

$D^*(2007)^0$

$I(J^P) = \frac{1}{2}(1^-)$
 I, J, P need confirmation.

J consistent with 1, value 0 ruled out (NGUYEN 77).

$D^*(2007)^0$ MASS

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|------|----------|
| 2006.85 ± 0.05 OUR FIT | Error includes scale factor of 1.1. | | |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 2006 ± 1.5 | ¹ GOLDHABER 77 | MRK1 | e^+e^- |
| ¹ From simultaneous fit to $D^*(2010)^+, D^*(2007)^0, D^+$, and D^0 . | | | |

$m_{D^*(2007)^0} - m_{D^0}$

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|---------------------------|------|-------------------------------|
| 142.014 ± 0.030 OUR FIT | Error includes scale factor of 1.5. | | | |
| 142.016 ± 0.030 OUR AVERAGE | Error includes scale factor of 1.5. | | | |
| 142.007 ± 0.015 ± 0.014 | 10k | ¹ TOMARADZE 15 | CLEO | $e^+e^- \rightarrow$ hadrons |
| 142.2 ± 0.3 ± 0.2 | 145 | ALBRECHT 95F | ARG | $e^+e^- \rightarrow$ hadrons |
| 142.12 ± 0.05 ± 0.05 | 1176 | BORTOLETTO92B | CLE2 | $e^+e^- \rightarrow$ hadrons |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 142.2 ± 2.0 | | SADROZINSKI 80 | CBAL | $D^{*0} \rightarrow D^0\pi^0$ |
| 142.7 ± 1.7 | | ² GOLDHABER 77 | MRK1 | e^+e^- |
| ¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration . This value comes from the average of the results for two decay modes, $D^0 \rightarrow K^- \pi^+$ and $D^0 \rightarrow K^- \pi^+ \pi^- \pi^+$. | | | | |
| ² From simultaneous fit to $D^*(2010)^+, D^*(2007)^0, D^+$, and D^0 . | | | | |

$D^*(2007)^0$ WIDTH

| VALUE (MeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|-------------------------|------|-------------------------------|
| <2.1 | 90 | ¹ ABACHI 88B | HRS | $D^{*0} \rightarrow D^+\pi^-$ |
| ¹ Assuming $m_{D^{*0}} = 2007.2 \pm 2.1$ MeV/ c^2 . | | | | |

$D^*(2007)^0$ DECAY MODES

$\bar{D}^*(2007)^0$ modes are charge conjugates of modes below.

| Mode | Fraction (Γ_i/Γ) |
|------------------------------|-------------------------------------|
| $\Gamma_1 \quad D^0 \pi^0$ | (64.7 \pm 0.9) % |
| $\Gamma_2 \quad D^0 \gamma$ | (35.3 \pm 0.9) % |
| $\Gamma_3 \quad D^0 e^+ e^-$ | (3.91 \pm 0.33) $\times 10^{-3}$ |

CONSTRAINED FIT INFORMATION

An overall fit to 2 branching ratios uses 5 measurements and one constraint to determine 2 parameters. The overall fit has a $\chi^2 = 2.5$ for 4 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$x_2 \begin{vmatrix} & -100 \\ & x_1 \end{vmatrix}$$

$D^*(2007)^0$ BRANCHING RATIOS

$\Gamma(D^0 \pi^0)/\Gamma(D^0 \gamma)$ Γ_1/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|-------------|--------------------|-------------|----------------|
|--------------|-------------|--------------------|-------------|----------------|

1.83 \pm 0.07 OUR FIT Error includes scale factor of 1.1.

1.85 \pm 0.07 OUR AVERAGE

| | | | | | |
|-----------------|------------|------|---------|----------|------------------------------------|
| 1.90 \pm 0.07 | \pm 0.05 | 4.9k | ABLIKIM | 15B BES3 | 10.6 $e^+ e^- \rightarrow$ hadrons |
|-----------------|------------|------|---------|----------|------------------------------------|

| | | | | | |
|-----------------|------------|--|------------|----------|------------------------------------|
| 1.74 \pm 0.02 | \pm 0.13 | | AUBERT, BE | 05G BABR | 10.6 $e^+ e^- \rightarrow$ hadrons |
|-----------------|------------|--|------------|----------|------------------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------------------|--|-------------------|----------|--|--------------------------------|
| 1.789 \pm 0.082 | | ¹ AAIJ | 22N LHCb | B ⁰ , B _s ⁰ | → $\bar{D}^{*0}(K\pi, \pi\pi)$ |
|-------------------|--|-------------------|----------|--|--------------------------------|

¹Statistical error only.

$\Gamma(D^0 e^+ e^-)/\Gamma(D^0 \gamma)$ Γ_3/Γ_2

| <u>VALUE (units 10^{-3})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|----------------|
|---|-------------|--------------------|-------------|----------------|

| | | | | |
|-----------------------------|-----|---------|-----------|---------------------|
| 11.08 \pm 0.76 \pm 0.49 | 421 | ABLIKIM | 21BD BES3 | 4.178 GeV $e^+ e^-$ |
|-----------------------------|-----|---------|-----------|---------------------|

$\Gamma(D^0 \pi^0)/\Gamma_{\text{total}}$ Γ_1/Γ

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|-------------|--------------------|-------------|----------------|
|--------------|-------------|--------------------|-------------|----------------|

0.647 \pm 0.009 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------------------------------|------|----------------------|----------|-------------------------------|
| 0.655 \pm 0.008 \pm 0.005 | 3.2k | ¹ ABLIKIM | 15B BES3 | $e^+ e^- \rightarrow$ hadrons |
|-------------------------------|------|----------------------|----------|-------------------------------|

| | | | | |
|-------------------------------|-----|-------------------------|----------|------------------------------------|
| 0.635 \pm 0.003 \pm 0.017 | 69k | ¹ AUBERT, BE | 05G BABR | 10.6 $e^+ e^- \rightarrow$ hadrons |
|-------------------------------|-----|-------------------------|----------|------------------------------------|

| | | | | | |
|-----------------------------|------|-----------------------|-----|------|-------------------------------|
| $0.596 \pm 0.035 \pm 0.028$ | 858 | ² ALBRECHT | 95F | ARG | $e^+ e^- \rightarrow$ hadrons |
| $0.636 \pm 0.023 \pm 0.033$ | 1097 | ² BUTLER | 92 | CLE2 | $e^+ e^- \rightarrow$ hadrons |

¹ Derived from the ratio $\Gamma(D^0 \pi^0) / \Gamma(D^0 \gamma)$ assuming that the branching fractions of $D^{*0} \rightarrow D^0 \pi^0$ and $D^{*0} \rightarrow D^0 \gamma$ decays sum to 100%.

² The BUTLER 92 and ALBRECHT 95F branching ratios are not independent, they have been constrained by the authors to sum to 100%.

| $\Gamma(D^0 \gamma) / \Gamma_{\text{total}}$ | | | | | Γ_2 / Γ |
|--|------|-------------|------|---------|---------------------|
| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT | |

0.353 ± 0.009 OUR FIT

0.381 ± 0.029 OUR AVERAGE

| | | | | | |
|-----------------------------|-----|-----------------------|-----|------|-------------------------------|
| $0.404 \pm 0.035 \pm 0.028$ | 456 | ¹ ALBRECHT | 95F | ARG | $e^+ e^- \rightarrow$ hadrons |
| $0.364 \pm 0.023 \pm 0.033$ | 621 | ¹ BUTLER | 92 | CLE2 | $e^+ e^- \rightarrow$ hadrons |
| $0.37 \pm 0.08 \pm 0.08$ | | ADLER | 88D | MRK3 | $e^+ e^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----------------------------|------|-------------------------|-----|------|------------------------------------|
| $0.345 \pm 0.008 \pm 0.005$ | 1.8k | ² ABLIKIM | 15B | BES3 | $e^+ e^- \rightarrow$ hadrons |
| $0.365 \pm 0.003 \pm 0.017$ | 68k | ² AUBERT, BE | 05G | BABR | $10.6 e^+ e^- \rightarrow$ hadrons |
| 0.47 ± 0.23 | | LOW | 87 | HRS | 29 GeV $e^+ e^-$ |
| 0.53 ± 0.13 | | BARTEL | 85G | JADE | $e^+ e^-$, hadrons |
| 0.47 ± 0.12 | | COLES | 82 | MRK2 | $e^+ e^-$ |
| 0.45 ± 0.15 | | GOLDHABER | 77 | MRK1 | $e^+ e^-$ |

¹ The BUTLER 92 and ALBRECHT 95F branching ratios are not independent, they have been constrained by the authors to sum to 100%.

² Derived from the ratio $\Gamma(D^0 \pi^0) / \Gamma(D^0 \gamma)$ assuming that the branching fractions of $D^{*0} \rightarrow D^0 \pi^0$ and $D^{*0} \rightarrow D^0 \gamma$ decays sum to 100%.

$D^*(2007)^0$ REFERENCES

| | | | | |
|-------------|------|-------------------|----------------------------------|-------------------------|
| AAIJ | 22N | PR D105 072005 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABLIKIM | 21BD | PR D104 112012 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 15B | PR D91 031101 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| TOMARADZE | 15 | PR D91 011102 | A. Tomaradze <i>et al.</i> | (NWES) |
| AUBERT, BE | 05G | PR D72 091101 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| ALBRECHT | 95F | ZPHY C66 63 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| BORTOLETTO | 92B | PRL 69 2046 | D. Bortoletto <i>et al.</i> | (CLEO Collab.) |
| BUTLER | 92 | PRL 69 2041 | F. Butler <i>et al.</i> | (CLEO Collab.) |
| ABACHI | 88B | PL B212 533 | S. Abachi <i>et al.</i> | (ANL, IND, MICH, PURD+) |
| ADLER | 88D | PL B208 152 | J. Adler <i>et al.</i> | (Mark III Collab.) |
| LOW | 87 | PL B183 232 | E.H. Low <i>et al.</i> | (HRS Collab.) |
| BARTEL | 85G | PL 161B 197 | W. Bartel <i>et al.</i> | (JADE Collab.) |
| COLES | 82 | PR D26 2190 | M.W. Coles <i>et al.</i> | (LBL, SLAC) |
| SADROZINSKI | 80 | Madison Conf. 681 | H.F.W. Sadrozinski <i>et al.</i> | (PRIN, CIT+) |
| GOLDHABER | 77 | PL 69B 503 | G. Goldhaber <i>et al.</i> | (Mark I Collab.) |
| NGUYEN | 77 | PRL 39 262 | H.K. Nguyen <i>et al.</i> | (LBL, SLAC) J |