

J/ ψ (1S)

$I^G(J^{PC}) = 0^-(1^{--})$

J/ ψ (1S) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3096.900±0.006 OUR AVERAGE				
3096.900±0.002±0.006		¹ ANASHIN 15	KEDR	$e^+ e^- \rightarrow$ hadrons
3096.89 ± 0.09	502	² ARTAMONOV 00	OLYA	$e^+ e^- \rightarrow$ hadrons
3096.91 ± 0.03 ± 0.01		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+ e^-$
3096.95 ± 0.1 ± 0.3	193	BAGLIN	SPEC	$\bar{p}p \rightarrow e^+ e^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3096.66 ± 0.19 ± 0.02	6.1k	⁴ AAIJ 15BI	LHCb	$p\bar{p} \rightarrow J/\psi X$
3096.917±0.010±0.007		AULCHENKO 03	KEDR	$e^+ e^- \rightarrow$ hadrons
3097.5 ± 0.3		GRIBUSHIN 96	FMPS	515 π^- Be → $2\mu X$
3098.4 ± 2.0	38k	LEMOIGNE 82	GOLI	185 π^- Be → $\gamma\mu^+\mu^- A$
3096.93 ± 0.09	502	⁵ ZHOLENTZ 80	REDE	$e^+ e^-$
3097.0 ± 1		⁶ BRANDELIK 79C	DASP	$e^+ e^-$

¹ Supersedes AULCHENKO 03.

² Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).

³ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.

⁴ From a sample of $\eta_c(1S)$ and J/ψ produced in b -hadron decays. Systematic uncertainties not estimated.

⁵ Superseded by ARTAMONOV 00.

⁶ From a simultaneous fit to $e^+ e^-$, $\mu^+ \mu^-$ and hadronic channels assuming $\Gamma(e^+ e^-) = \Gamma(\mu^+ \mu^-)$.

J/ ψ (1S) WIDTH

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
92.6 ± 1.7 OUR AVERAGE				
Error includes scale factor of 1.1.				
92.45 ± 1.40 ± 1.48		¹ ANASHIN 20	KEDR	$e^+ e^-$
96.1 ± 3.2	13k	² ADAMS 06A	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
84.4 ± 8.9		BAI 95B	BES	$e^+ e^-$
91 ± 11 ± 6		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+ e^-$
85.5 ± 6.1 - 5.8		⁴ HSUEH 92	RVUE	See γ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •				
92.94 ± 1.83		^{5,6} ANASHIN 18A	KEDR	$e^+ e^-$
94.1 ± 2.7		⁷ ANASHIN 10	KEDR	$3.097 e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
93.7 ± 3.5	7.8k	² AUBERT 04	BABR	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$

¹ Based on the same dataset as ANASHIN 18A and correlated to the values reported there

² Calculated by us from the reported values of $\Gamma(e^+ e^-) \times B(\mu^+ \mu^-)$ using $B(e^+ e^-) = (5.94 \pm 0.06)\%$ and $B(\mu^+ \mu^-) = (5.93 \pm 0.06)\%$.

³ The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].⁴ Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75B, BRANDELIK 79C.⁵ Using $\Gamma(e^+e^-)$ from ANASHIN 18A and $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ from PDG 16.⁶ Superseded by ANASHIN 20 that is based on the same dataset .⁷ Assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ and using $\Gamma(e^+e^-)/\Gamma_{\text{total}} = (5.94 \pm 0.06)\%$.

J/ ψ (1S) DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(87.7 \pm 0.5) %	
Γ_2 virtual $\gamma \rightarrow$ hadrons	(13.50 \pm 0.30) %	
Γ_3 ggg	(64.1 \pm 1.0) %	
Γ_4 γgg	(8.8 \pm 1.1) %	
Γ_5 e^+e^-	(5.971 \pm 0.032) %	
Γ_6 $e^+e^-\gamma$	[a] (8.8 \pm 1.4) $\times 10^{-3}$	
Γ_7 $\mu^+\mu^-$	(5.961 \pm 0.033) %	

Decays involving hadronic resonances

Γ_8 $\rho\pi$	(1.69 \pm 0.15) %	
Γ_9 $\rho^0\pi^0$	(5.6 \pm 0.7) $\times 10^{-3}$	
Γ_{10} $a_2(1320)^0\pi^+\pi^- \rightarrow 2(\pi^+\pi^-)\pi^0$	(2.8 \pm 0.6) $\times 10^{-3}$	
Γ_{11} $a_2(1320)^+\pi^-\pi^0 + \text{c.c} \rightarrow 2(\pi^+\pi^-)\pi^0$	(3.7 \pm 0.7) $\times 10^{-3}$	
Γ_{12} $a_2(1320)\rho$	(1.09 \pm 0.22) %	
Γ_{13} $\eta\pi^+\pi^-$	(3.8 \pm 0.7) $\times 10^{-4}$	
Γ_{14} $\eta\pi^+\pi^-\pi^0$	(1.17 \pm 0.20) %	
Γ_{15} $\eta\pi^+\pi^-3\pi^0$	(4.9 \pm 1.0) $\times 10^{-3}$	
Γ_{16} $\eta\rho$	(1.93 \pm 0.23) $\times 10^{-4}$	
Γ_{17} $\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-$	(1.2 \pm 0.4) $\times 10^{-4}$	
Γ_{18} $\eta\phi(2170) \rightarrow \eta K^*(892)^0\bar{K}^*(892)^0$	< 2.52 $\times 10^{-4}$	CL=90%
Γ_{19} $\eta K_S^\pm K_S^0\pi^\mp$	[b] (2.2 \pm 0.4) $\times 10^{-3}$	
Γ_{20} $\eta K^*(892)^0\bar{K}^*(892)^0$	(1.15 \pm 0.26) $\times 10^{-3}$	
Γ_{21} $\rho\eta'(958)$	(8.1 \pm 0.8) $\times 10^{-5}$	S=1.6
Γ_{22} $\rho^\pm\pi^\mp\pi^+\pi^-2\pi^0$	(2.8 \pm 0.8) %	
Γ_{23} $\rho^+\rho^-\pi^+\pi^-\pi^0$	(6 \pm 4) $\times 10^{-3}$	
Γ_{24} $\rho^+K^+K^-\pi^-\pi^+\pi^-\pi^0$	(3.5 \pm 0.8) $\times 10^{-3}$	
Γ_{25} $\rho^\mp K^\pm K_S^0$	(1.9 \pm 0.4) $\times 10^{-3}$	
Γ_{26} $h_1(1415)\eta' \rightarrow \gamma\eta\eta'$		
Γ_{27} $h_1(1595)\eta' \rightarrow \gamma\eta\eta'$		

Γ_{28}	$\rho(1450)\pi$			
Γ_{29}	$\rho(1450)\pi \rightarrow \pi^+ \pi^- \pi^0$	$(2.3 \pm 0.7) \times 10^{-3}$		
Γ_{30}	$\rho(1450)^\pm \pi^\mp \rightarrow K_S^0 K^\pm \pi^\mp$	$(3.5 \pm 0.6) \times 10^{-4}$		
Γ_{31}	$\rho(1450)^0 \pi^0 \rightarrow K^+ K^- \pi^0$	$(2.7 \pm 0.6) \times 10^{-4}$		
Γ_{32}	$\rho(1450)\eta'(958) \rightarrow \pi^+ \pi^- \eta'(958)$	$(3.3 \pm 0.7) \times 10^{-6}$		
Γ_{33}	$\rho(1700)\pi$			
Γ_{34}	$\rho(1700)\pi \rightarrow \pi^+ \pi^- \pi^0$	$(1.7 \pm 1.1) \times 10^{-4}$		
Γ_{35}	$\rho(2150)\pi$			
Γ_{36}	$\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0$	$(8 \pm 40) \times 10^{-6}$		
Γ_{37}	$\rho_3(1690)\pi \rightarrow \pi^+ \pi^- \pi^0$			
Γ_{38}	$\omega \pi^0$	$(4.5 \pm 0.5) \times 10^{-4}$	S=1.4	
Γ_{39}	$\omega \pi^0 \rightarrow \pi^+ \pi^- \pi^0$	$(1.7 \pm 0.8) \times 10^{-5}$		
Γ_{40}	$\omega \pi^+ \pi^-$	$(8.5 \pm 1.0) \times 10^{-3}$	S=1.3	
Γ_{41}	$\omega \pi^0 \pi^0$	$(3.4 \pm 0.8) \times 10^{-3}$		
Γ_{42}	$\omega 3\pi^0$	$(1.9 \pm 0.6) \times 10^{-3}$		
Γ_{43}	$\omega f_2(1270)$	$(4.3 \pm 0.6) \times 10^{-3}$		
Γ_{44}	$\omega \eta$	$(1.74 \pm 0.20) \times 10^{-3}$	S=1.6	
Γ_{45}	$\omega \pi^+ \pi^- \pi^0$	$(4.0 \pm 0.7) \times 10^{-3}$		
Γ_{46}	$\omega \pi^0 \eta$	$(3.4 \pm 1.7) \times 10^{-4}$		
Γ_{47}	$\omega \pi^+ \pi^+ \pi^- \pi^-$	$(8.5 \pm 3.4) \times 10^{-3}$		
Γ_{48}	$\omega \pi^+ \pi^- 2\pi^0$	$(3.3 \pm 0.5) \%$		
Γ_{49}	$\omega \eta' \pi^+ \pi^-$	$(1.12 \pm 0.13) \times 10^{-3}$		
Γ_{50}	$\omega \eta'(958)$	$(1.89 \pm 0.18) \times 10^{-4}$		
Γ_{51}	$\omega f_0(980)$	$(1.4 \pm 0.5) \times 10^{-4}$		
Γ_{52}	$\omega f_0(1710) \rightarrow \omega K \bar{K}$	$(4.8 \pm 1.1) \times 10^{-4}$		
Γ_{53}	$\omega f_1(1420)$	$(6.8 \pm 2.4) \times 10^{-4}$		
Γ_{54}	$\omega f'_2(1525)$	$< 2.2 \times 10^{-4}$	CL=90%	
Γ_{55}	$\omega X(1835) \rightarrow \omega p \bar{p}$	$< 3.9 \times 10^{-6}$	CL=95%	
Γ_{56}	$\omega X(1835), X \rightarrow \eta' \pi^+ \pi^-$	$< 6.2 \times 10^{-5}$		
Γ_{57}	$\omega K^+ K^-$	$(1.52 \pm 0.31) \times 10^{-3}$		
Γ_{58}	$\omega K^\pm K_S^0 \pi^\mp$	[b] $(3.4 \pm 0.5) \times 10^{-3}$		
Γ_{59}	$\omega K \bar{K}$	$(1.9 \pm 0.4) \times 10^{-3}$		
Γ_{60}	$\omega K^*(892) \bar{K} + \text{c.c.}$	$(6.1 \pm 0.9) \times 10^{-3}$		
Γ_{61}	$\eta' K^{*\pm} K^\mp$	$(1.48 \pm 0.13) \times 10^{-3}$		
Γ_{62}	$\eta' K^{*0} \bar{K}^0 + \text{c.c.}$	$(1.66 \pm 0.21) \times 10^{-3}$		
Γ_{63}	$\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.}$	$(2.16 \pm 0.31) \times 10^{-4}$		
Γ_{64}	$\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp$	$(1.51 \pm 0.23) \times 10^{-4}$		
Γ_{65}	$\eta' h_1(1415) \rightarrow \gamma \eta' \eta'$	$(4.7 \pm 1.1) \times 10^{-7}$		
Γ_{66}	$\bar{K} K^*(892) + \text{c.c.}$			
Γ_{67}	$\bar{K} K^*(892) + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp$	$(5.0 \pm 0.5) \times 10^{-3}$		
Γ_{68}	$K^+ K^*(892)^- + \text{c.c.}$	$(6.0 \pm 0.8) \times 10^{-3}$	S=2.9	

Γ_{69}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(2.69 \pm 0.13) \times 10^{-3}$
Γ_{70}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow$ $K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.0 \pm 0.4) \times 10^{-3}$
Γ_{71}	$K^0 \bar{K}^*(892)^0 + \text{c.c.}$	$(4.2 \pm 0.4) \times 10^{-3}$
Γ_{72}	$K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow$ $K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.2 \pm 0.4) \times 10^{-3}$
Γ_{73}	$\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}$	$(5.7 \pm 0.8) \times 10^{-3}$
Γ_{74}	$K^*(892)^\pm K^\mp \pi^0$	$(4.1 \pm 1.3) \times 10^{-3}$
Γ_{75}	$K^*(892)^+ K_S^0 \pi^- + \text{c.c.}$	$(2.0 \pm 0.5) \times 10^{-3}$
Γ_{76}	$K^*(892)^+ K_S^0 \pi^- + \text{c.c.} \rightarrow$ $K_S^0 K_S^0 \pi^+ \pi^-$	$(6.7 \pm 2.2) \times 10^{-4}$
Γ_{77}	$K^*(892)^0 K^- \pi^+ + \text{c.c.} \rightarrow$ $K^+ K^- \pi^+ \pi^-$	$(3.8 \pm 0.5) \times 10^{-3}$
Γ_{78}	$K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(6.3 \pm 0.6) \times 10^{-6}$
Γ_{79}	$K^*(892)^0 K_S^0 \pi^0$	$(7 \pm 4) \times 10^{-4}$
Γ_{80}	$K^*(892)^\pm K^*(700)^\mp$	$(1.1 \pm 1.0) \times 10^{-3}$
Γ_{81}	$K^*(892)^0 \bar{K}^*(892)^0$	$(2.3 \pm 0.6) \times 10^{-4}$
Γ_{82}	$K^*(892)^\pm K^*(892)^\mp$	$(1.00 \pm 0.22) \times 10^{-3}$
Γ_{83}	$K_1(1400)^\pm K^\mp$	$(3.8 \pm 1.4) \times 10^{-3}$
Γ_{84}	$K^*(1410) \bar{K} + \text{c.c.}$	
Γ_{85}	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow$ $K^\pm K^\mp \pi^0$	$(7 \pm 4) \times 10^{-5}$
Γ_{86}	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(8 \pm 6) \times 10^{-5}$
Γ_{87}	$K_2^*(1430) \bar{K} + \text{c.c.}$	
Γ_{88}	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow$ $K^\pm K^\mp \pi^0$	$(1.0 \pm 0.5) \times 10^{-4}$
Γ_{89}	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(4.0 \pm 1.0) \times 10^{-4}$
Γ_{90}	$\bar{K}_2^*(1430) K + \text{c.c.}$	$< 4.0 \times 10^{-3} \text{ CL}=90\%$
Γ_{91}	$K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(2.69 \pm 0.25) \times 10^{-4}$
Γ_{92}	$K_2^*(1430)^0 K^- \pi^+ + \text{c.c.} \rightarrow$ $K^+ K^- \pi^+ \pi^-$	$(2.6 \pm 0.9) \times 10^{-3}$
Γ_{93}	$K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}$	$(3.6 \pm 1.8) \times 10^{-3}$
Γ_{94}	$\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}$	$(4.67 \pm 0.29) \times 10^{-3}$
Γ_{95}	$K_2^*(1430)^- K^*(892)^+ + \text{c.c.}$	$(3.4 \pm 2.9) \times 10^{-3}$
Γ_{96}	$K_2^*(1430)^- K^*(892)^+ +$ $\text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- +$ c.c.	$(4 \pm 4) \times 10^{-4}$
Γ_{97}	$K_2^*(1430)^0 \bar{K}_2^*(1430)^0$	$< 2.9 \times 10^{-3} \text{ CL}=90\%$

Γ_{98}	$\overline{K}_2(1770)^0 K^*(892)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(6.9 \pm 0.9) \times 10^{-4}$
Γ_{99}	$K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(1.10 \pm 0.60) \times 10^{-5}$
Γ_{100}	$K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(6.2 \pm 2.9) \times 10^{-6}$
Γ_{101}	$K_1(1270)^{\pm} K^{\mp}$	$< 3.0 \times 10^{-3}$ CL=90%
Γ_{102}	$K_1(1270) K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(8.5 \pm 2.5) \times 10^{-7}$
Γ_{103}	$a_2(1320)^{\pm} \pi^{\mp}$	$[b] < 4.3 \times 10^{-3}$ CL=90%
Γ_{104}	$\phi \pi^0$	3×10^{-6} or 1×10^{-7}
Γ_{105}	$\phi \pi^+ \pi^-$	$(9.4 \pm 1.5) \times 10^{-4}$ S=1.7
Γ_{106}	$\phi \pi^0 \pi^0$	$(5.0 \pm 1.0) \times 10^{-4}$
Γ_{107}	$\phi 2(\pi^+ \pi^-)$	$(1.60 \pm 0.32) \times 10^{-3}$
Γ_{108}	$\phi \eta$	$(7.4 \pm 0.6) \times 10^{-4}$ S=1.2
Γ_{109}	$\phi \eta'(958)$	$(4.6 \pm 0.5) \times 10^{-4}$ S=2.2
Γ_{110}	$\phi \eta \eta'$	$(2.32 \pm 0.17) \times 10^{-4}$
Γ_{111}	$\phi f_0(980)$	$(3.2 \pm 0.9) \times 10^{-4}$ S=1.9
Γ_{112}	$\phi f_0(980) \rightarrow \phi \pi^+ \pi^-$	$(2.60 \pm 0.34) \times 10^{-4}$
Γ_{113}	$\phi f_0(980) \rightarrow \phi \pi^0 \pi^0$	$(1.8 \pm 0.5) \times 10^{-4}$
Γ_{114}	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-$	$(4.5 \pm 1.0) \times 10^{-6}$
Γ_{115}	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 p^0 \pi^0$	$(1.7 \pm 0.6) \times 10^{-6}$
Γ_{116}	$\phi f_0(980) \eta \rightarrow \eta \phi \pi^+ \pi^-$	$(3.2 \pm 1.0) \times 10^{-4}$
Γ_{117}	$\phi a_0(980)^0 \rightarrow \phi \eta \pi^0$	$(4.4 \pm 1.4) \times 10^{-6}$
Γ_{118}	$\phi f_2(1270)$	$(3.2 \pm 0.6) \times 10^{-4}$
Γ_{119}	$\phi f_1(1285)$	$(2.6 \pm 0.5) \times 10^{-4}$
Γ_{120}	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-$	$(9.4 \pm 2.8) \times 10^{-7}$
Γ_{121}	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi 3\pi^0$	$(2.1 \pm 2.2) \times 10^{-7}$
Γ_{122}	$\phi \eta(1405) \rightarrow \phi \eta \pi^+ \pi^-$	$(2.0 \pm 1.0) \times 10^{-5}$
Γ_{123}	$\phi f'_2(1525)$	$(8 \pm 4) \times 10^{-4}$ S=2.7
Γ_{124}	$\phi X(1835) \rightarrow \phi p \bar{p}$	$< 2.1 \times 10^{-7}$ CL=90%
Γ_{125}	$\phi X(1835) \rightarrow \phi \eta \pi^+ \pi^-$	$< 2.8 \times 10^{-4}$ CL=90%
Γ_{126}	$\phi X(1870) \rightarrow \phi \eta \pi^+ \pi^-$	$< 6.13 \times 10^{-5}$ CL=90%
Γ_{127}	$\phi K \bar{K}$	$(1.77 \pm 0.16) \times 10^{-3}$ S=1.3
Γ_{128}	$\phi f_0(1710) \rightarrow \phi K \bar{K}$	$(3.6 \pm 0.6) \times 10^{-4}$
Γ_{129}	$\phi K^+ K^-$	$(8.3 \pm 1.1) \times 10^{-4}$
Γ_{130}	$\phi K_S^0 K_S^0$	$(5.9 \pm 1.5) \times 10^{-4}$
Γ_{131}	$\phi K^{\pm} K_S^0 \pi^{\mp}$	$[b] (7.2 \pm 0.8) \times 10^{-4}$
Γ_{132}	$\phi K^*(892) \bar{K} + \text{c.c.}$	$(2.18 \pm 0.23) \times 10^{-3}$
Γ_{133}	$b_1(1235)^{\pm} \pi^{\mp}$	$[b] (3.0 \pm 0.5) \times 10^{-3}$
Γ_{134}	$b_1(1235)^0 \pi^0$	$(2.3 \pm 0.6) \times 10^{-3}$
Γ_{135}	$f'_2(1525) K^+ K^-$	$(1.06 \pm 0.35) \times 10^{-3}$

Γ_{136}	$\Delta(1232)^+ \bar{p}$	$< 1 \times 10^{-4}$	CL=90%
Γ_{137}	$\Delta(1232)^{++} \bar{p}\pi^-$	$(1.6 \pm 0.5) \times 10^{-3}$	
Γ_{138}	$\Delta(1232)^{++} \bar{\Delta}(1232)^{--}$	$(1.10 \pm 0.29) \times 10^{-3}$	
Γ_{139}	$\bar{\Sigma}(1385)^0 p K^-$	$(5.1 \pm 3.2) \times 10^{-4}$	
Γ_{140}	$\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.}$	$< 8.2 \times 10^{-6}$	CL=90%
Γ_{141}	$\Sigma(1385)^- \bar{\Sigma}^+(\text{or c.c.})$	[b] $(3.1 \pm 0.5) \times 10^{-4}$	
Γ_{142}	$\Sigma(1385)^- \bar{\Sigma}(1385)^+(\text{or c.c.})$	[b] $(1.16 \pm 0.05) \times 10^{-3}$	
Γ_{143}	$\Sigma(1385)^0 \bar{\Sigma}(1385)^0$	$(1.07 \pm 0.08) \times 10^{-3}$	
Γ_{144}	$\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda}$	$< 4.1 \times 10^{-6}$	CL=90%
Γ_{145}	$\bar{\Lambda}(1520) \Lambda + \text{c.c.}$	$< 1.80 \times 10^{-3}$	CL=90%
Γ_{146}	$\Xi^0 \bar{\Xi}^0$	$(1.17 \pm 0.04) \times 10^{-3}$	
Γ_{147}	$\Xi(1530)^- \bar{\Xi}^+ + \text{c.c.}$	$(3.18 \pm 0.08) \times 10^{-4}$	
Γ_{148}	$\Xi(1530)^0 \bar{\Xi}^0$	$(3.2 \pm 1.4) \times 10^{-4}$	
Γ_{149}	$\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.}$	[c] $< 1.1 \times 10^{-5}$	CL=90%
Γ_{150}	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	[c] $< 2.1 \times 10^{-5}$	CL=90%
Γ_{151}	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	[c] $< 1.6 \times 10^{-5}$	CL=90%
Γ_{152}	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	[c] $< 5.6 \times 10^{-5}$	CL=90%
Γ_{153}	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	[c] $< 1.1 \times 10^{-5}$	CL=90%

Decays into stable hadrons

Γ_{154}	$2(\pi^+ \pi^-) \pi^0$	$(4.2 \pm 0.4) \%$	S=2.1
Γ_{155}	$3(\pi^+ \pi^-) \pi^0$	$(2.9 \pm 0.6) \%$	
Γ_{156}	$\pi^+ \pi^- 3\pi^0$	$(1.9 \pm 0.9) \%$	
Γ_{157}	$\pi^+ \pi^- 4\pi^0$	$(6.5 \pm 1.3) \times 10^{-3}$	
Γ_{158}	$\rho^\pm \pi^\mp \pi^0 \pi^0$	$(1.41 \pm 0.22) \%$	
Γ_{159}	$\rho^+ \rho^- \pi^0$	$(6.0 \pm 1.1) \times 10^{-3}$	
Γ_{160}	$\pi^+ \pi^- \pi^0$	$(2.10 \pm 0.08) \%$	S=1.6
Γ_{161}	$2(\pi^+ \pi^- \pi^0)$	$(1.61 \pm 0.20) \%$	
Γ_{162}	$\pi^+ \pi^- \pi^0 K^+ K^-$	$(1.52 \pm 0.27) \%$	S=1.4
Γ_{163}	$\pi^+ \pi^-$	$(1.47 \pm 0.14) \times 10^{-4}$	
Γ_{164}	$2(\pi^+ \pi^-)$	$(3.20 \pm 0.25) \times 10^{-3}$	S=1.2
Γ_{165}	$3(\pi^+ \pi^-)$	$(4.3 \pm 0.4) \times 10^{-3}$	
Γ_{166}	$2(\pi^+ \pi^-) 3\pi^0$	$(6.2 \pm 0.9) \%$	
Γ_{167}	$4(\pi^+ \pi^-) \pi^0$	$(9.0 \pm 3.0) \times 10^{-3}$	
Γ_{168}	$2(\pi^+ \pi^-) \eta$	$(2.29 \pm 0.28) \times 10^{-3}$	
Γ_{169}	$3(\pi^+ \pi^-) \eta$	$(7.2 \pm 1.5) \times 10^{-4}$	
Γ_{170}	$2(\pi^+ \pi^- \pi^0) \eta$	$(1.6 \pm 0.5) \times 10^{-3}$	
Γ_{171}	$\pi^+ \pi^- \pi^0 \pi^0 \eta$	$(2.4 \pm 0.5) \times 10^{-3}$	
Γ_{172}	$\rho^\pm \pi^\mp \pi^0 \eta$	$(1.9 \pm 0.8) \times 10^{-3}$	
Γ_{173}	$K^+ K^-$	$(2.86 \pm 0.21) \times 10^{-4}$	
Γ_{174}	$K_S^0 K_L^0$	$(1.95 \pm 0.11) \times 10^{-4}$	S=2.4
Γ_{175}	$K_S^0 K_S^0$	$< 1.4 \times 10^{-8}$	CL=95%
Γ_{176}	$K \bar{K} \pi$	$(6.1 \pm 1.0) \times 10^{-3}$	

Γ_{177}	$K^+ K^- \pi^0$	$(2.88 \pm 0.12) \times 10^{-3}$	
Γ_{178}	$K_S^0 K^\pm \pi^\mp$	$(5.6 \pm 0.5) \times 10^{-3}$	
Γ_{179}	$K_S^0 K_L^0 \pi^0$	$(2.06 \pm 0.26) \times 10^{-3}$	
Γ_{180}	$K^*(892)^0 \bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0 K_L^0 \pi^0$	$(1.21 \pm 0.18) \times 10^{-3}$	
Γ_{181}	$K_2^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0 K_L^0 \pi^0$	$(4.3 \pm 1.3) \times 10^{-4}$	
Γ_{182}	$K^+ K^- \pi^+ \pi^-$	$(7.0 \pm 1.0) \times 10^{-3}$	
Γ_{183}	$K^+ K^- \pi^0 \pi^0$	$(2.13 \pm 0.22) \times 10^{-3}$	
Γ_{184}	$K_S^0 K_L^0 \pi^+ \pi^-$	$(3.8 \pm 0.6) \times 10^{-3}$	
Γ_{185}	$K_S^0 K_L^0 \pi^0 \pi^0$	$(1.9 \pm 0.4) \times 10^{-3}$	
Γ_{186}	$K_S^0 K_L^0 \eta$	$(1.45 \pm 0.33) \times 10^{-3}$	
Γ_{187}	$K_S^0 K_S^0 \pi^+ \pi^-$	$(1.68 \pm 0.19) \times 10^{-3}$	
Γ_{188}	$K^\mp K_S^0 \pi^\pm \pi^0$	$(5.7 \pm 0.5) \times 10^{-3}$	
Γ_{189}	$K^+ K^- 2(\pi^+ \pi^-)$	$(3.1 \pm 1.3) \times 10^{-3}$	
Γ_{190}	$K^+ K^- \pi^+ \pi^- \eta$	$(4.7 \pm 0.7) \times 10^{-3}$	
Γ_{191}	$2(K^+ K^-)$	$(7.2 \pm 0.8) \times 10^{-4}$	
Γ_{192}	$K^+ K^- K_S^0 K_S^0$	$(4.2 \pm 0.7) \times 10^{-4}$	
Γ_{193}	$p\bar{p}$	$(2.120 \pm 0.029) \times 10^{-3}$	
Γ_{194}	$p\bar{p}\pi^0$	$(1.19 \pm 0.08) \times 10^{-3}$	S=1.1
Γ_{195}	$p\bar{p}\pi^+ \pi^-$	$(6.0 \pm 0.5) \times 10^{-3}$	S=1.3
Γ_{196}	$p\bar{p}\pi^+ \pi^- \pi^0$	[d] $(2.3 \pm 0.9) \times 10^{-3}$	S=1.9
Γ_{197}	$p\bar{p}\eta$	$(2.00 \pm 0.12) \times 10^{-3}$	
Γ_{198}	$p\bar{p}\rho$	< 3.1 $\times 10^{-4}$	CL=90%
Γ_{199}	$p\bar{p}\omega$	$(9.8 \pm 1.0) \times 10^{-4}$	S=1.3
Γ_{200}	$p\bar{p}\eta'(958)$	$(1.29 \pm 0.14) \times 10^{-4}$	S=2.0
Γ_{201}	$p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0 \eta$	$(6.8 \pm 1.8) \times 10^{-5}$	
Γ_{202}	$p\bar{p}\phi$	$(5.19 \pm 0.33) \times 10^{-5}$	
Γ_{203}	$p\bar{n}\pi^-$	$(2.12 \pm 0.09) \times 10^{-3}$	
Γ_{204}	$n\bar{n}$	$(2.09 \pm 0.16) \times 10^{-3}$	
Γ_{205}	$n\bar{n}\pi^+ \pi^-$	$(4 \pm 4) \times 10^{-3}$	
Γ_{206}	$nN(1440)$	seen	
Γ_{207}	$nN(1520)$	seen	
Γ_{208}	$nN(1535)$	seen	
Γ_{209}	$\Lambda\bar{\Lambda}$	$(1.89 \pm 0.09) \times 10^{-3}$	S=2.8
Γ_{210}	$\Lambda\bar{\Lambda}\pi^0$	$(3.8 \pm 0.4) \times 10^{-5}$	
Γ_{211}	$\Lambda\bar{\Lambda}\pi^+ \pi^-$	$(4.3 \pm 1.0) \times 10^{-3}$	
Γ_{212}	$\Lambda\bar{\Lambda}\eta$	$(1.62 \pm 0.17) \times 10^{-4}$	
Γ_{213}	$\Lambda\bar{\Sigma}^-\pi^+ (\text{or c.c.})$	[b] $(8.3 \pm 0.7) \times 10^{-4}$	S=1.2
Γ_{214}	$pK^-\bar{\Lambda}+\text{c.c.}$	$(8.6 \pm 1.1) \times 10^{-4}$	
Γ_{215}	$pK^-\bar{\Sigma}^0$	$(2.9 \pm 0.8) \times 10^{-4}$	
Γ_{216}	$\bar{\Lambda}nK_S^0 + \text{c.c.}$	$(6.5 \pm 1.1) \times 10^{-4}$	
Γ_{217}	$\Lambda\bar{\Sigma} + \text{c.c.}$	$(2.83 \pm 0.23) \times 10^{-5}$	

Γ_{218}	$\Sigma^+ \bar{\Sigma}^-$	$(1.07 \pm 0.04) \times 10^{-3}$	
Γ_{219}	$\Sigma^0 \bar{\Sigma}^0$	$(1.172 \pm 0.032) \times 10^{-3}$	S=1.4
Γ_{220}	$\Sigma^+ \bar{\Sigma}^- \eta$	$(6.3 \pm 0.4) \times 10^{-5}$	
Γ_{221}	$\Xi^- \bar{\Xi}^+$	$(9.7 \pm 0.8) \times 10^{-4}$	S=1.4

Radiative decays

Γ_{222}	$\gamma \eta_c(1S)$	$(1.7 \pm 0.4) \%$	S=1.5
Γ_{223}	$\gamma \eta_c(1S) \rightarrow 3\gamma$	$(3.8 \pm 1.3) \times 10^{-6}$	S=1.1
Γ_{224}	$\gamma \eta_c(1S) \rightarrow \gamma \eta \eta \eta'$	$(4.9 \pm 0.8) \times 10^{-5}$	
Γ_{225}	3γ	$(1.16 \pm 0.22) \times 10^{-5}$	
Γ_{226}	4γ	$< 9 \times 10^{-6}$	CL=90%
Γ_{227}	5γ	$< 1.5 \times 10^{-5}$	CL=90%
Γ_{228}	$\gamma \pi^0$	$(3.56 \pm 0.17) \times 10^{-5}$	
Γ_{229}	$\gamma \pi^0 \pi^0$	$(1.15 \pm 0.05) \times 10^{-3}$	
Γ_{230}	$\gamma 2\pi^+ 2\pi^-$	$(2.8 \pm 0.5) \times 10^{-3}$	S=1.9
Γ_{231}	$\gamma f_2(1270) f_2(1270)$	$(9.5 \pm 1.7) \times 10^{-4}$	
Γ_{232}	$\gamma f_2(1270) f_2(1270)$ (non resonant)	$(8.2 \pm 1.9) \times 10^{-4}$	
Γ_{233}	$\gamma \pi^+ \pi^- 2\pi^0$	$(8.3 \pm 3.1) \times 10^{-3}$	
Γ_{234}	$\gamma K_S^0 K_S^0$	$(8.1 \pm 0.4) \times 10^{-4}$	
Γ_{235}	$\gamma(K\bar{K}\pi) [J^{PC} = 0-+]$	$(7 \pm 4) \times 10^{-4}$	S=2.1
Γ_{236}	$\gamma K^+ K^- \pi^+ \pi^-$	$(2.1 \pm 0.6) \times 10^{-3}$	
Γ_{237}	$\gamma K^*(892) \bar{K}^*(892)$	$(4.0 \pm 1.3) \times 10^{-3}$	
Γ_{238}	$\gamma \eta$	$(1.085 \pm 0.018) \times 10^{-3}$	
Γ_{239}	$\gamma \eta \pi^0$	$(2.14 \pm 0.31) \times 10^{-5}$	
Γ_{240}	$\gamma f_0(500) \rightarrow \gamma \pi \pi$		
Γ_{241}	$\gamma f_0(500) \rightarrow \gamma K \bar{K}$		
Γ_{242}	$\gamma f_0(500) \rightarrow \gamma \eta \eta$		
Γ_{243}	$\gamma a_0(980)^0 \rightarrow \gamma \eta \pi^0$	$< 2.5 \times 10^{-6}$	CL=95%
Γ_{244}	$\gamma a_2(1320)^0 \rightarrow \gamma \eta \pi^0$	$< 6.6 \times 10^{-6}$	CL=95%
Γ_{245}	$\gamma \eta \pi \pi$	$(6.1 \pm 1.0) \times 10^{-3}$	
Γ_{246}	$\gamma \eta_2(1870) \rightarrow \gamma \eta \pi^+ \pi^-$	$(6.2 \pm 2.4) \times 10^{-4}$	
Γ_{247}	$\gamma \eta'(958)$	$(5.25 \pm 0.07) \times 10^{-3}$	S=1.3
Γ_{248}	$\gamma f_0(980) \rightarrow \gamma \pi \pi$		
Γ_{249}	$\gamma f_0(980) \rightarrow \gamma K \bar{K}$		
Γ_{250}	$\gamma \rho \rho$	$(4.5 \pm 0.8) \times 10^{-3}$	
Γ_{251}	$\gamma \rho \omega$	$< 5.4 \times 10^{-4}$	CL=90%
Γ_{252}	$\gamma \rho \phi$	$< 8.8 \times 10^{-5}$	CL=90%
Γ_{253}	$\gamma \omega \omega$	$(1.61 \pm 0.33) \times 10^{-3}$	
Γ_{254}	$\gamma \phi \phi$	$(4.0 \pm 1.2) \times 10^{-4}$	S=2.1
Γ_{255}	$\gamma \eta(1405/1475) \rightarrow \gamma K \bar{K} \pi$	$(2.8 \pm 0.6) \times 10^{-3}$	S=1.6
Γ_{256}	$\gamma \eta(1405/1475) \rightarrow \gamma \gamma \rho^0$	$(7.8 \pm 2.0) \times 10^{-5}$	S=1.8
Γ_{257}	$\gamma \eta(1405/1475) \rightarrow \gamma \eta \pi^+ \pi^-$	$(3.0 \pm 0.5) \times 10^{-4}$	
Γ_{258}	$\gamma \eta(1405/1475) \rightarrow \gamma \rho^0 \rho^0$	$(1.7 \pm 0.4) \times 10^{-3}$	S=1.3

Γ_{259}	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi$	< 8.2	$\times 10^{-5}$	CL=95%
Γ_{260}	$\gamma\eta(1405) \rightarrow \gamma\gamma\gamma$	< 2.63	$\times 10^{-6}$	CL=90%
Γ_{261}	$\gamma\eta(1475) \rightarrow \gamma\gamma\gamma$	< 1.86	$\times 10^{-6}$	CL=90%
Γ_{262}	$\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0$	$(1.3 \pm 0.9) \times 10^{-4}$		
Γ_{263}	$\gamma\eta(1760) \rightarrow \gamma\omega\omega$	$(1.98 \pm 0.33) \times 10^{-3}$		
Γ_{264}	$\gamma\eta(1760) \rightarrow \gamma\gamma\gamma$	< 4.80	$\times 10^{-6}$	CL=90%
Γ_{265}	$\gamma\eta(2225)$	$(3.14 \pm 0.50) \times 10^{-4}$		
Γ_{266}	$\gamma f_2(1270)$	$(1.63 \pm 0.12) \times 10^{-3}$	S=1.3	
Γ_{267}	$\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0$	$(2.58 \pm 0.60) \times 10^{-5}$		
Γ_{268}	$\gamma f_1(1285)$	$(6.1 \pm 0.8) \times 10^{-4}$		
Γ_{269}	$\gamma f_0(1370) \rightarrow \gamma\pi\pi$			
Γ_{270}	$\gamma f_0(1370) \rightarrow \gamma K\bar{K}$	$(4.2 \pm 1.5) \times 10^{-4}$		
Γ_{271}	$\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0$	$(1.1 \pm 0.4) \times 10^{-5}$		
Γ_{272}	$\gamma f_0(1370) \rightarrow \gamma\eta\eta$			
Γ_{273}	$\gamma f_0(1370) \rightarrow \gamma\eta\eta'$			
Γ_{274}	$\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi$	$(7.9 \pm 1.3) \times 10^{-4}$		
Γ_{275}	$\gamma f_0(1500) \rightarrow \gamma\pi\pi$	$(1.09 \pm 0.24) \times 10^{-4}$		
Γ_{276}	$\gamma f_0(1500) \rightarrow \gamma\eta\eta$	$(1.7 \pm 0.6) \times 10^{-5}$		
Γ_{277}	$\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0$	$(1.59 \pm 0.24) \times 10^{-5}$		
Γ_{278}	$\gamma f_0(1500) \rightarrow \gamma\eta\eta'$			
Γ_{279}	$\gamma f_1(1510) \rightarrow \gamma\eta\pi^+\pi^-$	$(4.5 \pm 1.2) \times 10^{-4}$		
Γ_{280}	$\gamma f'_2(1525)$	$(5.7 \pm 0.8) \times 10^{-4}$	S=1.5	
Γ_{281}	$\gamma f'_2(1525) \rightarrow \gamma K_S^0 K_S^0$	$(8.0 \pm 0.7) \times 10^{-5}$		
Γ_{282}	$\gamma f'_2(1525) \rightarrow \gamma\eta\eta$	$(3.4 \pm 1.4) \times 10^{-5}$		
Γ_{283}	$\gamma f_2(1565) \rightarrow \gamma\eta\eta'$			
Γ_{284}	$\gamma f_2(1640) \rightarrow \gamma\omega\omega$	$(2.8 \pm 1.8) \times 10^{-4}$		
Γ_{285}	$\gamma f_0(1710) \rightarrow \gamma\pi\pi$	$(3.8 \pm 0.5) \times 10^{-4}$		
Γ_{286}	$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	$(9.5 \pm 1.0) \times 10^{-4}$	S=1.5	
Γ_{287}	$\gamma f_0(1710) \rightarrow \gamma\omega\omega$	$(3.1 \pm 1.0) \times 10^{-4}$		
Γ_{288}	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$	$(2.4 \pm 1.2) \times 10^{-4}$		
Γ_{289}	$\gamma f_0(1710) \rightarrow \gamma\eta\eta'$			
Γ_{290}	$\gamma f_0(1710) \rightarrow \gamma\omega\phi$	$(2.5 \pm 0.6) \times 10^{-4}$		
Γ_{291}	$\gamma f_0(1770) \rightarrow \gamma K_S^0 K_S^0$	$(1.11 \pm 0.20) \times 10^{-5}$		
Γ_{292}	$\gamma f_2(1810) \rightarrow \gamma\eta\eta$	$(5.4 \pm 3.5) \times 10^{-5}$		
Γ_{293}	$\gamma\eta_1(1855) \rightarrow \gamma\eta\eta'$	$(2.7 \pm 0.4) \times 10^{-6}$		
Γ_{294}	$\gamma f_0(1770) \rightarrow \gamma\eta\eta'$			
Γ_{295}	$\gamma f_2(1910) \rightarrow \gamma\omega\omega$	$(2.0 \pm 1.4) \times 10^{-4}$		

Γ_{296}	$\gamma f_2(1950) \rightarrow \gamma K^*(892) \bar{K}^*(892)$	$(7.0 \pm 2.2) \times 10^{-4}$
Γ_{297}	$\gamma f_2(2010) \rightarrow \gamma \eta \eta'$	
Γ_{298}	$\gamma f_0(2020) \rightarrow \gamma \pi \pi$	
Γ_{299}	$\gamma f_0(2020) \rightarrow \gamma K \bar{K}$	
Γ_{300}	$\gamma f_0(2020) \rightarrow \gamma \eta \eta$	
Γ_{301}	$\gamma f_0(2020) \rightarrow \gamma \eta' \eta'$	$(2.63 \pm 0.32) \times 10^{-4}$
Γ_{302}	$\gamma f_0(2020) \rightarrow \gamma \eta \eta'$	
Γ_{303}	$\gamma f_4(2050)$	$(2.7 \pm 0.7) \times 10^{-3}$
Γ_{304}	$\gamma f_4(2050) \rightarrow \gamma \eta \eta'$	
Γ_{305}	$\gamma f_0(2100) \rightarrow \gamma \eta \eta$	$(1.13 \pm 0.60) \times 10^{-4}$
Γ_{306}	$\gamma f_0(2100) \rightarrow \gamma K \bar{K}$	
Γ_{307}	$\gamma f_0(2100) \rightarrow \gamma \pi \pi$	$(6.2 \pm 1.0) \times 10^{-4}$
Γ_{308}	$\gamma f_0(2200)$	
Γ_{309}	$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$(5.9 \pm 1.3) \times 10^{-4}$
Γ_{310}	$\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0$	$(2.72 \pm 0.19) \times 10^{-4}$
Γ_{311}	$\gamma f_0(2200) \rightarrow \gamma \pi \pi$	
Γ_{312}	$\gamma f_0(2200) \rightarrow \gamma \eta \eta$	
Γ_{313}	$\gamma f_J(2220)$	
Γ_{314}	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	$< 3.9 \times 10^{-5}$ CL=90%
Γ_{315}	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	$< 4.1 \times 10^{-5}$ CL=90%
Γ_{316}	$\gamma f_J(2220) \rightarrow \gamma p \bar{p}$	$(1.5 \pm 0.8) \times 10^{-5}$
Γ_{317}	$\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0$	$(4.9 \pm 0.7) \times 10^{-5}$
Γ_{318}	$\gamma f_0(2330) \rightarrow \gamma \pi \pi$	
Γ_{319}	$\gamma f_0(2330) \rightarrow \gamma \eta \eta$	
Γ_{320}	$\gamma f_0(2330) \rightarrow \gamma \eta' \eta'$	$(6.1 \pm 4.0) \times 10^{-6}$
Γ_{321}	$\gamma f_0(2330) \rightarrow \gamma \eta \eta'$	
Γ_{322}	$\gamma f_2(2340) \rightarrow \gamma \eta \eta$	$(5.6 \pm 2.4) \times 10^{-5}$
Γ_{323}	$\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0$	$(5.5 \pm 4.0) \times 10^{-5}$
Γ_{324}	$\gamma f_2(2340) \rightarrow \gamma \eta' \eta'$	$(8.7 \pm 0.9) \times 10^{-6}$
Γ_{325}	$\gamma f_0(2470) \rightarrow \gamma \eta' \eta'$	$(8.2 \pm 4.0) \times 10^{-7}$
Γ_{326}	$\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta'$	$(2.7 \pm 0.6) \times 10^{-4}$ S=1.6
Γ_{327}	$\gamma X(1835) \rightarrow \gamma p \bar{p}$	$(7.7 \pm 1.5) \times 10^{-5}$
Γ_{328}	$\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta$	$(3.3 \pm 2.0) \times 10^{-5}$
Γ_{329}	$\gamma X(1835) \rightarrow \gamma \gamma \phi(1020)$	
Γ_{330}	$\gamma X(1835) \rightarrow \gamma \gamma \gamma$	$< 3.56 \times 10^{-6}$ CL=90%
Γ_{331}	$\gamma X(1835) \rightarrow \gamma 3(\pi^+ \pi^-)$	$(2.4 \pm 0.7) \times 10^{-5}$

Γ_{332}	$\gamma X(2370) \rightarrow \gamma K^+ K^- \eta'$	$(1.8 \pm 0.7) \times 10^{-5}$		
Γ_{333}	$\gamma X(2370) \rightarrow \gamma K_S^0 K_S^0 \eta'$	$(1.2 \pm 0.5) \times 10^{-5}$		
Γ_{334}	$\gamma X(2370) \rightarrow \gamma \eta \eta \eta'$	$< 9.2 \times 10^{-6}$	CL=90%	
Γ_{335}	$\gamma p \bar{p}$	$(3.8 \pm 1.0) \times 10^{-4}$		
Γ_{336}	$\gamma p \bar{p} \pi^+ \pi^-$	$< 7.9 \times 10^{-4}$	CL=90%	
Γ_{337}	$\gamma \Lambda \bar{\Lambda}$	$< 1.3 \times 10^{-4}$	CL=90%	
Γ_{338}	$\gamma A^0 \rightarrow \gamma \text{invisible}$	[e] $< 1.7 \times 10^{-6}$	CL=90%	
Γ_{339}	$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	[f] $< 7.8 \times 10^{-7}$	CL=90%	

Dalitz decays

Γ_{340}	$\pi^0 e^+ e^-$	$(7.6 \pm 1.4) \times 10^{-7}$		
Γ_{341}	$\eta e^+ e^-$	$(1.42 \pm 0.08) \times 10^{-5}$		
Γ_{342}	$\eta'(958) e^+ e^-$	$(6.59 \pm 0.18) \times 10^{-5}$		
Γ_{343}	$X(1835) e^+ e^-$, $X \rightarrow \pi^+ \pi^- \eta'$	$(3.58 \pm 0.25) \times 10^{-6}$		
Γ_{344}	$X(2120) e^+ e^-$, $X \rightarrow \pi^+ \pi^- \eta'$	$(8.2 \pm 1.3) \times 10^{-7}$		
Γ_{345}	$X(2370) e^+ e^-$, $X \rightarrow \pi^+ \pi^- \eta'$	$(1.08 \pm 0.17) \times 10^{-6}$		
Γ_{346}	$\eta U \rightarrow \eta e^+ e^-$	[g] $< 9.11 \times 10^{-7}$	CL=90%	
Γ_{347}	$\eta'(958) U \rightarrow \eta'(958) e^+ e^-$	[g] $< 2.0 \times 10^{-7}$	CL=90%	
Γ_{348}	$\phi e^+ e^-$	$< 1.2 \times 10^{-7}$	CL=90%	

Weak decays

Γ_{349}	$D^- e^+ \nu_e + \text{c.c.}$	$< 7.1 \times 10^{-8}$	CL=90%	
Γ_{350}	$\bar{D}^0 e^+ e^- + \text{c.c.}$	$< 8.5 \times 10^{-8}$	CL=90%	
Γ_{351}	$D_s^- e^+ \nu_e + \text{c.c.}$	$< 1.3 \times 10^{-6}$	CL=90%	
Γ_{352}	$D_s^{*-} e^+ \nu_e + \text{c.c.}$	$< 1.8 \times 10^{-6}$	CL=90%	
Γ_{353}	$D^- \pi^+ + \text{c.c.}$	$< 7.5 \times 10^{-5}$	CL=90%	
Γ_{354}	$\bar{D}^0 \bar{K}^0 + \text{c.c.}$	$< 1.7 \times 10^{-4}$	CL=90%	
Γ_{355}	$\bar{D}^0 \bar{K}^{*0} + \text{c.c.}$	$< 2.5 \times 10^{-6}$	CL=90%	
Γ_{356}	$D_s^- \pi^+ + \text{c.c.}$	$< 1.3 \times 10^{-4}$	CL=90%	
Γ_{357}	$D_s^- \rho^+ + \text{c.c.}$	$< 1.3 \times 10^{-5}$	CL=90%	

Charge conjugation (C), Parity (P), Lepton Family number (LF) violating modes

Γ_{358}	$\gamma \gamma$	$C < 2.7 \times 10^{-7}$	CL=90%	
Γ_{359}	$\gamma \phi$	$C < 1.4 \times 10^{-6}$	CL=90%	
Γ_{360}	$e^\pm \mu^\mp$	$LF < 1.6 \times 10^{-7}$	CL=90%	
Γ_{361}	$e^\pm \tau^\mp$	$LF < 7.5 \times 10^{-8}$	CL=90%	
Γ_{362}	$\mu^\pm \tau^\mp$	$LF < 2.0 \times 10^{-6}$	CL=90%	
Γ_{363}	$\Lambda_c^+ e^- + \text{c.c.}$	$< 6.9 \times 10^{-8}$	CL=90%	

Other decays

Γ_{364}	invisible	$< 7 \times 10^{-4}$	CL=90%	
----------------	-----------	----------------------	--------	--

- [a] For $E_\gamma > 100$ MeV.
- [b] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [c] $\Theta(1540)$ is a hypothetical pentaquark state of 1.54 GeV/c 2 mass and a width of less than 25 MeV/c 2 .
- [d] Includes $p\bar{p}\pi^+\pi^-\gamma$ and excludes $p\bar{p}\eta$, $p\bar{p}\omega$, $p\bar{p}\eta'$.
- [e] For a narrow state A with mass less than 960 MeV.
- [f] For a narrow scalar or pseudoscalar A^0 with mass 0.21 – 3.0 GeV.
- [g] For a dark photon U with mass between 100 and 2100 MeV.

J/ ψ (1S) PARTIAL WIDTHS

$\Gamma(\text{hadrons})$

Γ_1

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
81.37± 1.36±1.30	¹ ANASHIN	20	KEDR e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •			
74.1 ± 8.1	BAI	95B	BES e^+e^-
59 ± 24	BALDINI...	75	FRAG e^+e^-
59 ± 14	BOYARSKI	75	MRK1 e^+e^-
50 ± 25	ESPOSITO	75B	FRAM e^+e^-

¹ Based on the same dataset as ANASHIN 18A and correlated to the values reported there

$\Gamma(e^+e^-)$

Γ_5

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.53 ±0.10 OUR AVERAGE				
5.550±0.056±0.089		^{1,2} ANASHIN	18A	KEDR e^+e^-
5.36 $^{+0.29}_{-0.28}$		³ HSUEH	92	RVUE See γ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.58 ±0.05 ±0.08		⁴ ABLIKIM	16Q	BES3 $e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.71 ±0.16	13k	⁵ ADAMS	06A	CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.57 ±0.19	7.8k	⁵ AUBERT	04	BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.14 ±0.39		BAI	95B	BES e^+e^-
4.72 ±0.35		ALEXANDER	89	RVUE See γ mini-review
4.4 ±0.6		³ BRANDELIK	79c	DASP e^+e^-
4.6 ±0.8		⁶ BALDINI...	75	FRAG e^+e^-
4.8 ±0.6		BOYARSKI	75	MRK1 e^+e^-
4.6 ±1.0		ESPOSITO	75B	FRAM e^+e^-

¹ From the cross sections of $e^+e^- \rightarrow e^+e^-$ and $e^+e^- \rightarrow$ hadrons near the J/ ψ (1S) peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

⁴ Using $B(J/\psi \rightarrow \mu^+\mu^-) = (5.973 \pm 0.007 \pm 0.037)\%$ from ABLIKIM 13R.

⁵ Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.

⁶ Assuming equal partial widths for e^+e^- and $\mu^+\mu^-$.

$\Gamma(\mu^+ \mu^-)$				Γ_7
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.13 \pm 0.52	BAI	95B	BES $e^+ e^-$	
4.8 \pm 0.6	BOYARSKI	75	MRK1 $e^+ e^-$	
5 \pm 1	ESPOSITO	75B	FRAM $e^+ e^-$	

$\Gamma(\gamma\gamma)$				Γ_{358}
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.4	90	BRANDELIK	79C	DASP $e^+ e^-$

$J/\psi(1S) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into $e^+ e^-$ and with the total width is obtained from the integrated cross section into channel(I) in the $e^+ e^-$ annihilation.

$\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$				$\Gamma_1 \Gamma_5 / \Gamma$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.884 \pm 0.048 \pm 0.078	1,2 ANASHIN	18A	KEDR $e^+ e^-$	
4 \pm 0.8	3 BALDINI-...	75	FRAG $e^+ e^-$	
3.9 \pm 0.8	3 ESPOSITO	75B	FRAM $e^+ e^-$	

¹ From the cross sections of $e^+ e^- \rightarrow e^+ e^-$ and $e^+ e^- \rightarrow$ hadrons near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ Data redundant with branching ratios or partial widths above.

$\Gamma(e^+ e^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$				$\Gamma_5 \Gamma_5 / \Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
333.1 \pm 6.6 \pm 4.0	1,2 ANASHIN	18A	KEDR $e^+ e^-$	
332.3 \pm 6.4 \pm 4.8	ANASHIN	10	KEDR $3.097 e^+ e^- \rightarrow e^+ e^-$	
350 \pm 20	BRANDELIK	79C	DASP $e^+ e^-$	
320 \pm 70	3 BALDINI-...	75	FRAG $e^+ e^-$	
340 \pm 90	3 ESPOSITO	75B	FRAM $e^+ e^-$	
360 \pm 100	3 FORD	75	SPEC $e^+ e^-$	

¹ From the cross sections of $e^+ e^- \rightarrow e^+ e^-$ and $e^+ e^- \rightarrow$ hadrons near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ Data redundant with branching ratios or partial widths above.

$\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$				$\Gamma_7 \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
333 \pm 4 OUR AVERAGE				
333.4 \pm 2.5 \pm 4.4	ABLIKIM	16Q	BES3 $3.773 e^+ e^- \rightarrow \mu^+ \mu^- \gamma$	
331.8 \pm 5.2 \pm 6.3	ANASHIN	10	KEDR $3.097 e^+ e^- \rightarrow \mu^+ \mu^-$	
338.4 \pm 5.8 \pm 7.1	13k	ADAMS	06A CLEO $e^+ e^- \rightarrow \mu^+ \mu^- \gamma$	
330.1 \pm 7.7 \pm 7.3	7.8k	AUBERT	04 BABR $e^+ e^- \rightarrow \mu^+ \mu^- \gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

510 ± 90	DASP	75 DASP	$e^+ e^-$
380 ± 50	¹ ESPOSITO	75B FRAM	$e^+ e^-$

¹ Data redundant with branching ratios or partial widths above.

$\Gamma(\eta\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$$\Gamma_{13}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.4 OUR AVERAGE				
2.34 $\pm 0.43 \pm 0.16$	49	LEES	18	BABR $e^+ e^- \rightarrow \eta\pi^+\pi^-\gamma$
2.22 $\pm 0.96 \pm 0.02$	9	¹ AUBERT	07AU	BABR $10.6 e^+ e^- \rightarrow \eta\pi^+\pi^-\gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+\pi^-\pi^0)] = 0.51 \pm 0.22 \pm 0.03 \text{ eV}$ which we divide by our best value $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.02 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$$\Gamma_{14}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
64.8 $\pm 11.1 \pm 0.4$				
64.8 $\pm 11.1 \pm 0.4$	200	¹ LEES	21C	BABR $e^+ e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$

¹ LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] = 21.1 \pm 1.7 \pm 3.2 \text{ eV}$ which we divide by our best value $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta\pi^+\pi^-3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$$\Gamma_{15}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
26.9 $\pm 5.7 \pm 0.1$				
26.9 $\pm 5.7 \pm 0.1$	101	¹ LEES	21C	BABR $e^+ e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-3\pi^0\gamma\gamma)$

¹ LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 10.6 \pm 1.6 \pm 1.6 \text{ eV}$ which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta K^\pm K_S^0\pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$$\Gamma_{19}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
7.3 $\pm 1.4 \pm 0.4$				
7.3 $\pm 1.4 \pm 0.4$	44	LEES	17D	BABR $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$\Gamma(\rho^\pm\pi^\mp\pi^+\pi^-2\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$$\Gamma_{22}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
155 $\pm 26 \pm 36$				
155 $\pm 26 \pm 36$	14k	LEES	21	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

$\Gamma(\rho^+\rho^-\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$$\Gamma_{23}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
32 $\pm 13 \pm 15$				
32 $\pm 13 \pm 15$	14k	LEES	21	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

$\Gamma(\rho^\mp K^\pm K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$$\Gamma_{25}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.4 $\pm 1.0 \pm 1.9$				
10.4 $\pm 1.0 \pm 1.9$	130	LEES	17D	BABR $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{40}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
53.6±5.0±0.4	788	1 AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 47.8 \pm 3.1 \pm 3.2$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\omega\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{41}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
27.8±3.5±0.2	398	1 LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

¹ LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 24.8 \pm 1.8 \pm 2.5$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\omega 3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{42}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.5±3.1±0.1	89	1 LEES	21C BABR	$e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$

¹ LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \omega 3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 9.4 \pm 2.3 \pm 1.5$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\omega\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{45}\Gamma_5/\Gamma$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.2±0.3±0.2	170	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\pi^0\gamma$

 $\Gamma(\omega\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{44}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
16.9±7.6±0.2	1 LEES	21C BABR	$e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$

¹ Different final state as in AUBERT 06. LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \omega\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 4.9 \pm 2.1 \pm 0.7$ eV which we divide by our best values $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.21) \times 10^{-2}$, $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

 $\Gamma(\omega\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{46}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.90±0.96±0.01	27	1 LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

¹ LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^0\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 1.7 \pm 0.8 \pm 0.3$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\omega\pi^+\pi^-2\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{48}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
185±30±1	14k	¹ LEES	21 BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-2\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 165 \pm 9 \pm 25 \text{ eV}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\omega K\bar{K}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{59}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.70±1.98±0.03	24	¹ AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega K^+K^-\gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega K\bar{K}) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 3.3 \pm 1.3 \pm 1.2 \text{ eV}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^+K^*(892)^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{68}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
29.0±1.7±1.3		AUBERT	08S BABR	$10.6 e^+e^- \rightarrow K^+K^*(892)^-\gamma$

$$\Gamma(K^+K^*(892)^- + \text{c.c.} \rightarrow K^+K^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{69}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.96±0.85±0.70	155	AUBERT	08S BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^0\gamma$

$$\Gamma(K^+K^*(892)^- + \text{c.c.} \rightarrow K^0K^\pm\pi^\mp + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{70}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
16.76±1.70±1.00	89	AUBERT	08S BABR	$10.6 e^+e^- \rightarrow K_S^0K^\pm\pi^\mp\gamma$

$$\Gamma(K^0\bar{K}^*(892)^0 + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{71}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
26.6±2.5±1.5		AUBERT	08S BABR	$10.6 e^+e^- \rightarrow K^0\bar{K}^*(892)^0\gamma$

$$\Gamma(K^0\bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0K^\pm\pi^\mp + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{72}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
17.70±1.70±1.00	94	AUBERT	08S BABR	$10.6 e^+e^- \rightarrow K_S^0K^\pm\pi^\mp\gamma$

$$\Gamma(\bar{K}^*(892)^0K^+\pi^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{73}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
42.6±4.8±7.2	99	¹ LEES	17D BABR	$e^+e^- \rightarrow K_S^0K^\pm\pi^\mp\pi^0\gamma$

¹ Dividing by 1/6 to account for $B(K^*(892)^0 \rightarrow K_S^0\pi^0) = 1/6$.

$$\Gamma(K^*(892)^\pm K^\mp\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{74}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
22.8±2.8±6.8	80	¹ LEES	17D BABR	$e^+e^- \rightarrow K_S^0K^\pm\pi^\mp\pi^0\gamma$

¹ Dividing by 1/4 to account for $B(K^*(892)^\pm \rightarrow K_S^0\pi^\pm) = 1/4$.

$$\Gamma(K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{75} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.0 ± 2.8 OUR AVERAGE				
9.2 ± 1.2 ± 3.2	64	1 LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$
14.8 ± 4.8 ± 1.2	53	2 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by 1/2 to take into account $B(K^*(892)^\pm \rightarrow K^\pm \pi^\mp) = 1/2$.

² Dividing by 1/4 to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$$\Gamma(K^*(892)^+ K_S^0 \pi^- + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{76} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.7 ± 1.2 ± 0.3	53	LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

$$\Gamma(K^*(892)^0 K_S^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{79} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.60 ± 0.75 ± 2.25	34	1 LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

¹ Dividing by 2/3 to account for $B(K^*(892)^0 \rightarrow K^+ \pi^-) = 2/3$.

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{81} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.28 ± 0.34 ± 0.07	47 ± 12	1 LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$$1.28 \pm 0.40 \pm 0.11 \quad 25 \pm 8 \quad ^{1,2} \text{ AUBERT } 07AK \text{ BABR} \quad 10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$$

¹ Dividing by $(2/3)^2$ to take twice into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3 B(K^{*0} \rightarrow K\pi)$.

² Superseded by LEES 12F.

$$\Gamma(K^*(892)^\pm K^*(892)^\mp) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{82} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.80 ± 0.48 ± 0.32	1 ± 5	1 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by $(1/4)^2$ to take twice into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$$\Gamma(K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{93} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.1 ± 9.8 ± 0.5	35	1,2 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by 1/4 to take into account $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K\pi)$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)] / [\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 10.0 \pm 4.8 \pm 0.8 \text{ eV}$ which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K_2^*(1430)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{92} / \Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.65 ± 0.80 ± 0.44	1094	ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

$$\Gamma(\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{94} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

25.8±1.4±0.6 710 1,2,3 LEES 12F BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

33 ± 4 ± 1 317 2,4 AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 12.89 \pm 0.54 \pm 0.41$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3 B(K^{*0} \rightarrow K\pi)$.

³ The $K_2^*(1430)$ cannot be distinguished from the $K_0^*(1430)$.

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 16.4 \pm 1.1 \pm 1.4$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K_2^*(1430)^- K^*(892)^+ + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{95} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

18.6±16.1±0.4 8 ± 8 1,2 LEES 14H BABR $e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by $(1/4)^2$ to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ and $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K\pi)$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \bar{K}_2^*(1430)^- K^*(892)^+ + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 9.28 \pm 8.0 \pm 0.32$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K_2^*(1430)^- K^*(892)^+ + \text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{96} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

2.32±2.00±0.08 8 ± 8 1 LEES 14H BABR $e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by 1/4 to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$$\Gamma(\bar{K}_2(1770)^0 K^*(892)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{98} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

3.8±0.4±0.3 110 ± 14 1 AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3$.

$$\Gamma(\phi \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{105} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

4.48±0.35 OUR AVERAGE

4.46 $\pm 0.49 \pm 0.05$ 181 1 LEES 12F BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

4.51 $\pm 0.48 \pm 0.05$ 254 ± 23 2 SHEN 09 BELL 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.3 \pm 0.7 \pm 0.1$

103^3 AUBERT,BE 06D BABR $10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.19 \pm 0.23 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² SHEN 09 reports $4.50 \pm 0.41 \pm 0.26$ eV from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)]$ assuming $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.6) \times 10^{-2}$, which we rescale to our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Superseded by LEES 12F. AUBERT,BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.61 \pm 0.30 \pm 0.18$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$\Gamma_{106}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.77±0.57±0.03	45	¹ LEES 12F	BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.13 \pm 0.88 \pm 0.03$ 23^2 AUBERT,BE 06D BABR $10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.36 \pm 0.27 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by LEES 12F. AUBERT,BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.54 \pm 0.40 \pm 0.16$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi 2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$\Gamma_{107}\Gamma_5/\Gamma$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.96±0.19±0.01	35	¹ AUBERT 06D	BABR	$10.6 e^+ e^- \rightarrow \phi 2(\pi^+\pi^-) \gamma$

¹ AUBERT 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi 2(\pi^+\pi^-)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = (0.47 \pm 0.09 \pm 0.03) \times 10^{-2}$ keV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$\Gamma_{108}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.1±2.7±0.4	6	¹ AUBERT 07AU	BABR	$10.6 e^+ e^- \rightarrow \phi\eta\gamma$

¹ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi) = 0.84 \pm 0.37 \pm 0.05$ eV.

$\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{112}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.44±0.19 OUR AVERAGE				
1.40±0.25±0.02	57 ± 9	¹ LEES	12F BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
1.48±0.27±0.09	60 ± 11	² SHEN	09 BELL	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.02±0.24±0.01	20 ± 5	³ AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.69 \pm 0.11 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
² Multiplied by 2/3 to take into account the $\phi\pi^+\pi^-$ mode only. Using $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.6)\%$.				
³ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.50 \pm 0.11 \pm 0.04$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(\phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{113}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.98±0.26±0.01				
16 ± 4	¹ LEES	12F BABR	$10.6 e^+e^- \rightarrow \pi^0\pi^0K^+K^-\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.96±0.40±0.01	7.0 ± 2.8	² AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow \pi^0\pi^0K^+K^-\gamma$
¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.48 \pm 0.12 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.47 \pm 0.19 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(\phi f_2(1270)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{118}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.79±0.32^{+0.02}_{-0.06}				
61	1,2,3 LEES	12F BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.08±0.73 ^{+0.04} _{-0.14}	44	^{2,4} AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = 1.51 \pm 0.25 \pm 0.10$ eV which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
² Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$.				
³ Using $\pi^+\pi^-$ invariant mass between 1.1 and 1.5 GeV. May include other sources such as $f_0(1370)$.				

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = 3.44 \pm 0.55 \pm 0.28 \text{ eV}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f'_2(1525)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{123}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.2±3.2±0.2	11	1,2 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹ Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K\bar{K})$ and using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f'_2(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = 7.2 \pm 2.8 \pm 0.3 \text{ eV}$ which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{129}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.60±0.62±0.05	163	1 LEES	12F BABR	$e^+ e^- \rightarrow K^+ K^- K^+ K^- \gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.26 \pm 0.26 \pm 0.16 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi K_S^0 K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{130}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.26±0.84±0.04	29	1 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹ LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi K_S^0 K_S^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 1.6 \pm 0.4 \pm 0.1 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(f'_2(1525) K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{135}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.8±1.9±0.1	16	1,2 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹ Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K\bar{K})$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow f'_2(1525) K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = 5.12 \pm 1.68 \pm 0.20 \text{ eV}$ which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+ \pi^-)\pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{154}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
303±5±18	4990	AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-)\pi^0 \gamma$

$\Gamma(\pi^+\pi^-3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{156}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
104 ± 50 OUR AVERAGE				Error includes scale factor of 4.3.
55.4 ± 15.9 ± 0.5	14k	1 LEES	21 BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$
150.0 ± 4.0 ± 15.0	2.3k	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] = 19.2 \pm 4.5 \pm 3.2$ eV which we divide by our best value $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}} = 0.3468 \pm 0.0030$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\pi^+\pi^-4\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{157}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
35.8 ± 4.4 ± 5.4	340	LEES	21C BABR	$e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$

 $\Gamma(\rho^\pm\pi^\mp\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{158}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
78.0 ± 9.0 ± 8.0	1.2k	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

 $\Gamma(\rho^+\rho^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{159}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
33.0 ± 5.0 ± 3.3	529	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

 $\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{160}\Gamma_5/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.1248 ± 0.0019 ± 0.0026		LEES	21B BABR	$10.5 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.122 ± 0.005 ± 0.008	AUBERT,B	04N BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
-----------------------	----------	----------	---

 $\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{161}\Gamma_5/\Gamma$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.9 ± 0.5 ± 1.0	761	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$

 $\Gamma(\pi^+\pi^-\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{162}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
107.0 ± 4.3 ± 6.4	768	AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\pi^0\gamma$

 $\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{164}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.4 ± 0.9 ± 0.4		LEES	12E BABR	$10.6 e^+e^- \rightarrow 2\pi^+2\pi^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

19.5 ± 1.4 ± 1.3	270	¹ AUBERT	05D BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)\gamma$
------------------	-----	---------------------	----------	---

¹ Superseded by LEES 12E.

 $\Gamma(3(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{165}\Gamma_5/\Gamma$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.37 ± 0.16 ± 0.14	496	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow 3(\pi^+\pi^-)\gamma$

$$\Gamma(2(\pi^+\pi^-)3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{166}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
345±10±50	14k	LEES	21	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

$$\Gamma(2(\pi^+\pi^-)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{168}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1±2.4±0.1	85	1 AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)\eta\gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-)\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 5.16 \pm 0.85 \pm 0.39 \text{ eV}$ which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(2(\pi^+\pi^-\pi^0)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{170}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.1±2.6±1.4	14k	LEES	21	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

$$\Gamma(\pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{171}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1± 2.7 OUR AVERAGE				

26.1±17.9±0.3	14k	¹ LEES	21	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$
12.8± 1.8±2.0	203	LEES	18E	BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+\pi^-\pi^0)] = 6 \pm 4 \pm 1 \text{ eV}$ which we divide by our best value $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.02 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\rho^\pm\pi^\mp\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{172}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.5±4.1±1.6	168	LEES	18E	BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

$$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{173}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

1.78±0.11±0.05	462	¹ LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$
1.94±0.11±0.05	462	² LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$
1.42±0.23±0.08	51	³ LEES	13Q	BABR $e^+e^- \rightarrow K^+K^-\gamma$

¹ $\sin\phi > 0$.

² $\sin\phi < 0$.

³ Interference with non-resonant K^+K^- production not taken into account.

$$\Gamma(K_S^0 K_L^0 \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{179}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.4±1.3±0.6	182	LEES	17A	BABR $e^+e^- \rightarrow K_S^0 K_L^0 \pi^0\gamma$

$$\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{180}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.7±0.9±0.4	106	LEES	17A	BABR $e^+e^- \rightarrow K_S^0 K_L^0 \pi^0\gamma$

$$\Gamma(K_2^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{181} \Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.4±0.7±0.1	37	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

$$\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{182} \Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
37.94±0.81±1.10	3.1k	LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$$36.3 \pm 1.3 \pm 2.1 \quad 1.5k \quad ^1 \text{AUBERT} \quad 07AK \text{BABR} \quad 10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$$

$$33.6 \pm 2.7 \pm 2.7 \quad 233 \quad ^2 \text{AUBERT} \quad 05D \text{BABR} \quad 10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$$

¹ Superseded by LEES 12F.

² Superseded by AUBERT 07AK.

$$\Gamma(K^+ K^- \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{183} \Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.75±0.81±0.90	388	LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$$13.6 \pm 1.1 \pm 1.3 \quad 203 \quad ^1 \text{AUBERT} \quad 07AK \text{BABR} \quad 10.6 e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$$

¹ Superseded by LEES 12F.

$$\Gamma(K_S^0 K_L^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{184} \Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.8±2.3±2.1	248	LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_L^0 \gamma$

$$\Gamma(K_S^0 K_L^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{185} \Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.3±2.3±0.5	47	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \pi^0 \gamma$

$$\Gamma(K_S^0 K_L^0 \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{186} \Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.0±1.8±0.4	45	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \eta \gamma$

$$\Gamma(K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{187} \Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.3±0.9±0.5	133	LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

$$\Gamma(K^\mp K_S^0 \pi^\pm \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{188} \Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
31.7±1.9±1.8	393	LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$$\Gamma(K^+ K^- 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{189} \Gamma_5/\Gamma$$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.75±0.23±0.17	205	AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$

$\Gamma(K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{190} \Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
25.9±3.9±0.1	73	1 AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 10.2 \pm 1.3 \pm 0.8$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(K^+ K^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{191} \Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.00±0.33±0.29	287 ± 24	LEES	12F BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-)\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.11±0.39±0.30	156 ± 15	¹ AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-)\gamma$
4.0 ± 0.7 ± 0.6	38	² AUBERT	05D BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-)\gamma$

¹ Superseded by LEES 12F.

² Superseded by AUBERT 07AK.

$\Gamma(K^+ K^- K_S^0 K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{192} \Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.3±0.4±0.1	29	LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

$\Gamma(p\bar{p}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{193} \Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.9±0.6 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below.		

11.3±0.4±0.3 821 ¹ LEES 13O BABR $e^+ e^- \rightarrow p\bar{p}\gamma$

12.9±0.4±0.4 918 ² LEES 13Y BABR $e^+ e^- \rightarrow p\bar{p}\gamma$


9.7±1.7 3 ARMSTRONG 93B E760 $\bar{p}p \rightarrow e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

12.0±0.6±0.5 438 ⁴ AUBERT 06B BABR $e^+ e^- \rightarrow p\bar{p}\gamma$

WEIGHTED AVERAGE
11.9±0.6 (Error scaled by 1.8)

Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.



$$\Gamma(p\bar{p}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} (\text{eV})$$

¹ ISR photon reconstructed in the detector

² ISR photon undetected³ Using $\Gamma_{\text{total}} = 85.5^{+6.1}_{-5.8}$ MeV.⁴ Superseded by LEES 130

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{209}\Gamma_5/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
10.7±0.9±0.7	AUBERT	07BD BABR	$10.6 e^+ e^- \rightarrow \Lambda\bar{\Lambda}\gamma$
$\Gamma(\Sigma^0\bar{\Sigma}^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{219}\Gamma_5/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
6.4±1.2±0.6	AUBERT	07BD BABR	$10.6 e^+ e^- \rightarrow \Sigma^0\bar{\Sigma}^0\gamma$

J/ $\psi(1S)$ BRANCHING RATIOS

For the first four branching ratios, see also the partial widths, and (partial widths) $\times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ above.

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.877±0.005 OUR AVERAGE			
0.878±0.005	BAI	95B	BES $e^+ e^-$
0.86 ± 0.02	BOYARSKI	75	MRK1 $e^+ e^-$

$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.135±0.003	1,2 SETH	04	RVUE $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.17 ± 0.02	1 BOYARSKI	75	MRK1 $e^+ e^-$

¹ Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.² Using $B(J/\psi \rightarrow \ell^+\ell^-) = (5.90 \pm 0.09)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C.

$\Gamma(ggg)/\Gamma_{\text{total}}$	Γ_3/Γ		
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN
64.1±1.0	6 M	¹ BESSON	08 CLEO $\psi(2S) \rightarrow \pi^+\pi^- + \text{hadrons}$
1 Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the PDG 08 values of $B(\ell^+\ell^-)$, $B(\text{virtual } \gamma \rightarrow \text{hadrons})$, and $B(\gamma\eta_c)$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 08.			

$\Gamma(\gamma gg)/\Gamma_{\text{total}}$	Γ_4/Γ		
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN
8.79±1.05	200 k	¹ BESSON	08 CLEO $\psi(2S) \rightarrow \pi^+\pi^- \gamma + \text{hadrons}$
1 Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the value of $\Gamma(ggg)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(ggg)/\Gamma_{\text{total}}$ measurement of BESSON 08.			

$\Gamma(\gamma gg)/\Gamma(ggg)$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>
$13.7 \pm 0.1 \pm 0.7$	6 M

Γ_4/Γ_3

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>
5.971 ± 0.032 OUR AVERAGE	
5.983 $\pm 0.007 \pm 0.037$	720k
5.945 $\pm 0.067 \pm 0.042$	15k
5.90 $\pm 0.05 \pm 0.10$	
6.09 ± 0.33	
5.92 $\pm 0.15 \pm 0.20$	
6.9 ± 0.9	

Γ_5/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
BAI	95B	BES $e^+ e^-$
COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
BOYARSKI	75	MRK1 $e^+ e^-$

$\Gamma(e^+ e^- \gamma)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>
$8.8 \pm 1.3 \pm 0.4$

¹ For $E_\gamma > 100$ MeV.

Γ_6/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
¹ ARMSTRONG	96	$\bar{p}p \rightarrow e^+ e^- \gamma$

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>
5.961 ± 0.033 OUR AVERAGE	
5.973 $\pm 0.007 \pm 0.038$	770k
5.960 $\pm 0.065 \pm 0.050$	17k
5.84 $\pm 0.06 \pm 0.10$	
6.08 ± 0.33	
5.90 $\pm 0.15 \pm 0.19$	
6.9 ± 0.9	

Γ_7/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
BAI	95B	BES $e^+ e^-$
COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
BOYARSKI	75	MRK1 $e^+ e^-$

$\Gamma(e^+ e^-)/\Gamma(\mu^+ \mu^-)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.0016 ± 0.0031 OUR AVERAGE			
1.0022 $\pm 0.0044 \pm 0.0048$	¹ AULCHENKO	14	KEDR $3.097 e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
1.0017 $\pm 0.0017 \pm 0.0033$	² ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.002 $\pm 0.021 \pm 0.013$	³ ANASHIN	10	KEDR $3.097 e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
0.997 $\pm 0.012 \pm 0.006$	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.011 $\pm 0.013 \pm 0.016$	BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.00 ± 0.07	BAI	95B	BES $e^+ e^-$
1.00 ± 0.05	BOYARSKI	75	MRK1 $e^+ e^-$
0.91 ± 0.15	ESPOSITO	75B	FRAM $e^+ e^-$
0.93 ± 0.10	FORD	75	SPEC $e^+ e^-$

Γ_5/Γ_7


¹ From 235.3k $J/\psi \rightarrow e^+ e^-$ and 156.6k $J/\psi \rightarrow \mu^+ \mu^-$ observed events.

² Not independent of the corresponding measurements of $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$.

³ Not independent of the corresponding measurements of $\Gamma(e^+ e^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$.

HADRONIC DECAYS $\Gamma(\rho\pi)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_8/Γ
1.69 ± 0.15 OUR AVERAGE				Error includes scale factor of 2.4. See the ideogram below.	
2.18 ± 0.19		1,2 AUBERT,B	04N BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$	
2.184 ± 0.005 ± 0.201	220k	2,3 BAI	04H BES	$e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$	
2.091 ± 0.021 ± 0.116		2,4 BAI	04H BES	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	
1.21 ± 0.20		BAI	96D BES	$e^+ e^- \rightarrow \rho\pi$	
1.42 ± 0.01 ± 0.19		COFFMAN	88 MRK3	$e^+ e^-$	
1.3 ± 0.3	150	FRANKLIN	83 MRK2	$e^+ e^-$	
1.6 ± 0.4	183	ALEXANDER	78 PLUT	$e^+ e^-$	
1.33 ± 0.21		BRANDELIK	78B DASP	$e^+ e^-$	
1.0 ± 0.2	543	BARTEL	76 CNTR	$e^+ e^-$	
1.3 ± 0.3	153	JEAN-MARIE	76 MRK1	$e^+ e^-$	

WEIGHTED AVERAGE
1.69±0.15 (Error scaled by 2.4)¹ From the ratio of $\Gamma(e^+ e^-) B(\pi^+ \pi^- \pi^0)$ and $\Gamma(e^+ e^-) B(\mu^+ \mu^-)$ (AUBERT 04).² Not independent of their $B(\pi^+ \pi^- \pi^0)$.³ From $J/\psi \rightarrow \pi^+ \pi^- \pi^0$ events directly.⁴ Obtained comparing the rates for $\pi^+ \pi^- \pi^0$ and $\mu^+ \mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+ \mu^-) = 5.88 \pm 0.10\%$.

$\Gamma(\rho\pi)/\Gamma(\pi^+\pi^-\pi^0)$				Γ_8/Γ_{160}
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.142±0.011±0.026	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.331±0.033	20k	² LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model.
² From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho^0\pi^0)/\Gamma(\rho\pi)$				Γ_9/Γ_8
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.328±0.005±0.027		COFFMAN	88	MRK3 e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.35 ± 0.08		ALEXANDER	78	PLUT e^+e^-
0.32 ± 0.08		BRANDELIK	78B	DASP e^+e^-
0.39 ± 0.11		BARTEL	76	CNTR e^+e^-
0.37 ± 0.09		JEAN-MARIE	76	MRK1 e^+e^-

$\Gamma(\rho^+K^+K^-\pi^- + c.c \rightarrow K^+K^-\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$				Γ_{24}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.53±0.16±0.81	485	ANASHIN	22	KEDR $J/\psi \rightarrow K^+K^-\pi^+\pi^-\pi^0$

$\Gamma(a_2(1320)^0\pi^+\pi^- \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$				Γ_{10}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.84±0.08±0.60	1317	ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(a_2(1320)^+\pi^-\pi^0 + c.c \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$				Γ_{11}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.67±0.09±0.73	1628	ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(a_2(1320)\rho)/\Gamma_{\text{total}}$				Γ_{12}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
10.9±2.2 OUR AVERAGE				
11.7±0.7±2.5	7584	AUGUSTIN	89	$J/\psi \rightarrow \rho^0\rho^\pm\pi^\mp$
8.4±4.5	36	VANNUCCI	77	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$				Γ_{13}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.78±0.68	471	¹ ABLIKIM	19Q BES3	$e^+e^- \rightarrow J/\psi \rightarrow \eta\pi^+\pi^-$
¹ From an energy scan of $e^+e^- \rightarrow J/\psi \rightarrow \eta\pi^+\pi^-$ assuming PDG 16 values for $\Gamma(e^+e^-)$, $\Gamma(\mu^+\mu^-)$, and $\Gamma(\text{total})$.				

$\Gamma(\eta\rho)/\Gamma_{\text{total}}$				Γ_{16}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.193±0.023 OUR AVERAGE				
0.194±0.017±0.029	299	JOUSSET	90	$J/\psi \rightarrow \text{hadrons}$
0.193±0.013±0.029		COFFMAN	88	$e^+e^- \rightarrow \pi^+\pi^-\eta$

$\Gamma(\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.20 \pm 0.14 \pm 0.37$	471	ABLIKIM	15H BES3	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

$\Gamma(\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.52 \times 10^{-4}$	90	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+\pi^-\bar{K}^-\pi^+$

$\Gamma(\eta K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$21.8 \pm 2.2 \pm 3.4$	232 ± 23	ABLIKIM	08E BES2	$e^+e^- \rightarrow J/\psi$

$\Gamma(\eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.15 \pm 0.13 \pm 0.22$	209	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+\pi^-\bar{K}^-\pi^+$

$\Gamma(\rho\eta'(958))/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
8.1 ± 0.8 OUR AVERAGE		Error includes scale factor of 1.6.		
$7.90 \pm 0.19 \pm 0.49$	3476	¹ ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'$
$8.3 \pm 3.0 \pm 1.2$	19	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$11.4 \pm 1.4 \pm 1.6$		COFFMAN	88 MRK3	$J/\psi \rightarrow \pi^+\pi^-\eta'$

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+\pi^-\eta'$.

$\Gamma(\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{29}/Γ_{160}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$10.9 \pm 1.7 \pm 2.7$	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.80 ± 0.27	20k	² LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
-----------------	-----	-------------------	----------	--------------------------------------

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho(1450)^\pm\pi^\mp \rightarrow K_S^0 K^\pm\pi^\mp)/\Gamma(K_S^0 K^\pm\pi^\mp)$ Γ_{30}/Γ_{178}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$6.3 \pm 0.8 \pm 0.6$	4k	¹ LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm\pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(\rho(1450)^0\pi^0 \rightarrow K^+K^-\pi^0)/\Gamma(K^+K^-\pi^0)$ Γ_{31}/Γ_{177}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$9.3 \pm 2.0 \pm 0.6$	2k	¹ LEES	17C BABR	$J/\psi \rightarrow K^+K^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(\rho(1450)\eta'(958) \rightarrow \pi^+\pi^-\eta'(958))/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.28 \pm 0.55 \pm 0.44$	119	¹ ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'$

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+\pi^-\eta'$.

$$\Gamma(\rho(1700)\pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0)$$

Γ_{34}/Γ_{160}

<i>VALUE</i> (units 10^{-3})	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
$8 \pm 2 \pm 5$	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

- • • We do not use the following data for averages, fits, limits, etc. • • •

22 ± 6 20k $^2\text{LEES}$ 17c BABR $J/\psi \rightarrow \pi^+ \pi^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

²From a Dalitz plot analysis in a Veneziano model.

$$\Gamma(\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0)$$

Γ₃₆/Γ₁₆₀

<i>VALUE</i> (units 10^{-4})	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
4± 1±20	20k	1 LEES	17c BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

600 ± 250 20k 2 LEES 17c BABC $J/\psi \rightarrow \pi^+ \pi^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

$$\Gamma(\rho_3(1690)\pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0)$$

Γ₃₇/Γ₁₆₀

VALUE (units 10^{-3}) EVTS DOCUMENT_ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •


4.0 ± 0.8 29k $^{+1}_{-1}$ LEES 176 BABAR $1/2^+$ $\pi^+ \pi^- \pi^0$

¹ From a Dalitz plot analysis in a Veneziano model.

$$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$$

139 / 140

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.45 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
0.538 ± 0.012 ± 0.065	2090	1 ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\pi^0$
0.360 ± 0.028 ± 0.054	222	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.482 ± 0.019 ± 0.064		COFFMAN	88 MRK3	$e^+e^- \rightarrow \pi^0\pi^+\pi^-\pi^0$



¹ Using $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$.

$$\Gamma(\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0) \quad \Gamma_{39}/\Gamma_{160}$$


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$8\pm3\pm2$	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model and significance 4.9σ .

$$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{40}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8.5 ± 1.0 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
$10.6\pm1.2\pm0.1$	3531	¹ ANASHIN	22 KEDR	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.0 ± 1.6	18058	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.8 ± 1.6	215	BURMESTER	77D PLUT	e^+e^-
6.8 ± 1.9	348	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

WEIGHTED AVERAGE
 8.5 ± 1.0 (Error scaled by 1.3)



$$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}} (\text{units } 10^{-3})$$

¹ ANASHIN 22 reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (0.946 \pm 0.016 \pm 0.108) \times 10^{-2}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\omega\pi^0\pi^0)/\Gamma_{\text{total}} \quad \Gamma_{41}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.4\pm0.3\pm0.7$	509	AUGUSTIN	89 DM2	$J/\psi \rightarrow \pi^+\pi^- 3\pi^0$

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.3±0.6 OUR AVERAGE				
4.3±0.2±0.6	5860	AUGUSTIN	89	DM2 $e^+ e^-$
4.0±1.6	70	BURMESTER	77D	PLUT $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.9±0.8	81	VANNUCCI	77	MRK1 $e^+ e^- \rightarrow 2(\pi^+ \pi^-)\pi^0$


Γ_{43}/Γ

$\Gamma(\omega\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.74 ±0.20 OUR AVERAGE				
2.352±0.273	5k	1 ABLIKIM	06F	BES2 $J/\psi \rightarrow \omega\eta$
1.44 ±0.40 ±0.14	13	2 AUBERT	06D	BABR $10.6 e^+ e^- \rightarrow \omega\eta\gamma$
1.43 ±0.10 ±0.21	378	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
1.71 ±0.08 ±0.20		COFFMAN	88	MRK3 $e^+ e^- \rightarrow 3\pi\eta$

Γ_{44}/Γ

WEIGHTED AVERAGE
1.74±0.20 (Error scaled by 1.6)



$\Gamma(\omega\eta)/\Gamma_{\text{total}}$ (units 10^{-3})

¹ Using $B(\eta \rightarrow 2\gamma) = (39.43 \pm 0.26)\%$, $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 22.6 \pm 0.4\%$, $B(\eta \rightarrow \pi^+ \pi^- \gamma) = 4.68 \pm 0.11\%$, and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$.

² Using $\Gamma(J/\psi \rightarrow e^+ e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

$\Gamma(\omega\pi^+\pi^-\pi^-\pi^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
85±34				
85	140	VANNUCCI	77	MRK1 $e^+ e^- \rightarrow 3(\pi^+ \pi^-)\pi^0$

Γ_{47}/Γ

$\Gamma(\omega\eta'\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{49}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.12 \pm 0.02 \pm 0.13$	14k	¹ ABLIKIM	19AC BES3	$J/\psi \rightarrow \omega\eta'\pi^+\pi^-$

¹ Using the decays $\omega \rightarrow \pi^+\pi^-\pi^0$ and $\eta' \rightarrow \eta\pi^+\pi^-$.

 $\Gamma(\omega\eta'(958))/\Gamma_{\text{total}}$ Γ_{50}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.89 ± 0.18 OUR AVERAGE				
$2.08 \pm 0.30 \pm 0.14$	137	¹ ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'$
2.26 ± 0.43	218	² ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\eta'$
$1.8 \begin{array}{l} +1.0 \\ -0.8 \end{array} \pm 0.3$	6	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$1.66 \pm 0.17 \pm 0.19$		COFFMAN	88 MRK3	$e^+e^- \rightarrow 3\pi\eta'$

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+\pi^-\eta'$.

² Using $B(\eta' \rightarrow \pi^+\pi^-\eta) = (44.3 \pm 1.5)\%$, $B(\eta' \rightarrow \pi^+\pi^-\gamma) = 29.5 \pm 1.0\%$, $B(\eta \rightarrow 2\gamma) = 39.43 \pm 0.26\%$, and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

 $\Gamma(\omega f_0(980))/\Gamma_{\text{total}}$ Γ_{51}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.41 \pm 0.27 \pm 0.47$		¹ AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

 $\Gamma(\omega f_0(1710) \rightarrow \omega K\bar{K})/\Gamma_{\text{total}}$ Γ_{52}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.8 \pm 1.1 \pm 0.3$		^{1,2} FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

¹ Includes unknown branching fraction $f_0(1710) \rightarrow K\bar{K}$.

² Addition of $f_0(1710) \rightarrow K^+K^-$ and $f_0(1710) \rightarrow K^0\bar{K}^0$ branching ratios.

 $\Gamma(\omega f_1(1420))/\Gamma_{\text{total}}$ Γ_{53}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$6.8 \begin{array}{l} +1.9 \\ -1.6 \end{array} \pm 1.7$	$111 \begin{array}{l} +31 \\ -26 \end{array}$	BECKER	87 MRK3	$e^+e^- \rightarrow \text{hadrons}$

 $\Gamma(\omega f'_2(1525))/\Gamma_{\text{total}}$ Γ_{54}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.2 \times 10^{-4}$	90	¹ VANNUCCI	77 MRK1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.8 \times 10^{-4}$	90	¹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
-----------------------	----	----------------------	--------	-------------------------------------

¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.

 $\Gamma(\omega X(1835) \rightarrow \omega p\bar{p})/\Gamma_{\text{total}}$ Γ_{55}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.9 \times 10^{-6}$	95	ABLIKIM	13P BES3	$J/\psi \rightarrow \gamma\pi^0p\bar{p}$

$\Gamma(\omega X(1835), X \rightarrow \eta' \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$<6.2 \times 10^{-5}$	¹ ABLIKIM	19AC BES3	$J/\psi \rightarrow \omega \eta' \pi^+ \pi^-$

¹ Using the decays $\omega \rightarrow \pi^+ \pi^- \pi^0$ and $\eta' \rightarrow \eta \pi^+ \pi^-$.

$\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.52 \pm 0.30 \pm 0.01$	276	¹ ANASHIN	22	KEDR $J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

¹ ANASHIN 22 reports $[\Gamma(J/\psi(1S) \rightarrow \omega K^+ K^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = (0.136 \pm 0.008 \pm 0.026) \times 10^{-2}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
34 ± 5 OUR AVERAGE				
$37.7 \pm 0.8 \pm 5.8$	1972 ± 41	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$
$29.5 \pm 1.4 \pm 7.0$	879 ± 41	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\omega K \bar{K})/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
19 ± 4 OUR AVERAGE				
$19.8 \pm 2.1 \pm 3.9$		¹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
16 ± 10	22	FELDMAN	77 MRK1	$e^+ e^-$

¹ Addition of $\omega K^+ K^-$ and $\omega K^0 \bar{K}^0$ branching ratios.

$\Gamma(\omega K^*(892) \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
61 ± 9 OUR AVERAGE				
$62.0 \pm 6.8 \pm 10.6$	899 ± 98	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$
$65.3 \pm 10.2 \pm 13.5$	176 ± 28	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
$53 \pm 14 \pm 14$	530 ± 140	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\eta' K^{*\pm} K^\mp)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.48 ± 0.13 OUR AVERAGE			
$1.50 \pm 0.02 \pm 0.19$	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$
$1.47 \pm 0.03 \pm 0.17$	² ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K^+ K^- \pi^0$.
² From $\eta' K_S^0 K^\pm \pi^\mp$.

$\Gamma(\eta' K^{*0} \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.66 \pm 0.03 \pm 0.21$	1 ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K_S^0 K^\pm \pi^\mp$.

$\Gamma(\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{63}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.16±0.12±0.29	1.1k	1 ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K_S^0 K^\pm \pi^\mp$.

 $\Gamma(\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp)/\Gamma_{\text{total}}$ Γ_{64}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.51±0.09±0.21	1.0k	1 ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K^+ K^- \pi^0$.

 $\Gamma(\eta' h_1(1415) \rightarrow \gamma \eta' \eta')/\Gamma_{\text{total}}$ Γ_{65}/Γ

<u>VALUE</u> (units 10^{-7})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.69±0.80^{+0.74}_{-1.82}	1 ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta' \eta'$, and ($\eta' X$), with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

 $\Gamma(h_1(1415)\eta' \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$ Γ_{26}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.08±0.01 ^{+0.01} _{-0.02}	1 ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave.

 $\Gamma(h_1(1595)\eta' \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$ Γ_{27}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.16±0.02 ^{+0.03} _{-0.01}	1 ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave.

 $\Gamma(\bar{K} K^*(892)+\text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$ Γ_{67}/Γ_{178}

<u>VALUE</u> (%)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
90.5±0.9±3.8	4k	1 LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$


¹ From a Dalitz plot analysis in an isobar model.

 $\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{68}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.0^{+0.8}_{-1.0} OUR AVERAGE				Error includes scale factor of 2.9. See the ideogram below.
8.07±0.04 ^{+0.38} _{-0.61}	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$
4.57±0.17±0.70	2285	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
5.26±0.13±0.53		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$, $K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ± 0.6	24	FRANKLIN	83	MRK2	$J/\psi \rightarrow K^+ K^- \pi^0$
3.2 ± 0.6	48	VANNUCCI	77	MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
4.1 ± 1.2	39	BRAUNSCH...	76	DASP	$J/\psi \rightarrow K^\pm X$



$$\Gamma(K^+ K^*(892)^- + c.c. \rightarrow K^+ K^- \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{69}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.69 \pm 0.01^{+0.13}_{-0.20}$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

$$\Gamma(K^+ K^*(892)^- + c.c. \rightarrow K^+ K^- \pi^0) / \Gamma(K^+ K^- \pi^0) \quad \Gamma_{69}/\Gamma_{177}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$92.4 \pm 1.5 \pm 3.4$	2k	¹ LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$$\Gamma(K^0 \bar{K}^*(892)^0 + c.c.) / \Gamma_{\text{total}} \quad \Gamma_{71}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.2 ± 0.4 OUR AVERAGE				
$3.96 \pm 0.15 \pm 0.60$	1192	JOUSSET	90 DM2	$J/\psi \rightarrow$ hadrons
$4.33 \pm 0.12 \pm 0.45$		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
2.7 ± 0.6	45	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
5.73±0.14±0.82	1 ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	2 ABLIKIM	06c BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
------	-----------	----------	---

¹ Obtained from $J/\psi \rightarrow K^*(892) K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-$ taking the value 2/3 for the probability of the $K^*(892)^0 \rightarrow K^+ \pi^-$ decay.

² A $K_0^*(700)$ is observed by ABLIKIM 06c in the $K^+ \pi^-$ mass spectrum of the $\bar{K}^*(892)^0 K^+ \pi^-$ final state against the $\bar{K}^*(892)$. A corresponding branching fraction of the $J/\psi(1S)$ is not presented.

$\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.81±0.10±0.54	1559	ANASHIN	22	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{78}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
6.28±0.16±0.59	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(K^*(892)^\pm K^*(700)^\mp)/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.09±0.18±0.94	655	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(K^*(892)^\pm K^*(892)^\mp)/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.00±0.19±0.11	323	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
3.8±0.8±1.2	1 BAI	99c BES	$e^+ e^-$

¹ Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

$\Gamma(K^*(1410) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+ K^- \pi^0)$ Γ_{85}/Γ_{177}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.3±1.1±0.7	2k	¹ LEES	17c BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K^*(1410)\bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$	Γ_{86}/Γ_{178}			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.5 \pm 0.9$	4k	1 LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+ K^- \pi^0)$	Γ_{88}/Γ_{177}			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$3.5 \pm 1.3 \pm 0.9$	2k	1 LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$	Γ_{89}/Γ_{178}			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$7.1 \pm 1.3 \pm 1.2$	4k	1 LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(\bar{K}_2^*(1430)K + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{90}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<40 \times 10^{-4}$	90	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow K^0 \bar{K}_2^{*0}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<66 \times 10^{-4}$	90	BRAUNSCH... 76	DASP	$e^+ e^- \rightarrow K^\pm \bar{K}_2^{*\mp}$

$\Gamma(K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$	Γ_{91}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.69 \pm 0.04 \pm 0.25$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

$\Gamma(\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{94}/Γ			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
6.7 \pm 2.6	40	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

$\Gamma(K_2^*(1430)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$	Γ_{97}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<29 \times 10^{-4}$	90	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

$\Gamma(K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$	Γ_{99}/Γ			
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.1 \pm 0.1 \pm 0.6$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

$\Gamma(K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$	Γ_{100}/Γ			
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.2 \pm 0.7 \pm 2.8$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

$\Gamma(K_1(1270)^{\pm} K^{\mp})/\Gamma_{\text{total}}$				Γ_{101}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<3.0 \times 10^{-3}$	90	1 BAI	99c BES	$e^+ e^-$	

¹ Assuming $B(K_1(1270) \rightarrow K\rho) = 0.42 \pm 0.06$

$\Gamma(K_1(1270)K_S^0 \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$				Γ_{102}/Γ	
<u>VALUE (units 10^{-7})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$8.54^{+1.07+2.35}_{-1.20-2.13}$		ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$	

$\Gamma(a_2(1320)^{\pm} \pi^{\mp})/\Gamma_{\text{total}}$				Γ_{103}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<43 \times 10^{-4}$	90	BRAUNSCH...	76 DASP	$e^+ e^-$	

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$				Γ_{104}/Γ	
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.94 $\pm 0.16 \pm 0.16$	0.8k	1 ABLIKIM	15K BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma\gamma$	
0.124 $\pm 0.033 \pm 0.030$	35 ± 9	2 ABLIKIM	15K BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.4	90	3 ABLIKIM	05B BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \phi\gamma\gamma$
<6.8	90	COFFMAN	88 MRK3	$e^+ e^- \rightarrow K^+ K^- \pi^0$

¹ Corresponding to one of the two fit solutions with $\delta = (-95.9 \pm 1.5)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi\pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

² Corresponding to one of the two fit solutions with $\delta = (-152.1 \pm 7.7)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi\pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

³ Superseded by ABLIKIM 15K.

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$				Γ_{105}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.94 ± 0.15 OUR AVERAGE	Error includes scale factor of 1.7.				
1.09 $\pm 0.02 \pm 0.13$	ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$		
0.78 $\pm 0.03 \pm 0.12$	FALVAR	88 DM2	$J/\psi \rightarrow \text{hadrons}$		
2.1 ± 0.9	23 FELDMAN	77 MRK1	$e^+ e^-$		

$\Gamma(\phi 2(\pi^+\pi^-))/\Gamma_{\text{total}}$				Γ_{107}/Γ	
<u>VALUE (units 10^{-4})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
16.0 $\pm 1.0 \pm 3.0$		FALVAR	88 DM2	$J/\psi \rightarrow \text{hadrons}$	

$\Gamma(\phi\eta)/\Gamma_{\text{total}}$				Γ_{108}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.74 ± 0.06 OUR AVERAGE	Error includes scale factor of 1.2.				
0.71 $\pm 0.10 \pm 0.05$	99 ± 14	1 ZHU	23 BELL	$e^+ e^- \rightarrow \gamma(nS) \rightarrow \phi\eta\gamma$	
0.898 $\pm 0.024 \pm 0.089$		ABLIKIM	05B BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadr}$	


$0.64 \pm 0.04 \pm 0.11$	346	JOUSSET	90	DM2	$J/\psi \rightarrow$ hadrons
$0.661 \pm 0.045 \pm 0.078$		COFFMAN	88	MRK3	$e^+ e^- \rightarrow K^+ K^- \eta$

¹ From a fit to the combined $\phi\eta$ invariant mass spectrum with a Gaussian function for the J/ψ signals and a second-order polynomial function for the backgrounds.

$\Gamma(\phi\eta'(958))/\Gamma_{\text{total}}$

Γ_{109}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
4.6 ± 0.5 OUR AVERAGE					Error includes scale factor of 2.2. See the ideogram below.
$5.10 \pm 0.03 \pm 0.32$	31k	ABLIKIM	16M	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow$ hadrons
$5.46 \pm 0.31 \pm 0.56$		ABLIKIM	05B	BES2	$e^+ e^- \rightarrow J/\psi \rightarrow$ hadrons
$4.1 \pm 0.3 \pm 0.8$	167	JOUSSET	90	DM2	$J/\psi \rightarrow$ hadrons
$3.08 \pm 0.34 \pm 0.36$		COFFMAN	88	MRK3	$e^+ e^- \rightarrow K^+ K^- \eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 13	90	VANNUCCI	77	MRK1	$e^+ e^-$



$\Gamma(\phi\eta\eta')/\Gamma_{\text{total}}$

Γ_{110}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.32±0.06±0.16	2.2k	¹ ABLIKIM	19AN BES3	$e^+ e^- \rightarrow J/\psi \rightarrow$ hadrons

¹ Including contributions from intermediate resonances. Evidence for an intermediate resonance at $M \approx 2$ GeV and $\Gamma \approx 150$ MeV decaying to $\phi\eta'$ with $J^P = 1^+$ or $J^P = 1^-$, and $B(J/\psi \rightarrow \eta X) \times B(X \rightarrow \phi\eta') \approx 10^{-4}$.

$\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$ Γ_{111}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.2±0.9 OUR AVERAGE	Error includes scale factor of 1.9.			
4.6±0.4±0.8		¹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
2.6±0.6	50	¹ GIDAL	81 MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$. $\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{114}/Γ

<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.50±0.80±0.61	355	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

 $\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0 p^0\pi^0)/\Gamma_{\text{total}}$ Γ_{115}/Γ

<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.67±0.50±0.24	70	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

 $\Gamma(\phi f_0(980)\eta \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{116}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.23±0.75±0.73	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$

 $\Gamma(\phi a_0(980)^0 \rightarrow \phi\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{117}/Γ

<u>VALUE</u> (units 10^{-6})		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.37±1.35		¹ ABLIKIM	18D BES3	$J/\psi \rightarrow \phi\eta\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.0 ± 2.7 ± 2.5 ² ABLIKIM 11D BES3 $J/\psi \rightarrow \phi\eta\pi^0$ ¹ Assuming constructive interference between $a_0(980) - f_0(980)$ mixing and electromagnetic decay. Destructive interference gives a value of $(4.93 \pm 1.77) \times 10^{-6}$ for this branching fraction.² Assuming $a_0(980) - f_0(980)$ mixing and isospin breaking via γ^* and $K^* K$ loops. $\Gamma(\phi f_2(1270))/\Gamma_{\text{total}}$ Γ_{118}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.45	90	FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
< 0.37	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

 $\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$ Γ_{119}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.6±0.5 OUR AVERAGE				

3.4±1.8±1.5	1.1k	¹ ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$
3.2±0.6±0.4		JOUSSET	90 DM2	$J/\psi \rightarrow \phi 2(\pi^+\pi^-)$
2.1±0.5±0.4	25	² JOUSSET	90 DM2	$J/\psi \rightarrow \phi\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6±0.2±0.1 16 BECKER 87 MRK3 $J/\psi \rightarrow \phi K\bar{K}\pi$ ¹ ABLIKIM 15H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta\pi^+\pi^-)] = (1.20 \pm 0.6 \pm 0.14) \times 10^{-4}$ which we divide by our best value $B(f_1(1285) \rightarrow \eta\pi^+\pi^-) = (35 \pm 15) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.² We attribute to the $f_1(1285)$ the signal observed in the $\pi^+\pi^-\eta$ invariant mass distribution at 1297 MeV.

$\Gamma(\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{120}/Γ

<u>VALUE</u> (units 10^{-7})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$9.36 \pm 2.31 \pm 1.54$	78	ABLIKIM	15P	$J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow \phi 3\pi^0)/\Gamma_{\text{total}}$ Γ_{121}/Γ

<u>VALUE</u> (units 10^{-7})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.08 \pm 1.63 \pm 1.47$	9	ABLIKIM	15P	$J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{122}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.01 \pm 0.58 \pm 0.82$	172	1	ABLIKIM	15H	$e^+ e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 17 90 ² FALVARD 88 DM2 $J/\psi \rightarrow$ hadrons

¹ With 3.6 σ significance.

² Includes unknown branching fraction $\eta(1405) \rightarrow \eta\pi\pi$.

$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$ Γ_{123}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8 ± 4 OUR AVERAGE				Error includes scale factor of 2.7.
$12.3 \pm 0.6 \pm 2.0$	1, ² FALVARD 88 DM2 $J/\psi \rightarrow$ hadrons			
4.8 ± 1.8	46 ¹ GIDAL 81 MRK2 $J/\psi \rightarrow K^+ K^- K^+ K^-$			

¹ Re-evaluated using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.

² Including interference with $f_0(1710)$.

$\Gamma(\phi X(1835) \rightarrow \phi p\bar{p})/\Gamma_{\text{total}}$ Γ_{124}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.1 \times 10^{-7}$	90	1 ABLIKIM	16K	$J/\psi \rightarrow p\bar{p}K_S^0 K_L^0, p\bar{p}K^+ K^-$

¹ Upper limit applies to any $p\bar{p}$ mass enhancement near threshold.

$\Gamma(\phi X(1835) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{125}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.8 \times 10^{-4}$	90	ABLIKIM	15H	$e^+ e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$


$\Gamma(\phi X(1870) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{126}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.13 \times 10^{-5}$	90	ABLIKIM	15H	$e^+ e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

$\Gamma(\phi K\bar{K})/\Gamma_{\text{total}}$ Γ_{127}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
17.7 ± 1.6 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
$16.6 \pm 1.9 \pm 1.2$	163 ± 19	LEES	12F BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-)\gamma$
$21.4 \pm 0.4 \pm 2.2$		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
$48 \pm 20 \pm 6$	9.0 ± 3.7	1, ² HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$

14.6 \pm 0.8 \pm 2.1	³	FALVARD FELDMAN	88 77	DM2 MRK1	J/ ψ \rightarrow hadrons $e^+ e^-$
18 \pm 8	14				



¹ We have multiplied $K^+ K^-$ measurement by 2 to obtain $K\bar{K}$.

² Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

³ Addition of $\phi K^+ K^-$ and $\phi K^0 \bar{K}^0$ branching ratios.

$\Gamma(\phi f_0(1710) \rightarrow \phi K\bar{K})/\Gamma_{total}$

Γ_{128}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.6 \pm 0.2 \pm 0.6	1,2 FALVARD	88	DM2

¹ Including interference with $f'_2(1525)$.

² Includes unknown branching fraction $f_0(1710) \rightarrow K\bar{K}$.

$\Gamma(\phi K^\pm K_S^0 \pi^\mp)/\Gamma_{total}$

Γ_{131}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.2 \pm 0.8 OUR AVERAGE				
7.4 \pm 0.6 \pm 1.4	227 \pm 19	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$
7.4 \pm 0.9 \pm 1.1		FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
7 \pm 0.6 \pm 1.0	163 \pm 15	BECKER	87 MRK3	$e^+ e^- \rightarrow$ hadrons

$\Gamma(\phi K^*(892)\bar{K} + c.c.)/\Gamma_{total}$

Γ_{132}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
21.8 \pm 2.3 OUR AVERAGE				
20.8 \pm 2.7 \pm 3.9	195 \pm 25	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K_S^0 K^\pm \pi^\mp$
29.6 \pm 3.7 \pm 4.7	238 \pm 30	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K^+ K^- \pi^0$

$20.7 \pm 2.4 \pm 3.0$
 $20 \pm 3 \pm 3$ 155 ± 20

FALVARD 88 DM2 $J/\psi \rightarrow$ hadrons
BECKER 87 MRK3 $e^+ e^- \rightarrow$ hadrons

$\Gamma(b_1(1235)^{\pm} \pi^{\mp})/\Gamma_{\text{total}}$

VALUE (units 10^{-4}) EVTS
 30 ± 5 OUR AVERAGE

31 ± 6 4600
 29 ± 7 87

DOCUMENT ID	TECN	COMMENT
AUGUSTIN 89	DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
BURMESTER 77D	PLUT	$e^+ e^-$

Γ_{133}/Γ

$\Gamma(b_1(1235)^0 \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4}) EVTS
 $23 \pm 3 \pm 5$ 229

DOCUMENT ID	TECN	COMMENT
AUGUSTIN 89	DM2	$e^+ e^-$

Γ_{134}/Γ

$\Gamma(\Delta(1232)^+ \bar{p})/\Gamma_{\text{total}}$

VALUE CL%
 $<0.1 \times 10^{-3}$ 90

DOCUMENT ID	TECN	COMMENT
HENRARD 87	DM2	$e^+ e^-$

Γ_{136}/Γ

$\Gamma(\Delta(1232)^{++} \bar{p} \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3}) EVTS
 $1.58 \pm 0.23 \pm 0.40$ 332

DOCUMENT ID	TECN	COMMENT
EATON 84	MRK2	$e^+ e^-$

Γ_{137}/Γ

$\Gamma(\Delta(1232)^{++} \bar{\Delta}(1232)^{--})/\Gamma_{\text{total}}$

VALUE (units 10^{-3}) EVTS
 $1.10 \pm 0.09 \pm 0.28$ 233

DOCUMENT ID	TECN	COMMENT
EATON 84	MRK2	$e^+ e^-$

Γ_{138}/Γ

$\Gamma(\bar{\Sigma}(1385)^0 p K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3}) EVTS
 $0.51 \pm 0.26 \pm 0.18$ 89

DOCUMENT ID	TECN	COMMENT
EATON 84	MRK2	$e^+ e^-$

Γ_{139}/Γ

$\Gamma(\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE CL%
 $<0.82 \times 10^{-5}$ 90

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.2 \times 10^{-3}$ 90

DOCUMENT ID	TECN	COMMENT
ABLIKIM 13F	BES3	$J/\psi \rightarrow p \bar{p} \pi^+ \pi^- \gamma \gamma$

Γ_{140}/Γ

$\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ (\text{or c.c.}))/\Gamma_{\text{total}}$

VALUE (units 10^{-3}) EVTS
 0.31 ± 0.05 OUR AVERAGE

$0.30 \pm 0.03 \pm 0.07$ 74 ± 8
 $0.34 \pm 0.04 \pm 0.07$ 77 ± 9
 $0.29 \pm 0.11 \pm 0.10$ 26
 $0.31 \pm 0.11 \pm 0.11$ 28

DOCUMENT ID	TECN	COMMENT
HENRARD 87	DM2	$e^+ e^- \rightarrow \Sigma^{*-}$
HENRARD 87	DM2	$e^+ e^- \rightarrow \Sigma^+$
EATON 84	MRK2	$e^+ e^- \rightarrow \Sigma^{*-}$
EATON 84	MRK2	$e^+ e^- \rightarrow \Sigma^+$

Γ_{141}/Γ

$\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ (\text{or c.c.}))/\Gamma_{\text{total}}$

VALUE (units 10^{-3}) EVTS
 1.16 ± 0.05 OUR AVERAGE

$1.096 \pm 0.012 \pm 0.071$ 43k
 $1.258 \pm 0.014 \pm 0.078$ 53k
 $1.23 \pm 0.07 \pm 0.30$ 0.8k

DOCUMENT ID	TECN	COMMENT
ABLIKIM 16L	BES3	$J/\psi \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$
ABLIKIM 16L	BES3	$J/\psi \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
ABLIKIM 12P	BES2	$J/\psi \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$

Γ_{142}/Γ

1.50 ± 0.08	± 0.38	1k	ABLIKIM	12P	BES2	$J/\psi \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
1.00 ± 0.04	± 0.21	0.6k	HENRARD	87	DM2	$e^+ e^- \rightarrow \Sigma^{*-}$
1.19 ± 0.04	± 0.25	0.7k	HENRARD	87	DM2	$e^+ e^- \rightarrow \Sigma^{*+}$
0.86 ± 0.18	± 0.22	56	EATON	84	MRK2	$e^+ e^- \rightarrow \Sigma^{*-}$
1.03 ± 0.24	± 0.25	68	EATON	84	MRK2	$e^+ e^- \rightarrow \Sigma^{*+}$

$\Gamma(\Sigma(1385)^0 \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$

Γ_{143}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.071 \pm 0.009 \pm 0.082$	103k	ABLIKIM	17E	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadrons}$

$\Gamma(\Lambda(1520)\bar{\Lambda} + \text{c.c.} \rightarrow \gamma\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$

Γ_{144}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.1 \times 10^{-6}$	90	ABLIKIM	12B	$J/\psi \rightarrow \Lambda\bar{\Lambda}\gamma$

$\Gamma(\bar{\Lambda}(1520)\Lambda + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{145}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.80 \times 10^{-3}$	90	LU	19	$B^+ \rightarrow \bar{p}\Lambda K^+ K^+$

$\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$

Γ_{146}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.17 ± 0.04 OUR AVERAGE				
$1.165 \pm 0.004 \pm 0.043$	135k	ABLIKIM	17E	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
$1.20 \pm 0.12 \pm 0.21$	206	ABLIKIM	080	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\Xi(1530)^- \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{147}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.318 ± 0.008 OUR AVERAGE				
$0.317 \pm 0.002 \pm 0.008$	70k	ABLIKIM	20	$e^+ e^- \rightarrow J/\psi$
$0.59 \pm 0.09 \pm 0.12$	75	HENRARD	87	$DM2$

$\Gamma(\Xi(1530)^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$

Γ_{148}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.32 \pm 0.12 \pm 0.07$	24 ± 9	HENRARD	87	$DM2$

$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{149}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-5}$	90	BAI	04G	$BES2$

$\Gamma(\Theta(1540)K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$

Γ_{150}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.1 \times 10^{-5}$	90	BAI	04G	$BES2$

$\Gamma(\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$

Γ_{151}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.6 \times 10^{-5}$	90	BAI	04G	$BES2$

$\Gamma(\overline{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{152}/Γ
$<5.6 \times 10^{-5}$	90	BAI	04G	BES2	$e^+ e^-$

$\Gamma(\overline{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{153}/Γ
$<1.1 \times 10^{-5}$	90	BAI	04G	BES2	$e^+ e^-$


— STABLE HADRONS —

$\Gamma(2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$

Γ_{154}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{154}/Γ
4.2 ± 0.4 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.	
5.44 ± 0.07 ± 0.33	23K	ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$	
3.5 ± 0.8 ± 0.1	14k	¹ LEES	21	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-)3\pi^0\gamma$	
4.73 ± 0.44	228k	ABLIKIM	19Q	BES3 $J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$	
3.25 ± 0.49	46055	AUGUSTIN	89	DM2 $J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$	
3.17 ± 0.42	147	FRANKLIN	83	MRK2 $e^+ e^- \rightarrow \text{hadrons}$	
3.64 ± 0.52	1500	BURMESTER	77D	PLUT $e^+ e^-$	
4 ± 1	675	JEAN-MARIE	76	MRK1 $e^+ e^-$	

WEIGHTED AVERAGE
4.2 ± 0.4 (Error scaled by 2.1)



$\Gamma(2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}} (\text{units } 10^{-2})$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0)] = (14.8 \pm 2.6 \pm 2.2) \times 10^{-3} \text{ keV}$ which we divide by our best values $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04 \text{ keV}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0) =$

$(18.24 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² From an energy scan of $e^+ e^- \rightarrow J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$, assuming PDG 16 values for $\Gamma(e^+ e^-)$, $\Gamma(\mu^+ \mu^-)$, and $\Gamma(\text{total})$, and for a phase difference between strong and electromagnetic amplitudes of $(84.9 \pm 3.6)^\circ$. An alternative solution is $(4.85 \pm 0.45)\%$ with a phase of $(-84.7 \pm 3.1)^\circ$.

$\Gamma(3(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$


VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{155}/Γ
0.029±0.006 OUR AVERAGE					
0.028±0.009	11	FRANKLIN	83	MRK2 $e^+ e^- \rightarrow \text{hadrons}$	
0.029±0.007	181	JEAN-MARIE	76	MRK1 $e^+ e^-$	

$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{160}/Γ
21.0 ±0.8 OUR AVERAGE					
21.37±0.04 ^{+0.64} _{-0.62}	1.8M	1,2 ABLIKIM	12H BES3	$e^+ e^- \rightarrow J/\psi$	
23.0 ±2.0 ±0.4	256	³ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow J/\psi \pi^+ \pi^- \gamma$	
21.84±0.05±2.01	220k	1,4 BAI	04H BES	$e^+ e^-$	
20.91±0.21±1.16		4,5 BAI	04H BES	$e^+ e^-$	
15 ±2	168	FRANKLIN	83	MRK2 $e^+ e^-$	

WEIGHTED AVERAGE
21.0±0.8 (Error scaled by 1.6)

Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.



¹ From $J/\psi \rightarrow \pi^+ \pi^- \pi^0$ events directly.

² The quoted systematic error includes a contribution of 1.23% (added in quadrature) from the uncertainty on the number of J/ψ events.

³ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] = (18.6 \pm 1.2 \pm 1.1) \times 10^{-3} \text{ keV}$ which we divide by our best value $\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}} = 0.808 \pm 0.013 \text{ keV}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Mostly $\rho\pi$, see also $\rho\pi$ subsection.

⁵ Obtained comparing the rates for $\pi^+\pi^-\pi^0$ and $\mu^+\mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$.

$\Gamma(\pi^+\pi^-\pi^0 K^+ K^-)/\Gamma_{\text{total}}$

Γ_{162}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.52 ± 0.27 OUR AVERAGE	Error includes scale factor of 1.4.			
$1.74 \pm 0.08 \pm 0.24$	2616	ANASHIN	22	KEDR $J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1.2 ± 0.3	309	VANNUCCI	77	MRK1 $e^+ e^-$

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{163}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.47 ± 0.14 OUR AVERAGE				
$1.47 \pm 0.13 \pm 0.13$	140	¹ METREVELI	12	$\psi(2S) \rightarrow 2(\pi^+\pi^-)$
$1.58 \pm 0.20 \pm 0.15$	84	BALTRUSAIT..85D	MRK3	$e^+ e^-$
1.0 ± 0.5	5	BRANDELIK	78B	DASP $e^+ e^-$
1.6 ± 1.6	1	VANNUCCI	77	MRK1 $e^+ e^-$

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$

Γ_{164}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.20 ± 0.25 OUR AVERAGE	Error includes scale factor of 1.2.			
$2.88 \pm 0.14 \pm 0.24$	2654	ANASHIN	22	$J/\psi \rightarrow 2(\pi^+\pi^-)$
$3.53 \pm 0.12 \pm 0.29$	1107	¹ ABLIKIM	05H	BES2 $e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \pi^+ \pi^-$, $J/\psi \rightarrow 2(\pi^+\pi^-)$
4.0 ± 1.0	76	JEAN-MARIE	76	MRK1 $e^+ e^-$

¹ Computed using $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$

Γ_{165}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
40 ± 20	32	JEAN-MARIE	76	MRK1 $e^+ e^-$

$\Gamma(4(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

Γ_{167}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
90 ± 30	13	JEAN-MARIE	76	MRK1 $e^+ e^-$

$\Gamma(2(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}$

Γ_{168}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.29 ± 0.28 OUR AVERAGE				
$3.1 \pm 1.5 \pm 0.1$	14k	¹ LEES	21	BABR $e^+ e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$
$2.26 \pm 0.08 \pm 0.27$	4.8k	ABLIKIM	05C	BES2 $e^+ e^- \rightarrow 2(\pi^+\pi^-)\eta$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] \times [B(\eta \rightarrow 3\pi^0)] = (5.6 \pm 2.6 \pm 0.8) \times 10^{-3}$ keV which we divide by our best values

$\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.53 \pm 0.10$ keV, $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.21) \times 10^{-2}$.

Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(3(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.24 \pm 0.96 \pm 1.11$	616	ABLIKIM	05C BES2	$e^+ e^- \rightarrow 3(\pi^+\pi^-)\eta$

Γ_{169}/Γ

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.86 \pm 0.09 \pm 0.19$	1k	¹ METREVELI	12	$\psi(2S) \rightarrow \pi^+\pi^- K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$2.39 \pm 0.24 \pm 0.22$	107	² BALTRUSAIT..85D	MRK3	e^+e^-
2.2 ± 0.9	6	² BRANDELIK	79c DASP	e^+e^-

Γ_{173}/Γ

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Interference with non-resonant K^+K^- production not taken into account.


$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$

Γ_{174}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.95 ± 0.11 OUR AVERAGE				Error includes scale factor of 2.4. See the ideogram below.
1.93 $\pm 0.01 \pm 0.05$	110k	ABLIKIM	17AH BES3	$J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^- X$
2.62 $\pm 0.15 \pm 0.14$	0.3k	¹ METREVELI	12	$\psi(2S) \rightarrow \pi^+\pi^- K_S^0 K_L^0$
1.82 $\pm 0.04 \pm 0.13$	2.1k	² BAI	04A BES2	$J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.18 $\pm 0.12 \pm 0.18$		JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
1.01 $\pm 0.16 \pm 0.09$	74	BALTRUSAIT..85D	MRK3	e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6868 \pm 0.0027$.



$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$

Γ_{174}/Γ

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$

Γ_{175}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-8}$	95	¹ ABLIKIM	17AH BES3	$J/\psi \rightarrow K_S^0 K_S^0 \rightarrow \pi^+\pi^-\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1 \times 10^{-6}$	95	¹ BAI	04D	BES	$e^+ e^-$
$<5.2 \times 10^{-6}$	90	¹ BALTRUSAIT..85C	..85C	MRK3	$e^+ e^-$

¹ Forbidden by CP.

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS
61 ± 10 OUR AVERAGE	
55.2 ± 12.0	25
78.0 ± 21.0	126

Γ_{176}/Γ

DOCUMENT ID	TECN	COMMENT
FRANKLIN	83	MRK2 $e^+ e^- \rightarrow K^+ K^- \pi^0$
VANNUCCI	77	MRK1 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS
2.88 ± 0.01 ± 0.12	183k

Γ_{177}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

$\Gamma(K^+ K^- \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$

VALUE (%)	EVTS
12.0 ± 0.3 ± 0.9	23k

$\Gamma_{177}/\Gamma_{160}$

DOCUMENT ID	TECN	COMMENT
LEES	17C BABR	$J/\psi \rightarrow h^+ h^- \pi^0$

$\Gamma(K_S^0 K^\pm \pi^\mp)/\Gamma(\pi^+ \pi^- \pi^0)$

VALUE (%)	EVTS
26.5 ± 0.5 ± 2.1	24k

$\Gamma_{178}/\Gamma_{160}$

DOCUMENT ID	TECN	COMMENT
LEES	17C BABR	$J/\psi \rightarrow h^0 h^+ h^-$

$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS
7.04 ± 0.26 ± 0.92	2671

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.2 ± 2.3	205
-----------	-----

Γ_{182}/Γ

DOCUMENT ID	TECN	COMMENT
ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$
VANNUCCI	77 MRK1	$e^+ e^-$

$\Gamma(K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS
31 ± 13	30

Γ_{189}/Γ

DOCUMENT ID	TECN	COMMENT
VANNUCCI	77 MRK1	$e^+ e^-$

$\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS
• • • We do not use the following data for averages, fits, limits, etc. • • •	

$1.4^{+0.5}_{-0.4} \pm 0.2$	$11.0^{+4.3}_{-3.5}$	¹ HUANG	03	BELL	$B^+ \rightarrow 2(K^+ K^-) K^+$
-----------------------------	----------------------	--------------------	----	------	----------------------------------

0.7 ± 0.3		VANNUCCI	77	MRK1	$e^+ e^-$
-----------	--	----------	----	------	-----------

¹ Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS
2.120 ± 0.029 OUR AVERAGE	

Γ_{193}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	12C BES3	$e^+ e^-$
¹ WU	06 BELL	$B^+ \rightarrow p\bar{p} K^+$
BAI	04E BES2	$e^+ e^- \rightarrow J/\psi$
BALDINI	98 FENI	$e^+ e^-$

1.91 ± 0.04 ± 0.30	PALLIN	87	DM2	$e^+ e^-$
2.16 ± 0.07 ± 0.15	EATON	84	MRK2	$e^+ e^-$
2.5 ± 0.4	BRANDELIK	79c	DASP	$e^+ e^-$
2.0 ± 0.5	BESCH	78	BONA	$e^+ e^-$
2.2 ± 0.2	² PERUZZI	78	MRK1	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.3	48	ANTONELLI	93	SPEC	$e^+ e^-$
---------------	----	-----------	----	------	-----------

¹ WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] = (2.21 \pm 0.13 \pm 0.10) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assuming angular distribution $(1+\cos^2\theta)$.

$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$

Γ_{194}/Γ


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.19 ± 0.08 OUR AVERAGE	Error includes scale factor of 1.1.			
1.33 $\pm 0.02 \pm 0.11$	11k	ABLIKIM	09B	BES2 $e^+ e^-$
1.13 $\pm 0.09 \pm 0.09$	685	EATON	84	MRK2 $e^+ e^-$
1.4 ± 0.4		BRANDELIK	79c	DASP $e^+ e^-$
1.00 ± 0.15	109	PERUZZI	78	MRK1 $e^+ e^-$

$\Gamma(p\bar{p}\pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{195}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ± 0.5 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
6.46 $\pm 0.17 \pm 0.43$	1435	EATON	84	MRK2 $e^+ e^-$
3.8 ± 1.6	48	BESCH	81	BONA $e^+ e^-$
5.5 ± 0.6	533	PERUZZI	78	MRK1 $e^+ e^-$

WEIGHTED AVERAGE
6.0 ± 0.5 (Error scaled by 1.3)



$\Gamma(p\bar{p}\pi^+ \pi^-)/\Gamma_{\text{total}}$ (units 10^{-3})

$\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{196}/Γ

Including $p\bar{p}\pi^+\pi^-\gamma$ and excluding ω, η, η'

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.9 OUR AVERAGE	Error includes scale factor of 1.9.			
$3.36 \pm 0.65 \pm 0.28$	364	EATON	84	MRK2 e^+e^-
1.6 ± 0.6	39	PERUZZI	78	MRK1 e^+e^-

$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$

Γ_{197}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.00 ± 0.12 OUR AVERAGE				

$1.91 \pm 0.02 \pm 0.17$	13k	¹ ABLIKIM	09	BES2 e^+e^-
$2.03 \pm 0.13 \pm 0.15$	826	EATON	84	MRK2 e^+e^-
2.5 ± 1.2		BRANDELIK	79C	DASP e^+e^-
2.3 ± 0.4	197	PERUZZI	78	MRK1 e^+e^-

¹ From the combination of $p\bar{p}\eta \rightarrow p\bar{p}\gamma\gamma$ and $p\bar{p}\eta \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ channels.

$\Gamma(p\bar{p}\rho)/\Gamma_{\text{total}}$

Γ_{198}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.31 \times 10^{-3}$	90	EATON	84	MRK2 $e^+e^- \rightarrow \text{hadrons}\gamma$


$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$

Γ_{199}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.98 ± 0.10 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			

$0.90 \pm 0.02 \pm 0.09$	2670	ABLIKIM	13P	BES3 e^+e^-
$0.98 \pm 0.03 \pm 0.14$	2449	ABLIKIM	08	BES2 e^+e^-
$1.10 \pm 0.17 \pm 0.18$	486	EATON	84	MRK2 e^+e^-
1.6 ± 0.3	77	PERUZZI	78	MRK1 e^+e^-

WEIGHTED AVERAGE
 0.98 ± 0.10 (Error scaled by 1.3)



$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ (units 10^{-3})

$\Gamma(p\bar{p}\eta'(958))/\Gamma_{\text{total}}$ Γ_{200}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.129±0.014 OUR AVERAGE		Error includes scale factor of 2.0.		
0.126±0.002±0.007	16k	¹ ABLIKIM	19N	BES3 $e^+ e^-$
0.200±0.023±0.028	265 ± 31	² ABLIKIM	09	BES2 $e^+ e^-$
0.68 ± 0.23 ± 0.17	19	EATON	84	MRK2 $e^+ e^-$
1.8 ± 0.6	19	PERUZZI	78	MRK1 $e^+ e^-$

¹ From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\gamma$ channels.

² From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\gamma\rho^0$ channels.

 $\Gamma(p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta)/\Gamma_{\text{total}}$ Γ_{201}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.8±1.2±1.3	ABLIKIM	14N	$e^+ e^- \rightarrow J/\psi$

 $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{202}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.519±0.033 OUR AVERAGE				
0.523±0.006±0.033	14k	ABLIKIM	16K	BES3 $J/\psi \rightarrow p\bar{p}K_S^0 K_L^0$, $p\bar{p}K^+ K^-$
0.45 ± 0.13 ± 0.07		FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$

 $\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{203}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.12±0.09 OUR AVERAGE				
2.36±0.02±0.21	59k	ABLIKIM	06K	$J/\psi \rightarrow p\pi^-\bar{n}$
2.47±0.02±0.24	55k	ABLIKIM	06K	$J/\psi \rightarrow \bar{p}\pi^+n$
2.02±0.07±0.16	1288	EATON	84	MRK2 $e^+ e^- \rightarrow p\pi^-$
1.93±0.07±0.16	1191	EATON	84	MRK2 $e^+ e^- \rightarrow \bar{p}\pi^+$
1.7 ± 0.7	32	BESCH	81	BONA $e^+ e^- \rightarrow p\pi^-$
1.6 ± 1.2	5	BESCH	81	BONA $e^+ e^- \rightarrow \bar{p}\pi^+$
2.16±0.29	194	PERUZZI	78	MRK1 $e^+ e^- \rightarrow p\pi^-$
2.04±0.27	204	PERUZZI	78	MRK1 $e^+ e^- \rightarrow \bar{p}\pi^+$

 $\Gamma(n\bar{n})/\Gamma_{\text{total}}$ Γ_{204}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.09±0.16 OUR AVERAGE				
2.07±0.01±0.17	36k	ABLIKIM	12C	$e^+ e^-$
2.31±0.49	79	BALDINI	98	FENI $e^+ e^-$
1.8 ± 0.9		BESCH	78	BONA $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.90±0.55	40	ANTONELLI	93	SPEC $e^+ e^-$

 $\Gamma(n\bar{n}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{205}/Γ


<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.8±3.6	5	BESCH	81	BONA $e^+ e^-$

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.89 ± 0.09 OUR AVERAGE				Error includes scale factor of 2.8. See the ideogram below.
1.943 ± 0.003 ± 0.033	441k	ABLIKIM	17L	BES3 $e^+ e^-$
2.03 ± 0.03 ± 0.15	8887	ABLIKIM	06	BES2 $J/\psi \rightarrow \Lambda\bar{\Lambda}$
1.96 $\begin{array}{l} +0.47 \\ -0.44 \end{array}$ ± 0.04	46	1 WU	06	BELL $B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
1.08 ± 0.06 ± 0.24	631	BAI	98G	BES $e^+ e^-$
1.38 ± 0.05 ± 0.20	1847	PALLIN	87	DM2 $e^+ e^-$
1.58 ± 0.08 ± 0.19	365	EATON	84	MRK2 $e^+ e^-$
2.6 ± 1.6	5	BESCH	81	BONA $e^+ e^-$
1.1 ± 0.2	196	PERUZZI	78	MRK1 $e^+ e^-$

¹WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.00^{+0.34}_{-0.29} \pm 0.34) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

WEIGHTED AVERAGE
1.89±0.09 (Error scaled by 2.8)



$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.78±0.27±0.30		323	¹ ABLIKIM	13F	BES3 $J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.4	90	² ABLIKIM	07H	BES2	$e^+ e^- \rightarrow \psi(2S)$
23 ± 7	± 8	11	BAI	98G	BES $e^+ e^-$
22 ± 5	± 5	19	HENRARD	87	DM2 $e^+ e^-$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

$\Gamma(\Lambda \bar{\Lambda} \pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{211}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.30 $\pm 0.13 \pm 0.99$	2.4k	ABLIKIM	12P	BES2 J/ψ

$\Gamma(\Lambda \bar{\Lambda} \eta)/\Gamma_{\text{total}}$

Γ_{212}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
16.2 ± 1.7 OUR AVERAGE				
15.7 $\pm 0.80 \pm 1.54$	454	¹ ABLIKIM	13F	BES3 $J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
26.2 $\pm 6.0 \pm 4.4$	44	² ABLIKIM	07H	BES2 $e^+ e^- \rightarrow \psi(2S)$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

$\Gamma(\Lambda \bar{\Sigma}^+ \pi^+ (\text{or c.c.}))/\Gamma_{\text{total}}$

Γ_{213}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.83 ± 0.07 OUR AVERAGE		Error includes scale factor of 1.2.		
0.770 $\pm 0.051 \pm 0.083$	335	¹ ABLIKIM	07H	BES2 $e^+ e^- \rightarrow \bar{\Lambda} \Sigma^+ \pi^-$
0.747 $\pm 0.056 \pm 0.076$	254	¹ ABLIKIM	07H	BES2 $e^+ e^- \rightarrow \Lambda \bar{\Sigma}^+ \pi^+$
0.90 $\pm 0.06 \pm 0.16$	225 ± 15	HENRARD	87	DM2 $e^+ e^- \rightarrow \bar{\Lambda} \Sigma^+ \pi^-$
1.11 $\pm 0.06 \pm 0.20$	342 ± 18	HENRARD	87	DM2 $e^+ e^- \rightarrow \Lambda \bar{\Sigma}^+ \pi^+$
1.53 $\pm 0.17 \pm 0.38$	135	EATON	84	MRK2 $e^+ e^- \rightarrow \bar{\Lambda} \Sigma^+ \pi^-$
1.38 $\pm 0.21 \pm 0.35$	118	EATON	84	MRK2 $e^+ e^- \rightarrow \Lambda \bar{\Sigma}^+ \pi^+$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\Sigma^+ \rightarrow \pi^0 p) = 51.6\%$.

$\Gamma(p K^- \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{214}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.86 ± 0.11 OUR AVERAGE				
0.84 $^{+0.17}_{-0.15} \pm 0.02$	45	¹ LU	19	BELL $B^+ \rightarrow \bar{p} \Lambda K^+ K^+$
0.89 $\pm 0.07 \pm 0.14$	307	EATON	84	MRK2 $e^+ e^-$

¹ LU 19 reports $(8.32^{+1.63}_{-1.45} \pm 0.49) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow p K^- \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)]$ assuming $B(B^+ \rightarrow J/\psi(1S) K^+) = (1.026 \pm 0.031) \times 10^{-3}$, which we rescale to our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p K^- \bar{\Sigma}^0)/\Gamma_{\text{total}}$

Γ_{215}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.29 $\pm 0.06 \pm 0.05$	90	EATON	84	MRK2 $e^+ e^-$

$\Gamma(\bar{\Lambda} n K_S^0 + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{216}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.46 \pm 0.20 \pm 1.07$	1058	¹ ABLIKIM	08C BES2	$e^+ e^- \rightarrow J/\psi$

¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.

$\Gamma(\Lambda \Sigma + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{217}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.83 ± 0.23 OUR AVERAGE					
$2.74 \pm 0.24 \pm 0.22$		234 ± 21	¹ ABLIKIM	12B BES3	$J/\psi \rightarrow \Lambda \bar{\Sigma}^0$
$2.92 \pm 0.22 \pm 0.24$		308 ± 24	² ABLIKIM	12B BES3	$J/\psi \rightarrow \bar{\Lambda} \Sigma^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<18			² HENRARD	87 DM2	$J/\psi \rightarrow \bar{\Lambda} \Sigma^0$
<15	90		PERUZZI	78 MRK1	$e^+ e^- \rightarrow \Lambda X$

¹ ABLIKIM 12B quotes $B(J/\psi \rightarrow \Lambda \bar{\Sigma}^0)$ which we multiply by 2.

² ABLIKIM 12B and HENRARD 87 quote results for $B(J/\psi \rightarrow \bar{\Lambda} \Sigma^0)$ which we multiply by 2.

$\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}$

Γ_{218}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.07 ± 0.04 OUR AVERAGE				
$1.061 \pm 0.004 \pm 0.036$	87k	ABLIKIM	21AT BES3	$J/\psi \rightarrow p\pi^0 \bar{p}\pi^0$
$1.50 \pm 0.10 \pm 0.22$	399	ABLIKIM	080 BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$

Γ_{219}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.172 ± 0.032 OUR AVERAGE				
$1.164 \pm 0.004 \pm 0.023$	111k	ABLIKIM	17L BES3	$J/\psi \rightarrow \Sigma^0 \bar{\Sigma}^0$
$1.33 \pm 0.04 \pm 0.11$	1.7k	ABLIKIM	06 BES2	$J/\psi \rightarrow \Sigma^0 \bar{\Sigma}^0$
$1.06 \pm 0.04 \pm 0.23$	884	PALLIN	87 DM2	$e^+ e^- \rightarrow \Sigma^0 \bar{\Sigma}^0$
$1.58 \pm 0.16 \pm 0.25$	90	EATON	84 MRK2	$e^+ e^- \rightarrow \Sigma^0 \bar{\Sigma}^0$
1.3 ± 0.4	52	PERUZZI	78 MRK1	$e^+ e^- \rightarrow \Sigma^0 \bar{\Sigma}^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.4 ± 2.6	3	BESCH	81 BONA	$e^+ e^- \rightarrow \Sigma^+ \bar{\Sigma}^-$

$\Gamma(\Sigma^+ \bar{\Sigma}^- \eta)/\Gamma_{\text{total}}$

Γ_{220}/Γ


VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.34 \pm 0.21 \pm 0.37$	1821	ABLIKIM	22AY BES3	$J/\psi \rightarrow \Sigma^+ \bar{\Sigma}^- \eta$



$\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$

Γ_{221}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.97 ± 0.08 OUR AVERAGE				
Error includes scale factor of 1.4. See the ideogram below.				
$1.040 \pm 0.006 \pm 0.074$	43k	ABLIKIM	16L BES3	$J/\psi \rightarrow \Xi^- \bar{\Xi}^+$
$0.90 \pm 0.03 \pm 0.18$	961	ABLIKIM	12P BES2	$J/\psi \rightarrow \Xi^- \bar{\Xi}^+$
$0.70 \pm 0.06 \pm 0.12$	132	HENRARD	87 DM2	$e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$
$1.14 \pm 0.08 \pm 0.20$	194	EATON	84 MRK2	$e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$
1.4 ± 0.5	51	PERUZZI	78 MRK1	$e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$



$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$

Γ_{222}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.7 ± 0.4 OUR AVERAGE	Error includes scale factor of 1.5.			
2.00±0.31±0.02		¹ MITCHELL 09	CLEO	$e^+ e^- \rightarrow \gamma X$
1.27±0.36		GAISER 86	CBAL	$J/\psi \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen		ANASHIN 14	KEDR	$J/\psi \rightarrow \gamma\eta_c$
0.79±0.20	273 ± 43	² AUBERT 06E	BABR	$B^\pm \rightarrow K^\pm \chi_{c\bar{c}}$
seen	16	BALTRUSAITIS 84	MRK3	$J/\psi \rightarrow 2\phi\gamma$

¹ MITCHELL 09 reports $(1.98 \pm 0.09 \pm 0.30) \times 10^{-2}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (35.04 \pm 0.07 \pm 0.77) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Calculated by the authors using an average of $B(J/\psi \rightarrow \gamma\eta_c) \times B(\eta_c \rightarrow K\bar{K}\pi)$ from BALTRUSAITIS 86, BISELLO 91, BAI 04 and $B(\eta_c \rightarrow K\bar{K}\pi) = (8.5 \pm 1.8)\%$ from AUBERT 06E.

$\Gamma(\gamma\eta_c(1S) \rightarrow 3\gamma)/\Gamma_{\text{total}}$

Γ_{223}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8^{+1.3}_{-1.0} OUR AVERAGE	Error includes scale factor of 1.1.			
4.5±1.2±0.6	33 ± 9	ABLIKIM 13I	BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
$1.2^{+2.7}_{-1.1} \pm 0.3$	$1.2^{+2.8}_{-1.1}$	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$\Gamma(\gamma\eta_c(1S) \rightarrow \gamma\eta\eta\eta')/\Gamma_{\text{total}}$ Γ_{224}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.86 \pm 0.62 \pm 0.45$	137	ABLIKIM	21c	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta\eta'$

 $\Gamma(3\gamma)/\Gamma_{\text{total}}$ Γ_{225}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
11.6 ± 2.2 OUR AVERAGE					
$11.3 \pm 1.8 \pm 2.0$		113 ± 18	ABLIKIM	13l	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
$12 \pm 3 \pm 2$		$24.2^{+7.2}_{-6.0}$	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<55	90		PARTRIDGE	80	CBAL $e^+ e^-$

 $\Gamma(4\gamma)/\Gamma_{\text{total}}$ Γ_{226}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<9 \times 10^{-6}$	90	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

 $\Gamma(5\gamma)/\Gamma_{\text{total}}$ Γ_{227}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<15 \times 10^{-6}$	90	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

 $\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$ Γ_{228}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.56 ± 0.17 OUR AVERAGE				
$3.59 \pm 0.20 \pm 0.03$	1.6k	1 ABLIKIM	180	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$
$3.63 \pm 0.36 \pm 0.13$		PEDLAR	09	$J/\psi \rightarrow \pi^0 \gamma$
$3.13^{+0.65}_{-0.47}$	586	ABLIKIM	06E	$J/\psi \rightarrow \pi^0 \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.6 \pm 1.1 \pm 0.7$	BLOOM	83	CBAL	$e^+ e^-$
7.3 ± 4.7	10	BRANDELIK	79C	DASP $e^+ e^-$

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] = (3.57 \pm 0.12 \pm 0.16) \times 10^{-5}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\pi^0 \rightarrow 2\gamma) = (98.823 \pm 0.034) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.


 $\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{229}/Γ

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.15 ± 0.05	1 ABLIKIM	15AE	BES3 $J/\psi \rightarrow \gamma\pi^0\pi^0$

¹ The uncertainty is systematic as statistical is negligible. $\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ Γ_{230}/Γ

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.8 ± 0.5 OUR AVERAGE			Error includes scale factor of 1.9. See the ideogram below.
4.32 $\pm 0.14 \pm 0.73$	¹ BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
2.08 $\pm 0.13 \pm 0.35$	² BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$

3.05 \pm 0.08 \pm 0.45	²	BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
4.85 \pm 0.45 \pm 1.20	³	BURKE	82	MRK2 e^+e^-



¹ 4π mass less than 3.0 GeV.

² 4π mass less than 2.0 GeV.

³ 4π mass less than 2.5 GeV.

$\Gamma(\gamma f_2(1270) f_2(1270))/\Gamma_{\text{total}}$

Γ_{231}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.5 \pm 0.7 \pm 1.6	646 \pm 45	ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

$\Gamma(\gamma f_2(1270) f_2(1270)(\text{non resonant}))/\Gamma_{\text{total}}$

Γ_{232}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.2 \pm 0.8 \pm 1.7	¹ ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

¹ Subtracting contribution from intermediate $\eta_c(1S)$ decays.

$\Gamma(\gamma \pi^+ \pi^- 2\pi^0)/\Gamma_{\text{total}}$

Γ_{233}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
8.3 \pm 0.2 \pm 3.1	¹ BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$

¹ 4π mass less than 2.0 GeV.

$\Gamma(\gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

Γ_{234}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.1 \pm 0.4	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma(K\bar{K}\pi)[J^PC=0^-+])/\Gamma_{\text{total}}$ Γ_{235}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.7 ± 0.4 OUR AVERAGE	Error includes scale factor of 2.1.		
0.58 ± 0.03 ± 0.20	¹ BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
2.1 ± 0.1 ± 0.7	² BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$

¹ For a broad structure around 1800 MeV.² For a broad structure around 2040 MeV.
 $\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{236}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.1 ± 0.1 ± 0.6	1516	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

 $\Gamma(\gamma K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{237}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.0 ± 0.3 ± 1.3	320	¹ BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

¹ Summed over all charges.
 $\Gamma(\gamma\eta)/\Gamma_{\text{total}}$ Γ_{238}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.085 ± 0.018 OUR AVERAGE				
1.067 ± 0.005 ± 0.023	87.9k	ABLIKIM	21AM BES3	$e^+ e^- \rightarrow J/\psi$
1.12 ± 0.05 ± 0.01	18.6k	¹ ABLIKIM	18O BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$
1.101 ± 0.029 ± 0.022		PEDLAR	09 CLE3	$J/\psi \rightarrow \eta \gamma$
1.123 ± 0.089	11k	ABLIKIM	06E BES2	$J/\psi \rightarrow \eta \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.88 ± 0.08 ± 0.11		BLOOM	83 CBAL	$e^+ e^-$
0.82 ± 0.10		BRANDELIK	79C DASP	$e^+ e^-$
1.3 ± 0.4	21	BARTEL	77 CNTR	$e^+ e^-$

¹ ABLIKIM 18O reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = (4.42 \pm 0.04 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

 $\Gamma(\gamma\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{239}/Γ

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
21.4 ± 1.8 ± 2.5	596	ABLIKIM	16P BES3	$J/\psi \rightarrow 5\gamma$

 $\Gamma(\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{243}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 2.5 \times 10^{-6}$	95	ABLIKIM	16P BES3	$J/\psi \rightarrow 5\gamma$

 $\Gamma(\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{244}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 6.6 \times 10^{-6}$	95	ABLIKIM	16P BES3	$J/\psi \rightarrow 5\gamma$

$\Gamma(\gamma\eta\pi\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})

6.1 ±1.0 OUR AVERAGE

$5.85 \pm 0.3 \pm 1.05$

$7.8 \pm 1.2 \pm 2.4$

¹ Broad enhancement at 1700 MeV.

Γ_{245}/Γ

DOCUMENT ID

TECN

COMMENT

¹ EDWARDS

83B

CBAL

$J/\psi \rightarrow \eta\pi^+\pi^-$

¹ EDWARDS

83B

CBAL

$J/\psi \rightarrow \eta 2\pi^0$

$\Gamma(\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})

6.2±2.2±0.9

DOCUMENT ID

TECN

COMMENT

BAI

99

BES

$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})

5.25±0.07 OUR AVERAGE

DOCUMENT ID

TECN

COMMENT

Error includes scale factor of 1.3. See the ideogram below.

$5.27 \pm 0.03 \pm 0.05$

36k

19T

BES

$J/\psi \rightarrow \gamma\eta'$

$5.43 \pm 0.23 \pm 0.09$

5.0k

¹ ABLIKIM

180

BES3

$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

$4.77 \pm 0.22 \pm 0.06$

² ABLIKIM

11

BES3

$J/\psi \rightarrow \eta'\gamma$

$5.24 \pm 0.12 \pm 0.11$

PEDLAR

09

CLE3

$J/\psi \rightarrow \eta'\gamma$

5.55 ± 0.44

35k

ABLIKIM

06E

BES2

$J/\psi \rightarrow \eta'\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.50 \pm 0.14 \pm 0.53$

BOLTON

92B

MRK3

$J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \gamma\gamma$

$4.30 \pm 0.31 \pm 0.71$

BOLTON

92B

MRK3

$J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \pi^+\pi^-\pi^0$

$4.04 \pm 0.16 \pm 0.85$

622

AUGUSTIN

90

DM2

$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

$4.39 \pm 0.09 \pm 0.66$

2420

AUGUSTIN

90

DM2

$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

$4.1 \pm 0.3 \pm 0.6$

BLOOM

83

CBAL

$e^+e^- \rightarrow 3\gamma + \text{hadrons}$

2.9 ± 1.1

6

BRANDELIK

79C

DASP

$e^+e^- \rightarrow 3\gamma$

2.4 ± 0.7

57


BARTEL

76

CNTR

$e^+e^- \rightarrow 2\gamma\rho$

WEIGHTED AVERAGE
5.25±0.07 (Error scaled by 1.3)



χ^2

0.1

0.5

4.4

0.0

5.0

(Confidence Level = 0.168)

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ (units 10^{-3})

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] = (1.26 \pm 0.02 \pm 0.05) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\eta'(958) \rightarrow \gamma\gamma) = (2.307 \pm 0.033) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² ABLIKIM 11 reports $(4.84 \pm 0.03 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] / [B(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)]$ assuming $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (43.2 \pm 0.7) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$, which we rescale to our best values $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (42.5 \pm 0.5) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\gamma f_0(500) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

Γ_{240}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.5 ± 2.0 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(500) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

Γ_{241}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

5 ± 5 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(500) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$

Γ_{242}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4 ± 3 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(980) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

Γ_{248}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.3 ± 0.2 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(980) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

Γ_{249}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.8 ± 0.3 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma\rho\rho)/\Gamma_{\text{total}}$

Γ_{250}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

4.5 ±0.8 OUR AVERAGE

$4.7 \pm 0.3 \pm 0.9$

¹ BALTRUSAIT..86B MRK3 $J/\psi \rightarrow 4\pi\gamma$

$3.75 \pm 1.05 \pm 1.20$

² BURKE 82 MRK2 $J/\psi \rightarrow 4\pi\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.09 90 ³ BISELLO 89B $J/\psi \rightarrow 4\pi\gamma$

¹ 4π mass less than 2.0 GeV.

² 4π mass less than 2.0 GeV. We have multiplied $2\rho^0$ measurement by 3 to obtain 2ρ .

³ 4π mass in the range 2.0–25 GeV.

$\Gamma(\gamma\rho\omega)/\Gamma_{\text{total}}$

VALUE	CL%
$<5.4 \times 10^{-4}$	90

Γ_{251}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	08A BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\gamma\rho\phi)/\Gamma_{\text{total}}$

VALUE	CL%
$<8.8 \times 10^{-5}$	90

Γ_{252}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	08A BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\gamma\omega\omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS
1.61 ± 0.33 OUR AVERAGE	

6.0 $\pm 4.8 \pm 1.8$	
$1.41 \pm 0.2 \pm 0.42$	120 ± 17
$1.76 \pm 0.09 \pm 0.45$	

Γ_{253}/Γ

DOCUMENT ID	TECN	COMMENT
-------------	------	---------

ABLIKIM	08A BES2	$J/\psi \rightarrow \gamma\omega\pi^+\pi^-$
BISELLO	87 SPEC	$e^+ e^-$, hadrons γ
BALTRUSAIT..85C	MRK3	$e^+ e^- \rightarrow$ hadrons γ

$\Gamma(\gamma\phi\phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS
4.0 ± 1.2 OUR AVERAGE	Error includes scale factor of 2.1. See the ideogram below.


7.5 $\pm 0.6 \pm 1.2$	168
$3.4 \pm 0.8 \pm 0.6$	33 ± 7
$3.1 \pm 0.7 \pm 0.4$	

Γ_{254}/Γ

DOCUMENT ID	TECN	COMMENT
-------------	------	---------

BAI	90B MRK3	$J/\psi \rightarrow \gamma 4K$
¹ BISELLO	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
¹ BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

WEIGHTED AVERAGE
 4.0 ± 1.2 (Error scaled by 2.1)



$\Gamma(\gamma\phi\phi)/\Gamma_{\text{total}}$ (units 10^{-4})

¹ $\phi\phi$ mass less than 2.9 GeV, η_C excluded.


$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$

Γ_{255}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.6 OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.		
1.66 ± 0.1	1,2 BAI	00D BES	$J/\psi \rightarrow \gamma K^{\pm} K_S^0 \pi^{\mp}$
3.8 ± 0.3	3 AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
4.0 ± 0.7	3 EDWARDS	82E CBAL	$J/\psi \rightarrow K^+ K^- \pi^0 \gamma$
4.3 ± 1.7	3,4 SCHARRE	80 MRK2	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.78 ± 0.21 ± 0.33	3,5,6 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
0.83 ± 0.13 ± 0.18	3,7,8 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
0.66 ^{+0.17 + 0.24} _{-0.16 - 0.15}	3,6,9 BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$
1.03 ^{+0.21 + 0.26} _{-0.18 - 0.19}	3,8,10 BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$

WEIGHTED AVERAGE

2.8 ± 0.6 (Error scaled by 1.6)



$$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}} (\text{units } 10^{-3})$$

¹ Interference with the $J/\psi(1S)$ radiative transition to the broad $K\bar{K}\pi$ pseudoscalar state around 1800 is $(0.15 \pm 0.01 \pm 0.05) \times 10^{-3}$.

² Interference with $J/\psi \rightarrow \gamma f_1(1420)$ is $(-0.03 \pm 0.01 \pm 0.01) \times 10^{-3}$.

³ Includes unknown branching fraction $\eta(1405) \rightarrow K\bar{K}\pi$.

⁴ Corrected for spin-zero hypothesis for $\eta(1405)$.

⁵ From fit to the $a_0(980)\pi$ $0^- +$ partial wave.

⁶ $a_0(980)\pi$ mode.

⁷ From fit to the $K^*(892)K$ $0^- +$ partial wave.

⁸ $K^* K$ mode.

⁹ From $a_0(980)\pi$ final state.

¹⁰ From $K^*(890)K$ final state.

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0)/\Gamma_{\text{total}}$ Γ_{256}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.78±0.20 OUR AVERAGE	Error includes scale factor of 1.8.		
1.07±0.17±0.11	¹ BAI 04J	BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
0.64±0.12±0.07	¹ COFFMAN 90	MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

¹ Includes unknown branching fraction $\eta(1405) \rightarrow \gamma\rho^0$.

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{257}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.0 ±0.5 OUR AVERAGE				
2.6 ±0.7 ±0.4		BAI 99	BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
3.38±0.33±0.64		¹ BOLTON 92B	MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
7.0 ±0.6 ±1.1	261	² AUGUSTIN 90	DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

¹ Via $a_0(980)\pi$.

² Includes unknown branching fraction to $\eta\pi^+\pi^-$.

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$ Γ_{258}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.7 ±0.4 OUR AVERAGE	Error includes scale factor of 1.3.		
2.1 ±0.4	BUGG 95	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
1.36±0.38	^{1,2} BISELLO 89B	DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits.

² Includes unknown branching fraction to $\rho^0\rho^0$.

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi)/\Gamma_{\text{total}}$ Γ_{259}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<82	95		BAI 04J	BES2	$J/\psi \rightarrow \gamma\gamma K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
7.03±0.92±0.91	1.3k	¹ ABLIKIM 18I	BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$	
10.36±1.51±1.54	1.9k	² ABLIKIM 18I	BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$	

¹ Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

² Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

 $\Gamma(\gamma\eta(1405) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{260}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.63 \times 10^{-6}$	90	ABLIKIM 180	BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

 $\Gamma(\gamma\eta(1475) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{261}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.86 \times 10^{-6}$	90	ABLIKIM 180	BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

$\Gamma(\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$ Γ_{262}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.13 ± 0.09	1,2 BISELLO	89B	DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits.² Includes unknown branching fraction to $\rho^0\rho^0$. $\Gamma(\gamma\eta(1760) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$ Γ_{263}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.98 \pm 0.08 \pm 0.32$	1045	ABLIKIM	06H	BES

 $\Gamma(\gamma\eta(1760) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{264}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<4.80 \times 10^{-6}$	90	ABLIKIM	180	BES3

 $\Gamma(\gamma\eta(2225))/\Gamma_{\text{total}}$ Γ_{265}/Γ


<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.14^{+0.50}_{-0.19} \text{ OUR AVERAGE}$				

$2.40 \pm 0.10^{+2.47}_{-0.18}$	1,2 ABLIKIM	16N	BES3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$4.4 \pm 0.4 \pm 0.8$	196	2 ABLIKIM	08I	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$3.3 \pm 0.8 \pm 0.5$		2 BAI	90B	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$2.7 \pm 0.6 \pm 0.6$		2 BAI	90B	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$2.4^{+1.5}_{-1.0}$	3,4 BISELLO	89B	DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma\phi\phi$ that also finds significant signals for $\eta(2100)$, 0^-+ phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen 0^-+ state $X(2500)$ ($M = 2470^{+15+101}_{-19-23}$ MeV, $\Gamma = 230^{+64+56}_{-35-33}$ MeV).² Includes unknown branching fraction to $\phi\phi$.³ Estimated by us from various fits.⁴ Includes unknown branching fraction to $\rho^0\rho^0$. $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{266}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.63 \pm 0.12 \text{ OUR AVERAGE}$		Error includes scale factor of 1.3. See the ideogram below.		

$2.07 \pm 0.16^{+0.02}_{-0.07}$	2.4k	1,2 DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
$1.63 \pm 0.26^{+0.02}_{-0.06}$		3 ABLIKIM	06v	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
$1.42 \pm 0.21^{+0.02}_{-0.05}$		4 ABLIKIM	06v	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$
$1.33 \pm 0.05 \pm 0.20$		5 AUGUSTIN	87	$J/\psi \rightarrow \gamma\pi^+\pi^-$
$1.36 \pm 0.09 \pm 0.23$		5 BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$
$1.48 \pm 0.25 \pm 0.30$	178	EDWARDS	82B	$CBAL \quad e^+ e^- \rightarrow 2\pi^0\gamma$
2.0 ± 0.7	35	ALEXANDER	78	$PLUT \quad e^+ e^-$
1.2 ± 0.6	30	BRANDELIK	78B	$DASP \quad e^+ e^- \rightarrow \pi^+\pi^-\gamma$



¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.744 \pm 0.052 \pm 0.122) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.371 \pm 0.010 \pm 0.222) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.200 \pm 0.027 \pm 0.174) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ Estimated using $B(f_2(1270) \rightarrow \pi\pi) = 0.843 \pm 0.012$. The errors do not contain the uncertainty in the $f_2(1270)$ decay.

⁶ Restated by us to take account of spread of E1, M2, E3 transitions.

$\Gamma(\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$	Γ_{267}/Γ		
VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$2.58^{+0.08+0.59}_{-0.09-0.20}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_1(1285))/\Gamma_{\text{total}}$	Γ_{268}/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.61 ± 0.08 OUR AVERAGE			
0.69 $\pm 0.16 \pm 0.20$	¹ BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\rho^0$
0.61 $\pm 0.04 \pm 0.21$	² BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$

$0.45 \pm 0.09 \pm 0.17$	³ BAI	99	BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$0.625 \pm 0.063 \pm 0.103$	⁴ BOLTON	92	MRK3	$J/\psi \rightarrow \gamma f_1(1285)$
$0.70 \pm 0.08 \pm 0.16$	⁵ BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

¹ Assuming $B(f_1(1285) \rightarrow \rho^0 \gamma) = 0.055 \pm 0.013$.

² Assuming $\Gamma(f_1(1285) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}} = 0.090 \pm 0.004$.

³ Assuming $\Gamma(f_1(1285) \rightarrow \eta\pi\pi)/\Gamma_{\text{total}} = 0.5 \pm 0.18$.

⁴ Obtained summing the sequential decay channels

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi\pi\pi\pi) = (1.44 \pm 0.39 \pm 0.27) \times 10^{-4}$;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980)\pi, a_0(980) \rightarrow \eta\pi) = (3.90 \pm 0.42 \pm 0.87) \times 10^{-4}$;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980)\pi, a_0(980) \rightarrow K\bar{K}) = (0.66 \pm 0.26 \pm 0.29) \times 10^{-4}$;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma\rho^0) = (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}$.

⁵ Using $B(f_1(1285) \rightarrow a_0(980)\pi) = 0.37$, and including unknown branching ratio for $a_0(980) \rightarrow \eta\pi$.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

Γ_{269}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

38 ± 10	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------------	-----------	----	------	---

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

Γ_{270}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

4.19 ± 0.73 ± 1.34	478	¹ DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
---------------------------	-----	--------------------	----	--------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.3 ± 0.4	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	----	------	---

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

Γ_{271}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

$1.07^{+0.08}_{-0.07} {}^{+0.36}_{-0.34}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
---	---------	-----------	---

$\Gamma(\gamma f_0(1370) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$

Γ_{272}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.5 ± 1.0	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	----	------	---

$\Gamma(\gamma f_0(1370) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$

Γ_{273}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.9 ± 0.3	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	----	------	---

$\Gamma(\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$

Γ_{274}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.79±0.13 OUR AVERAGE			
0.68±0.04±0.24	BAI	00D BES	$J/\psi \rightarrow \gamma K^{\pm} K_S^0 \pi^{\mp}$
0.76±0.15±0.21	1,2 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
0.87±0.14 ^{+0.14} _{-0.11}	1 BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$

¹ Included unknown branching fraction $f_1(1420) \rightarrow K\bar{K}\pi$.

² From fit to the $K^*(892)K$ 1^{++} partial wave.

$\Gamma(\gamma f_0(1500) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

Γ_{275}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.09±0.24 OUR AVERAGE				
1.21±0.29±0.24	174	1 DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
1.00±0.03±0.45		2 ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1.02±0.09±0.45		2 ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.90±0.17		SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
5.7 ± 0.8	3,4 BUGG		95 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi\pi$.

³ Including unknown branching ratio for $f_0(1500) \rightarrow \pi^+\pi^-\pi^+\pi^-$.

⁴ Assuming that $f_0(1500)$ decays only to two S -wave dipions.

$\Gamma(\gamma f_0(1500) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$

Γ_{276}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.65^{+0.26+0.51}_{-0.31-1.40}				
5.5k	1 ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.1 ± 0.4		SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

Γ_{277}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
1.59±0.16^{+0.18}_{-0.56}			
	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.7 ± 0.3		SARANTSEV	21 RVUE
			$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(1500) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$

Γ_{278}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
18.1±1.1 ^{+1.9} _{-1.3}	1 ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
12 ± 5	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta' P$ -wave.			

$\Gamma(\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{279}/Γ

VALUE (units 10^{-4})

$4.5 \pm 1.0 \pm 0.7$

DOCUMENT ID

BAI

TECN

99

COMMENT

$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$

Γ_{280}/Γ

VALUE (units 10^{-4})

CL% EVTS

DOCUMENT ID

TECN


COMMENT

5.7	+0.8	OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.
8.1	± 0.9	± 0.2	750 1,2 DOBBS 15 $J/\psi \rightarrow \gamma K\bar{K}$
3.85 ± 0.17	$+1.91$	-0.73	3 BAI 03G BES $J/\psi \rightarrow \gamma K\bar{K}$
3.6 ± 0.4	$+1.4$	-0.4	3 BAI 96C BES $J/\psi \rightarrow \gamma K^+ K^-$
5.6 ± 1.4	± 0.9		3 AUGUSTIN 88 DM2 $J/\psi \rightarrow \gamma K^+ K^-$
4.5 ± 0.4	± 0.9		3 AUGUSTIN 88 DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
6.8 ± 1.6	± 1.4		3 BALTRUSAIT...87 MRK3 $J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.4	90	4	⁴ BRANDELIK	79C	DASP	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
<2.3	90	3	ALEXANDER	78	PLUT	$e^+ e^- \rightarrow K^+ K^- \gamma$

WEIGHTED AVERAGE
5.7+0.8-0.5 (Error scaled by 1.5)



$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$ (units 10^{-4})

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = (7.09 \pm 0.46 \pm 0.67) \times 10^{-4}$ which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.888$.

⁴ Assuming isotropic production and decay of the $f'_2(1525)$ and isospin.

$\Gamma(\gamma f_2(1565) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$

Γ_{283}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.32 \pm 0.05^{+0.12}_{-0.02}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta' P$ -wave.

$\Gamma(\gamma f'_2(1525) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

Γ_{281}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$7.99^{+0.03+0.69}_{-0.04-0.50}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
----------------------------------	---------	-----------	---

$\Gamma(\gamma f'_2(1525) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$

Γ_{282}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$3.42^{+0.43+1.37}_{-0.51-1.30}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$
----------------------------------	------	----------------------	----------	-------------------------------------

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_2(1640) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$

Γ_{284}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$0.28 \pm 0.05 \pm 0.17$	141	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$
--------------------------	-----	---------	---------	---

$\Gamma(\gamma f_0(1710) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

Γ_{285}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.8 ± 0.5 OUR AVERAGE

$3.72 \pm 0.30 \pm 0.43$	483	¹ DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
$3.96 \pm 0.06 \pm 1.12$		² ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
$3.99 \pm 0.15 \pm 2.64$		² ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.2	³ SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	------------------------	---------	---

$2.5 \pm 1.6 \pm 0.8$	BAI	98H BES	$J/\psi \rightarrow \gamma\pi^0\pi^0$
-----------------------	-----	---------	---------------------------------------

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi\pi$.

³ There is a further $(2.4 \pm 0.8) \times 10^{-4}$ scalar contribution at 1765 MeV.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

Γ_{286}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

$9.5^{+1.0}_{-0.5}$ OUR AVERAGE			Error includes scale factor of 1.5. See the ideogram below.
---	--	--	---


$8.00^{+0.12+1.24}_{-0.08-0.40}$		¹ ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
----------------------------------	--	----------------------	-----------	---

$11.76 \pm 0.54 \pm 0.94$	1.2k	² DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
---------------------------	------	--------------------	----	--------------------------------------

$9.62 \pm 0.29^{+3.51}_{-1.86}$		³ BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
---------------------------------	--	------------------	---------	--------------------------------------

$5.0 \pm 0.8^{+1.8}_{-0.4}$	1,4 BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
$9.2 \pm 1.4 \pm 1.4$	1 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
$10.4 \pm 1.2 \pm 1.6$	1 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$9.6 \pm 1.2 \pm 1.8$	1 BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.3 ± 0.8	5 SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$1.6 \pm 0.2^{+0.6}_{-0.2}$	1,6 BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
< 0.8	7 BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$
$1.6 \pm 0.4 \pm 0.3$	8 BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$
3.8 ± 1.6	9 EDWARDS	82D CBAL	$e^+e^- \rightarrow \eta\eta\gamma$

WEIGHTED AVERAGE
9.5+1.0-0.5 (Error scaled by 1.5)



¹ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied $K^+ K^-$ measurement by 2, and $K_S^0 K_S^0$ by 4 to obtain $K\bar{K}$ result.

² Using CLEO-c data but not authored by the CLEO Collaboration.

³ Includes unknown branching ratio to $K^+ K^-$ or $K_S^0 K_S^0$.

⁴ Assuming $J^P = 2^+$ for $f_0(1710)$.

⁵ There is a further $(6 \pm 2) \times 10^{-4}$ scalar contribution at 1765 MeV.

⁶ Assuming $J^P = 0^+$ for $f_0(1710)$.

⁷ Includes unknown branching fraction to $\rho^0 \rho^0$.

⁸ Includes unknown branching fraction to $\pi^+ \pi^-$.

⁹ Includes unknown branching fraction to $\eta\eta$.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$

Γ_{287}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.31 \pm 0.06 \pm 0.08$	180	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$

Γ_{288}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.35^{+0.13+1.24}_{-0.11-0.74}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.2 ± 0.4	² SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	------------------------	----	------	---

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

² There is a further $(0.7 \pm 0.1) \times 10^{-4}$ scalar contribution at 1765 MeV.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$

Γ_{289}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.5 ± 2.5	¹ SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	------------------------	----	------	---

¹ There is a further $(2.5 \pm 1.1) \times 10^{-5}$ scalar contribution at 1765 MeV.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma\omega\phi)/\Gamma_{\text{total}}$

Γ_{290}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	-------------	--------------------	-------------	----------------

2.5 ± 0.6 OUR AVERAGE

$2.00 \pm 0.08^{+1.38}_{-1.64}$	1.3k	ABLIKIM	13J BES3	$J/\psi \rightarrow \gamma\omega\phi$
---------------------------------	------	---------	----------	---------------------------------------

$2.61 \pm 0.27 \pm 0.65$	95	ABLIKIM	06J BES2	$J/\psi \rightarrow \gamma\omega\phi$
--------------------------	----	---------	----------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.1 ± 0.1	¹ SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	------------------------	----	------	---

¹ There is a further $(2.2 \pm 0.4) \times 10^{-4}$ scalar contribution at 1765 MeV.

$\Gamma(\gamma f_0(1770) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

Γ_{291}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	--------------------	-------------	----------------

$1.11 \pm 0.06^{+0.19}_{-0.32}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
---	---------	-----------	---

$\Gamma(\gamma f_0(1770) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$

Γ_{294}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.11 \pm 0.01^{+0.04}_{-0.03}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P -wave.

$\Gamma(\gamma f_2(1810) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{292}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
5.40\pm0.60\pm3.42 -0.67 -2.35	5.5k	¹ ABLIKIM	13N $J/\psi \rightarrow \gamma\eta\eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

 $\Gamma(\gamma\eta_1(1855) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$ Γ_{293}/Γ

<u>VALUE</u> (units 10^{-6})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.70\pm0.41\pm0.16 -0.35	¹ ABLIKIM	22AI BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$

¹ From a Breit-Wigner fit involving 9 resonances and the resonating exotic $\eta_1(1855) \rightarrow \eta\eta' P$ -wave. For analysis details see ABLIKIM 22AS.

 $\Gamma(\gamma f_2(1910) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$ Γ_{295}/Γ

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.20\pm0.04\pm0.13	151	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

 $\Gamma(\gamma f_2(1950) \rightarrow \gamma K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{296}/Γ

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.7\pm0.1\pm0.2	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

 $\Gamma(\gamma f_2(2010) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$ Γ_{297}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			

$0.71 \pm 0.06 \pm 0.10$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
--------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta' P$ -wave.

 $\Gamma(\gamma f_0(2020) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{298}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
42 \pm 10	SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$	

 $\Gamma(\gamma f_0(2020) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{299}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
55 \pm 25	SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$	

 $\Gamma(\gamma f_0(2020) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{300}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10 \pm 10	SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$	

$\Gamma(\gamma f_0(2020) \rightarrow \gamma \eta' \eta')/\Gamma_{\text{total}}$ Γ_{301}/Γ

<u>VALUE</u> (units 10^{-4})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.63 \pm 0.06^{+0.31}_{-0.46}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta' \eta'$, and ($\eta' X$), with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

 $\Gamma(\gamma f_0(2020) \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$ Γ_{302}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			

$2.28 \pm 0.12^{+0.29}_{-0.20}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave.

 $\Gamma(\gamma f_4(2050))/\Gamma_{\text{total}}$ Γ_{303}/Γ

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.7 \pm 0.5 \pm 0.5$	¹ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$

¹ Assuming branching fraction $f_4(2050) \rightarrow \pi \pi / \text{total} = 0.167$.

 $\Gamma(\gamma f_4(2050) \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$ Γ_{304}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			

$0.06 \pm 0.01^{+0.03}_{-0.01}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave.

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{305}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.13^{+0.09+0.64}_{-0.10-0.28}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

1.8 ± 1.5	SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K\bar{K}, \eta \eta, \omega \phi)$
---------------	-------------------	---

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{307}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$6.24 \pm 0.48 \pm 0.87$	744	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

2.0 ± 0.8	SARANTSEV 21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K\bar{K}, \eta \eta, \omega \phi)$
---------------	-------------------	---

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2100) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

Γ_{306}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
32 \pm 20	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(2200))/\Gamma_{\text{total}}$

Γ_{308}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
1.5	¹ AUGUSTIN 88	DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
¹ Includes unknown branching fraction to $K_S^0 K_S^0$.			

$\Gamma(\gamma f_0(2200) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

Γ_{311}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
5 \pm 2	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

Γ_{309}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.86 \pm 0.49 \pm 1.20	490	¹ DOBBS 15		$J/\psi \rightarrow \gamma K\bar{K}$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.5 \pm 0.5	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$	

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

Γ_{310}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.72 \pm 0.08 \pm 0.17	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(2200) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$

Γ_{312}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.7 \pm 0.4	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$

Γ_{313}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
>300			¹ BAI 96B	BES	$e^+ e^- \rightarrow \gamma\bar{p}p, K\bar{K}$
>250	99.9		² HASAN 96	SPEC	$\bar{p}p \rightarrow \pi^+\pi^-$
< 2.3	95		³ AUGUSTIN 88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$
< 1.6	95		³ AUGUSTIN 88	DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$12.4^{+6.4}_{-5.2} \pm 2.8$		23	³ BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$8.4^{+3.4}_{-2.8} \pm 1.6$		93	³ BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

¹ Using BARNES 93.

² Using BAI 96B.

³ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

Γ_{314}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 3.9	90	1,2 DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

14 ± 8 ± 4	BAI	98H BES	$J/\psi \rightarrow \gamma\pi^0\pi^0$
8.4 ± 2.6 ± 3.0	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+\pi^-$ and $\pi^0\pi^0$ are $2.6/5.2 \times 10^{-5}$ and $1.3/1.9 \times 10^{-5}$, respectively.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

Γ_{315}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 4.1	90	1,2 DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.6	³ DEL-AMO-SA..100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+K^-$
< 2.9	³ DEL-AMO-SA..100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
6.6 ± 2.9 ± 2.4	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+K^-$
10.8 ± 4.0 ± 3.2	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for K^+K^- and $K_S^0K_S^0$ are $1.7/3.1 \times 10^{-5}$ and $1.2/2.0 \times 10^{-5}$, respectively.

³ For spin 2 and helicity 0; other combinations lead to more stringent upper limits.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$

Γ_{316}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.5±0.6±0.5	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}$

$\Gamma(\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

Γ_{317}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.95±0.21^{+0.66}_{-0.72}	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.1	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-----------	-----------	---------	---

$\Gamma(\gamma f_0(2330) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

Γ_{318}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			

4 ± 2 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(2330) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$

Γ_{319}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			

1.5 ± 0.4 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(2330) \rightarrow \gamma\eta'\eta')/\Gamma_{\text{total}}$ Γ_{320}/Γ

<u>VALUE</u> (units 10^{-6})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$6.09 \pm 0.64^{+4.00}_{-1.68}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta'\eta'$, and ($\eta' X$), with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$ Γ_{321}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			

$0.10 \pm 0.02^{+0.01}_{-0.02}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta' P$ -wave.

 $\Gamma(\gamma f_0(2470) \rightarrow \gamma\eta'\eta')/\Gamma_{\text{total}}$ Γ_{325}/Γ

<u>VALUE</u> (units 10^{-7})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.18 \pm 1.77^{+3.73}_{-2.23}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta'\eta'$, and ($\eta' X$), with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

 $\Gamma(\gamma f_2(2340) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{322}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.60^{+0.62+2.37}_{-0.65-2.07}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

 $\Gamma(\gamma f_2(2340) \rightarrow \gamma\eta'\eta')/\Gamma_{\text{total}}$ Γ_{324}/Γ

<u>VALUE</u> (units 10^{-6})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.67 \pm 0.70^{+0.61}_{-1.67}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta'\eta'$, and ($\eta' X$), with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

 $\Gamma(\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{323}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.54^{+0.34+3.82}_{-0.40-1.49}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma X(1835) \rightarrow \gamma\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{326}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.7 $^{+0.6}_{-0.8}$ OUR AVERAGE				Error includes scale factor of 1.6.
$3.93 \pm 0.38^{+0.31}_{-0.84}$		¹ ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
$2.2 \pm 0.4 \pm 0.4$	264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$2.87 \pm 0.09^{+0.49}_{-0.52}$	4265	² ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

- ¹ From a fit of the measured $\pi^+\pi^-\eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold with a Flatte formula in addition to known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the $X(1835)$; fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.
- ² From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two states $\gamma X(2120)$ and $\gamma X(2370)$, for $M(\pi^+\pi^-\eta') < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$.

 $\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{327}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.77 $^{+0.15}_{-0.09}$ OUR AVERAGE				
$0.90^{+0.04+0.27}_{-0.11-0.55}$		¹ ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$
$1.14^{+0.43+0.42}_{-0.30-0.26}$	231	² ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p\bar{p}$
$0.70 \pm 0.04^{+0.19}_{-0.08}$		BAI	03F BES2	$J/\psi \rightarrow \gamma p\bar{p}$

- ¹ From the fit including final state interaction effects in isospin 0 *S*-wave according to SIBIRTSEV 05A.
- ² From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.

 $\Gamma(\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta)/\Gamma_{\text{total}}$ Γ_{328}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.31 $^{+0.33+1.96}_{-0.30-1.29}$		ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$

 $\Gamma(\gamma X(1835) \rightarrow \gamma\gamma\phi(1020))/\Gamma_{\text{total}}$ Γ_{329}/Γ

<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.77 \pm 0.35 \pm 0.25$	305	¹ ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
$8.09 \pm 1.99 \pm 1.36$	1.3k	² ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$

- ¹ Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.
- ² Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

$\Gamma(\gamma X(1835) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{330}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.56 \times 10^{-6}$	90	ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$

 $\Gamma(\gamma X(1835) \rightarrow \gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{331}/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.44 \pm 0.36^{+0.60}_{-0.74}$	0.6k	ABLIKIM	13U	BES3 $J/\psi \rightarrow \gamma 3(\pi^+ \pi^-)$

 $\Gamma(\gamma X(2370) \rightarrow \gamma K^+ K^- \eta')/\Gamma_{\text{total}}$ Γ_{332}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.79 \pm 0.23 \pm 0.65$	ABLIKIM	20Q	BES3 $J/\psi \rightarrow \gamma K^+ K^- \eta'$

 $\Gamma(\gamma X(2370) \rightarrow \gamma K_S^0 K_S^0 \eta')/\Gamma_{\text{total}}$ Γ_{333}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.18 \pm 0.32 \pm 0.39$	ABLIKIM	20Q	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$

 $\Gamma(\gamma X(2370) \rightarrow \gamma \eta \eta \eta')/\Gamma_{\text{total}}$ Γ_{334}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<9.2	90	ABLIKIM	21C	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta \eta'$

 $\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{335}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.38 \pm 0.07 \pm 0.07$	49	EATON	84	MRK2	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.11	90	PERUZZI	78	MRK1	$e^+ e^-$
---------	----	---------	----	------	-----------

 $\Gamma(\gamma p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{336}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<0.79 \times 10^{-3}$	90	EATON	84	MRK2 $e^+ e^-$

 $\Gamma(\gamma \Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{337}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<0.13 \times 10^{-3}$	90	HENRARD	87	DM2 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.16 \times 10^{-3}$	90	BAI	98G	BES	$e^+ e^-$
------------------------	----	-----	-----	-----	-----------

 $\Gamma(\gamma A^0 \rightarrow \gamma \text{invisible})/\Gamma_{\text{total}}$ Γ_{338}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.7 \times 10^{-6}$	90	88M	¹ ABLIKIM	20K	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<6.3 \times 10^{-6}$	90	3.7M	² INSLER	10	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
-----------------------	----	------	---------------------	----	--

¹ For a narrow state, A^0 , with mass $m_{A^0} < 1.2$ GeV. The limit varies with m_{A^0} , reaching its largest value of 1.7×10^{-6} at 1.2 GeV and being 7.0×10^{-7} for $m_{A^0} = 0$.

² The limit varies with mass m_{A^0} of a narrow state A^0 and is 4.3×10^{-6} for $m_{A^0} = 0$, reaches its largest value of 6.3×10^{-6} at $m_{A^0} = 500$ MeV, and is 3.6×10^{-6} at $m_{A^0} = 960$ MeV.

$\Gamma(\gamma A^0 \rightarrow \gamma\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{339}/Γ (narrow state A^0 with 0.2 GeV $< m_{A^0} < 3$ GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.8 \times 10^{-7}$	90	1 ABLIKIM	22H BES3	$J/\psi \rightarrow \gamma\mu^+\mu^-$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<0.5 \times 10^{-5}$	90	2 ABLIKIM	16E BES3	$J/\psi \rightarrow \gamma\mu^+\mu^-$
$<2.1 \times 10^{-5}$	90	3 ABLIKIM	12 BES3	$J/\psi \rightarrow \gamma\mu^+\mu^-$

¹ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3.0 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(1.2\text{--}778.0) \times 10^{-9}$.

² For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(2.8\text{--}495.3) \times 10^{-8}$.

³ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.21–3.00 GeV. The measured 90% CL limit as a function of m_{A^0} ranges from 4×10^{-7} to 2.1×10^{-5} .

 DALITZ DECAYS

 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{340}/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.56 \pm 1.32 \pm 0.50$	39	ABLIKIM	14I BES3	$J/\psi \rightarrow \pi^0 e^+ e^-$

 $\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_{341}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.42 \pm 0.04 \pm 0.07$	2.47k	1,2 ABLIKIM	19A BES3	$J/\psi \rightarrow \eta e^+ e^-$

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

$1.16 \pm 0.07 \pm 0.06$	320	1 ABLIKIM	14I BES3	$J/\psi \rightarrow \eta e^+ e^-$
--------------------------	-----	-----------	----------	-----------------------------------

¹ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decays.

² Approximation of the transition form factor squared as an incoherent sum of the ρ -meson and one-pole non-resonant amplitudes gives the pole mass $m(\Lambda) = 2.56 \pm 0.04 \pm 0.03$ GeV. Supersedes ABLIKIM 14I.

 $\Gamma(\eta'(958)e^+e^-)/\Gamma_{\text{total}}$ Γ_{342}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.59 \pm 0.07 \pm 0.17$	8.9k	1 ABLIKIM	19H BES3	$J/\psi \rightarrow \eta'(958)e^+e^-$

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

$5.81 \pm 0.16 \pm 0.31$	1.4k	1,2 ABLIKIM	14I BES3	$J/\psi \rightarrow \eta'(958)e^+e^-$
--------------------------	------	-------------	----------	---------------------------------------

¹ Using both $\eta' \rightarrow \gamma\pi^+\pi^-$ and $\eta' \rightarrow \pi^+\pi^-\eta$ decays.

² Superseded by ABLIKIM 19H.

 $\Gamma(X(1835)e^+e^-, X \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{343}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.58 \pm 0.19 \pm 0.16$	1364	1 ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'e^+e^-$

¹ Assuming constructive interference. Destructive interference gives a value of $(4.43 \pm 0.23 \pm 0.19) \times 10^{-6}$ for this branching fraction.

 $\Gamma(X(2120)e^+e^-, X \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{344}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.82 \pm 0.12 \pm 0.06$	310	ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'e^+e^-$

$\Gamma(X(2370)e^+e^-, X \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}$	Γ_{345}/Γ				
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	■
$1.08 \pm 0.14 \pm 0.10$	397	ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'e^+e^-$	

$\Gamma(\eta U \rightarrow \eta e^+e^-)/\Gamma_{\text{total}}$	Γ_{346}/Γ				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	■
$<9.11 \times 10^{-7}$	90	¹ ABLIKIM	19A BES3	$J/\psi \rightarrow \eta e^+e^-$	

¹ For a dark photon U with mass between 10 and 2400 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.9×10^{-8} to 91.1×10^{-8} .

$\Gamma(\eta'(958)U \rightarrow \eta'(958)e^+e^-)/\Gamma_{\text{total}}$	Γ_{347}/Γ				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	■
$<2.0 \times 10^{-7}$	90	¹ ABLIKIM	19H BES3	$J/\psi \rightarrow \eta'(958)e^+e^-$	

¹ For a dark photon U with mass between 100 and 2100 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.8×10^{-8} to 2.0×10^{-7} . The corresponding limits on the branching fraction $J/\psi \rightarrow \eta' U$ range from 5.7×10^{-8} to 7.4×10^{-7} .

$\Gamma(\phi e^+e^-)/\Gamma_{\text{total}}$	Γ_{348}/Γ				
VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT	■
<1.2	90	¹ ABLIKIM	19AB BES3	$J/\psi \rightarrow \phi e^+e^-$	

¹ Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ and $B(\psi(2S) \rightarrow \pi^+\pi^-J/\psi) = (34.49 \pm 0.30)\%$.

WEAK DECAYS

$\Gamma(D^-e^+\nu_e + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{349}/Γ				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	■
$<7.1 \times 10^{-8}$	90	ABLIKIM	21Q BES3	$e^+e^- \rightarrow J/\psi$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.2 \times 10^{-5}$ 90 ABLIKIM 06M BES2 $e^+e^- \rightarrow J/\psi$

$\Gamma(\bar{D}^0e^+e^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{350}/Γ				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	■
$<8.5 \times 10^{-8}$	90	¹ ABLIKIM	17AF BES3	$e^+e^- \rightarrow J/\psi$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.1 \times 10^{-5}$ 90 ABLIKIM 06M BES2 $e^+e^- \rightarrow J/\psi$

¹ Using D^0 decays to $K^-\pi^+$, $K^-\pi^+\pi^0$, and $K^-\pi^+\pi^+\pi^-$.

$\Gamma(D_s^-e^+\nu_e + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{351}/Γ				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	■
$<1.3 \times 10^{-6}$	90	ABLIKIM	14R BES3	$e^+e^- \rightarrow J/\psi$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<3.6 \times 10^{-5}$ 90 ¹ ABLIKIM 06M BES2 $e^+e^- \rightarrow J/\psi$

¹ Using $B(D_s^- \rightarrow \phi\pi^-) = 4.4 \pm 0.5\%$.

$\Gamma(D_s^{*-} e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$< 1.8 \times 10^{-6}$	90

$\Gamma(D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$< 7.5 \times 10^{-5}$	90

$\Gamma(\bar{D}^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$< 1.7 \times 10^{-4}$	90

$\Gamma(\bar{D}^0 \bar{K}^{*0} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$< 2.5 \times 10^{-6}$	90

$\Gamma(D_s^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$< 1.3 \times 10^{-4}$	90

$\Gamma(D_s^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$< 1.3 \times 10^{-5}$	90

— CHARGE CONJUGATION (C), PARITY (P), —
— LEPTON FAMILY NUMBER (LF) VIOLATING MODES —

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE	CL%
$< 2.7 \times 10^{-7}$	90
• • • We do not use the following data for averages, fits, limits, etc. • • •	
$< 0.5 \times 10^{-5}$	90
$< 1.6 \times 10^{-4}$	90
$< 2.2 \times 10^{-5}$	90
$< 50 \times 10^{-5}$	90

¹ WICHT 08 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [\mathcal{B}(B^+ \rightarrow J/\psi(1S) K^+)] < 0.16 \times 10^{-6}$ which we divide by our best value $\mathcal{B}(B^+ \rightarrow J/\psi(1S) K^+) = 1.020 \times 10^{-3}$.

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$

VALUE	CL%
$< 1.4 \times 10^{-6}$	90

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$

VALUE	CL%
$< 1.6 \times 10^{-7}$	90

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

VALUE	CL%
$< 1.1 \times 10^{-6}$	90

Γ_{352}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	14R	$e^+ e^- \rightarrow J/\psi$

Γ_{353}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	08J	$e^+ e^- \rightarrow J/\psi$

Γ_{354}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	08J	$e^+ e^- \rightarrow J/\psi$

Γ_{355}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	14K	$e^+ e^- \rightarrow J/\psi$

Γ_{356}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	08J	$e^+ e^- \rightarrow J/\psi$

Γ_{357}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	14K	$e^+ e^- \rightarrow J/\psi$

Γ_{358}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	14Q	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •		
ADAMS	08	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
¹ WICHT	08	$B^\pm \rightarrow K^\pm \gamma\gamma$
ABLIKIM	07J	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
BARTEL	77	$e^+ e^-$

Γ_{359}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	14Q	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

Γ_{360}/Γ

DOCUMENT ID	TECN	COMMENT
ABLIKIM	13L	$e^+ e^- \rightarrow J/\psi$

$\Gamma(e^\pm\tau^\mp)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{361}/Γ
$<7.5 \times 10^{-8}$	90	ABLIKIM	21M	BES3 $e^+e^- \rightarrow J/\psi$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<8.3 \times 10^{-6}$	90	¹ ABLIKIM	04	BES $e^+e^- \rightarrow J/\psi$	

¹ Superseded by ABLIKIM 21M. $\Gamma(\mu^\pm\tau^\mp)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{362}/Γ
$<2.0 \times 10^{-6}$	90	ABLIKIM	04	BES $e^+e^- \rightarrow J/\psi$	

 $\Gamma(\Lambda_c^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{363}/Γ
$<6.9 \times 10^{-8}$	90	ABLIKIM	19AF	BES3 $e^+e^- \rightarrow J/\psi \rightarrow pK^-\pi^+e^- (+ \text{ c.c.})$	

OTHER DECAYS

 $\Gamma(\text{invisible})/\Gamma(e^+e^-)$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{364}/Γ_5
$<6.6 \times 10^{-2}$	90	LEES	13I	BABR $B \rightarrow K^{(*)}J/\psi$	

 $\Gamma(\text{invisible})/\Gamma(\mu^+\mu^-)$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{364}/Γ_7
$<1.2 \times 10^{-2}$	90	ABLIKIM	08G	BES2 $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$	

J/ ψ (1S) REFERENCES

ZHU	23	PR D107 012006	W. Zhu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	22AI	PRL 129 192002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D107 079901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AY	PR D106 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22B	PRL 129 022002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22C	PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22H	PR D105 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	22	EPJ C82 938	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
ABLIKIM	21AM	PR D104 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AT	JHEP 2111 226	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21C	PR D103 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21M	PR D103 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21Q	JHEP 2106 157	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	21	PR D103 092001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	21C	PR D104 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
ABLIKIM	20	PR D101 012004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20K	PR D101 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20Q	EPJ C80 746	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	20	JHEP 2007 112	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
ABLIKIM	19A	PR D99 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D104 099901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AB	PR D99 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AC	PR D99 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AF	PR D99 072006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AN	PR D99 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)

ABLIKIM	19H	PR D99 012013	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19N	PR D99 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19Q	PL B791 375	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19T	PRL 122 142002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18AB	PR D98 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18D	PRL 121 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	18A	JHEP 1805 119	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	18E	PR D98 112015	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AH	PR D96 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	17D	PR D95 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	16E	PR D93 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16K	PR D93 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16M	PR D93 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16N	PR D93 112011	M. Ablikim	(BESIII Collab.)
ABLIKIM	16P	PR D94 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16Q	PL B761 98	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15K	PR D91 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	14I	PR D89 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14K	PR D89 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14N	PR D90 052009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14Q	PR D90 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14R	PR D90 112014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
AULCHENKO	14	PL B731 227	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13L	PR D87 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13P	PR D87 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	12	PR D85 092012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12C	PR D86 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12P	CP C36 1031	M. Ablikim <i>et al.</i>	(BES II Collab.)
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)
ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)

ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ANASHIN	10	PL B685 134	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DEL-AMO-SA...	10O	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
INSLER	10	PR D81 091101	J. Insler <i>et al.</i>	(CLEO Collab.)
ABLIKIM	09	PL B676 25	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	09B	PR D80 052004	M. Ablikim <i>et al.</i>	(BES II Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08	EPJ C53 15	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08A	PR D77 012001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08G	PRL 100 192001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08J	PL B663 297	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08O	PR D78 092005	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSON	08	PR D78 032012	D. Besson <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07J	PR D76 117101	M. Ablikim <i>et al.</i>	(BES Collab.)
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Fermilab E835 Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABLIKIM	06	PL B632 181	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06F	PR D73 052007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06K	PRL 97 062001	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	06M	PL B639 418	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	06A	PR D73 051103	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05B	PR D71 032003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05C	PL B610 192	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
LI	05C	PR D71 111103	Z. Li <i>et al.</i>	(CLEO Collab.)
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer	
ABLIKIM	04	PL B598 172	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	04	PR D69 011103	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	04	PL B578 16	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04A	PR D69 012003	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04E	PL B591 42	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04H	PR D70 012005	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
SETH	04	PR D69 097503	K.K. Seth	
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03D	PL B561 49	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)

BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00B	PL B472 200	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98G	PL B424 213	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALDINI	98	PL B444 111	R. Baldini <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	96	PR D54 7067	T.A. Armstrong <i>et al.</i>	(E760 Collab.)
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96D	PR D54 1221	J.Z. Bai <i>et al.</i>	(BES Collab.)
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 and E706 Collab.)
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)
BAI	95B	PL B355 374	J.Z. Bai <i>et al.</i>	(BES Collab.)
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ANTONELLI	93	PL B301 317	A. Antonelli <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)
COFFMAN	92	PRL 68 282	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
HSUEH	92	PR D45 2181	S. Hsueh, S. Palestini	(FNAL, TORI)
BISELLLO	91	NP B350 1	D. Bisello <i>et al.</i>	(DM2 Collab.)
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
BISELLLO	90	PL B241 617	D. Bisello <i>et al.</i>	(DM2 Collab.)
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
JOUSSET	90	PR D41 1389	J. Jousset <i>et al.</i>	(DM2 Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)
BISELLLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
COFFMAN	88	PR D38 2695	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)
BAGLIN	87	NP B286 592	C. Baglin <i>et al.</i>	(LAPP, CERN, GENO, LYON+)
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)
BISELLLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
HENRARD	87	NP B292 670	P. Henrard <i>et al.</i>	(CLER, FRAS, LALO+)
PALLIN	87	NP B292 653	D. Pallin <i>et al.</i>	(CLER, FRAS, LALO, PADO)
BALTRUSAIT...	86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)
BISELLLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
BALTRUSAIT...	85C	PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
BALTRUSAIT...	85D	PR D32 566	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
		Translated from YAF 41 733.		
BALTRUSAIT...	84	PRL 52 2126	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
EATON	84	PR D29 804	M.W. Eaton <i>et al.</i>	(LBL, SLAC)
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
EDWARDS	83B	PRL 51 859	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)
EDWARDS	82B	PR D25 3065	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
Also		ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
BESCH	81	ZPHY C8 1	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)

GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)
PARTRIDGE	80	PRL 44 712	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
		Translated from YAF 34 1471.		
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
ALEXANDER	78	PL 72B 493	G. Alexander <i>et al.</i>	(DESY, HAMB, SIEG+)
BESCH	78	PL 78B 347	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)
BRANDELIK	78B	PL 74B 292	R. Brandelik <i>et al.</i>	(DASP Collab.)
PERUZZI	78	PR D17 2901	I. Peruzzi <i>et al.</i>	(SLAC, LBL)
BARTEL	77	PL 66B 489	W. Bartel <i>et al.</i>	(DESY, HEIDP)
BURMESTER	77D	PL 72B 135	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
VANNUCCI	77	PR D15 1814	F. Vannucci <i>et al.</i>	(SLAC, LBL)
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)
BRAUNSCH...	76	PL 63B 487	W. Braunschweig <i>et al.</i>	(DASP Collab.)
JEAN-MARIE	76	PRL 36 291	B. Jean-Marie <i>et al.</i>	(SLAC, LBL) IG
BALDINI...	75	PL 58B 471	R. Baldini-Celio <i>et al.</i>	(FRAS, ROMA)
BOYARSKI	75	PRL 34 1357	A.M. Boyarski <i>et al.</i>	(SLAC, LBL) JPC
DASP	75	PL 56B 491	W. Braunschweig <i>et al.</i>	(DASP Collab.)
ESPOSITO	75B	LNC 14 73	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
FORD	75	PRL 34 604	R.L. Ford <i>et al.</i>	(SLAC, PENN)