



$$I(J^P) = \frac{1}{2}(0^-)$$

K_S^0 MEAN LIFE

For earlier measurements, beginning with BOLDT 58B, see our 1986 edition, Physics Letters **170B** 130 (1986).

OUR FIT is described in the note on “*CP* violation in K_L decays” in the K_L^0 Particle Listings. The result labeled “OUR FIT Assuming *CPT*” [“OUR FIT Not assuming *CPT*”] includes all measurements except those with the comment “Not assuming *CPT*” [“Assuming *CPT*”]. Measurements with neither comment do not assume *CPT* and enter both fits.

VALUE (10^{-10} s)	EVTS	DOCUMENT ID	TECN	COMMENT
0.8954 ± 0.0004 OUR FIT		Error includes scale factor of 1.1. Assuming <i>CPT</i>		
0.89564 ± 0.00033 OUR FIT		Not assuming <i>CPT</i>		
0.89589 ± 0.00070		1,2 ABOUZAID	11 KTEV	Not assuming <i>CPT</i>
0.89623 ± 0.00047		1,3 ABOUZAID	11 KTEV	Assuming <i>CPT</i>
0.89562 ± 0.00029 ± 0.00043	20M	4 AMBROSINO	11 KLOE	Not assuming <i>CPT</i>
0.89598 ± 0.00048 ± 0.00051	16M	LAI	02C NA48	
0.8971 ± 0.0021		BERTANZA	97 NA31	
0.8941 ± 0.0014 ± 0.0009		SCHWINGEN...95	E773	Assuming <i>CPT</i>
0.8929 ± 0.0016		GIBBONS	93 E731	Assuming <i>CPT</i>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.8965 ± 0.0007		5 ALAVI-HARATI03	KTEV	Assuming <i>CPT</i>
0.8958 ± 0.0013		6 ALAVI-HARATI03	KTEV	Not assuming <i>CPT</i>
0.8920 ± 0.0044	214k	GROSSMAN	87 SPEC	
0.905 ± 0.007		7 ARONSON	82B SPEC	
0.881 ± 0.009	26k	ARONSON	76 SPEC	
0.8926 ± 0.0032 ± 0.0002		8 CARITHERS	75 SPEC	
0.8937 ± 0.0048	6M	GEWENIGER	74B ASPK	
0.8958 ± 0.0045	50k	9 SKJEGGEST...	72 HBC	
0.856 ± 0.008	19994	10 DONALD	68B HBC	
0.872 ± 0.009	20000	9,10 HILL	68 DBC	

¹ The two ABOUZAID 11 values use the same full KTeV dataset from 1996, 1997, and 1999. The first enters the “assuming *CPT*” fit and the second enters the “not assuming *CPT*” fit.

² ABOUZAID 11 fit has Δm , τ_s , ϕ_ϵ , $\text{Re}(\epsilon'/\epsilon)$, and $\text{Im}(\epsilon'/\epsilon)$ as free parameters. See $\text{Im}(\epsilon'/\epsilon)$ in the “ K_L^0 CP violation” section for correlation information.

³ ABOUZAID 11 fit has Δm and τ_s free but constrains ϕ_ϵ to the Superweak value, i.e. assumes *CPT*. This τ_s value is correlated with their $\Delta m = m_{K_L^0} - m_{K_S^0}$ measurement in the K_L^0 listings. The correlation coefficient $\rho(\tau_s, \Delta m) = -0.670$.

⁴ Fit to the proper time distribution.

⁵ This ALAVI-HARATI 03 fit has Δm and τ_s free but constrains ϕ_{+-} to the Superweak value, i.e. assumes *CPT*. This τ_s value is correlated with their $\Delta m = m_{K_L^0} - m_{K_S^0}$ measurement in the K_L^0 listings. The correlation coefficient $\rho(\tau_s, \Delta m) = -0.396$. Superseded by ABOUZAID 11.

- ⁶This ALAVI-HARATI 03 fit has Δm , ϕ_{+-} , and τ_{K_S} free. See ϕ_{+-} in the “ K_L CP violation” section for correlation information. Superseded by ABOUZAID 11.
- ⁷ARONSON 82 find that K_S^0 mean life may depend on the kaon energy.
- ⁸CARITHERS 75 measures the Δm dependence of the total decay rate (inverse mean life) to be $\Gamma(K_S^0) = [(1.122 \pm 0.004) + 0.16(\Delta m - 0.5348)/\Delta m] 10^{10} / \text{s}$, or, in terms of mean life, CARITHERS 75 measures $\tau_s = (0.8913 \pm 0.0032) - 0.238 [\Delta m - 0.5348] (10^{-10} \text{ s})$. We have adjusted the measurement to use our best values of ($\Delta m = 0.5293 \pm 0.0009$) (10^{10} s^{-1}). Our first error is their experiment's error and our second error is the systematic error from using our best values.
- ⁹HILL 68 has been changed by the authors from the published value (0.865 ± 0.009) because of a correction in the shift due to η_{+-} . SKJEGGESTAD 72 and HILL 68 give detailed discussions of systematics encountered in this type of experiment.
- ¹⁰Pre-1971 experiments are excluded from the average because of disagreement with later more precise experiments.

K_S^0 DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Hadronic modes		
Γ_1 $\pi^0 \pi^0$	(30.69 ± 0.05) %	
Γ_2 $\pi^+ \pi^-$	(69.20 ± 0.05) %	
Γ_3 $\pi^+ \pi^- \pi^0$	($3.5 \begin{array}{l} +1.1 \\ -0.9 \end{array} \right) \times 10^{-7}$	
Modes with photons or $\ell\bar{\ell}$ pairs		
Γ_4 $\pi^+ \pi^- \gamma$	[a,b] (1.79 ± 0.05) $\times 10^{-3}$	
Γ_5 $\pi^+ \pi^- e^+ e^-$	(4.79 ± 0.15) $\times 10^{-5}$	
Γ_6 $\pi^0 \gamma \gamma$	[a] (4.9 ± 1.8) $\times 10^{-8}$	
Γ_7 $\gamma \gamma$	(2.63 ± 0.17) $\times 10^{-6}$	S=3.1
Semileptonic modes		
Γ_8 $\pi^\pm e^\mp \nu_e$	[c] (7.04 ± 0.08) $\times 10^{-4}$	
Γ_9 $\pi^\pm \mu^\mp \nu_\mu$	[c,d] (4.56 ± 0.20) $\times 10^{-4}$	
CP violating (CP) and $\Delta S = 1$ weak neutral current (S1) modes		
Γ_{10} $3\pi^0$	CP	< 2.6×10^{-8} CL=90%
Γ_{11} $\mu^+ \mu^-$	S1	< 2.1×10^{-10} CL=90%
Γ_{12} $e^+ e^-$	S1	< 9×10^{-9} CL=90%
Γ_{13} $\pi^0 e^+ e^-$	S1	[a] ($3.0 \begin{array}{l} +1.5 \\ -1.2 \end{array} \right) \times 10^{-9}$
Γ_{14} $\pi^0 \mu^+ \mu^-$	S1	($2.9 \begin{array}{l} +1.5 \\ -1.2 \end{array} \right) \times 10^{-9}$

[a] See the Particle Listings below for the energy limits used in this measurement.

[b] Most of this radiative mode, the low-momentum γ part, is also included in the parent mode listed without γ 's.

- [c] The value is for the sum of the charge states or particle/antiparticle states indicated.
 [d] Not a measurement. Calculated as $0.666 \cdot B(\pi^\pm e^\mp \nu_e)$.
-

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 5 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 0.1$ for 2 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-100		
x_8	-6	4	
x_9	-1	-3	0
	x_1	x_2	x_8

K_S^0 DECAY RATES

$\Gamma(\pi^\pm e^\mp \nu_e)$

Γ_8

VALUE (10^6 s^{-1})	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
8.1 ± 1.6	75	¹ AKHMETSHIN 99	CMD2	Tagged K_S^0 using $\phi \rightarrow K_L^0 K_S^0$
7.50 ± 0.08		² PDG	98	
seen		BURGUN	72	HBC $K^+ p \rightarrow K^0 p \pi^+$
9.3 ± 2.5		AUBERT	65	HLBC $\Delta S = \Delta Q$, CP cons. not assumed

¹ AKHMETSHIN 99 is from a measured branching ratio $B(K_S^0 \rightarrow \pi e \nu_e) = (7.2 \pm 1.4) \times 10^{-4}$ and $\tau_{K_S^0} = (0.8934 \pm 0.0008) \times 10^{-10} \text{ s}$. Not independent of measured branching ratio.

² PDG 98 from K_L^0 measurements, assuming that $\Delta S = \Delta Q$ in K^0 decay so that $\Gamma(K_S^0 \rightarrow \pi^\pm e^\mp \nu_e) = \Gamma(K_L^0 \rightarrow \pi^\pm e^\mp \nu_e)$.

$\Gamma(\pi^\pm \mu^\mp \nu_\mu)$

Γ_9

VALUE (10^6 s^{-1})	DOCUMENT ID
---------------------------------	-------------

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

5.25 ± 0.07	¹ PDG	98
-----------------	------------------	----

¹ PDG 98 from K_L^0 measurements, assuming that $\Delta S = \Delta Q$ in K^0 decay so that $\Gamma(K_S^0 \rightarrow \pi^\pm \mu^\mp \nu_\mu) = \Gamma(K_L^0 \rightarrow \pi^\pm \mu^\mp \nu_\mu)$.

K_S^0 BRANCHING RATIOS**Hadronic modes** **$\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
0.3069 ± 0.0005 OUR FIT			

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.335 ± 0.014	1066	BROWN	63	HLBC
0.288 ± 0.021	198	CHRETIEN	63	HLBC
0.30 ± 0.035		BROWN	61	HLBC

 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.6920 ± 0.0005 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.670 ± 0.010	3447	DOYLE	69	HBC $\pi^- p \rightarrow \Lambda K^0$
-------------------	------	-------	----	---------------------------------------

 $\Gamma(\pi^+\pi^-)/\Gamma(\pi^0\pi^0)$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.255 ± 0.005 OUR FIT				

 2.2549 ± 0.0054

¹ AMBROSINO 06c KLOE

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.2555 $\pm 0.0012 \pm 0.0054$		2 AMBROSINO	06c	KLOE
2.236 $\pm 0.003 \pm 0.015$	766k	² ALOISIO	02B	KLOE
2.11 ± 0.09	1315	EVERHART	76	WIRE $\pi^- p \rightarrow \Lambda K^0$
2.169 ± 0.094	16k	COWELL	74	OSPK $\pi^- p \rightarrow \Lambda K^0$
2.16 ± 0.08	4799	HILL	73	DBC $K^+ d \rightarrow K^0 pp$
2.22 ± 0.10	3068	³ ALITTI	72	HBC $K^+ p \rightarrow \pi^+ p K^0$
2.22 ± 0.08	6380	MORSE	72B	DBC $K^+ n \rightarrow K^0 p$
2.10 ± 0.11	701	⁴ NAGY	72	HLBC $K^+ n \rightarrow K^0 p$
2.22 ± 0.095	6150	⁵ BALTAY	71	HBC $K p \rightarrow K^0$ neutrals
2.282 ± 0.043	7944	⁶ MOFFETT	70	OSPK $K^+ n \rightarrow K^0 p$
2.12 ± 0.17	267	⁴ BOZOKI	69	HLBC
2.285 ± 0.055	3016	⁶ GOBBI	69	OSPK $K^+ n \rightarrow K^0 p$
2.10 ± 0.06	3700	MORFIN	69	HLBC $K^+ n \rightarrow K^0 p$

¹ This result combines AMBROSINO 06c KLOE 2001-02 data with ALOISIO 02B KLOE 2000 data. $K_S^0 \rightarrow \pi^+\pi^-$ fully inclusive.

² Includes radiative decays $\pi^+\pi^-\gamma$.

³ The directly measured quantity is $K_S^0 \rightarrow \pi^+\pi^-/\text{all } K^0 = 0.345 \pm 0.005$.

⁴ NAGY 72 is a final result which includes BOZOKI 69.

⁵ The directly measured quantity is $K_S^0 \rightarrow \pi^+\pi^-/\text{all } \bar{K}^0 = 0.345 \pm 0.005$.

⁶ MOFFETT 70 is a final result which includes GOBBI 69.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$				Γ_3/Γ
<u>VALUE (units 10^{-7})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.5^{+1.1}_{-0.9}$ OUR AVERAGE				
$4.7^{+2.2+1.7}_{-1.7-1.5}$		¹ BATLEY	05	NA48
$2.5^{+1.3+0.5}_{-1.0-0.6}$	500k	² ADLER	97B	CPLR
$4.8^{+2.2}_{-1.6}\pm 1.1$		³ ZOU	96	E621
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$4.1^{+2.5+0.5}_{-1.9-0.6}$		⁴ ADLER	96E	CPLR Sup. by ADLER 97B
$3.9^{+5.4+0.9}_{-1.8-0.7}$		⁵ THOMSON	94	E621 Sup. by ZOU 96

¹ BATLEY 05 is obtained by measuring the interference parameters in K_S , $K_L \rightarrow \pi^+\pi^-\pi^0$: $\text{Re}(\lambda) = 0.038 \pm 0.008 \pm 0.006$ and $\text{Im}(\lambda) = -0.013 \pm 0.005 \pm 0.004$; the correlation coeff. between $\text{Re}(\lambda)$ and $\text{Im}(\lambda)$ is 0.66 (statistical only).

² ADLER 97B find the CP -conserving parameters $\text{Re}(\lambda) = (28 \pm 7 \pm 3) \times 10^{-3}$, $\text{Im}(\lambda) = (-10 \pm 8 \pm 2) \times 10^{-3}$. They estimate $B(K_S^0 \rightarrow \pi^+\pi^-\pi^0)$ from $\text{Re}(\lambda)$ and the K_L^0 decay parameters. See also ANGELOPOULOS 98C.

³ ZOU 96 is from the measured quantities $|\rho_{+-0}| = 0.039^{+0.009}_{-0.006} \pm 0.005$ and $\phi_\rho = (-9 \pm 18)^\circ$.

⁴ ADLER 96E is from the measured quantities $\text{Re}(\lambda) = 0.036 \pm 0.010^{+0.002}_{-0.003}$ and $\text{Im}(\lambda)$ consistent with zero. Note that the quantity λ is the same as ρ_{+-0} used in other footnotes.

⁵ THOMSON 94 calculates this branching ratio from their measurements $|\rho_{+-0}| = 0.035^{+0.019}_{-0.011} \pm 0.004$ and $\phi_\rho = (-59 \pm 48)^\circ$ where $|\rho_{+-0}| e^{i\phi_\rho} = A(K_S^0 \rightarrow \pi^+\pi^-\pi^0, I=2)/A(K_L^0 \rightarrow \pi^+\pi^-\pi^0)$.

— Modes with photons or $\ell\bar{\ell}$ pairs —

$\Gamma(\pi^+\pi^-\gamma)/\Gamma(\pi^+\pi^-)$				Γ_4/Γ_2
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.59 ± 0.08 OUR AVERAGE				
2.56 ± 0.09	1286	RAMBERG	93	E731 $p_\gamma > 50 \text{ MeV}/c$
2.68 ± 0.15		¹ TAUREG	76	SPEC $p_\gamma > 50 \text{ MeV}/c$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
7.10 ± 0.22	3723	RAMBERG	93	E731 $p_\gamma > 20 \text{ MeV}/c$
3.0 ± 0.6	29	² BOBISUT	74	HLBC $p_\gamma > 40 \text{ MeV}/c$
2.8 ± 0.6		³ BURGUN	73	HBC $p_\gamma > 50 \text{ MeV}/c$

¹ TAUREG 76 find direct emission contribution < 0.06 , CL = 90%.

² BOBISUT 74 not included in average because p_γ cut differs. Estimates direct emission contribution to be 0.5 or less, CL = 95%.

³ BURGUN 73 estimates that direct emission contribution is 0.3 ± 0.6 .

$\Gamma(\pi^+\pi^-e^+e^-)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
4.79±0.15 OUR AVERAGE				
4.83±0.11±0.14	23k	¹ BATLEY	11	NA48 2002 data
4.69±0.30	676	² LAI	03C	NA48 1998+1999 data
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.71±0.23±0.22	620	^{2,3} LAI	03C	NA48 1999 data
4.5 ± 0.7 ± 0.4	56	LAI	00B	NA48 1998 data

¹ BATLEY 11 reports $[\Gamma(K_S^0 \rightarrow \pi^+\pi^-e^+e^-)/\Gamma_{\text{total}}] / [B(K_L^0 \rightarrow \pi^+\pi^-\pi^0)] / [B(\pi^0 \rightarrow e^+e^-\gamma)] = (3.28 \pm 0.06 \pm 0.04) \times 10^{-2}$ which we multiply by our best values $B(K_L^0 \rightarrow \pi^+\pi^-\pi^0) = (12.54 \pm 0.05) \times 10^{-2}$, $B(\pi^0 \rightarrow e^+e^-\gamma) = (1.174 \pm 0.035) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values. Also a limit on the absolute value of the interference between bremsstrahlung and E1 transition is given : $< 4 \times 10^{-7}$ at 90% C.L.

² Uses normalization $BR(K_L \rightarrow \pi^+\pi^-\pi^0)*BR(\pi^0 \rightarrow e^+e^-) = (1.505 \pm 0.047) \times 10^{-3}$ from our 2000 Edition.

³ Second error is 0.16(syst)±0.15(norm) combined in quadrature.

$\Gamma(\pi^0\gamma\gamma)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-8})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
4.9±1.6±0.9	17	¹ LAI	04	NA48	$m_{\gamma\gamma}^2/m_K^2 > 0.2$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<33	90	LAI	03B	NA48	$m_{\gamma\gamma}^2/m_K^2 > 0.2$

¹ Spectrum also measured and found consistent with the one generated by a constant matrix element.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.63 ±0.17 OUR AVERAGE			Error includes scale factor of 3.1.		
2.26 ± 0.12 ± 0.06	711	¹ AMBROSINO	08C	KLOE	$\phi \rightarrow K_S^0 K_L^0$
2.713±0.063±0.005	7.5k	² LAI	03	NA48	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.58 ± 0.36 ± 0.22	149	LAI	00	NA48	
2.2 ± 1.1	16	³ BARR	95B	NA31	
2.4 ± 0.9	35	⁴ BARR	95B	NA31	
< 13	90	BALATS	89	SPEC	
2.4 ± 1.2	19	BURKHARDT	87	NA31	
<133	90	BARMIN	86B	XEBC	

¹ AMBROSINO 08C reports $(2.26 \pm 0.12 \pm 0.06) \times 10^{-6}$ from a measurement of $[\Gamma(K_S^0 \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(K_S^0 \rightarrow \pi^0\pi^0)]$ assuming $B(K_S^0 \rightarrow \pi^0\pi^0) = (30.69 \pm 0.05) \times 10^{-2}$.

² LAI 03 reports $[\Gamma(K_S^0 \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] / [B(K_S^0 \rightarrow \pi^0\pi^0)] = (8.84 \pm 0.18 \pm 0.10) \times 10^{-6}$ which we multiply by our best value $B(K_S^0 \rightarrow \pi^0\pi^0) = (30.69 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ BARR 95B result is calculated using $B(K_L \rightarrow \gamma\gamma) = (5.86 \pm 0.17) \times 10^{-4}$.

⁴ BARR 95B quotes this as the combined BARR 95B + BURKHARDT 87 result after rescaling BURKHARDT 87 to use same branching ratios and lifetimes as BARR 95B.

Semileptonic modes $\Gamma(\pi^\pm e^\mp \nu_e)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

7.04 ±0.08 OUR FIT**7.04 ±0.08 OUR AVERAGE**

7.046 ± 0.18 ± 0.16	¹ BATLEY	07D NA48	$K^0(\bar{K}^0)(t) \rightarrow \pi e \nu$
6.91 ± 0.34 ± 0.15	624 ² ALOISIO	02 KLOE	Tagged K_S^0 using $\phi \rightarrow K_L^0 K_S^0$

• • • We use the following data for averages but not for fits. • • •

7.05 ± 0.09	13k ³ AMBROSINO	06E KLOE	Not fitted
-------------	----------------------------	----------	------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.2 ± 1.4	75 AKHMETSHIN 99	CMD2	Tagged K_S^0 using $\phi \rightarrow K_L^0 K_S^0$
-----------	------------------	------	---

¹ Reconstructed from $K^0(\bar{K}^0)(t) \rightarrow \pi e \nu$ distributions using PDG values of $B(K_L^0 \rightarrow \pi e \nu) = 0.4053 \pm 0.0015$, $\tau_L = (5.114 \pm 0.021) \times 10^{-8}$ s and $\tau_S = (0.8958 \pm 0.0005) \times 10^{-10}$ s.

² Uses the PDG 00 value for $B(K_S^0 \rightarrow \pi^+ \pi^-)$.

³ Obtained by imposing $\sum_i B(K_S^0 \rightarrow i) = 1$, where i runs over all the four branching ratios $\pi^+ \pi^-$, $\pi^0 \pi^0$, $\pi e \nu$, and $\pi \mu \nu$. Input value of $B(K_S^0 \rightarrow \pi^+ \pi^-) / B(K_S^0 \rightarrow \pi^0 \pi^0)$ from AMBROSINO 06C is used. To derive $\Gamma(K_S^0 \rightarrow \pi^+ \mu \nu) / \Gamma(K_S^0 \rightarrow \pi^+ e \nu)$, lepton universality is assumed, radiative corrections from ANDRE 07 are used, and phase space integrals are taken from KTeV, ALEXOPOULOS 04A. This branching fraction enters our fit via their $\Gamma(\pi^\pm e^\mp \nu_e) / \Gamma(\pi^+ \pi^-)$ branching ratio measurement.

 $\Gamma(\pi^\pm \mu^\mp \nu_\mu)/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

4.56±0.20 OUR FIT**4.56±0.11±0.17**

7223	¹ BABUSCI	20 KLOE	direct measurement
------	----------------------	---------	--------------------

¹ Value obtained by normalizing to the KLOE measurement $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.196 \pm 0.051)\%$. Also comparison with the PDG 18 based derived value leads to a lepton flavor universality test $|V_{us} f_+(0)|_{K_S^0 \rightarrow \pi \mu \nu}^2 / |V_{us} f_+(0)|_{K_S^0 \rightarrow \pi e \nu}^2 = 0.975 \pm 0.044$.

 $\Gamma(\pi^\pm e^\mp \nu_e)/\Gamma(\pi^+ \pi^-)$ Γ_8/Γ_2

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
---	-------------	--------------------	-------------

10.18±0.12 OUR FIT**10.19±0.11±0.07**

13k	AMBROSINO 06E	KLOE
-----	---------------	------

— CP violating (CP) and $\Delta S = 1$ weak neutral current (S1) modes — $\Gamma(3\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

Violates CP conservation.

<u>VALUE (units 10^{-7})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	-------------	--------------------	-------------	----------------

< 0.26	90	590M	¹ BABUSCI	13C KLOE	$\phi \rightarrow K_L^0 K_S^0$
--------	----	------	----------------------	----------	--------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.2	90	37.8M	AMBROSINO 05B	KLOE
-------	----	-------	---------------	------

< 7.4	90	4.9M	² LAI	05A NA48
-------	----	------	------------------	----------

<140	90	7M	ACHASOV	99D SND
------	----	----	---------	---------

<190	90	17300	³ ANGELOPO...	98B	CPLR
<370	90		BARMIN	83	HLBC

¹ BABUSCI 13C uses 1.7 fb^{-1} of data of $\phi \rightarrow K_L^0 K_S^0$ decays with K_L^0 interaction in the calorimeter, collected from 2004 to 2005. No candidate events were found in the data with an expected background of $0.04^{+0.15}_{-0.03}$ events. Upper limit is obtained by normalizing to $K_S^0 \rightarrow 2\pi^0$ decays.

² LAI 05A value is obtained from their bound on $|\eta_{000}|$ (not assuming *CPT*) and $B(K_L^0 \rightarrow 3\pi^0) = 0.211 \pm 0.003$, and PDG 04 values for K_L^0 and K_S^0 lifetimes. If *CPT* is assumed then $B(K_S^0 \rightarrow 3\pi^0)_{CPT} < 2.3 \times 10^{-7}$ at 90% CL

³ ANGELOPOULOS 98B is from $\text{Im}(\eta_{000}) = -0.05 \pm 0.12 \pm 0.05$, assuming $\text{Re}(\eta_{000}) = \text{Re}(\epsilon) = 1.635 \times 10^{-3}$ and using the value $B(K_L^0 \rightarrow \pi^0 \pi^0 \pi^0) = 0.2112 \pm 0.0027$.

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$

Γ_{11}/Γ

Test for $\Delta S = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE	CL%	DOCUMENT ID	TECN
<2.1 × 10⁻¹⁰	90	¹ AAIJ	20AE LHCb

• • • We do not use the following data for averages, fits, limits, etc. • • •

<8 × 10 ⁻¹⁰	90	² AAIJ	17BQ LHCb
<9 × 10 ⁻⁹	90	³ AAIJ	13G LHCb
<3.2 × 10 ⁻⁷	90	GJESDAL	73 ASPK
<7 × 10 ⁻⁶	90	HYAMS	69B OSPK

¹ AAIJ 20AE uses 8.6 fb^{-1} of LHCb data from 2011 to 2012 and 2016 to 2018. The result utilizes the normalization mode branching fraction $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05) \times 10^{-2}$ from PDG 18. Supersedes AAIJ 17BQ.

² AAIJ 17BQ uses 3.0 fb^{-1} of $p p$ collisions at $\sqrt{s} = 7$ and 8 TeV. The result utilizes the normalization mode branching fraction $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05) \times 10^{-2}$ from PDG 16. Supersedes AAIJ 13G.

³ AAIJ 13G uses 1.0 fb^{-1} of $p p$ collisions at $\sqrt{s} = 7$ TeV. They obtained $B(K_S^0 \rightarrow \mu^+ \mu^-) < 11 \times 10^{-9}$ at 95% C.L.

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

Γ_{12}/Γ

Test for $\Delta S = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.09	90	¹ AMBROSINO 09A	KLOE	$e^+ e^- \rightarrow \phi \rightarrow K_S^0 K_L^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.4	90	ANGELOPO...	97	CPLR
< 28	90	BLICK	94	CNTR Hyperon facility
<100	90	BARMIN	86	XEBC

¹ AMBROSINO 09A reports $< 0.09 \times 10^{-7}$ from a measurement of $[\Gamma(K_S^0 \rightarrow e^+ e^-)/\Gamma_{\text{total}}] / [B(K_S^0 \rightarrow \pi^+ \pi^-)]$ assuming $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05) \times 10^{-2}$.

$\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$

Γ_{13}/Γ

Test for $\Delta S = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units 10^{-9})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$3.0^{+1.5}_{-1.2} \pm 0.2$		7	¹ BATLEY	03	NA48 $m_{ee} > 0.165$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 140	90	LAI	01	NA48
< 1100	90	BARR	93B	NA31
< 45000	90	GIBBONS	88	E731

¹ BATLEY 03 extrapolate also to the full kinematical region using a constant form factor and a vector matrix element. The resulting branching ratio is $(5.8^{+2.9}_{-2.4}) \times 10^{-9}$.

$\Gamma(\pi^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$

Γ_{14}/Γ

Test for $\Delta S = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units 10^{-9})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.9^{+1.5}_{-1.2} \pm 0.2$	6	¹ BATLEY	04A	NA48/1 K_S^0 beam

¹ Background estimate is $0.22^{+0.18}_{-0.11}$ events. Branching ratio assumes a vector matrix element and unit form factor.

K_S^0 FORM FACTORS

For discussion, see note on $K_{\ell 3}$ form factors in the K^\pm section of the Particle Listings above. Because the semileptonic branching fraction is smaller in K_S^0 than K_L^0 by the ratio of the mean lives, the K_S^0 semileptonic form factor has so far been measured only in the K_{e3} mode using the linear expansion $f_+(t) = f_+(0) (1 + \lambda_+ t / m_{\pi^+}^2)$, which gives the vector form factor $f_+(t)$ relative to its value at $t = 0$.

λ_+ (LINEAR ENERGY DEPENDENCE OF f_+ IN K_{e3}^0 DECAY)

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN
3.39 ± 0.41	15k	AMBROSINO	06E KLOE

CP VIOLATION IN $K_S \rightarrow 3\pi$

Written 1996 by T. Nakada (Paul Scherrer Institute) and L. Wolfenstein (Carnegie-Mellon University).

The possible final states for the decay $K^0 \rightarrow \pi^+ \pi^- \pi^0$ have isospin $I = 0, 1, 2$, and 3 . The $I = 0$ and $I = 2$ states have $CP = +1$ and K_S can decay into them without violating CP symmetry, but they are expected to be strongly suppressed by centrifugal barrier effects. The $I = 1$ and $I = 3$ states, which

have no centrifugal barrier, have $CP = -1$ so that the K_S decay to these requires CP violation.

In order to see CP violation in $K_S \rightarrow \pi^+ \pi^- \pi^0$, it is necessary to observe the interference between K_S and K_L decay, which determines the amplitude ratio

$$\eta_{+-0} = \frac{A(K_S \rightarrow \pi^+ \pi^- \pi^0)}{A(K_L \rightarrow \pi^+ \pi^- \pi^0)} . \quad (1)$$

If η_{+-0} is obtained from an integration over the whole Dalitz plot, there is no contribution from the $I = 0$ and $I = 2$ final states and a nonzero value of η_{+-0} is entirely due to CP violation.

Only $I = 1$ and $I = 3$ states, which are $CP = -1$, are allowed for $K^0 \rightarrow \pi^0 \pi^0 \pi^0$ decays and the decay of K_S into $3\pi^0$ is an unambiguous sign of CP violation. Similarly to η_{+-0} , η_{000} is defined as

$$\eta_{000} = \frac{A(K_S \rightarrow \pi^0 \pi^0 \pi^0)}{A(K_L \rightarrow \pi^0 \pi^0 \pi^0)} . \quad (2)$$

If one assumes that CPT invariance holds and that there are no transitions to $I = 3$ (or to nonsymmetric $I = 1$ states), it can be shown that

$$\begin{aligned} \eta_{+-0} &= \eta_{000} \\ &= \epsilon + i \frac{\text{Im } a_1}{\text{Re } a_1} . \end{aligned} \quad (3)$$

With the Wu-Yang phase convention, a_1 is the weak decay amplitude for K^0 into $I = 1$ final states; ϵ is determined from CP violation in $K_L \rightarrow 2\pi$ decays. The real parts of η_{+-0} and η_{000} are equal to $\text{Re}(\epsilon)$. Since currently-known upper limits on $|\eta_{+-0}|$ and $|\eta_{000}|$ are much larger than $|\epsilon|$, they can be interpreted as upper limits on $\text{Im}(\eta_{+-0})$ and $\text{Im}(\eta_{000})$ and so as limits on the CP -violating phase of the decay amplitude a_1 .

CP-VIOLATION PARAMETERS IN K_S^0 DECAY

$$A_S = [\Gamma(K_S^0 \rightarrow \pi^- e^+ \nu_e) - \Gamma(K_S^0 \rightarrow \pi^+ e^- \bar{\nu}_e)] / \text{SUM}$$

Such asymmetry violates CP. If CPT is assumed then $A_S = 2 \operatorname{Re}(\epsilon)$.

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$-3.8 \pm 5.0 \pm 2.6$	83k	¹ ANASTASI	18A KLOE

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5 \pm 9.6 \pm 2.9	13k	AMBROSINO	06E KLOE
-------------------------	-----	-----------	----------

¹ ANASTASI 18A result is a combination of the new measurement and AMBROSINO 06E. The new ANASTASI 18A measurement using data collected from 2004–2005, which corresponds to an integrated luminosity of 1.63 fb^{-1} is $A_S = (-4.9 \pm 5.7 \pm 2.6) \times 10^{-3}$.

PARAMETERS FOR $K_S^0 \rightarrow 3\pi$ DECAY

$$\operatorname{Im}(\eta_{+-0})^2 = \Gamma(K_S^0 \rightarrow \pi^+ \pi^- \pi^0, \text{CP-violating}) / \Gamma(K_L^0 \rightarrow \pi^+ \pi^- \pi^0)$$

CPT assumed valid (i.e. $\operatorname{Re}(\eta_{+-0}) \simeq 0$).

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
--------------	------------	-------------	--------------------	-------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.23	90	601	¹ BARMIN	85 HLBC
<0.12	90	384	METCALF	72 ASPK

¹ BARMIN 85 find $\operatorname{Re}(\eta_{+-0}) = (0.05 \pm 0.17)$ and $\operatorname{Im}(\eta_{+-0}) = (0.15 \pm 0.33)$. Includes events of BALDO-CEOLIN 75.

$$\operatorname{Im}(\eta_{+-0}) = \operatorname{Im}(A(K_S^0 \rightarrow \pi^+ \pi^- \pi^0, \text{CP-violating}) / A(K_L^0 \rightarrow \pi^+ \pi^- \pi^0))$$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.002 \pm 0.009^{+0.002}_{-0.001}$	500k	¹ ADLER	97B CPLR	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.002 \pm 0.018 \pm 0.003$	137k	² ADLER	96D CPLR	Sup. by ADLER 97B
$-0.015 \pm 0.017 \pm 0.025$	272k	³ ZOU	94 SPEC	

¹ ADLER 97B also find $\operatorname{Re}(\eta_{+-0}) = -0.002 \pm 0.007^{+0.004}_{-0.001}$. See also ANGELOPOU-LOS 98C.

² The ADLER 96D fit also yields $\operatorname{Re}(\eta_{+-0}) = 0.006 \pm 0.013 \pm 0.001$ with a correlation $+0.66$ between real and imaginary parts. Their results correspond to $|\eta_{+-0}| < 0.037$ with 90% CL.

³ ZOU 94 use theoretical constraint $\operatorname{Re}(\eta_{+-0}) = \operatorname{Re}(\epsilon) = 0.0016$. Without this constraint they find $\operatorname{Im}(\eta_{+-0}) = 0.019 \pm 0.061$ and $\operatorname{Re}(\eta_{+-0}) = 0.019 \pm 0.027$.

$$\operatorname{Im}(\eta_{000})^2 = \Gamma(K_S^0 \rightarrow 3\pi^0) / \Gamma(K_L^0 \rightarrow 3\pi^0)$$

CPT assumed valid (i.e. $\operatorname{Re}(\eta_{000}) \simeq 0$). This limit determines branching ratio $\Gamma(3\pi^0)/\Gamma_{\text{total}}$ above.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.1	90	632	¹ BARMIN	83 HLBC
<0.28	90		² GJESDAL	74B SPEC Indirect meas.

¹ BARMIN 83 find $\operatorname{Re}(\eta_{000}) = (-0.08 \pm 0.18)$ and $\operatorname{Im}(\eta_{000}) = (-0.05 \pm 0.27)$. Assuming CPT invariance they obtain the limit quoted above.

² GJESDAL 74B uses $K_{2\pi}$, $K_{\mu 3}$, and K_{e3} decay results, unitarity, and CPT. Calculates $|\eta_{000}| = 0.26 \pm 0.20$. We convert to upper limit.

$\text{Im}(\eta_{000}) = \text{Im}(A(K_S^0 \rightarrow \pi^0 \pi^0 \pi^0)/A(K_L^0 \rightarrow \pi^0 \pi^0 \pi^0))$

$K_S^0 \rightarrow \pi^0 \pi^0 \pi^0$ violates CP conservation, in contrast to $K_S^0 \rightarrow \pi^+ \pi^- \pi^0$ which has a CP -conserving part.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.001 ± 0.016 OUR AVERAGE				
$0.000 \pm 0.009 \pm 0.013$	4.9M	¹ LAI	05A	NA48 Assumes CPT
$-0.05 \pm 0.12 \pm 0.05$	17300	² ANGELOPO...	98B	CPLR Assumes CPT

¹ LAI 05A assumes $\text{Re}(\eta_{000}) = \text{Re}(\epsilon) = 1.66 \times 10^{-3}$. The equivalent limit is $|\eta_{000}|_{CPT} < 0.025$ at 90% CL. Without assuming CPT invariance, they obtain $\text{Re}(\eta_{000}) = -0.002 \pm 0.011 \pm 0.015$ and $\text{Im}(\eta_{000}) = -0.003 \pm 0.013 \pm 0.017$ with a statistical correlation coefficient of 0.77 and an overall correlation coefficient of 0.57 between imaginary and real part. The equivalent limit is $|\eta_{000}| < 0.045$ at 90% CL

² ANGELOPOULOS 98B assumes $\text{Re}(\eta_{000}) = \text{Re}(\epsilon) = 1.635 \times 10^{-3}$. Without assuming CPT invariance, they obtain $\text{Re}(\eta_{000}) = 0.18 \pm 0.14 \pm 0.06$ and $\text{Im}(\eta_{000}) = 0.15 \pm 0.20 \pm 0.03$.

$|\eta_{000}| = |A(K_S^0 \rightarrow 3\pi^0)/A(K_L^0 \rightarrow 3\pi^0)|$

A non-zero value violates CP invariance.

VALUE	CL%	EVTS	DOCUMENT ID	TECN
<0.0088	90	590M	BABUSCI	13C KLOE
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.018	90	37.8M	AMBROSINO	05B KLOE
<0.045	90	4.9M	LAI	05A NA48

— DECAY-PLANE ASYMMETRY IN $\pi^+ \pi^- e^+ e^-$ DECAYS —

This is the CP -violating asymmetry

$$A = \frac{N_{\sin\phi\cos\phi>0.0} - N_{\sin\phi\cos\phi<0.0}}{N_{\sin\phi\cos\phi>0.0} + N_{\sin\phi\cos\phi<0.0}}$$

where ϕ is the angle between the $e^+ e^-$ and $\pi^+ \pi^-$ planes in the K_S^0 rest frame.

CP asymmetry A in $K_S^0 \rightarrow \pi^+ \pi^- e^+ e^-$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-0.4 ± 0.8 OUR AVERAGE			
-0.4 ± 0.8	¹ BATLEY	11	NA48 2002 data
-1.1 ± 4.1	LAI	03C	NA48 1998+1999 data
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.5 \pm 4.0 \pm 1.6$	LAI	03C	NA48 1999 data

¹ The result is used to set the limit $A < 1.5\%$ at 90% C.L.

K_S^0 REFERENCES

AAIJ	20AE	PRL 125 231801	R. Aaij <i>et al.</i>	(LHCb Collab.)
BABUSCI	20	PL B804 135378	D. Babusci <i>et al.</i>	(KLOE-2 Collab.)
ANASTASI	18A	JHEP 1809 021	A. Anastasi <i>et al.</i>	(KLOE-2 Collab.)
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)
AAIJ	17BQ	EPJ C77 678	R. Aaij <i>et al.</i>	(LHCb Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
AAIJ	13G	JHEP 1301 090	R. Aaij <i>et al.</i>	(LHCb Collab.)
BABUSCI	13C	PL B723 54	D. Babusci <i>et al.</i>	(KLOE-2 Collab.)
ABOUZAID	11	PR D83 092001	E. Abouzaid <i>et al.</i>	(FNAL KTeV Collab.)
AMBROSINO	11	EPJ C71 1604	F. Ambrosino <i>et al.</i>	(KLOE Collab.)

BATLEY	11	PL B694 301	J.R. Batley <i>et al.</i>	(CERN NA48/1 Collab.)
AMBROSINO	09A	PL B672 203	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AMBROSINO	08C	JHEP 0805 051	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
ANDRE	07	ANP 322 2518	T. Andre	(EFI)
BATLEY	07D	PL B653 145	J.R. Batley <i>et al.</i>	(CERN NA48 Collab.)
AMBROSINO	06C	EPJ C48 767	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AMBROSINO	06E	PL B636 173	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AMBROSINO	05B	PL B619 61	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
BATLEY	05	PL B630 31	J.R. Batley <i>et al.</i>	(NA48 Collab.)
LAI	05A	PL B610 165	A. Lai <i>et al.</i>	(CERN NA48 Collab.)
ALEXOPOU...	04A	PR D70 092007	T. Alexopoulos <i>et al.</i>	(FNAL KTeV Collab.)
BATLEY	04A	PL B599 197	J.R. Batley <i>et al.</i>	(NA48 Collab.)
LAI	04	PL B578 276	A. Lai <i>et al.</i>	(CERN NA48 Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
ALAVI-HARATI	03	PR D67 012005	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
Also		PR D70 079904 (errat.)	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
BATLEY	03	PL B576 43	J.R. Batley <i>et al.</i>	(CERN NA48 Collab.)
LAI	03	PL B551 7	A. Lai <i>et al.</i>	(CERN NA48 Collab.)
LAI	03B	PL B556 105	A. Lai <i>et al.</i>	(CERN NA48 Collab.)
LAI	03C	EPJ C30 33	A. Lai <i>et al.</i>	(CERN NA48 Collab.)
ALOISIO	02	PL B535 37	A. Aloisio <i>et al.</i>	(KLOE Collab.)
ALOISIO	02B	PL B538 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)
LAI	02C	PL B537 28	A. Lai <i>et al.</i>	(CERN NA48 Collab.)
LAI	01	PL B514 253	A. Lai <i>et al.</i>	(CERN NA48 Collab.)
LAI	00	PL B493 29	A. Lai <i>et al.</i>	(CERN NA48 Collab.)
LAI	00B	PL B496 137	A. Lai <i>et al.</i>	(CERN NA48 Collab.)
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i>	(PDG Collab.)
ACHASOV	99D	PL B459 674	M.N. Achasov <i>et al.</i>	
AKHMETSHIN	99	PL B456 90	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ANGELOPO...	98B	PL B425 391	A. Angelopoulos <i>et al.</i>	(CPLEAR Collab.)
ANGELOPO...	98C	EPJ C5 389	A. Angelopoulos <i>et al.</i>	(CPLEAR Collab.)
PDG	98	EPJ C3 1	C. Caso <i>et al.</i>	(PDG Collab.)
ADLER	97B	PL B407 193	R. Adler <i>et al.</i>	(CPLEAR Collab.)
ANGELOPO...	97	PL B413 232	A. Angelopoulos <i>et al.</i>	(CPLEAR Collab.)
BERTANZA	97	ZPHY C73 629	L. Bertanza (PISA, CERN, EDIN, MAINZ, ORSAY+)	
ADLER	96D	PL B370 167	R. Adler <i>et al.</i>	(CPLEAR Collab.)
ADLER	96E	PL B374 313	R. Adler <i>et al.</i>	(CPLEAR Collab.)
ZOU	96	PL B369 362	Y. Zou <i>et al.</i>	(RUTG, MINN, MICH)
BARR	95B	PL B351 579	G.D. Barr <i>et al.</i>	(CERN, EDIN, MAINZ, LALO+)
SCHWINGEN...	95	PRL 74 4376	B. Schwingenheuer <i>et al.</i>	(EFI, CHIC+)
BLICK	94	PL B334 234	A.M. Blick <i>et al.</i>	(SERP, JINR)
THOMSON	94	PL B337 411	G.B. Thomson <i>et al.</i>	(RUTG, MINN, MICH)
ZOU	94	PL B329 519	Y. Zou <i>et al.</i>	(RUTG, MINN, MICH)
BARR	93B	PL B304 381	G.D. Barr <i>et al.</i>	(CERN, EDIN, MAINZ, LALO+)
GIBBONS	93	PRL 70 1199	L.K. Gibbons <i>et al.</i>	(FNAL E731 Collab.)
Also		PR D55 6625	L.K. Gibbons <i>et al.</i>	(FNAL E731 Collab.)
RAMBERG	93	PRL 70 2525	E. Ramberg <i>et al.</i>	(FNAL E731 Collab.)
BALATS	89	SJNP 49 828	M.Y. Balats <i>et al.</i>	(ITEP)
		Translated from YAF 49 1332.		
GIBBONS	88	PRL 61 2661	L.K. Gibbons <i>et al.</i>	(FNAL E731 Collab.)
BURKHARDT	87	PL B199 139	H. Burkhardt <i>et al.</i>	(CERN, EDIN, MAINZ+)
GROSSMAN	87	PRL 59 18	N. Grossman <i>et al.</i>	(MINN, MICH, RUTG)
BARMIN	86	SJNP 44 622	V.V. Barmin <i>et al.</i>	(ITEP)
		Translated from YAF 44 965.		
BARMIN	86B	NC 96A 159	V.V. Barmin <i>et al.</i>	(ITEP, PADO)
PDG	86B	PL 170B 130	M. Aguilar-Benitez <i>et al.</i>	(CERN, CIT+)
BARMIN	85	NC 85A 67	V.V. Barmin <i>et al.</i>	(ITEP, PADO)
Also		SJNP 41 759	V.V. Barmin <i>et al.</i>	(ITEP)
		Translated from YAF 41 1187.		
BARMIN	83	PL 128B 129	V.V. Barmin <i>et al.</i>	(ITEP, PADO)
Also		SJNP 39 269	V.V. Barmin <i>et al.</i>	(ITEP, PADO)
		Translated from YAF 39 428.		
ARONSON	82	PRL 48 1078	S.H. Aronson <i>et al.</i>	(BNL, CHIC, STAN+)
ARONSON	82B	PRL 48 1306	S.H. Aronson <i>et al.</i>	(BNL, CHIC, PURD)
Also		PL 116B 73	E. Fischbach <i>et al.</i>	(PURD, BNL, CHIC)
Also		PR D28 476	S.H. Aronson <i>et al.</i>	(BNL, CHIC, PURD)
Also		PR D28 495	S.H. Aronson <i>et al.</i>	(BNL, CHIC, PURD)
ARONSON	76	NC 32A 236	S.H. Aronson <i>et al.</i>	(WISC, EFI, UCSD+)
EVERHART	76	PR D14 661	G.C. Everhart <i>et al.</i>	(PENN)
TAUREG	76	PL 65B 92	H. Taureg <i>et al.</i>	(HEIDH, CERN, DORT)
BALDO-...	75	NC 25A 688	M. Baldo-Ceolin <i>et al.</i>	(PADO, WISC)

CARITHERS	75	PRL 34 1244	W.C.J. Carithers <i>et al.</i>	(COLU, NYU)
BOBISUT	74	LNC 11 646	F. Bobisut <i>et al.</i>	(PADO)
COWELL	74	PR D10 2083	P.L. Cowell <i>et al.</i>	(STON, COLU)
GEWENIGER	74B	PL 48B 487	C. Geweniger <i>et al.</i>	(CERN, HEIDH)
GJESDAL	74B	PL 52B 119	S. Gjesdal <i>et al.</i>	(CERN, HEIDH)
BURGUN	73	PL 46B 481	G. Burgun <i>et al.</i>	(SACL, CERN)
GJESDAL	73	PL 44B 217	S. Gjesdal <i>et al.</i>	(CERN, HEIDH)
HILL	73	PR D8 1290	D.G. Hill <i>et al.</i>	(BNL, CMU)
ALITTI	72	PL 39B 568	J. Alitti, E. Lesquoy, A. Muller	(SACL)
BURGUN	72	NP B50 194	G. Burgun <i>et al.</i>	(SACL, CERN, OSLO)
METCALF	72	PL 40B 703	M. Metcalf <i>et al.</i>	(CERN, IPN, WIEN)
MORSE	72B	PRL 28 388	R. Morse <i>et al.</i>	(COLO, PRIN, UMD)
NAGY	72	NP B47 94	E. Nagy, F. Telbisz, G. Vesztergombi	(BUDA)
Also		PL 30B 498	G. Bozoki <i>et al.</i>	(BUDA)
SKJEGGEST...	72	NP B48 343	O. Skjeggestad <i>et al.</i>	(OSLO, CERN, SACL)
BALTAY	71	PRL 27 1678	C. Baltay <i>et al.</i>	(COLU)
Also		Thesis Nevis 187	W.A. Cooper	(COLU)
MOFFETT	70	BAPS 15 512	R. Moffett <i>et al.</i>	(ROCH)
BOZOKI	69	PL 30B 498	G. Bozoki <i>et al.</i>	(BUDA)
DOYLE	69	Thesis UCRL 18139	J.C. Doyle	(LRL)
GOBBI	69	PRl 22 682	B. Gobbi <i>et al.</i>	(ROCH)
HYAMS	69B	PL 29B 521	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
MORFIN	69	PRl 23 660	J.G. Morfin, D. Sinclair	(MICH)
DONALD	68B	PL 27B 58	R.A. Donald <i>et al.</i>	(LIVP, CERN, IPNP+)
HILL	68	PR 171 1418	D.G. Hill <i>et al.</i>	(BNL, CMU)
AUBERT	65	PL 17 59	B. Aubert <i>et al.</i>	(EPOL, ORSAY)
BROWN	63	PR 130 769	J.L. Brown <i>et al.</i>	(LRL, MICH)
CHRETIEN	63	PR 131 2208	M. Chretien <i>et al.</i>	(BRAN, BROW, HARV+)
BROWN	61	NC 19 1155	J.L. Brown <i>et al.</i>	(MICH)
BOLDT	58B	PRL 1 150	E. Boldt, D.O. Caldwell, Y. Pal	(MIT)

OTHER RELATED PAPERS

LITTENBERG	93	ARNPS 43 729 Rare and Radiative Kaon Decays	L.S. Littenberg, G. Valencia	(BNL, FNAL)
BATTISTON	92	PRPL 214 293 Status and Perspectives of K Decay Physics	R. Battiston <i>et al.</i>	(PGIA, CERN, TRSTT)
TRILLING	65B	UCRL 16473 Updated from 1965 Argonne Conference, page 115.	G.N. Trilling	(LRL)
CRAWFORD	62	CERN Conf. 827	F.S. Crawford	(LRL)
FITCH	61	NC 22 1160	V.L. Fitch, P.A. Piroue, R.B. Perkins	(PRIN+)
GOOD	61	PR 124 1223	R.H. Good <i>et al.</i>	(LRL)
BIRGE	60	Rochester Conf. 601	R.W. Birge <i>et al.</i>	(LRL, WISC)
MULLER	60	PRL 4 418	F. Muller <i>et al.</i>	(LRL, BNL)