

$\chi_{b2}(3P)$

$I^G(J^{PC}) = 0^+(2^{++})$

Observed in the radiative decay to  $\Upsilon(3S)$ , therefore  $C = +$ .  $J$  needs confirmation.

### $\chi_{b2}(3P)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

**10524.02 ± 0.57 ± 0.53** <sup>1</sup> SIRUNYAN 18N CMS  $p p \rightarrow \gamma \mu^+ \mu^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

10530 ± 5 ± 9 <sup>2</sup> AAD 12A ATLAS  $p p \rightarrow \gamma \mu^+ \mu^- X$

<sup>1</sup> Systematic error includes an additional 0.5 MeV for the uncertainty on the  $\Upsilon(3S)$  mass. Also measures  $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.60 \pm 0.64 \pm 0.17$  MeV. A total of 372  $\chi_{b1}(3P)$  and  $\chi_{b2}(3P)$  events was observed.

<sup>2</sup> The mass barycenter of the merged lineshapes from the  $J = 1$  and 2 states.

### $\chi_{b2}(3P)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \Upsilon(3S)\gamma$	seen

### $\chi_{b2}(3P)$ BRANCHING RATIOS

$\Gamma(\Upsilon(3S)\gamma)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$
--	-------------------

VALUE	DOCUMENT ID	TECN	COMMENT
seen	SIRUNYAN 18N CMS	$p p \rightarrow \gamma \mu^+ \mu^- X$	

### $\chi_{b2}(3P)$ REFERENCES

SIRUNYAN AAD	18N 12A	PRL 121 092002 PRL 108 152001	A.M. Sirunyan <i>et al.</i> G. Aad <i>et al.</i>	(CMS Collab.) (ATLAS Collab.)
-----------------	------------	----------------------------------	---	----------------------------------