

$\chi_{c0}(1P)$  $I^G(J^{PC}) = 0^+(0^{++})$  **$\chi_{c0}(1P)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3414.71 \pm 0.30</math> OUR AVERAGE</b>				
3413.0 $\pm 1.9 \pm 0.6$	933	<sup>1</sup> AAIJ	17BB LHCb	$p p \rightarrow b \bar{b} X \rightarrow 2(K^+ K^-)X$
3414.2 $\pm 0.5 \pm 2.3$	5.4k	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow \text{hadrons}$
3406 $\pm 7 \pm 6$	230	<sup>2</sup> ABE	07	BELL $e^+ e^- \rightarrow J/\psi(c\bar{c})$
3414.21 $\pm 0.39 \pm 0.27$		ABLIKIM	05G	BES2 $\psi(2S) \rightarrow \gamma \chi_{c0}$
3414.7 $\pm 0.7 \pm 0.2$		<sup>3</sup> ANDREOTTI	03	E835 $\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0 \pi^0$
3415.5 $\pm 0.4 \pm 0.4$	392	<sup>4</sup> BAGNASCO	02	E835 $\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi \gamma$
3417.4 $\pm 1.8 \pm 0.2$		<sup>3</sup> AMBROGIANI	99B	E835 $\bar{p}p \rightarrow e^+ e^- \gamma$
3414.1 $\pm 0.6 \pm 0.8$		BAI	99B	BES $\psi(2S) \rightarrow \gamma X$
3417.8 $\pm 0.4 \pm 4$		<sup>3</sup> GAISER	86	CBAL $\psi(2S) \rightarrow \gamma X$
3416 $\pm 3 \pm 4$		<sup>5</sup> TANENBAUM	78	MRK1 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3414.6 $\pm 1.1$	266	UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
3416.5 $\pm 3.0$		EISENSTEIN	01	CLE2 $e^+ e^- \rightarrow e^+ e^- \chi_{c0}$
3422 $\pm 10$		<sup>5</sup> BARTEL	78B	CNTR $e^+ e^- \rightarrow J/\psi 2\gamma$
3415 $\pm 9$		<sup>5</sup> BIDDICK	77	CNTR $e^+ e^- \rightarrow \gamma X$

<sup>1</sup> From a fit of the  $\phi\phi$  invariant mass with the width of  $\chi_{c0}(1P)$  fixed to the PDG 16 value.

<sup>2</sup> From a fit of the  $J/\psi$  recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

<sup>3</sup> Using mass of  $\psi(2S) = 3686.0$  MeV.

<sup>4</sup> Recalculated by ANDREOTTI 05A, using the value of  $\psi(2S)$  mass from AULCHENKO 03.

<sup>5</sup> Mass value shifted by us by amount appropriate for  $\psi(2S)$  mass = 3686 MeV and  $J/\psi(1S)$  mass = 3097 MeV.

 **$\chi_{c0}(1P)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>10.8 \pm 0.6</math> OUR FIT</b>				
<b><math>10.5 \pm 0.8</math> OUR AVERAGE</b> Error includes scale factor of 1.1.				
10.6 $\pm 1.9 \pm 2.6$	5.4k	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow \text{hadrons}$
$12.6^{+1.5}_{-1.6} {}^{+0.9}_{-1.1}$		ABLIKIM	05G	BES2 $\psi(2S) \rightarrow \gamma \chi_{c0}$
$8.6^{+1.7}_{-1.3} {}^{+0.1}_{-0.1}$		<sup>1</sup> ANDREOTTI	03	E835 $\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0 \pi^0$
9.7 $\pm 1.0$	392	<sup>1</sup> BAGNASCO	02	E835 $\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi \gamma$
$16.6^{+5.2}_{-3.7} {}^{+0.1}_{-0.1}$		AMBROGIANI	99B	E835 $\bar{p}p \rightarrow e^+ e^- \gamma$
14.3 $\pm 2.0 \pm 3.0$		BAI	98I	BES $\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
13.5 $\pm 3.3 \pm 4.2$		GAISER	86	CBAL $\psi(2S) \rightarrow \gamma X, \gamma \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$13.2 \pm 2.1$       266      UEHARA      13      BELL       $\gamma\gamma \rightarrow K_S^0 K_S^0$

<sup>1</sup> Recalculated by ANDREOTTI 05A.

### $\chi_{c0}(1P)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
<b>Hadronic decays</b>		
$\Gamma_1$ $2(\pi^+ \pi^-)$	$(2.34 \pm 0.18) \%$	
$\Gamma_2$ $\rho^0 \pi^+ \pi^-$	$(9.1 \pm 2.9) \times 10^{-3}$	
$\Gamma_3$ $\rho^0 \rho^0$		
$\Gamma_4$ $f_0(980) f_0(980)$	$(6.6 \pm 2.1) \times 10^{-4}$	
$\Gamma_5$ $\pi^+ \pi^- \pi^0 \pi^0$	$(3.3 \pm 0.4) \%$	
$\Gamma_6$ $\rho^+ \pi^- \pi^0 + \text{c.c.}$	$(2.9 \pm 0.4) \%$	
$\Gamma_7$ $4\pi^0$	$(3.3 \pm 0.4) \times 10^{-3}$	
$\Gamma_8$ $\pi^+ \pi^- K^+ K^-$	$(1.81 \pm 0.14) \%$	
$\Gamma_9$ $K_0^*(1430)^0 \bar{K}_0^*(1430)^0 \rightarrow \pi^+ \pi^- K^+ K^-$	$(9.8 \pm 4.0) \times 10^{-4}$	
$\Gamma_{10}$ $K_0^*(1430)^0 \bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-$	$(8.0 \pm 2.0) \times 10^{-4}$	
$\Gamma_{11}$ $K_1(1270)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-$	$(6.3 \pm 1.9) \times 10^{-3}$	
$\Gamma_{12}$ $K_1(1400)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-$	$< 2.7 \times 10^{-3}$	CL=90%
$\Gamma_{13}$ $f_0(980) f_0(980)$	$(1.6 \pm 1.0) \times 10^{-4}$	
$\Gamma_{14}$ $f_0(980) f_0(2200)$	$(7.9 \pm 2.0) \times 10^{-4}$	
$\Gamma_{15}$ $f_0(1370) f_0(1370)$	$< 2.7 \times 10^{-4}$	CL=90%
$\Gamma_{16}$ $f_0(1370) f_0(1500)$	$< 1.7 \times 10^{-4}$	CL=90%
$\Gamma_{17}$ $f_0(1370) f_0(1710)$	$(6.7 \pm 3.5) \times 10^{-4}$	
$\Gamma_{18}$ $f_0(1500) f_0(1370)$	$< 1.3 \times 10^{-4}$	CL=90%
$\Gamma_{19}$ $f_0(1500) f_0(1500)$	$< 5 \times 10^{-5}$	CL=90%
$\Gamma_{20}$ $f_0(1500) f_0(1710)$	$< 7 \times 10^{-5}$	CL=90%
$\Gamma_{21}$ $K^+ K^- \pi^+ \pi^- \pi^0$	$(8.6 \pm 0.9) \times 10^{-3}$	
$\Gamma_{22}$ $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(4.2 \pm 0.4) \times 10^{-3}$	
$\Gamma_{23}$ $K^+ K^- \pi^0 \pi^0$	$(5.6 \pm 0.9) \times 10^{-3}$	
$\Gamma_{24}$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	$(2.49 \pm 0.33) \%$	
$\Gamma_{25}$ $\rho^+ K^- K^0 + \text{c.c.}$	$(1.21 \pm 0.21) \%$	
$\Gamma_{26}$ $K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	$(4.6 \pm 1.2) \times 10^{-3}$	

$\Gamma_{27}$	$K_S^0 K_S^0 \pi^+ \pi^-$	$(5.7 \pm 1.1) \times 10^{-3}$	
$\Gamma_{28}$	$K^+ K^- \eta \pi^0$	$(3.0 \pm 0.7) \times 10^{-3}$	
$\Gamma_{29}$	$3(\pi^+ \pi^-)$	$(1.20 \pm 0.18) \%$	
$\Gamma_{30}$	$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	$(7.5 \pm 1.6) \times 10^{-3}$	
$\Gamma_{31}$	$K^*(892)^0 \bar{K}^*(892)^0$	$(1.7 \pm 0.6) \times 10^{-3}$	
$\Gamma_{32}$	$\pi \pi$	$(8.51 \pm 0.33) \times 10^{-3}$	
$\Gamma_{33}$	$\pi^0 \eta$	$< 1.8 \times 10^{-4}$	
$\Gamma_{34}$	$\pi^0 \eta'$	$< 1.1 \times 10^{-3}$	
$\Gamma_{35}$	$\pi^0 \eta_c$	$< 1.6 \times 10^{-3}$	CL=90%
$\Gamma_{36}$	$\eta \eta$	$(3.01 \pm 0.19) \times 10^{-3}$	
$\Gamma_{37}$	$\eta \eta'$	$(9.1 \pm 1.1) \times 10^{-5}$	
$\Gamma_{38}$	$\eta' \eta'$	$(2.17 \pm 0.12) \times 10^{-3}$	
$\Gamma_{39}$	$\omega \omega$	$(9.7 \pm 1.1) \times 10^{-4}$	
$\Gamma_{40}$	$\omega \phi$	$(1.41 \pm 0.13) \times 10^{-4}$	
$\Gamma_{41}$	$\omega K^+ K^-$	$(1.94 \pm 0.21) \times 10^{-3}$	
$\Gamma_{42}$	$K^+ K^-$	$(6.05 \pm 0.31) \times 10^{-3}$	
$\Gamma_{43}$	$K_S^0 K_S^0$	$(3.16 \pm 0.17) \times 10^{-3}$	
$\Gamma_{44}$	$\pi^+ \pi^- \eta$	$< 2.0 \times 10^{-4}$	CL=90%
$\Gamma_{45}$	$\pi^+ \pi^- \eta'$	$< 4 \times 10^{-4}$	CL=90%
$\Gamma_{46}$	$\bar{K}^0 K^+ \pi^- + \text{c.c.}$	$< 9 \times 10^{-5}$	CL=90%
$\Gamma_{47}$	$K^+ K^- \pi^0$	$< 6 \times 10^{-5}$	CL=90%
$\Gamma_{48}$	$K^+ K^- \eta$	$< 2.3 \times 10^{-4}$	CL=90%
$\Gamma_{49}$	$K^+ K^- K_S^0 K_S^0$	$(1.4 \pm 0.5) \times 10^{-3}$	
$\Gamma_{50}$	$K_S^0 K_S^0 K_S^0 K_S^0$	$(5.8 \pm 0.5) \times 10^{-4}$	
$\Gamma_{51}$	$K^+ K^- K^+ K^-$	$(2.82 \pm 0.29) \times 10^{-3}$	
$\Gamma_{52}$	$K^+ K^- \phi$	$(9.7 \pm 2.5) \times 10^{-4}$	
$\Gamma_{53}$	$\bar{K}^0 K^+ \pi^- \phi + \text{c.c.}$	$(3.7 \pm 0.6) \times 10^{-3}$	
$\Gamma_{54}$	$K^+ K^- \pi^0 \phi$	$(1.90 \pm 0.35) \times 10^{-3}$	
$\Gamma_{55}$	$\phi \pi^+ \pi^- \pi^0$	$(1.18 \pm 0.15) \times 10^{-3}$	
$\Gamma_{56}$	$\phi \phi$	$(8.0 \pm 0.7) \times 10^{-4}$	
$\Gamma_{57}$	$\phi \phi \eta$	$(8.4 \pm 1.0) \times 10^{-4}$	
$\Gamma_{58}$	$p \bar{p}$	$(2.21 \pm 0.08) \times 10^{-4}$	
$\Gamma_{59}$	$p \bar{p} \pi^0$	$(7.0 \pm 0.7) \times 10^{-4}$	S=1.3
$\Gamma_{60}$	$p \bar{p} \eta$	$(3.5 \pm 0.4) \times 10^{-4}$	
$\Gamma_{61}$	$p \bar{p} \omega$	$(5.2 \pm 0.6) \times 10^{-4}$	
$\Gamma_{62}$	$p \bar{p} \phi$	$(6.0 \pm 1.4) \times 10^{-5}$	
$\Gamma_{63}$	$p \bar{p} \pi^+ \pi^-$	$(2.1 \pm 0.7) \times 10^{-3}$	S=1.4
$\Gamma_{64}$	$p \bar{p} \pi^0 \pi^0$	$(1.04 \pm 0.28) \times 10^{-3}$	
$\Gamma_{65}$	$p \bar{p} K^+ K^- (\text{non-resonant})$	$(1.22 \pm 0.26) \times 10^{-4}$	
$\Gamma_{66}$	$p \bar{p} K_S^0 K_S^0$	$< 8.8 \times 10^{-4}$	CL=90%
$\Gamma_{67}$	$p \bar{n} \pi^-$	$(1.27 \pm 0.11) \times 10^{-3}$	
$\Gamma_{68}$	$\bar{p} n \pi^+$	$(1.37 \pm 0.12) \times 10^{-3}$	

$\Gamma_{69}$	$p\bar{n}\pi^-\pi^0$	$(2.34 \pm 0.21) \times 10^{-3}$	
$\Gamma_{70}$	$\bar{p}n\pi^+\pi^0$	$(2.21 \pm 0.18) \times 10^{-3}$	
$\Gamma_{71}$	$\Lambda\bar{\Lambda}$	$(3.59 \pm 0.15) \times 10^{-4}$	
$\Gamma_{72}$	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(1.18 \pm 0.13) \times 10^{-3}$	
$\Gamma_{73}$	$\Lambda\bar{\Lambda}\pi^+\pi^-$ (non-resonant)	$< 5 \times 10^{-4}$	CL=90%
$\Gamma_{74}$	$\Lambda\bar{\Lambda}\eta$	$(2.3 \pm 0.4) \times 10^{-4}$	
$\Gamma_{75}$	$\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.}$	$< 5 \times 10^{-4}$	CL=90%
$\Gamma_{76}$	$\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.}$	$< 5 \times 10^{-4}$	CL=90%
$\Gamma_{77}$	$K^+\bar{p}\Lambda + \text{c.c.}$	$(1.25 \pm 0.12) \times 10^{-3}$	S=1.3
$\Gamma_{78}$	$nK_S^0\bar{\Lambda} + \text{c.c.}$	$(6.6 \pm 0.5) \times 10^{-4}$	
$\Gamma_{79}$	$K^*(892)^+\bar{p}\Lambda + \text{c.c.}$	$(4.8 \pm 0.9) \times 10^{-4}$	
$\Gamma_{80}$	$K^+\bar{p}\Lambda(1520) + \text{c.c.}$	$(2.9 \pm 0.7) \times 10^{-4}$	
$\Gamma_{81}$	$\Lambda(1520)\bar{\Lambda}(1520)$	$(3.1 \pm 1.2) \times 10^{-4}$	
$\Gamma_{82}$	$\Sigma^0\bar{\Sigma}^0$	$(4.68 \pm 0.32) \times 10^{-4}$	
$\Gamma_{83}$	$\Sigma^+\bar{p}K_S^0 + \text{c.c.}$	$(3.52 \pm 0.27) \times 10^{-4}$	
$\Gamma_{84}$	$\Sigma^0\bar{p}K^+ + \text{c.c.}$	$(3.03 \pm 0.20) \times 10^{-4}$	
$\Gamma_{85}$	$\Sigma^+\bar{\Sigma}^-$	$(4.6 \pm 0.8) \times 10^{-4}$	S=2.6
$\Gamma_{86}$	$\Sigma^-\bar{\Sigma}^+$	$(5.1 \pm 0.5) \times 10^{-4}$	
$\Gamma_{87}$	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	$(1.6 \pm 0.6) \times 10^{-4}$	
$\Gamma_{88}$	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$	$(2.3 \pm 0.7) \times 10^{-4}$	
$\Gamma_{89}$	$K^-\Lambda\bar{\Xi}^+ + \text{c.c.}$	$(1.94 \pm 0.35) \times 10^{-4}$	
$\Gamma_{90}$	$\Xi^0\bar{\Xi}^0$	$(4.5 \pm 0.5) \times 10^{-4}$	S=1.7
$\Gamma_{91}$	$\Xi^-\bar{\Xi}^+$	$(4.45 \pm 0.19) \times 10^{-4}$	
$\Gamma_{92}$	$\eta_c\pi^+\pi^-$	$< 7 \times 10^{-4}$	CL=90%

### Radiative decays

$\Gamma_{93}$	$\gamma J/\psi(1S)$	$(1.40 \pm 0.05) \%$	
$\Gamma_{94}$	$\gamma\rho^0$	$< 9 \times 10^{-6}$	CL=90%
$\Gamma_{95}$	$\gamma\omega$	$< 8 \times 10^{-6}$	CL=90%
$\Gamma_{96}$	$\gamma\phi$	$< 6 \times 10^{-6}$	CL=90%
$\Gamma_{97}$	$\gamma\gamma$	$(2.04 \pm 0.09) \times 10^{-4}$	
$\Gamma_{98}$	$e^+e^- J/\psi(1S)$	$(1.33 \pm 0.29) \times 10^{-4}$	
$\Gamma_{99}$	$\mu^+\mu^- J/\psi(1S)$	$< 1.9 \times 10^{-5}$	CL=90%

## CONSTRAINED FIT INFORMATION

A multiparticle fit to  $\chi_{c1}(1P)$ ,  $\chi_{c0}(1P)$ ,  $\chi_{c2}(1P)$ , and  $\psi(2S)$  with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 248 measurements to determine 49 parameters. The overall fit has a  $\chi^2 = 379.8$  for 199 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ .

$x_2$	24									
$x_8$	9	2								
$x_{30}$	5	1	28							
$x_{32}$	8	2	10	3						
$x_{36}$	4	1	5	1	14					
$x_{42}$	8	2	8	3	18	11				
$x_{43}$	7	2	8	2	18	10	14			
$x_{51}$	5	1	5	2	9	5	7	7		
$x_{56}$	7	2	6	2	9	5	7	7	4	
$x_{58}$	3	1	4	1	3	-1	7	7	3	3
$x_{71}$	7	2	9	2	23	13	18	18	8	9
$x_{93}$	5	1	6	2	17	11	13	12	6	6
$x_{97}$	-8	-2	-2	-3	14	9	10	10	3	1
$\Gamma$	-26	-6	-19	-10	-15	-7	-14	-12	-10	-13
	$x_1$	$x_2$	$x_8$	$x_{30}$	$x_{32}$	$x_{36}$	$x_{42}$	$x_{43}$	$x_{51}$	$x_{56}$
$x_{71}$	9									
$x_{93}$	-19	16								
$x_{97}$	6	15	13							
$\Gamma$	-4	-13	-9	-38						
	$x_{58}$	$x_{71}$	$x_{93}$	$x_{97}$						

### $\chi_{c0}(1P)$ PARTIAL WIDTHS

$$\text{—— } \chi_{c0}(1P) \Gamma(i) \Gamma(\gamma J/\psi(1S)) / \Gamma(\text{total}) \text{ ——}$$

$$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}} \quad \Gamma_{58}\Gamma_{93}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>33.6 \pm 2.3</math> OUR FIT</b>				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$26.6 \pm 2.6 \pm 1.4$       392      1,2 BAGNASCO      02      E835       $\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi \gamma$

$48.7^{+11.3}_{-8.9} \pm 2.4$       1,2 AMBROGIANI      99B      E835       $\bar{p}p \rightarrow \gamma J/\psi$

<sup>1</sup> Calculated by us using  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$ .

<sup>2</sup> Values in  $(\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}})$  and  $(\Gamma(p\bar{p}) / \Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}})$  are not independent. The latter is used in the fit since it is less correlated to the total width.

$\chi_{c0}(1P) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ 

$$\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_1 \Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>52 ± 4 OUR FIT</b>				
<b>49 ± 10 OUR AVERAGE</b>				Error includes scale factor of 1.8.
44.7 ± 3.6 ± 4.9	3.6k	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(\pi^+\pi^-)$
75 ± 13 ± 8		EISENSTEIN	01	CLE2 $e^+e^- \rightarrow e^+e^-\chi_{c0}$

$$\Gamma(\rho^0\rho^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_3 \Gamma_{97}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<12	90	<252	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(\pi^+\pi^-)$

$$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_8 \Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>40.0 ± 3.5 OUR FIT</b>				
<b>38.8 ± 3.7 ± 4.7</b>	1.7k	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+K^-\pi^+\pi^-$

$$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{21} \Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>26 ± 4 ± 4</b>	1094	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

$$\Gamma(K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{30} \Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>16 ± 4 OUR FIT</b>				
<b>16.7 ± 6.1 ± 3.0</b>	495 ± 182	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+K^-\pi^+\pi^-$

$$\Gamma(K^*(892)^0\bar{K}^*(892)^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{31} \Gamma_{97}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<6	90	<148	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+K^-\pi^+\pi^-$

$$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{32} \Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>18.8 ± 1.3 OUR FIT</b>				
<b>23 ± 5 OUR AVERAGE</b>				

29.7 $\pm$ 17.4 $\pm$ 4.8	103 $\pm$ 60 $\pm$ 42	<sup>1</sup> UEHARA	09	BELL $10.6 e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
22.7 $\pm$ 3.2 $\pm$ 3.5	129 $\pm$ 18	<sup>2</sup> NAKAZAWA	05	BELL $10.6 e^+e^- \rightarrow e^+e^-\pi^+\pi^-$

<sup>1</sup> We multiplied the measurement by 3 to convert from  $\pi^0\pi^0$  to  $\pi\pi$ . Interference with the continuum included.

<sup>2</sup> We have multiplied  $\pi^+\pi^-$  measurement by 3/2 to obtain  $\pi\pi$ .

$$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{36} \Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.4 ± 2.3 ± 1.2</b>	22	<sup>1</sup> UEHARA	10A	BELL $10.6 e^+e^- \rightarrow e^+e^-\eta\eta$

<sup>1</sup> Interference with the continuum not included.

$\Gamma(\omega\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{39}\Gamma_{97}/\Gamma$ 

<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
<3.9	90	<sup>1</sup> LIU	12B BELL	$\gamma\gamma \rightarrow 2(\pi^+\pi^-\pi^0)$
<sup>1</sup> Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$ .				

 $\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{40}\Gamma_{97}/\Gamma$ 

<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
<0.34	90	<sup>1</sup> LIU	12B BELL	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$
<sup>1</sup> Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$ .				

 $\Gamma(K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{42}\Gamma_{97}/\Gamma$ 

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>13.4±1.0 OUR FIT</b>				
<b>14.3±1.6±2.3</b>	$153 \pm 17$	NAKAZAWA 05	BELL	$e^+e^- \rightarrow e^+e^-K^+K^-$

 $\Gamma(K_S^0K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{43}\Gamma_{97}/\Gamma$ 

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>7.0 ± 0.5 OUR FIT</b>				
<b>8.7 ± 1.7 ± 0.9</b>	266	<sup>1</sup> UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0K_S^0$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
7.00±0.65±0.71	$134 \pm 12$	CHEN	07B BELL	$e^+e^- \rightarrow e^+e^-\chi_{c0}$
<sup>1</sup> Supersedes CHEN 07B.				

 $\Gamma(K^+K^-K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{51}\Gamma_{97}/\Gamma$ 

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>6.2±0.7 OUR FIT</b>				
<b>7.9±1.3±1.1</b>	$215 \pm 36$	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(K^+K^-)$

 $\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{56}\Gamma_{97}/\Gamma$ 

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.76±0.18 OUR FIT</b>				
<b>1.72±0.33±0.14</b>	$56 \pm 11$	<sup>1</sup> LIU	12B BELL	$\gamma\gamma \rightarrow 2(K^+K^-)$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
2.3 ± 0.9 ± 0.4	$23.6 \pm 9.6$	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(K^+K^-)$
<sup>1</sup> Supersedes UEHARA 08. Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ .				

 **$\chi_{c0}(1P)$  BRANCHING RATIOS****———— HADRONIC DECAYS ————** $\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>
<b>0.0234±0.0018 OUR FIT</b>	

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-))$ VALUE**0.39±0.12 OUR FIT****0.39±0.12**DOCUMENT IDTECNCOMMENT $\Gamma_2/\Gamma_1$  $\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE**0.0091±0.0029 OUR FIT**DOCUMENT ID $\Gamma_2/\Gamma$  $\Gamma(f_0(980)f_0(980))/\Gamma_{\text{total}}$ VALUE (units  $10^{-4}$ )**6.6±2.1±0.1**EVTS $36 \pm 9$ DOCUMENT ID<sup>1</sup> ABLIKIMTECN

04G BES

COMMENT $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$  $\Gamma_4/\Gamma$ 

<sup>1</sup> ABLIKIM 04G reports  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980)f_0(980))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (6.5 \pm 1.6 \pm 1.3) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$ VALUE (%)**3.3±0.4±0.1**EVTS $1751.4$ DOCUMENT ID<sup>1</sup> HETECN

08B CLEO

COMMENT $\Gamma_5/\Gamma$ 

<sup>1</sup> HE 08B reports  $3.54 \pm 0.10 \pm 0.43 \pm 0.18 \%$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\rho^+\pi^-\pi^0+c.c.)/\Gamma_{\text{total}}$ VALUE (%)**2.9±0.4±0.1**EVTS $1358.5$ DOCUMENT ID<sup>1,2</sup> HETECN

08B CLEO

COMMENT $\Gamma_6/\Gamma$ 

<sup>1</sup> HE 08B reports  $3.04 \pm 0.18 \pm 0.42 \pm 0.16 \%$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \rho^+\pi^-\pi^0 + c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Calculated by us. We have added the values from HE 08B for  $\rho^+\pi^-\pi^0$  and  $\rho^-\pi^+\pi^0$  decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**3.3±0.4±0.1**EVTS $3296$ DOCUMENT ID<sup>1</sup> ABLIKIMTECN

11A BES3

COMMENT $\Gamma_7/\Gamma$ 

<sup>1</sup> ABLIKIM 11A reports  $(3.34 \pm 0.06 \pm 0.44) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}} \quad \Gamma_8/\Gamma$$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>
<b><math>18.1 \pm 1.4</math> OUR FIT</b>	

$$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma(\pi^+ \pi^- K^+ K^-) \quad \Gamma_{30}/\Gamma_8$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.41 \pm 0.09</math> OUR FIT</b>			
<b><math>0.41 \pm 0.10</math></b>	TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$

$$\Gamma(K_0^*(1430)^0 \bar{K}_0^*(1430)^0 \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma$$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>9.8^{+3.6}_{-2.8} \pm 0.2</math></b>	83	<sup>1</sup> ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

<sup>1</sup> ABLIKIM 05Q reports  $(10.44 \pm 2.37^{+3.05}_{-1.90}) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K_0^*(1430)^0 \bar{K}_0^*(1430)^0 \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K_0^*(1430)^0 \bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}} \quad \Gamma_{10}/\Gamma$$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>8.0^{+2.0}_{-2.4} \pm 0.2</math></b>	62	<sup>1</sup> ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

<sup>1</sup> ABLIKIM 05Q reports  $(8.49 \pm 1.66^{+1.32}_{-1.99}) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K_0^*(1430)^0 \bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K_1(1270)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}} \quad \Gamma_{11}/\Gamma$$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>6.3 \pm 1.9 \pm 0.1</math></b>	68	<sup>1</sup> ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

<sup>1</sup> ABLIKIM 05Q reports  $(6.66 \pm 1.31^{+1.60}_{-1.51}) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K_1(1270)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. The measurement assumes  $B(K_1(1270) \rightarrow K\rho(770)) = 42 \pm 6\%$ .

$$\Gamma(K_1(1400)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}} \quad \Gamma_{12}/\Gamma$$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;2.7</b>	90	<sup>1</sup> ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

<sup>1</sup> ABLIKIM 05Q reports  $< 2.85 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K_1(1400)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to

our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ . The measurement assumes  $B(K_1(1400) \rightarrow K^*(892)\pi) = 94 \pm 6\%$ .

### $\Gamma(f_0(980)f_0(980))/\Gamma_{\text{total}}$ $\Gamma_{13}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>16.2^{+10.4}_{-9.0} \pm 0.3</math></b>	28	<sup>1</sup> ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

<sup>1</sup> ABLIKIM 05Q reports  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980)f_0(980))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (1.59 \pm 0.50^{+0.89}_{-0.72}) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. One of the  $f_0(980)$  mesons is identified via decay to  $\pi^+ \pi^-$  while the other via  $K^+ K^-$  decay.

### $\Gamma(f_0(980)f_0(2200))/\Gamma_{\text{total}}$ $\Gamma_{14}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>7.9^{+2.0}_{-2.5} \pm 0.2</math></b>	77	<sup>1</sup> ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

<sup>1</sup> ABLIKIM 05Q reports  $(8.42 \pm 1.42^{+1.65}_{-2.29}) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980)f_0(2200))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. The  $f_0$  mesons are identified via  $f_0(980) \rightarrow \pi^+ \pi^-$  and  $f_0(2200) \rightarrow K^+ K^-$  decays.

### $\Gamma(f_0(1370)f_0(1370))/\Gamma_{\text{total}}$ $\Gamma_{15}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.7</b>	90	<sup>1</sup> ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

<sup>1</sup> ABLIKIM 05Q reports  $< 2.9 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1370))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ . One of the  $f_0(1370)$  mesons is identified via decay to  $\pi^+ \pi^-$  while the other via  $K^+ K^-$  decay. Both branching fractions for these  $f_0$  decays are implicitly included in the quoted result.

### $\Gamma(f_0(1370)f_0(1500))/\Gamma_{\text{total}}$ $\Gamma_{16}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.7</b>	90	<sup>1</sup> ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

<sup>1</sup> ABLIKIM 05Q reports  $< 1.8 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1500))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ . The  $f_0$  mesons are identified via  $f_0(1370) \rightarrow \pi^+ \pi^-$  and  $f_0(1500) \rightarrow K^+ K^-$  decays. Both branching fractions for these  $f_0$  decays are implicitly included in the quoted result.

$\Gamma(f_0(1370)f_0(1710))/\Gamma_{\text{total}}$  $\Gamma_{17}/\Gamma$ 

<u>VALUE</u> (units $10^{-4}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>6.7^{+3.5}_{-2.3} \pm 0.1</math></b>	61	<sup>1</sup> ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

<sup>1</sup> ABLIKIM 05Q reports  $(7.12 \pm 1.85^{+3.28}_{-1.68}) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1710))/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \approx (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. The  $f_0$  mesons are identified via  $f_0(1370) \rightarrow \pi^+\pi^-$  and  $f_0(1710) \rightarrow K^+K^-$  decays. Both branching fractions for these  $f_0$  decays are implicitly included in the quoted result.

 $\Gamma(f_0(1500)f_0(1370))/\Gamma_{\text{total}}$  $\Gamma_{18}/\Gamma$ 

<u>VALUE</u> (units $10^{-4}$ )	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;1.3</b>	90	<sup>1</sup> ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

<sup>1</sup> ABLIKIM 05Q reports  $< 1.4 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500)f_0(1370))/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \approx (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ . The  $f_0$  mesons are identified via  $f_0(1500) \rightarrow \pi^+\pi^-$  and  $f_0(1370) \rightarrow K^+K^-$  decays. Both branching fractions for these  $f_0$  decays are implicitly included in the quoted result.

 $\Gamma(f_0(1500)f_0(1500))/\Gamma_{\text{total}}$  $\Gamma_{19}/\Gamma$ 

<u>VALUE</u> (units $10^{-4}$ )	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.5</b>	90	<sup>1</sup> ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

<sup>1</sup> ABLIKIM 05Q reports  $< 0.55 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500)f_0(1500))/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \approx (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ . One of the  $f_0(1500)$  is identified via decay to  $\pi^+\pi^-$  while the other via  $K^+K^-$  decay. Both branching fractions for these  $f_0$  decays are implicitly included in the quoted result.

 $\Gamma(f_0(1500)f_0(1710))/\Gamma_{\text{total}}$  $\Gamma_{20}/\Gamma$ 

<u>VALUE</u> (units $10^{-4}$ )	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.7</b>	90	<sup>1</sup> ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

<sup>1</sup> ABLIKIM 05Q reports  $< 0.73 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500)f_0(1710))/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \approx (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ . The  $f_0$  mesons are identified via  $f_0(1500) \rightarrow \pi^+\pi^-$  and  $f_0(1710) \rightarrow K^+K^-$  decays. Both branching fractions for these  $f_0$  decays are implicitly included in the quoted result.

 $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{21}/\Gamma$ 

<u>VALUE</u> (units $10^{-3}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>8.61 \pm 0.13 \pm 0.94</math></b>	9.0k	<sup>1</sup> ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $\mathcal{B}(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.68 \pm 0.31)\%$ .

$\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_{22}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.22±0.10±0.43</b>	2.7k	1 ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.68 \pm 0.31)\%$ .

 $\Gamma(K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{23}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.56±0.09±0.01</b>	213.5	1 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $0.59 \pm 0.05 \pm 0.08 \pm 0.03$  % from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{24}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.49±0.33±0.05</b>	401.7	1 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $2.64 \pm 0.15 \pm 0.31 \pm 0.14$  % from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{25}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.21±0.21±0.02</b>	179.7	1 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $1.28 \pm 0.16 \pm 0.15 \pm 0.07$  % from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{26}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.46±0.12±0.01</b>	64.1	1 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $0.49 \pm 0.10 \pm 0.07 \pm 0.03$  % from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_{27}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.7±1.0±0.1</b>	$152 \pm 14$	1 ABLIKIM	050 BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$

<sup>1</sup> ABLIKIM 050 reports  $[\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (0.558 \pm 0.051 \pm 0.089) \times 10^{-3}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(K^+ K^- \eta\pi^0)/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{28}/\Gamma$
<b>0.30±0.07±0.01</b>	56.4	1 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$	

<sup>1</sup> HE 08B reports  $0.32 \pm 0.05 \pm 0.05 \pm 0.02$  % from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \eta\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{29}/\Gamma$
<b>12.0±1.8 OUR EVALUATION</b>		Treating systematic error as correlated.			


**19.6±2.1 OUR AVERAGE** Error includes scale factor of 3.3. See the ideogram below.

$20.8 \pm 0.1 \pm 0.7$	145K	1 ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$	
$11.7 \pm 1.0 \pm 1.9$		2 BAI	99B BES	$\psi(2S) \rightarrow \gamma\chi_{c0}$	
$12.5 \pm 2.9 \pm 0.5$		2 TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma\chi_{c0}$	

<sup>1</sup> ABLIKIM 22Q reports  $(2.080 \pm 0.006 \pm 0.068) \times 10^{-2}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow 3(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.2) \times 10^{-2}$ .

<sup>2</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+ \pi^-) = (32.6 \pm 0.5)\%$ .

WEIGHTED AVERAGE  
 $19.6 \pm 2.1$  (Error scaled by 3.3)



### $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID
<b>0.0075±0.0016 OUR FIT</b>	

### $\Gamma_{30}/\Gamma$

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}}$	$\Gamma_{31}/\Gamma$			
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.72^{+0.60}_{-0.54} \pm 0.04$	64	<sup>1</sup> ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.56 \pm 0.40 \pm 0.03$      $30 \pm 6$     <sup>2,3</sup> ABLIKIM    04H BES    Repl. by ABLIKIM 050

<sup>1</sup> ABLIKIM 05Q reports  $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (0.168 \pm 0.035^{+0.047}_{-0.040}) \times 10^{-3}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Assumes  $B(K^*(892)^0 \rightarrow K^- \pi^+) = 2/3$ .

<sup>3</sup> ABLIKIM 04H reports  $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (1.53 \pm 0.29 \pm 0.26) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$   
VALUE (units  $10^{-3}$ ) **8.51 ± 0.33 OUR FIT** DOCUMENT ID

$\Gamma(\pi^0 \eta_c)/\Gamma_{\text{total}}$	$CL\%$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_{35}/\Gamma$
$<1.6 \times 10^{-3}$	90	<sup>1</sup> ABLIKIM	15N	BES3	$\psi(2s) e^+ e^- \rightarrow \gamma \pi^0 \eta_c$
<sup>1</sup> Using $B(\eta_c \rightarrow K_c^0 K^\pm \pi^\mp) \times B(K_c^0 \rightarrow \pi^+ \pi^-) \times B(\pi^0 \rightarrow \gamma \gamma) = (1.66 \pm 0.11) \times 10^{-2}$ .					

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	DOCUMENT ID
<u>3.01 ± 0.19 OUR FIT</u>	

$\Gamma(\eta\eta)/\Gamma(\pi\pi)$	$\Gamma_{36}/\Gamma_{32}$		
VALUE	DOCUMENT ID	TECN	COMMENT

We have made the following plots of  $\chi^2$  versus fit parameters.

0.26 ± 0.09 ± 0.03      <sup>1</sup> ANDREOTTI 05C E835  $\bar{p}p \rightarrow 2 \text{ mesons}$   
 0.24 ± 0.10 ± 0.08      <sup>1</sup> BAI 03C BES  $\psi(2S) \rightarrow 5\gamma$

<sup>1</sup>We have multiplied  $\pi^0\pi^0$  measurement by 3 to obtain  $\pi\pi$

$\Gamma(\eta\eta')/\Gamma_{\text{total}}$	$\Gamma_{37}/\Gamma$				
<i>VALUE</i> (units $10^{-5}$ )	<i>CL%</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<b>9.1±1.1±0.2</b>		85	1 ABLIKIM	17AI BES3	$\psi(2S) \rightarrow \gamma\eta'\eta$

- • • We do not use the following data for averages, fits, limits, etc. • • •

$<24$	90	$35 \pm 13$	<sup>2</sup> ASNER	09	CLEO	$\psi(2S) \rightarrow \gamma\eta'\eta$
$<50$	90		<sup>3</sup> ADAMS	07	CLEO	$\psi(2S) \rightarrow \gamma\chi_{c0}$

<sup>1</sup> ABLIKIM 17AI reports  $(8.92 \pm 0.84 \pm 0.65) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm$

$0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ASNER 09 reports  $< 0.25 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

<sup>3</sup> Superseded by ASNER 09. ADAMS 07 reports  $< 0.5 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

### $\Gamma(\eta'\eta')/\Gamma_{\text{total}}$

### $\Gamma_{38}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.17 \pm 0.12</math> OUR AVERAGE</b>				

$2.23 \pm 0.13 \pm 0.05$       2.5k      <sup>1</sup> ABLIKIM      17AI BES3       $\psi(2S) \rightarrow \gamma\eta'\eta'$

$2.00 \pm 0.21 \pm 0.04$       0.4k      <sup>2</sup> ASNER      09 CLEO       $\psi(2S) \rightarrow \gamma\eta'\eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.60 \pm 0.41 \pm 0.03$       23      <sup>3</sup> ADAMS      07 CLEO       $\psi(2S) \rightarrow \gamma\chi_{c0}$

<sup>1</sup> ABLIKIM 17AI reports  $(2.19 \pm 0.03 \pm 0.14) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ASNER 09 reports  $(2.12 \pm 0.13 \pm 0.21) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Superseded by ASNER 09. ADAMS 07 reports  $(1.7 \pm 0.4 \pm 0.2) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 0.0922 \pm 0.0011 \pm 0.0046$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(\omega\omega)/\Gamma_{\text{total}}$

### $\Gamma_{39}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.97 \pm 0.11</math> OUR AVERAGE</b>				

$0.93 \pm 0.11 \pm 0.02$       991      <sup>1</sup> ABLIKIM      11K BES3       $\psi(2S) \rightarrow \gamma$  hadrons

$2.16 \pm 0.66 \pm 0.04$        $38.1 \pm 9.6$       <sup>2</sup> ABLIKIM      05N BES2       $\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma 6\pi$

<sup>1</sup> ABLIKIM 11K reports  $(0.95 \pm 0.03 \pm 0.11) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ABLIKIM 05N reports  $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (0.212 \pm 0.053 \pm 0.037) \times 10^{-3}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega\phi)/\Gamma_{\text{total}}$  $\Gamma_{40}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.41 \pm 0.13 \pm 0.03</math></b>	486	1 ABLIKIM	19J BES3	$\psi(2S) \rightarrow \gamma$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.18 \pm 0.22 \pm 0.02$	76	2,3 ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
--------------------------	----	-------------	----------	---------------------------------------

<sup>1</sup> ABLIKIM 19J reports  $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (13.83 \pm 0.70 \pm 1.01) \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ABLIKIM 11K reports  $(1.2 \pm 0.1 \pm 0.2) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Superseded by ABLIKIM 19J.

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{41}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.94 \pm 0.06 \pm 0.20</math></b>	1.4k	1 ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.68 \pm 0.31)\%$ .

 $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{42}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>
<b><math>6.05 \pm 0.31</math> OUR FIT</b>	

 $\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$  $\Gamma_{43}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>
<b><math>3.16 \pm 0.17</math> OUR FIT</b>	

 $\Gamma(K_S^0 K_S^0)/\Gamma(\pi\pi)$  $\Gamma_{43}/\Gamma_{32}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.371 \pm 0.023</math> OUR FIT</b>			

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.31 \pm 0.05 \pm 0.05$	1,2 CHEN	07B BELL	$e^+ e^- \rightarrow e^+ e^- \chi_{c0}$
--------------------------	----------	----------	---

<sup>1</sup> Using  $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  from the  $\pi^+\pi^-$  measurement of NAKAZAWA 05 rescaled by 3/2 to convert to  $\pi\pi$ .

<sup>2</sup> Not independent from other measurements.

 $\Gamma(K_S^0 K_S^0)/\Gamma(K^+ K^-)$  $\Gamma_{43}/\Gamma_{42}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.52 \pm 0.04</math> OUR FIT</b>			

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.49 \pm 0.07 \pm 0.08$	1,2 CHEN	07B BELL	$e^+ e^- \rightarrow e^+ e^- \chi_{c0}$
--------------------------	----------	----------	---

<sup>1</sup> Using  $\Gamma(K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  from NAKAZAWA 05.

<sup>2</sup> Not independent from other measurements.

$\Gamma(\pi^+\pi^-\eta)/\Gamma_{\text{total}}$  $\Gamma_{44}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.20	90	1 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.0	90	<sup>2</sup> ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$
------	----	----------------------	----------	---

<sup>1</sup> ATHAR 07 reports  $< 0.21 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

<sup>2</sup> ABLIKIM 06R reports  $< 1.1 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

 $\Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$  $\Gamma_{45}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	1 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>1</sup> ATHAR 07 reports  $< 0.38 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

 $\Gamma(\bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{46}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.09	90	1 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.7	90	<sup>2,3</sup> ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$
<0.7	90	<sup>3,4</sup> BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c0}$

<sup>1</sup> ATHAR 07 reports  $< 0.10 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

<sup>2</sup> ABLIKIM 06R reports  $< 0.70 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

<sup>3</sup> We have multiplied the  $K_S^0 K^+ \pi^-$  measurement by a factor of 2 to convert to  $\bar{K}^0 K^+ \pi^-$ .

<sup>4</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.4 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$ .

 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{47}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.06	90	1 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>1</sup> ATHAR 07 reports  $< 0.06 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

$\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$  $\Gamma_{48}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.23</b>	90	1 ATTHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>1</sup> ATTHAR 07 reports  $< 0.24 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

 $\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}$  $\Gamma_{49}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.41 ± 0.47 ± 0.03</b>	$16.8 \pm 4.8$	1 ABLIKIM	050	BES2 $\psi(2S) \rightarrow \gamma \chi_{c0}$

<sup>1</sup> ABLIKIM 050 reports  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (0.138 \pm 0.039 \pm 0.025) \times 10^{-3}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}$  $\Gamma_{50}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.8 ± 0.5 ± 0.1</b>	319	1 ABLIKIM	19AA	BES3 $\psi(2S) \rightarrow \gamma 4K_S^0$

<sup>1</sup> Using  $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$ . ABLIKIM 19AA reports  $[\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (5.64 \pm 0.33 \pm 0.37) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value..

 $\Gamma(K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{51}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>
<b>2.82 ± 0.29 OUR FIT</b>	

 $\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$  $\Gamma_{52}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.97 ± 0.25 ± 0.02</b>	38	1 ABLIKIM	06T	BES2 $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>1</sup> ABLIKIM 06T reports  $(1.03 \pm 0.22 \pm 0.15) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\bar{K}^0 K^+ \pi^- \phi + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{53}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>3.68 ± 0.30 ± 0.50</b>	ABLIKIM	15M	BES3 $\psi(2S) \rightarrow \gamma \chi_{c0}$

 $\Gamma(K^+ K^- \pi^0 \phi)/\Gamma_{\text{total}}$  $\Gamma_{54}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.90 ± 0.14 ± 0.32</b>	ABLIKIM	15M	BES3 $\psi(2S) \rightarrow \gamma \chi_{c0}$

$\Gamma(\phi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{55}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.18±0.07±0.13</b>	538	1 ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.68 \pm 0.31)\%$ .

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$  $\Gamma_{56}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>
<b>0.80±0.07 OUR FIT</b>	

 $\Gamma(\phi\phi\eta)/\Gamma_{\text{total}}$  $\Gamma_{57}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.4±0.7±0.6</b>	186.6	1 ABLIKIM	20B BES3	$\psi(2S) \rightarrow \gamma\phi\phi\eta$

<sup>1</sup> ABLIKIM 20B reports  $(8.41 \pm 0.74 \pm 0.62) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \phi\phi\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ .

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$  $\Gamma_{58}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>
<b>2.21±0.08 OUR FIT</b>	

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{59}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.70±0.07 OUR AVERAGE</b>	Error includes scale factor of 1.3.		
0.73±0.06±0.01	<sup>1</sup> ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$
0.56±0.12±0.01	<sup>2</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+h^-h^0$

<sup>1</sup> ONYISI 10 reports  $(7.76 \pm 0.37 \pm 0.51 \pm 0.39) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ATHAR 07 reports  $(0.59 \pm 0.10 \pm 0.08) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$  $\Gamma_{60}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.35±0.04 OUR AVERAGE</b>			
0.35±0.04±0.01	<sup>1</sup> ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$
0.37±0.11±0.01	<sup>2</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+h^-h^0$

<sup>1</sup> ONYISI 10 reports  $(3.73 \pm 0.38 \pm 0.28 \pm 0.19) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ATHAR 07 reports  $(0.39 \pm 0.11 \pm 0.04) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.52±0.06±0.01</b>	<sup>1</sup> ONYISI	10	CLE3 $\psi(2S) \rightarrow \gamma p\bar{p}X$
<sup>1</sup> ONYISI 10 reports $(5.57 \pm 0.48 \pm 0.42 \pm 0.14) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.			

### $\Gamma_{61}/\Gamma$

### $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.0±1.4±0.1</b>	$42 \pm 8$	<sup>1</sup> ABLIKIM	11F	$BES3 \psi(2S) \rightarrow \gamma p\bar{p}K^+ K^-$
<sup>1</sup> ABLIKIM 11F reports $(6.12 \pm 1.18 \pm 0.86) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

### $\Gamma_{62}/\Gamma$

### $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.1 ± 0.7 OUR EVALUATION</b>	Error includes scale factor of 1.4. Treating systematic error as correlated.		
<b>2.1 ± 1.0 OUR AVERAGE</b> Error includes scale factor of 2.0.			
$1.57 \pm 0.21 \pm 0.53$ <sup>1</sup> BAI      99B      BES $\psi(2S) \rightarrow \gamma \chi_{c0}$			
$4.20 \pm 1.15 \pm 0.18$ <sup>1</sup> TANENBAUM      78      MRK1 $\psi(2S) \rightarrow \gamma \chi_{c0}$			
<sup>1</sup> Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.4 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$ .			

### $\Gamma_{63}/\Gamma$

### $\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.104±0.028±0.002</b>	$39.5$	<sup>1</sup> HE	08B	$CLEO e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
<sup>1</sup> HE 08B reports $0.11 \pm 0.02 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

### $\Gamma_{64}/\Gamma$

### $\Gamma(p\bar{p}K^+K^- (\text{non-resonant}))/\Gamma_{\text{total}}$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.22±0.26±0.02</b>	$48 \pm 8$	<sup>1</sup> ABLIKIM	11F	$BES3 \psi(2S) \rightarrow \gamma p\bar{p}K^+ K^-$
<sup>1</sup> ABLIKIM 11F reports $(1.24 \pm 0.20 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}K^+ K^- (\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

### $\Gamma_{65}/\Gamma$

$\Gamma(p\bar{p}K_S^0 K_S^0)/\Gamma_{\text{total}}$  $\Gamma_{66}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;8.8</b>	90	<sup>1</sup> ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c0}\gamma$

<sup>1</sup> Using  $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.2 \pm 0.5)\%$

 $\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{67}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>12.7 \pm 1.1 \pm 0.2</math></b>	5150	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$
$11.2 \pm 3.1 \pm 0.2$		<sup>2</sup> ABLIKIM	06I BES2	$\psi(2S) \rightarrow \gamma p\pi^- X$

<sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.26 \pm 0.02 \pm 0.11) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ABLIKIM 06I reports  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.10 \pm 0.24 \pm 0.18) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$  $\Gamma_{68}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>13.7 \pm 1.2 \pm 0.3</math></b>	5808	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+$

<sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.34 \pm 0.03 \pm 0.11) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{69}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>23.4 \pm 2.0 \pm 0.5</math></b>	2480	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-\pi^0$

<sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (2.29 \pm 0.08 \pm 0.18) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{70}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>22.1 \pm 1.8 \pm 0.5</math></b>	2757	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+\pi^0$

<sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (2.16 \pm 0.07 \pm 0.16) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$  $\Gamma_{71}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>
<b><math>3.59 \pm 0.15</math> OUR FIT</b>	

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{72}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>118±12±2</b>		426	<sup>1</sup> ABLIKIM	12I	BES3 $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<400	90	<sup>2</sup> ABLIKIM	06D	BES2	$\psi(2S) \rightarrow \chi_{c0}\gamma$
------	----	----------------------	-----	------	--

<sup>1</sup> ABLIKIM 12I reports  $(119.0 \pm 6.4 \pm 11.4) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Using  $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.2 \pm 0.5)\%$

 $\Gamma(\Lambda\bar{\Lambda}\pi^+(\text{non-resonant}))/\Gamma_{\text{total}}$  $\Gamma_{73}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;50</b>	90	<sup>1</sup> ABLIKIM	12I	BES3 $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

<sup>1</sup> ABLIKIM 12I reports  $< 54 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+(\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

 $\Gamma(\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{75}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;50</b>	90	<sup>1</sup> ABLIKIM	12I	BES3 $\psi(2S) \rightarrow \gamma\Sigma(1385)^+\bar{\Lambda}\pi^-$

<sup>1</sup> ABLIKIM 12I reports  $< 55 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

 $\Gamma(\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{76}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;50</b>	90	<sup>1</sup> ABLIKIM	12I	BES3 $\psi(2S) \rightarrow \gamma\Sigma(1385)^-\bar{\Lambda}\pi^+$

<sup>1</sup> ABLIKIM 12I reports  $< 50 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

 $\Gamma(K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{77}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**1.25±0.12 OUR AVERAGE** Error includes scale factor of 1.3.

1.30±0.09±0.03	9k	<sup>1,2</sup> ABLIKIM	13D	BES3 $\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$
1.01±0.19±0.02		<sup>3</sup> ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+h^-h^0$

<sup>1</sup> ABLIKIM 13D reports  $(1.32 \pm 0.03 \pm 0.10) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))$

$= (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Using  $B(\Lambda \rightarrow p\pi^-) = 63.9\%$ .

<sup>3</sup> ATTHAR 07 reports  $(1.07 \pm 0.17 \pm 0.12) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ \bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(K^*(892)^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$

$\Gamma_{79}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.8±0.9±0.1</b>	254	<sup>1</sup> ABLIKIM	19AU BES3	$\psi(2S) \rightarrow \gamma K^*+\bar{p}\Lambda$

<sup>1</sup> ABLIKIM 19AU reports  $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (4.7 \pm 0.7 \pm 0.5) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(K^+\bar{p}\Lambda(1520)+\text{c.c.})/\Gamma_{\text{total}}$

$\Gamma_{80}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.9±0.7±0.1</b>	$62 \pm 12$	<sup>1</sup> ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

<sup>1</sup> ABLIKIM 11F reports  $(3.00 \pm 0.58 \pm 0.50) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+\bar{p}\Lambda(1520)+\text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(nK_S^0\bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$

$\Gamma_{78}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.7±0.3±0.4</b>	1284	<sup>1</sup> ABLIKIM	21AV BES3	$\psi(2S) \rightarrow \gamma nK_S^0\bar{\Lambda} + \text{c.c.}$

<sup>1</sup> ABLIKIM 21AV reports  $(6.65 \pm 0.26 \pm 0.41) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow nK_S^0\bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 0.0979 \pm 0.0020$ . Also uses  $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = (63.9 \pm 0.5)\%$  and  $B(K_S^0 \rightarrow \pi^+\pi^-) = (69.20 \pm 0.05)\%$ .

### $\Gamma(\Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}$

$\Gamma_{81}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.1±1.2±0.1</b>	$28 \pm 10$	<sup>1</sup> ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

<sup>1</sup> ABLIKIM 11F reports  $(3.18 \pm 1.11 \pm 0.53) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$

$\Gamma_{82}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.68±0.32 OUR AVERAGE</b>				
4.82±0.34±0.10	1046	<sup>1</sup> ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma\Sigma^0\bar{\Sigma}^0$
4.2 ± 0.7 ± 0.1	$78 \pm 10$	<sup>2</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma\Sigma^0\bar{\Sigma}^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.7 \pm 0.5 \pm 0.1$       243      <sup>3,4</sup> ABLIKIM      13H      BES3       $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

<sup>1</sup> ABLIKIM 18V reports  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (4.72 \pm 0.18 \pm 0.28) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> NAIK 08 reports  $(4.41 \pm 0.56 \pm 0.47) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> ABLIKIM 13H reports  $(4.78 \pm 0.34 \pm 0.39) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> Superseded by ABLIKIM 18v

### $\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}$

### $\Gamma_{85}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.6 ± 0.8 OUR AVERAGE</b>		Error includes scale factor of 2.6.		
$5.10 \pm 0.35 \pm 0.10$	747	<sup>1</sup> ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$
$3.1 \pm 0.7 \pm 0.1$	$39 \pm 7$	<sup>2</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.5 \pm 0.5 \pm 0.1$       148      <sup>3,4</sup> ABLIKIM      13H      BES3       $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

<sup>1</sup> ABLIKIM 18V reports  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (4.99 \pm 0.24 \pm 0.24) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> NAIK 08 reports  $(3.25 \pm 0.57 \pm 0.43) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> ABLIKIM 13H reports  $(4.54 \pm 0.42 \pm 0.30) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> Superseded by ABLIKIM 18v

### $\Gamma(\Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}$

### $\Gamma_{86}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.1 ± 0.2 ± 0.4</b>	2143	<sup>1</sup> ABLIKIM	20I BES3	$\psi(2S) \rightarrow \gamma \Sigma^- \bar{\Sigma}^+$

<sup>1</sup> ABLIKIM 20I reports  $(5.13 \pm 0.24 \pm 0.41) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (9.79 \pm 0.20) \times 10^{-2}$ .

$\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$  $\Gamma_{87}/\Gamma$ 

<u>VALUE</u> (units $10^{-5}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>16.2 \pm 5.8 \pm 0.3</math></b>	27	<sup>1</sup> ABLIKIM	12I	BES3 $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

<sup>1</sup> ABLIKIM 12I reports  $(16.4 \pm 5.7 \pm 1.6) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}]$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$  $\Gamma_{88}/\Gamma$ 

<u>VALUE</u> (units $10^{-5}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>23.2 \pm 6.5 \pm 0.5</math></b>	33	<sup>1</sup> ABLIKIM	12I	BES3 $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

<sup>1</sup> ABLIKIM 12I reports  $(23.5 \pm 6.2 \pm 2.3) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}]$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{89}/\Gamma$ 

<u>VALUE</u> (units $10^{-4}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.94 \pm 0.35 \pm 0.04</math></b>	57	<sup>1</sup> ABLIKIM	15I	BES3 $\psi(2S) \rightarrow \gamma K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$

<sup>1</sup> ABLIKIM 15I reports  $[\Gamma(\chi_{c0}(1P) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (1.90 \pm 0.30 \pm 0.16) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$  $\Gamma_{90}/\Gamma$ 

<u>VALUE</u> (units $10^{-4}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>4.5 \pm 0.5</math> OUR AVERAGE</b>				Error includes scale factor of 1.7.
$4.67 \pm 0.19 \pm 0.26$	1741	<sup>1</sup> ABLIKIM	220	BES3 $\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$

$3.1 \pm 0.8 \pm 0.1$	$23.3 \pm 4.9$	<sup>2</sup> NAIK	08	CLEO $\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$
-----------------------	----------------	-------------------	----	--

<sup>1</sup> ABLIKIM 220 reports  $(4.67 \pm 0.19 \pm 0.26) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.2) \times 10^{-2}]$ .

<sup>2</sup> NAIK 08 reports  $(3.34 \pm 0.70 \pm 0.48) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}]$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$  $\Gamma_{91}/\Gamma$ 

<u>VALUE</u> (units $10^{-4}$ )	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>4.45 \pm 0.19</math> OUR AVERAGE</b>					

$4.43 \pm 0.08 \pm 0.18$	4932	<sup>1</sup> ABLIKIM	220	BES3 $\psi(2S) \rightarrow \gamma \Xi^- \bar{\Xi}^+$
$4.8 \pm 0.7 \pm 0.1$	95	<sup>2</sup> NAIK	08	CLEO $\psi(2S) \rightarrow \gamma \Xi^+ \bar{\Xi}^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<10.3	90	<sup>3</sup> ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c0}\gamma$
<sup>1</sup> ABLIKIM 220 reports $(4.43 \pm 0.08 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.2) \times 10^{-2}$ .				
<sup>2</sup> NAIK 08 reports $(5.14 \pm 0.60 \pm 0.47) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				
<sup>3</sup> Using $\mathcal{B}(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.2 \pm 0.5)\%$				

$\Gamma(\eta_c\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{92}/\Gamma$
VALUE	CL %	DOCUMENT ID	TECN	COMMENT	
<b>&lt; <math>7 \times 10^{-4}</math></b>	90	1,2 ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$	
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>					
$< 41 \times 10^{-4}$	90	1,3 ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$	
<sup>1</sup> Using $1.06 \times 10^8$ $\psi(2S)$ mesons and $\mathcal{B}(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.68 \pm 0.31)\%$ .					
<sup>2</sup> From the $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ decays.					
<sup>3</sup> From the $\eta_c \rightarrow K^+ K^- \pi^0$ decays.					

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi\pi)/\Gamma_{\text{total}}$					$\Gamma_{58}/\Gamma \times \Gamma_{32}/\Gamma$
VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT		
<b>18.8 <math>\pm 1.0</math> OUR FIT</b>					
<b>15.3 <math>\pm 2.4 \pm 0.8</math></b>	<sup>1</sup> ANDREOTTI 03	E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0\pi^0$		

<sup>1</sup> We have multiplied  $\mathcal{B}(p\bar{p}) \cdot \mathcal{B}(\pi^0\pi^0)$  measurement by 3 to obtain  $\mathcal{B}(p\bar{p}) \cdot \mathcal{B}(\pi\pi)$ .

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi^0\eta)/\Gamma_{\text{total}}$					$\Gamma_{58}/\Gamma \times \Gamma_{33}/\Gamma$
VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT		
<b>&lt;0.4</b>	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0\eta$		

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi^0\eta')/\Gamma_{\text{total}}$					$\Gamma_{58}/\Gamma \times \Gamma_{34}/\Gamma$
VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT		
<b>&lt;2.5</b>	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0\eta$		

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\eta\eta)/\Gamma_{\text{total}}$					$\Gamma_{58}/\Gamma \times \Gamma_{36}/\Gamma$
VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT		
<b>6.7 <math>\pm 0.5</math> OUR FIT</b>					
<b>4.0 <math>\pm 1.2^{+0.5}_{-0.3}</math></b>	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \eta\eta$		

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\eta\eta')/\Gamma_{\text{total}}$					$\Gamma_{58}/\Gamma \times \Gamma_{37}/\Gamma$
VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT		
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>					
<b><math>2.1^{+2.3}_{-1.5}</math></b>	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0\eta$		

---

RADIATIVE DECAYS

---

 $\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$  $\Gamma_{93}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**1.40±0.05 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.25 \pm 0.16 \pm 2.15$	12k	<sup>1</sup> ABLIKIM	17U BES3	$e^+ e^- \rightarrow \gamma X$
$2.0 \pm 0.2 \pm 0.2$		<sup>2</sup> ADAM	05A CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$

<sup>1</sup> Not independent from  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))$  and the product  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))$  also measured in ABLIKIM 17U.

<sup>2</sup> Uses  $B(\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow \gamma \gamma J/\psi)$  from ADAM 05A and  $B(\psi(2S) \rightarrow \gamma \chi_{c0})$  from ATHAR 04.

 $\Gamma(\gamma \rho^0)/\Gamma_{\text{total}}$  $\Gamma_{94}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

< 9	90	$1.2 \pm 4.5$	<sup>1</sup> BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma \gamma \rho^0$
-----	----	---------------	----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<10	90	$6 \pm 12$	<sup>2</sup> ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma \gamma \rho^0$
-----	----	------------	----------------------	----------	---

<sup>1</sup> BENNETT 08A reports  $< 9.6 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma \rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

<sup>2</sup> ABLIKIM 11E reports  $< 10.5 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma \rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

 $\Gamma(\gamma \omega)/\Gamma_{\text{total}}$  $\Gamma_{95}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

< 8	90	$0.0 \pm 2.8$	<sup>1</sup> BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma \gamma \omega$
-----	----	---------------	----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13	90	$5 \pm 11$	<sup>2</sup> ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma \gamma \omega$
-----	----	------------	----------------------	----------	---

<sup>1</sup> BENNETT 08A reports  $< 8.8 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma \omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

<sup>2</sup> ABLIKIM 11E reports  $< 12.9 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma \omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

 $\Gamma(\gamma \phi)/\Gamma_{\text{total}}$  $\Gamma_{96}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

< 6	90	$0.1 \pm 1.6$	<sup>1</sup> BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma \gamma \phi$
-----	----	---------------	----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16	90	$15 \pm 7$	<sup>2</sup> ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma \gamma \phi$
-----	----	------------	----------------------	----------	---

<sup>1</sup> BENNETT 08A reports  $< 6.4 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

<sup>2</sup> ABLIKIM 11E reports  $< 16.2 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

## $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{97}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>2.04 \pm 0.09</math> OUR FIT</b>				

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7 90 <sup>1</sup> WICHT 08 BELL  $B^\pm \rightarrow K^\pm \gamma\gamma$

<sup>1</sup> WICHT 08 reports  $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c0} K^+)] < 0.11 \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c0} K^+) = 1.51 \times 10^{-4}$ .

## $\Gamma(e^+ e^- J/\psi(1S))/\Gamma_{\text{total}}$ $\Gamma_{98}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

$1.54 \pm 0.33 \pm 0.03$  56 <sup>1,2</sup> ABLIKIM 17I BES3  $\psi(2S) \rightarrow \gamma e^+ e^- J/\psi$

<sup>1</sup> ABLIKIM 17I reports  $(1.51 \pm 0.30 \pm 0.13) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow e^+ e^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Not independent from other measurements reported by ABLIKIM 17I

## $\Gamma(e^+ e^- J/\psi(1S))/\Gamma(\gamma J/\psi(1S))$ $\Gamma_{98}/\Gamma_{93}$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>9.5 \pm 1.9 \pm 0.7</math></b>	56	<sup>1</sup> ABLIKIM	17I	BES3 $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

<sup>1</sup> Uses  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (15.8 \pm 0.3 \pm 0.6) \times 10^{-4}$  from ABLIKIM 17N and accounts for common systematic errors.

## $\Gamma(\mu^+ \mu^- J/\psi(1S))/\Gamma(e^+ e^- J/\psi(1S))$ $\Gamma_{99}/\Gamma_{98}$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.14</b>	90	<9.5	ABLIKIM	19Z	BES3 $\psi(2S) \rightarrow \gamma \chi_c \rightarrow \gamma (\mu^+ \mu^- J/\psi)$

## $\Gamma(\gamma\gamma)/\Gamma(\gamma J/\psi(1S))$ $\Gamma_{97}/\Gamma_{93}$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>1.45 \pm 0.08</math> OUR FIT</b>			

### 2.0 ± 0.4 OUR AVERAGE

$2.2 \pm 0.4$  <sup>+0.1</sup> <sub>-0.2</sub> <sup>1</sup> ANDREOTTI 04 E835  $p\bar{p} \rightarrow \chi_{c0} \rightarrow \gamma\gamma$

$1.45 \pm 0.74$  <sup>2</sup> AMBROGIANI 00B E835  $\bar{p}p \rightarrow \chi_{c2} \rightarrow \gamma\gamma, \gamma J/\psi$

<sup>1</sup> The values of  $B(p\bar{p})B(\gamma\gamma)$  and  $B(\gamma\gamma)B(\gamma J/\psi)$  measured by ANDREOTTI 04 are not independent. The latter is used in the fit because of smaller systematics.

<sup>2</sup> Calculated by us using  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$ .

## $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$ $\Gamma_{58}/\Gamma \times \Gamma_{93}/\Gamma$

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>31.1 \pm 1.5</math> OUR FIT</b>				

### 28.2 ± 2.1 OUR AVERAGE

$28.0 \pm 1.9 \pm 1.3$  392 <sup>1,2,3</sup> BAGNASCO 02 E835  $\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi \gamma$

$29.3$  <sup>+5.7</sup> <sub>-4.7</sub>  $\pm 1.5$  89 <sup>1,2</sup> AMBROGIANI 99B  $\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi \gamma$

<sup>1</sup> Values in  $(\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}})$  and  $(\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}})$  are not independent. The latter is used in the fit since it is less correlated to the total width.

<sup>2</sup> Calculated by us using  $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$ .

<sup>3</sup> Recalculated by ANDREOTTI 05A.

$$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{58}/\Gamma \times \Gamma_{97}/\Gamma$$

VALUE (units $10^{-8}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.52 ± 0.27 OUR FIT</b>			

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.52 \pm 1.18^{+0.48}_{-0.72}$	<sup>1</sup> ANDREOTTI 04 E835	$p\bar{p} \rightarrow \chi_{c0} \rightarrow \gamma\gamma$
---------------------------------	--------------------------------	---

<sup>1</sup> The values of  $B(p\bar{p})B(\gamma\gamma)$  and  $B(\gamma\gamma)B(\gamma J/\psi)$  measured by ANDREOTTI 04 are not independent. The latter is used in the fit because of smaller systematics.

### $\chi_{c0}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

$$\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}} \quad \Gamma_{58}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{166}^{\psi(2S)}$$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**21.7 ± 0.9 OUR FIT**

**23.7 ± 1.0 OUR AVERAGE**

$23.7 \pm 0.8 \pm 0.9$	1222	ABLIKIM	13V BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$
$23.7 \pm 1.4 \pm 1.4$	383 ± 22	<sup>1</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
$23.6^{+3.7}_{-3.4} \pm 3.4$	$89.5^{+14}_{-13}$	BAI	04F BES	$\psi(2S) \rightarrow \gamma\chi_{c0}(1P) \rightarrow \gamma\bar{p}p$

<sup>1</sup> Calculated by us. NAIK 08 reports  $B(\chi_{c0} \rightarrow p\bar{p}) = (25.7 \pm 1.5 \pm 1.5 \pm 1.3) \times 10^{-5}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$ .

$$\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{58}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**6.25 ± 0.26 OUR FIT**

<b>4.6 ± 1.9</b>	<sup>1</sup> BAI	98I BES	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma\bar{p}p$
------------------	------------------	---------	---

<sup>1</sup> Calculated by us. The value for  $B(\chi_{c0} \rightarrow p\bar{p})$  reported in BAI 98I is derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.3 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

$$\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}} \quad \Gamma_{71}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**35.2 ± 1.3 OUR FIT**

**35.1 ± 1.4 OUR AVERAGE** Error includes scale factor of 1.1.

$35.6 \pm 1.0 \pm 1.0$	1486	ABLIKIM	21L BES3	$\psi(2S) \rightarrow \gamma p\pi^-\bar{p}\pi^+$
$31.2 \pm 3.3 \pm 2.0$	131	<sup>1</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$32.0 \pm 1.9 \pm 2.2$	369	<sup>2,3</sup> ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$
------------------------	-----	------------------------	----------	---

<sup>1</sup> Calculated by us. NAIK 08 reports  $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) = (33.8 \pm 3.6 \pm 2.2 \pm 1.7) \times 10^{-5}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$ .

<sup>2</sup> Superseded by ABLIKIM 21L

<sup>3</sup> Calculated by us. ABLIKIM 13H reports  $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) = (33.3 \pm 2.0 \pm 2.6) \times 10^{-5}$  from a measurement of  $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma\chi_{c0})$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.62 \pm 0.31)\%$ .

$$\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{71}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>10.1 \pm 0.4</math> OUR FIT</b>				

**$13.0^{+3.6}_{-3.5} \pm 2.5$**      $15.2^{+4.2}_{-4.0}$     <sup>1</sup> BAI    03E BES     $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

<sup>1</sup> BAI 03E reports [  $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) B(\psi(2S) \rightarrow \gamma\chi_{c0}) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)$  ]  $\times$   $[B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p\bar{p})] = (2.45^{+0.68}_{-0.65} \pm 0.46)\%$ . We calculate from this measurement the presented value using  $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$  and  $B(J/\psi \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$ .

$$\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{74}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.26 \pm 0.30 \pm 0.20</math></b>	67	ABLIKIM	22AO BES3	$\psi(2S) \rightarrow \gamma p\pi^-\bar{p}\pi^+\gamma\gamma$

$$\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{93}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{10}^{\psi(2S)}$$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.138 \pm 0.005</math> OUR FIT</b>				

<b><math>0.147 \pm 0.029</math> OUR AVERAGE</b>	Error includes scale factor of 4.6.
$0.158 \pm 0.003 \pm 0.006$	<sup>1</sup> ABLIKIM    4.8k $\psi(2S) \rightarrow \gamma\gamma J/\psi$
$0.024 \pm 0.015 \pm 0.205$	ABLIKIM    12k $e^+e^- \rightarrow \gamma X$
$0.069 \pm 0.018$	<sup>2</sup> OREGLIA    82    CBAL $\psi(2S) \rightarrow \gamma\chi_{c0}$
$0.4 \pm 0.3$	<sup>3</sup> BRANDELIK    79B    DASP $\psi(2S) \rightarrow \gamma\chi_{c0}$
$0.16 \pm 0.11$	<sup>3</sup> BARTEL    78B    CNTR $\psi(2S) \rightarrow \gamma\chi_{c0}$
$3.3 \pm 1.7$	<sup>4</sup> BIDDICK    77    CNTR $e^+e^- \rightarrow \gamma X$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>	
$0.151 \pm 0.003 \pm 0.010$	<sup>5</sup> ABLIKIM    4.3k $\psi(2S) \rightarrow \gamma\chi_{c0}$
$0.125 \pm 0.007 \pm 0.013$	<sup>6</sup> MENDEZ    560    CLEO $\psi(2S) \rightarrow \gamma\chi_{c0}$
$0.18 \pm 0.01 \pm 0.02$	<sup>7</sup> ADAM    172    CLEO    Repl. by MENDEZ 08

<sup>1</sup> Uses  $B(J/\psi \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$  and  $B(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$ .

<sup>2</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$ .

<sup>3</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$ .

<sup>4</sup> Assumes isotropic gamma distribution.

<sup>5</sup> Superseded by ABLIKIM 17N.

<sup>6</sup> Not independent from other measurements of MENDEZ 08.

<sup>7</sup> Not independent from other values reported by ADAM 05A.

$$\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\text{anything})$$

$$\Gamma_{93}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{10}^{\psi(2S)}$$

$$\Gamma_{93}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{10}^{\psi(2S)} = \Gamma_{93}/\Gamma \times \Gamma_{166}^{\psi(2S)}/(\Gamma_{12}^{\psi(2S)} + \Gamma_{13}^{\psi(2S)} + \Gamma_{14}^{\psi(2S)} + 0.343\Gamma_{167}^{\psi(2S)} + 0.190\Gamma_{168}^{\psi(2S)})$$

<u>VALUE</u> (units $10^{-2}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.224±0.009 OUR FIT</b>				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.201±0.011±0.021	560	<sup>1</sup> MENDEZ	08	CLEO $\psi(2S) \rightarrow \gamma\chi_{c0}$
0.31 ± 0.02 ± 0.03	172	ADAM	05A	CLEO    Repl. by MENDEZ 08

<sup>1</sup> Not independent from other measurements of MENDEZ 08.

$$\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = \frac{\Gamma_{93}/\Gamma}{\Gamma_{166}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

<u>VALUE</u> (units $10^{-2}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.397±0.015 OUR FIT</b>				

0.358±0.020±0.037	560	MENDEZ	08	CLEO $\psi(2S) \rightarrow \gamma\chi_{c0}$
-------------------	-----	--------	----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.55 ± 0.04 ± 0.06	172	<sup>1</sup> ADAM	05A	CLEO    Repl. by MENDEZ 08
--------------------	-----	-------------------	-----	----------------------------

<sup>1</sup> Not independent from other values reported by ADAM 05A.

$$\Gamma(\chi_{c0}(1P) \rightarrow \gamma\gamma) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) / \Gamma_{\text{total}} = \frac{\Gamma_{97}/\Gamma}{\Gamma_{166}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

<u>VALUE</u> (units $10^{-5}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.00±0.08 OUR FIT</b>				

#### **1.95±0.09 OUR AVERAGE**

1.93±0.08±0.05	3.5k	ABLIKIM	17AE	BES3 $\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow 3\gamma$
2.17±0.32±0.10	0.2k	ECKLUND	08A	CLEO $\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow 3\gamma$
3.7 ± 1.8 ± 1.0		LEE	85	CBAL $\psi(2S) \rightarrow \gamma\chi_{c0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.17±0.17±0.12	0.8k	<sup>1</sup> ABLIKIM	12A	BES3 $\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow 3\gamma$
----------------	------	----------------------	-----	---

<sup>1</sup> Superseded by ABLIKIM 17AE.

$$\Gamma(\chi_{c0}(1P) \rightarrow \pi\pi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) / \Gamma_{\text{total}} = \frac{\Gamma_{32}/\Gamma}{\Gamma_{166}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

<u>VALUE</u> (units $10^{-4}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.34±0.29 OUR FIT</b>				

#### **8.80±0.34 OUR AVERAGE**

9.11±0.08±0.65	17k	<sup>1</sup> ABLIKIM	10A	BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$
8.81±0.11±0.43	8.9k	<sup>2</sup> ASNER	09	CLEO $\psi(2S) \rightarrow \gamma\pi^+\pi^-$
8.13±0.19±0.89	2.8k	<sup>3</sup> ASNER	09	CLEO $\psi(2S) \rightarrow \gamma\pi^0\pi^0$

<sup>1</sup> Calculated by us. ABLIKIM 10A reports  $B(\chi_{c0} \rightarrow \pi^0\pi^0) = (3.23 \pm 0.03 \pm 0.23 \pm 0.14) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$ . We have multiplied the  $\pi^0\pi^0$  measurement by 3 to obtain  $\pi\pi$ .

<sup>2</sup> Calculated by us. ASNER 09 reports  $B(\chi_{c0} \rightarrow \pi^+\pi^-) = (6.37 \pm 0.08 \pm 0.31 \pm 0.32) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$ . We have multiplied the  $\pi^+\pi^-$  measurement by 3/2 to obtain  $\pi\pi$ .

<sup>3</sup> Calculated by us. ASNER 09 reports  $B(\chi_{c0} \rightarrow \pi^0\pi^0) = (2.94 \pm 0.07 \pm 0.32 \pm 0.15) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$ . We have multiplied the  $\pi^0\pi^0$  measurement by 3 to obtain  $\pi\pi$ .

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{32}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**24.0±0.8 OUR FIT****20.7±1.7 OUR AVERAGE**

23.9±2.7±4.1	97 ± 11	<sup>1</sup> BAI	03C BES	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma\pi^0\pi^0$
20.2±1.1±1.5	720 ± 32	<sup>2</sup> BAI	98I BES	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma\pi^+\pi^-$

<sup>1</sup> We have multiplied  $\pi^0\pi^0$  measurement by 3 to obtain  $\pi\pi$ .<sup>2</sup> Calculated by us. The value for  $B(\chi_{c0} \rightarrow \pi^+\pi^-)$  reported in BAI 98I is derived using  $B(\psi' \rightarrow \gamma\chi_{c0}) = (9.3 \pm 0.8)\%$  and  $B(\psi' \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D]. We have multiplied  $\pi^+\pi^-$  measurement by 3/2 to obtain  $\pi\pi$ .

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{36}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**2.95±0.18 OUR FIT****3.12±0.19 OUR AVERAGE**

3.23±0.09±0.23	2132	<sup>1</sup> ABLIKIM	10A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$
2.93±0.12±0.29	0.9k	<sup>2</sup> ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\eta\eta$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
2.86±0.46±0.37	48	<sup>3</sup> ADAMS	07 CLEO	$\psi(2S) \rightarrow \gamma\chi_{c0}$

<sup>1</sup> Calculated by us. ABLIKIM 10A reports  $B(\chi_{c0} \rightarrow \eta\eta) = (3.44 \pm 0.10 \pm 0.24 \pm 0.13) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$ .<sup>2</sup> Calculated by us. ASNER 09 reports  $B(\chi_{c0} \rightarrow \eta\eta) = (3.18 \pm 0.13 \pm 0.31 \pm 0.16) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$ .<sup>3</sup> Superseded by ASNER 09. Calculated by us. The value of  $B(\chi_{c0}(1P) \rightarrow \eta\eta)$  reported by ADAMS 07 was derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46)\%$  (ATHAR 04).

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{36}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**0.85 ±0.05 OUR FIT**

<b>0.578±0.241±0.158</b>	BAI	03C BES	$\psi(2S) \rightarrow \gamma\eta\eta$
--------------------------	-----	---------	---------------------------------------

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{42}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**5.92±0.28 OUR FIT**

<b>5.97±0.07±0.32</b>	8.1k	<sup>1</sup> ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma K^+K^-$
-----------------------	------	--------------------	---------	--------------------------------------

<sup>1</sup> Calculated by us. ASNER 09 reports  $B(\chi_{c0} \rightarrow K^+K^-) = (6.47 \pm 0.08 \pm 0.35 \pm 0.32) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$ .

$$\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

$$\Gamma_{42}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**1.71±0.08 OUR FIT****1.63±0.10±0.15**       $774 \pm 38$       <sup>1</sup> BAI      98I BES       $\psi(2S) \rightarrow \gamma K^+ K^-$ 

<sup>1</sup> Calculated by us. The value for  $B(\chi_{c0} \rightarrow K^+ K^-)$  reported by BAI 98I is derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.3 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

$$\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{43}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**3.10±0.16 OUR FIT****3.18±0.17 OUR AVERAGE**

$3.22 \pm 0.07 \pm 0.17$	2.1k	<sup>1</sup> ASNER	09	CLEO $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
$3.02 \pm 0.19 \pm 0.33$	322	ABLIKIM	050	BES2 $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

<sup>1</sup> Calculated by us. ASNER 09 reports  $B(\chi_{c0} \rightarrow K_S^0 K_S^0) = (3.49 \pm 0.08 \pm 0.18 \pm 0.17) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$ .

$$\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

$$\Gamma_{43}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**8.9±0.5 OUR FIT****5.6±0.8±1.3**       $1$  BAI      99B BES       $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$ 

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c0} \rightarrow K_S^0 K_S^0)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.3 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

$$\Gamma(\chi_{c0}(1P) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

$$\Gamma_1/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**6.6±0.5 OUR FIT****6.9±2.4 OUR AVERAGE** Error includes scale factor of 3.8.

$4.4 \pm 0.1 \pm 0.9$	<sup>1</sup> BAI	99B	BES	$\psi(2S) \rightarrow \gamma \chi_{c0}$
$9.3 \pm 0.9$	<sup>2</sup> TANENBAUM	78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$

<sup>1</sup> Calculated by us. The value for  $B(\chi_{c0} \rightarrow 2\pi^+ 2\pi^-)$  reported in BAI 99B is derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.3 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

<sup>2</sup> The value  $B(\psi(1S) \rightarrow \gamma \chi_{c0}) \times B(\chi_{c0} \rightarrow 2\pi^+ 2\pi^-)$  reported in TANENBAUM 78 is derived using  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (4.6 \pm 0.7)\%$ . Calculated by us using  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .

$$\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_8 / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

<u>VALUE</u> (units $10^{-3}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.78 ± 0.14 OUR FIT</b>			
<b>1.64 ± 0.05 ± 0.2</b>	ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$

$$\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

$$\Gamma_8 / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}$$

<u>VALUE</u> (units $10^{-3}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.1 ± 0.4 OUR FIT</b>			
<b>5.8 ± 1.6 OUR AVERAGE</b>	Error includes scale factor of 2.3.		

4.22 ± 0.20 ± 0.97	BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c0}$
7.4 ± 1.0	<sup>1</sup> TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$

<sup>1</sup> The reported value is derived using  $B(\psi(2S) \rightarrow \pi^+ \pi^- J/\psi) \times B(J/\psi \rightarrow \ell^+ \ell^-) = (4.6 \pm 0.7)\%$ . Calculated by us using  $B(J/\psi \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .

$$\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{51} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}$$

<u>VALUE</u> (units $10^{-4}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.76 ± 0.28 OUR FIT</b>				
<b>3.20 ± 0.11 ± 0.41</b>	278	<sup>1</sup> ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c0} \rightarrow 2K^+ 2K^-)$  reported by ABLIKIM 06T was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4)\%$ .

$$\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) /$$

$$\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \Gamma_{51} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}$$

<u>VALUE</u> (units $10^{-4}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.0 ± 0.8 OUR FIT</b>				
<b>6.1 ± 0.8 ± 0.9</b>	<sup>1</sup> BAI	99B BES	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$	

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c0} \rightarrow 2K^+ 2K^-)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.3 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

$$\Gamma(\chi_{c0}(1P) \rightarrow \phi\phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{56} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}$$

<u>VALUE</u> (units $10^{-4}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.78 ± 0.07 OUR FIT</b>				
<b>0.78 ± 0.08 OUR AVERAGE</b>				

0.77 ± 0.03 ± 0.08	612	<sup>1</sup> ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
0.86 ± 0.19 ± 0.12	26	<sup>2</sup> ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c0} \rightarrow \phi\phi)$  reported by ABLIKIM 11K was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31)\%$ .

<sup>2</sup> Calculated by us. The value of  $B(\chi_{c0} \rightarrow \phi\phi)$  reported by ABLIKIM 06T was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4)\%$ .

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{56}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.25±0.21 OUR FIT</b>			

**2.6 ±1.0 ±1.1**      <sup>1</sup> BAI      99B BES       $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c0} \rightarrow \phi\phi)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.3 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+\bar{p}K_S^0 + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{83}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>3.45±0.17±0.19</b>	493	<sup>1</sup> ABLIKIM	19BB BES3	$\psi(2S) \rightarrow \gamma\Sigma^+\bar{p}K_S^0 + \text{c.c.}$

<sup>1</sup> Calculated by us. ABLIKIM 19BB reports  $B(\chi_c^0 \rightarrow \Sigma^+\bar{p}K_S^0 + \text{c.c.}) = (3.52 \pm 0.19 \pm 0.21) \times 10^{-4}$  using  $B(\psi(2S) \rightarrow \gamma\chi_c^0) = (9.79 \pm 0.20)\%$  and other branching fractions from PDG 18.

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0\bar{p}K^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{84}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.97±0.12±0.14</b>	871	<sup>1</sup> ABLIKIM	20AE BES3	$\psi(2S) \rightarrow \gamma\Sigma^0\bar{p}K^+ + \text{c.c.}$

<sup>1</sup> Calculated by us. ABLIKIM 20AE reports  $B(\chi_c^0 \rightarrow \Sigma^0\bar{p}K^+ + \text{c.c.}) = (3.03 \pm 0.12 \pm 0.15) \times 10^{-4}$  using  $B(\psi(2S) \rightarrow \gamma\chi_c^0) = (9.79 \pm 0.20)\%$  and other branching fractions from PDG 20.

## $\chi_{c0}(1P)$ REFERENCES

ABLIKIM	22AO PR D106 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22O JHEP 2206 074	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22Q PR D106 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AV JHEP 2111 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21L PR D103 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20AE PR D102 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20B PR D101 012012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20I PR D101 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	20 PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)
ABLIKIM	19AA PR D99 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AU PR D100 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19BB PR D100 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19J PR D99 012015	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19Z PR D99 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18V PR D97 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	18 PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)
AAIJ	17BB EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17AE PR D96 092007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AI PR D96 112006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17I PRL 118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17N PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17U PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16 CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
ABLIKIM	15I PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15M PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15N PR D91 112018	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13B PR D87 012002	M. Ablikim <i>et al.</i>	(BESIII Collab.)

ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
ABLIKIM	12A	PR D85 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LIU	12B	PRL 108 232001	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
ABLIKIM	10A	PR D81 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)
ASNER	09	PR D79 072007	D.M. Asner <i>et al.</i>	(CLEO Collab.)
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)
BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
ADAMS	07	PR D75 071101	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
CHEN	07B	PL B651 15	W.T. Chen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05N	PL B630 7	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
ANDREOTTI	05C	PR D72 112002	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
NAKAZAWA	05	PL B615 39	H. Nakazawa <i>et al.</i>	(BELLE Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ABLIKIM	04G	PR D70 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ANDREOTTI	04	PL B584 16	M. Andreotti <i>et al.</i>	(E835 Collab.)
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)
ANDREOTTI	03	PRL 91 091801	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
BAGNASCO	02	PL B533 237	S. Bagnasco <i>et al.</i>	(FNAL E835 Collab.)
EISENSTEIN	01	PRL 87 061801	B.I. Eisenstein <i>et al.</i>	(CLEO Collab.)
AMBROGIANI	00B	PR D62 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
AMBROGIANI	99B	PRL 83 2902	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)
Also		Private Comm.	G. Trilling	(LBL, UCB)
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)