

$\eta_c(2S)$

$I^G(J^{PC}) = 0^+(0^{-+})$

Quantum numbers are quark model predictions.

## $\eta_c(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3637.7±1.1 OUR AVERAGE</b>				Error includes scale factor of 1.2.
3643.4±2.3±4.4	569	ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$
3635.1±3.7±2.9	106	XU	18 BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
3633.6±1.7±0.6	106	<sup>1</sup> AAIJ	17ADLHCb	$p p \rightarrow B^+ X \rightarrow p \bar{p} K^+ X$
3636.4±4.1±0.7	365	<sup>2</sup> AAIJ	17BBLHCb	$p p \rightarrow b \bar{b} X \rightarrow 2(K^+ K^-) X$
3637.0±5.7±3.4	178	<sup>3,4</sup> LEES	14E BABR	$\gamma \gamma \rightarrow K^+ K^- \pi^0$
3635.1±5.8±2.1	47	<sup>3,5</sup> LEES	14E BABR	$\gamma \gamma \rightarrow K^+ K^- \eta$
3646.9±1.6±3.6	57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
3637.6±2.9±1.6	127 ± 18	<sup>6</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K\pi, K K \pi^0$
3638.5±1.5±0.8	624	<sup>3</sup> DEL-AMO-SA..11M BABR		$\gamma \gamma \rightarrow K_S^0 K^\pm \pi^\mp$
3640.5±3.2±2.5	1201	<sup>3</sup> DEL-AMO-SA..11M BABR		$\gamma \gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
3636.1 <sup>+3.9 +0.7</sup> <sub>-4.2 -2.0</sub>	128	<sup>7</sup> VINOKUROVA 11	BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
3626 ± 5 ± 6	311	<sup>8</sup> ABE	07 BELL	$e^+ e^- \rightarrow J/\psi(c\bar{c})$
3645.0±5.5 <sup>+4.9</sup> <sub>-7.8</sub>	121 ± 27	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
3642.9±3.1±1.5	61	ASNER	04 CLEO	$\gamma \gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3639 ± 7	98 ± 52	<sup>9</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X c\bar{c}$
3630.8±3.4±1.0	112 ± 24	<sup>10</sup> AUBERT	04D BABR	$\gamma \gamma \rightarrow \eta_c(2S) \rightarrow K \bar{K} \pi$
3654 ± 6 ± 8	39 ± 11	<sup>11</sup> CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
3594 ± 5		<sup>12</sup> EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

<sup>1</sup>AAIJ 17AD report  $m_{\psi(2S)} - m_{\eta_c(2S)} = 52.5 \pm 1.7 \pm 0.6$  MeV. We use the current value  $m_{\psi(2S)} = 3686.097 \pm 0.025$  MeV to obtain the quoted mass.

<sup>2</sup>From a fit of the  $\phi\phi$  invariant mass with the width of  $\eta_c(2S)$  fixed to the PDG 16 value.

<sup>3</sup>Ignoring possible interference with continuum.

<sup>4</sup>With a width fixed to 11.3 MeV.

<sup>5</sup>With a width fixed to 11.3 MeV. Using both  $\eta \rightarrow \gamma \gamma$  and  $\eta \rightarrow \pi^+ \pi^- \pi^0$  decays.

<sup>6</sup>From a simultaneous fit to  $K_S^0 K^\pm \pi^\mp$  and  $K^+ K^- \pi^0$  decay modes.

<sup>7</sup>Accounts for interference with non-resonant continuum.

<sup>8</sup>From a fit of the  $J/\psi$  recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

<sup>9</sup>From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

<sup>10</sup>Superseded by DEL-AMO-SANCHEZ 11M.

<sup>11</sup>Superseded by VINOKUROVA 11.

<sup>12</sup>Assuming mass of  $\psi(2S) = 3686$  MeV.

## $\eta_c(2S)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>13.9 ± 2.6 OUR AVERAGE</b>						
19.8 ± 3.9 ± 3.1		569	ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$	■
9.9 ± 4.8 ± 2.9		57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	
16.9 ± 6.4 ± 4.8		127 ± 18	<sup>1</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K\pi,$ $K K\pi^0$	
13.4 ± 4.6 ± 3.2		624	<sup>2</sup> DEL-AMO-SA..11M BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$		
6.6 ± 8.4 ± 2.6 5.1 – 0.9		128	<sup>3</sup> VINOKUROVA 11 BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$		
6.3 ± 12.4 ± 4.0		61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 23	90	98 ± 52	<sup>4</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$	
22 ± 14		121 ± 27	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$	
17.0 ± 8.3 ± 2.5		112 ± 24	<sup>5</sup> AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$	
< 55	90	39 ± 11	<sup>6</sup> CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$	
< 8.0	95		<sup>7</sup> EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$	

<sup>1</sup> From a simultaneous fit to  $K_S^0 K^\pm \pi^\mp$  and  $K^+ K^- \pi^0$  decay modes.

<sup>2</sup> Ignoring possible interference with continuum.

<sup>3</sup> Accounts for interference with non-resonant continuum.

<sup>4</sup> From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

<sup>5</sup> Superseded by DEL-AMO-SANCHEZ 11M.

<sup>6</sup> For a mass value of  $3654 \pm 6$  MeV. Superseded by VINOKUROVA 11.

<sup>7</sup> For a mass value of  $3594 \pm 5$  MeV

## $\eta_c(2S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ hadrons	not seen	
$\Gamma_2$ $K\bar{K}\pi$	( 1.9 ± 1.2 ) %	
$\Gamma_3$ $K\bar{K}\eta$	( 5 ± 4 ) × 10 <sup>-3</sup>	
$\Gamma_4$ $2\pi^+ 2\pi^-$	< 2.1 %	90%
$\Gamma_5$ $\rho^0 \rho^0$	< 1.9 × 10 <sup>-3</sup>	90%
$\Gamma_6$ $3\pi^+ 3\pi^-$	( 1.3 ± 0.9 ) %	
$\Gamma_7$ $K^+ K^- \pi^+ \pi^-$	< 1.4 %	90%
$\Gamma_8$ $K^{*0} \bar{K}^{*0}$	< 2.9 × 10 <sup>-3</sup>	90%
$\Gamma_9$ $K^+ K^- \pi^+ \pi^- \pi^0$	( 1.4 ± 1.0 ) %	
$\Gamma_{10}$ $K^+ K^- 2\pi^+ 2\pi^-$	< 1.4 %	90%
$\Gamma_{11}$ $K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$	( 1.0 ± 0.8 ) %	
$\Gamma_{12}$ $2K^+ 2K^-$	< 1.3 × 10 <sup>-3</sup>	90%

$\Gamma_{13}$	$\phi\phi$	< 1.1	$\times 10^{-3}$	90%
$\Gamma_{14}$	$p\bar{p}$	< 2.0	$\times 10^{-3}$	90%
$\Gamma_{15}$	$p\bar{p}\pi^+\pi^-$	seen		
$\Gamma_{16}$	$\gamma\gamma$	( $1.6 \pm 1.0$ )	$\times 10^{-4}$	
$\Gamma_{17}$	$\gamma J/\psi(1S)$	< 1.4	%	90%
$\Gamma_{18}$	$\pi^+\pi^-\eta$	< 6	$\times 10^{-3}$	90%
$\Gamma_{19}$	$\pi^+\pi^-\eta'$	( $2.6 \pm 1.9$ )	$\times 10^{-3}$	
$\Gamma_{20}$	$\pi^+\pi^-\eta_c(1S)$	< 25	%	90%

 **$\eta_c(2S)$  PARTIAL WIDTHS** **$\Gamma(\gamma\gamma)$**  **$\Gamma_{16}$** 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
$0.44 \pm 0.14$	106	<sup>1</sup> XU	18	BELL $e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
$1.3 \pm 0.6$		<sup>2</sup> ASNER	04	CLEO $\gamma\gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$

<sup>1</sup> Assuming that the branching fraction into  $\eta'\pi^+\pi^-$  is the same as for  $\eta_c(1S)$ .<sup>2</sup> They measure  $\Gamma(\eta_c(2S)\gamma\gamma) B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma) B(\eta_c(1S) \rightarrow K\bar{K}\pi)$ . The value for  $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$  is derived assuming that the branching fractions for  $\eta_c(2S)$  and  $\eta_c(1S)$  decays to  $K_S K\pi$  are equal and using  $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$  keV. **$\Gamma(\gamma\gamma) \times \Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$**  **$\Gamma_{16}\Gamma_{19}/\Gamma$** 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$5.6^{+1.2}_{-1.1} \pm 1.1$	106	XU	18	BELL $e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$

 **$\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$**  **$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$**  **$\Gamma_2\Gamma_{16}/\Gamma$** 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$41 \pm 4 \pm 6$	624	<sup>1</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$

<sup>1</sup> Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M. **$\Gamma(2\pi^+2\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$**  **$\Gamma_4\Gamma_{16}/\Gamma$** 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.5$	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+\pi^-)$

 **$\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$**  **$\Gamma_7\Gamma_{16}/\Gamma$** 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.0$	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+K^-\pi^+\pi^-$

 **$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$**  **$\Gamma_9\Gamma_{16}/\Gamma$** 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$30 \pm 6 \pm 5$	1201	<sup>1</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

<sup>1</sup> Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\Gamma(2K^+2K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{12}\Gamma_{16}/\Gamma$
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<2.9	90	UEHARA	08	BELL	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+ K^-)$
$\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{20}\Gamma_{16}/\Gamma$
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<133	90	LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$	

### $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{14}/\Gamma \times \Gamma_{16}/\Gamma$
<u>VALUE (units <math>10^{-8}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 5.6	90	1,2,3 AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 8.0	90	1,2,4 AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$	
<12.0	90	2,4 AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$	

<sup>1</sup> Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.

<sup>2</sup> For a total width  $\Gamma=5$  MeV.

<sup>3</sup> For the resonance mass region  $3589$ – $3599$  MeV/ $c^2$ .

<sup>4</sup> For the resonance mass region  $3575$ – $3660$  MeV/ $c^2$ .

### $\eta_c(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>not seen</b>		ABREU	980 DLPH	$e^+ e^- \rightarrow e^+ e^- + \text{hadrons}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
seen	<sup>1</sup>	EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$	

<sup>1</sup> For a mass value of  $3594 \pm 5$  MeV

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$					$\Gamma_2/\Gamma$
<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>1.9±0.4±1.1</b>	$59 \pm 12$	<sup>1</sup> AUBERT	08AB BABR	$B \rightarrow \eta_c(2S) K \rightarrow K\bar{K}\pi K$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
seen	$127 \pm 18$	ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K\bar{K}\pi$	
seen	$39 \pm 11$	<sup>2</sup> CHOI	02 BELL	$B \rightarrow KK_S K^- \pi^+$	

<sup>1</sup> Derived from a measurement of  $[B(B^+ \rightarrow \eta_c(2S)K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6^{+2.0}_{-1.9} \pm 2.5)\%$  and using  $B(B^+ \rightarrow \eta_c(2S)K^+) = (3.4 \pm 1.8) \times 10^{-4}$ , and  $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77^{+0.55}_{-0.66}) \times 10^{-5}$ .

<sup>2</sup> For a mass value of  $3654 \pm 6$  MeV

$\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$					$\Gamma_3/\Gamma_2$
<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>27.3±7.0±9.0</b>	225	<sup>1</sup> LEES	14E BABR	$\gamma\gamma \rightarrow K^+ K^- \gamma\gamma$	
<sup>1</sup> LEES 14E reports $B(\eta_c(2S) \rightarrow K^+ K^- \eta)/B(\eta_c(2S) \rightarrow K^+ K^- \pi^0) = 0.82 \pm 0.21 \pm 0.27$ , which we divide by 3 to account for isospin symmetry.					

### $\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$

$\Gamma_4/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
not seen	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$

### $\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$

$\Gamma_5/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
not seen	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

### $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$

$\Gamma_7/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
not seen	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$

### $\Gamma(K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$

$\Gamma_8/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
not seen	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

### $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K \bar{K} \pi)$

$\Gamma_9/\Gamma_2$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.73 ± 0.17 ± 0.17</b>	1201	<sup>1</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

<sup>1</sup> We have multiplied the value of  $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$  reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain  $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K \bar{K} \pi)$ . Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

### $\Gamma(K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

$\Gamma_{11}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
seen	$57 \pm 17$	ABLIKIM	13K	BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$

### $\Gamma(2K^+ 2K^-)/\Gamma_{\text{total}}$

$\Gamma_{12}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
not seen	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$

### $\Gamma(\phi\phi)/\Gamma_{\text{total}}$

$\Gamma_{13}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
not seen	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

### $\Gamma(p\bar{p})/\Gamma_{\text{total}}$

$\Gamma_{14}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
seen	106	<sup>1</sup> AAIJ	17AD LHCb	$p\bar{p} \rightarrow B^+ X \rightarrow p\bar{p} K^+ X$

<sup>1</sup> AAIJ 17AD report a 6.4 standard deviation signal, with  $B(B^+ \rightarrow \eta_c(2S) K^+ \rightarrow p\bar{p} K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p} K^+) = (1.58 \pm 0.33 \pm 0.09) \times 10^{-2}$ .

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{15}/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	110	<sup>1</sup> CHILIKIN	19	BELL $e^+e^- \rightarrow \gamma(4S)$	

<sup>1</sup> CHILIKIN 19 reports signals in  $B^+ \rightarrow \eta_c(2S)K^+$  and  $B^0 \rightarrow \eta_c(2S)K_S^0$  with 12.3 and 5.9 standard deviations, respectively.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{16}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<4 \times 10^{-4}$	90	<sup>1</sup> WICHT	08	BELL $B^\pm \rightarrow K^\pm\gamma\gamma$	
not seen		AMBROGIANI	01	E835 $\bar{p}p \rightarrow \gamma\gamma$	
$<0.01$	90	LEE	85	CBAL $\psi' \rightarrow \text{photons}$	

<sup>1</sup> WICHT 08 reports  $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c(2S)K^+)] < 0.18 \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \eta_c(2S)K^+) = 4.4 \times 10^{-4}$ .

$\Gamma(\pi^+\pi^-\eta_c(1S))/\Gamma(K\bar{K}\pi)$					$\Gamma_{20}/\Gamma_2$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;3.33</b>	90	<sup>1</sup> LEES	12AE BABR	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$	

<sup>1</sup> We divided the reported limit by 3 to take into account isospin relations.

### $\eta_c(2S)$ CROSS-PARTICLE BRANCHING RATIOS

$\Gamma(\eta_c(2S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$					$\Gamma_3/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<11.8 \times 10^{-6}$	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+K^-\eta$	

<sup>1</sup> CRONIN-HENNESSY 10 reports a limit of  $< 5.9 \times 10^{-6}$  for the decay  $\eta_c(2S) \rightarrow K^+K^-\eta$  which we multiply by 2 account for isospin symmetry. It assumes  $\Gamma(\eta_c(2S)) = 14$  MeV. It also gives the analytic dependence of limits on width.

$\Gamma(\eta_c(2S) \rightarrow 2\pi^+2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;14.6 \times 10^{-6}</b>	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 2\pi^+2\pi^-$	

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$\Gamma(\eta_c(2S) \rightarrow \rho^0\rho^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$					$\Gamma_5/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;12.7 \times 10^{-7}</b>	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma 2\pi^+2\pi^-$	

$$\Gamma(\eta_c(2S) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \\ \Gamma_6/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.2±1.0±1.2</b>	569	ABLIKIM	22Q	BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13.2 90 <sup>1</sup> CRONIN-HEN..10 CLEO  $\psi(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \\ \Gamma_7/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;9.6 × 10<sup>-6</sup></b>	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \\ \Gamma_8/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;19.6 × 10<sup>-7</sup></b>	90	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \\ \Gamma_9/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;43.0 × 10<sup>-6</sup></b>	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \\ \Gamma_{10}/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;9.7 × 10<sup>-6</sup></b>	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- 2\pi^+ 2\pi^-$

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \\ \Gamma_{11}/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.03±2.10±0.7</b>	60	ABLIKIM	13K	BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 15.2 90 <sup>1</sup> CRONIN-HEN..10 CLEO  $\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{13}/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.8 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

$$\Gamma(\eta_c(2S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{14}/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.4 \times 10^{-6}$	90	ABLIKIM	13V BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$

$$\Gamma(\eta_c(2S) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{17}/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<9.7 \times 10^{-6}$	90	33	1 ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma J/\psi$

<sup>1</sup> Uses  $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$  and  $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$ .

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{18}/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<4.3 \times 10^{-6}$	90	1 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta$

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{19}/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<14.2 \times 10^{-6}$	90	1 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta'$

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{20}/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.7 \times 10^{-4}$	90	1 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta_c(1S)$

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

## $\eta_c(2S)$ REFERENCES

ABLIKIM	22Q	PR D106 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
CHILIKIN	19	PR D100 012001	K. Chilikin <i>et al.</i>	(BELLE Collab.)
XU	18	PR D98 072001	Q.N. Xu <i>et al.</i>	(BELLE Collab.)
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)

ABLIKIM	13K	PR D87 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	11H	PR D84 091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
VINOKUROVA	11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
CRONIN-HEN...	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)
AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05C	PR D72 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)