

# $\psi_2(3823)$

$I^G(JPC) = 0^-(2^{--})$   
 $I, J, P$  need confirmation.

was  $\psi(3823)$ ,  $X(3823)$

Seen by BHARDWAJ 13 in  $B \rightarrow \chi_{c1}\gamma K$  and ABLIKIM 15S in  $e^+e^- \rightarrow \pi^+\pi^-\gamma\chi_{c1}$  decays as a narrow peak in the invariant mass distribution of the  $\chi_{c1}\gamma$  system. Properties consistent with the  $\psi_2(1^3D_2)$   $c\bar{c}$  state.

## $\psi_2(3823)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3823.5 ± 0.5 OUR AVERAGE</b>		Error includes scale factor of 1.4.		
3823.12 ± 0.43 ± 0.13	120	ABLIKIM	22R	$BES3 e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
3824.08 ± 0.53 ± 0.14	137	<sup>1</sup> AAIJ	20S	$LHCb B^+ \rightarrow J/\psi\pi^+\pi^-K^+$
3823.1 ± 1.8 ± 0.7	33 ± 10	<sup>2</sup> BHARDWAJ	13	$BELL B^\pm \rightarrow \chi_{c1}\gamma K^\pm$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
3821.7 ± 1.3 ± 0.7	19 ± 5	<sup>3</sup> ABLIKIM	15S	$BES3 e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$

<sup>1</sup> Using the measured  $m_{\psi_2(3823)} - m_{\psi(2S)} = 137.98 \pm 0.53 \pm 0.14$  MeV.

<sup>2</sup> From a simultaneous fit to  $B^\pm \rightarrow (\chi_{c1}\gamma)K^\pm$  and  $B^0 \rightarrow (\chi_{c1}\gamma)K_S^0$  with significance  $4.0\sigma$  including systematics. Corrected for the measured  $\psi(2S)$  mass using  $B \rightarrow \psi(2S)K \rightarrow (\gamma\chi_{c1})K$  decays.

<sup>3</sup> From a simultaneous unbinned maximum likelihood fit of  $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$  data (the  $\pi^+\pi^-$  recoil mass) taken at  $\sqrt{s}$  values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both  $\psi(2S) \rightarrow \chi_{c1}\gamma$  and  $\psi_2(3823) \rightarrow \chi_{c1}\gamma$  together, with floating mass scale offset for  $\psi(2S)$ , floating  $\psi_2(3823)$  mass, and zero  $\psi_2(3823)$  width, resulting in a significance of  $5.9\sigma$  when including systematic uncertainties. Superseded by ABLIKIM 22R.

## $m_{\psi_2(3823)} - m_{\psi(2S)}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
137.98 ± 0.53 ± 0.14	137	<sup>1</sup> AAIJ	20S	$LHCb B^+ \rightarrow J/\psi\pi^+\pi^-K^+$
<sup>1</sup> AAIJ 20S also reports $m_{\chi_{c1}(3872)} - m_{\psi_2(3823)} = 47.50 \pm 0.53 \pm 0.13$ MeV.				

## $\psi_2(3823)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 2.9	90	120	ABLIKIM	22R	$BES3 e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
< 5.2	90		<sup>1</sup> AAIJ	20S	$LHCb B^+ \rightarrow J/\psi\pi^+\pi^-K^+$
< 16	90		<sup>2</sup> ABLIKIM	15S	$BES3 e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
< 24	90		<sup>3</sup> BHARDWAJ	13	$BELL B^\pm \rightarrow \chi_{c1}\gamma K^\pm$

<sup>1</sup> AAIJ 20S also provides a limit of < 6.6 MeV with 95% CL.

<sup>2</sup> From a fit of  $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$  data (the  $\pi^+\pi^-$  recoil mass) taken at  $\sqrt{s}$  values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to a Breit-Wigner function with the mass fixed from the likelihood fit above, Gaussian resolution smearing, and floating width.

<sup>3</sup> From a simultaneous fit to  $B^\pm \rightarrow (\chi_{c1}\gamma)K^\pm$  and  $B^0 \rightarrow (\chi_{c1}\gamma)K_S^0$  with significance  $4.0\sigma$  including systematics.

## $\psi_2(3823)$ DECAY MODES

Branching fractions are given relative to the one **DEFINED AS 1**.

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1 J/\psi(1S)\pi^+\pi^-$	<0.06	90%
$\Gamma_2 J/\psi(1S)\pi^0\pi^0$	<0.11	90%
$\Gamma_3 J/\psi(1S)\pi^0$	<0.030	90%
$\Gamma_4 J/\psi(1S)\eta$	<0.14	90%
$\Gamma_5 \chi_{c0}\gamma$	<0.24	90%
$\Gamma_6 \chi_{c1}\gamma$	<b>DEFINED AS 1</b>	
$\Gamma_7 \chi_{c2}\gamma$	$0.28^{+0.14}_{-0.11}$	

## $\psi_2(3823)$ BRANCHING RATIOS

$$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	<sup>1</sup> ABLIKIM	210	BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
seen	$137 \pm 26$	AAIJ	20S	LHCb $B^+ \rightarrow J/\psi\pi^+\pi^-K^+$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

$$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(\chi_{c1}\gamma) \quad \Gamma_1/\Gamma_6$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.06</b>	90	<sup>1</sup> ABLIKIM	210	BES3 $e^+e^- \rightarrow \pi^+\pi^-X$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(\chi_{c1}\gamma) \quad \Gamma_2/\Gamma_6$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.11</b>	90	<sup>1</sup> ABLIKIM	210	BES3 $e^+e^- \rightarrow \pi^+\pi^-X$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

$$\Gamma(J/\psi(1S)\pi^0)/\Gamma(\chi_{c1}\gamma) \quad \Gamma_3/\Gamma_6$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.03</b>	90	<sup>1</sup> ABLIKIM	210	BES3 $e^+e^- \rightarrow \pi^+\pi^-X$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

$$\Gamma(J/\psi(1S)\eta)/\Gamma(\chi_{c1}\gamma) \quad \Gamma_4/\Gamma_6$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.14</b>	90	<sup>1</sup> ABLIKIM	210	BES3 $e^+e^- \rightarrow \pi^+\pi^-X$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$   $\Gamma_4/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>4.4<sup>+2.5</sup><sub>-1.9</sub><sup>±0.9</sup></b>	<sup>1</sup> AAIJ	22D LHCb	$B^+ \rightarrow J/\psi(1S)\eta K^+$

<sup>1</sup> Using the branching ratio for  $B^+ \rightarrow \psi_2(3823)K^+$  with  $\psi_2(3823) \rightarrow J/\psi(1S)\pi^+\pi^-$  from AAIJ 20S.

 $\Gamma(\chi_{c0}\gamma)/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	<sup>1</sup> ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
----------	----------------------	----------	----------------------------------

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

 $\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	120	<sup>1</sup> ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
seen	$63 \pm 9$	<sup>2</sup> ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
seen	$16 \pm 5$	<sup>3</sup> ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^0\pi^0X$
seen	$33 \pm 10$	<sup>4</sup> BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1}\gamma K^\pm$

<sup>1</sup> From a fit to the  $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$  cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak  $R$  with mass  $4417.5 \pm 26.2 \pm 3.5$  MeV and width  $245 \pm 48 \pm 13$  MeV, which leads to  $\Gamma(e^+e^-)B(R \rightarrow \pi^+\pi^-\psi_2(3823))B(\psi_2(3823) \rightarrow \chi_{c1}\gamma) = 0.57 \pm 0.08$  eV.

<sup>2</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ . Signal has a  $11.8\sigma$  significance.

<sup>3</sup> From a fit of the invariant  $\pi^0\pi^0$  recoil-mass distribution. Signal has a  $4.3\sigma$  significance.

<sup>4</sup> BHARDWAJ 13 reports  $B(B^\pm \rightarrow \psi_2(3823)K^\pm) \times B(\psi_2(3823) \rightarrow \gamma\chi_{c1}) = (9.7 \pm 2.8 \pm 1.1) \times 10^{-6}$  with statistical significance  $3.8\sigma$ .

 $\Gamma(\chi_{c0}\gamma)/\Gamma(\chi_{c1}\gamma)$   $\Gamma_5/\Gamma_6$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.24	90	<sup>1</sup> ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

 $\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	<sup>1</sup> ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
not seen	<sup>2</sup> ABLIKIM	15S BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$
not seen	<sup>3</sup> BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c2}\gamma K^\pm$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ . Signal has a  $3.2\sigma$  significance.

<sup>2</sup> From a simultaneous unbinned maximum likelihood fit of  $e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$  data (the  $\pi^+\pi^-$  recoil mass) taken at  $\sqrt{s}$  values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to

simulated events including both  $\psi(2S) \rightarrow \chi_{c2}\gamma$  and  $\psi_2(3823) \rightarrow \chi_{c2}\gamma$  together, with floating mass scale offset for  $\psi(2S)$ ,  $\psi_2(3823)$  mass floating (fixed to that above), and zero  $\psi_2(3823)$  width.

<sup>3</sup> BHARDWAJ 13 reports  $B(B^\pm \rightarrow \psi_2(3823)K^\pm) \times B(\psi_2(3823) \rightarrow \gamma\chi_{c2}) < 3.6 \times 10^{-6}$  at 90% CL.

$\Gamma(\chi_{c2}\gamma)/\Gamma(\chi_{c1}\gamma)$	$\Gamma_7/\Gamma_6$				
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.28<sup>+0.14</sup><sub>-0.11</sub><sup>±0.02</sup></b>		9 ± 4	<sup>1</sup> ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.42	90	<sup>2</sup> ABLIKIM	15S BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$
<0.41	90	BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1/c2}\gamma K^\pm$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

<sup>2</sup> From a simultaneous unbinned maximum likelihood fit of  $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}(2)\gamma$  data (the  $\pi^+\pi^-$  recoil mass) taken at  $\sqrt{s}$  values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both  $\psi(2S) \rightarrow \chi_{c1}(2)\gamma$  and  $\psi_2(3823) \rightarrow \chi_{c1}(2)\gamma$  together, with floating mass scale offset for  $\psi(2S)$ ,  $\psi_2(3823)$  mass floating (fixed to that above), and zero  $\psi_2(3823)$  width.

## $\psi_2(3823)$ REFERENCES

AAIJ	22D	JHEP 2204 046	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	22R	PRL 129 102003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21O	PR D103 L091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	20S	JHEP 2008 123	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BHARDWAJ	13	PRL 111 032001	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)