

**$\Lambda(1405) S_{01}$**  $I(J^P) = 0(\frac{1}{2}^-)$  Status: \*\*\*\***NOTE ON THE  $\Lambda(1405)$** 

Revised March 1998 by R.H. Dalitz (Oxford University).

It is generally accepted that the  $\Lambda(1405)$  is a well-established  $J^P = 1/2^-$  resonance. It is assigned to the lowest  $L = 1$  supermultiplet of the 3-quark system and paired with the  $J^P = 3/2^-$   $\Lambda(1520)$ . Lying about 30 MeV below the  $N\bar{K}$  threshold, the  $\Lambda(1405)$  can be observed directly only as a resonance bump in the  $(\Sigma\pi)^0$  subsystem in final states of production experiments. It was first reported by ALSTON 61B in the reaction  $K^-p \rightarrow \Sigma\pi\pi\pi$  at 1.15 GeV/ $c$  and has since been seen in at least eight other experiments. However, only two of them had enough events for a detailed analysis: THOMAS 73, with about 400  $\Sigma^\pm\pi^\mp$  events from  $\pi^-p \rightarrow K^0(\Sigma\pi)^0$  at 1.69 GeV/ $c$ ; and HEMINGWAY 85, with 766  $\Sigma^+\pi^-$  and 1106  $\Sigma^-\pi^+$  events from  $K^-p \rightarrow (\Sigma\pi\pi)^+\pi^-$  at 4.2 GeV/ $c$ , after the selections  $1600 \leq M(\Sigma\pi\pi)^+ \leq 1720$  MeV and momentum transfer  $\leq 1.0$  (GeV/ $c$ )<sup>2</sup> to purify the  $\Lambda(1405) \rightarrow (\Sigma\pi)^0$  sample. These experiments agree on a mass of about 1395–1400 MeV and a width of about 60 MeV. (Hemingway's mass of  $1391 \pm 1$  MeV is from his best, but unacceptably poor, Breit-Wigner fit.)

The Byers-Fenster tests on these data allow any spin and either parity: neither  $J$  nor  $P$  has yet been determined *directly*. The early indications for  $J^P = 1/2^-$  came from finding  $\text{Re } A_{I=0}$  to be large and negative in a constant-scattering-length analysis of low-energy  $N\bar{K}$  reaction data (see KIM 65, SAKITT 65, and earlier references cited therein). The first multichannel energy-dependent K-matrix analysis (KIM 67) strengthened the case for a resonance around 1400–1420 MeV strongly coupled to the  $I = 0$   $S$ -wave  $N\bar{K}$  system.

THOMAS 73 and HEMINGWAY 85 both found the  $\Lambda(1405)$  bump to be asymmetric and not well fitted by a Breit-Wigner resonance function with constant parameters. The asymmetry involves a rapid fall in intensity as the  $N\bar{K}$  threshold energy is approached from below. This is readily understood as due to a strong coupling of the  $\Lambda(1405)$  to the  $S$ -wave  $N\bar{K}$  channel (see DALITZ 81). This striking  $S$ -shaped cusp behavior at a new threshold is characteristic of  $S$ -wave coupling; the other below-threshold hyperon, the  $\Sigma(1385)$ , has no such threshold distortion because its  $N\bar{K}$  coupling is  $P$ -wave. For the  $\Lambda(1405)$ , this asymmetry is the *sole direct evidence* that  $J^P = 1/2^-$ .

Following the early work cited above, a considerable literature has developed on proper procedures for phenomenological extrapolation below the  $N\bar{K}$  threshold, partly in order to strengthen the evidence for the spin-parity of the  $\Lambda(1405)$ , and partly to provide an estimate for the amplitude  $f(N\bar{K})$  in the unphysical domain below the  $N\bar{K}$  threshold; the latter is needed for the evaluation of the dispersion relation for  $N\bar{K}$  and  $NK$  forward scattering amplitudes. For recent reviews, see MILLER 84 and BARRETT 89. In most recent work, the  $(\Sigma\pi)^0$  production spectrum is included in the data fitted (see, *e.g.*, CHAO 73, MARTIN 81).

It is now accepted that the data can be fitted only with an  $S$ -wave pole in the reaction amplitudes below  $N\bar{K}$  threshold (see, however, FINK 90), but there is still controversy about the physical origin of this pole (for a review, see DALITZ 81 and DALITZ 82). Two extreme possibilities are: (a) an  $L = 1$  SU(3)-singlet  $uds$  state coupled with the  $S$ -wave meson-baryon systems; or (b) an unstable  $N\bar{K}$  bound state, analogous to the (stable) deuteron in the  $NN$  system. The problem with (a) is that the  $\Lambda(1405)$  mass is so much lower than that of

its partner, the  $\Lambda(1520)$ . This requires, in the QCD-inspired quark model, rather large spin-orbit couplings, whether or not one uses relativistic kinetic energies. CAPSTICK 86 and CAPSTICK 89 conclude that a proper QCD calculation leads only to small energy splittings, whereas LEINWEBER 90, using QCD sum rules, obtains a good fit to this splitting.

On the other hand, the problem with (b) is that then another  $J^P = 1/2^- \Lambda$  is needed to replace the  $\Lambda(1405)$  in the  $L = 1$  supermultiplet, and it would have to lie close to the  $\Lambda(1520)$ , a region already well explored by  $N\bar{K}$  experiments without result. Intermediate structures are possible; for example, the cloudy bag model allows the configurations (a) and (b) to mix and finds the intensity of (a) in the  $\Lambda(1405)$  to be only 14% (VEIT 84, VEIT 85, JENNINGS 86). Such models naturally predict a second  $1/2^- \Lambda$  close to the  $\Lambda(1520)$ .

The determination of the mass and width of the resonance from  $(\Sigma\pi)^0$  data is usually based on the “Watson approximation,” which states that the production rate  $R(\Sigma\pi)$  of the  $(\Sigma\pi)^0$  state has a mass dependence proportional to  $(\sin^2\delta_{\Sigma\pi})/q$ ,  $q$  being the  $\Sigma\pi$  c.m. momentum, in a  $\Sigma\pi$  mass range where  $\delta_{\Sigma\pi}$  is not far from  $\pi/2$  and only the  $\Sigma\pi$  channel is open, *i.e.*, between the  $\Sigma\pi$  and the  $N\bar{K}$  thresholds. Then  $q R(\Sigma\pi)$  is proportional to  $\sin^2\delta_{\Sigma\pi}$ , and the mass  $M$  may be defined as the energy at which  $\sin^2\delta_{\Sigma\pi} = 1$ . The width  $\Gamma$  may be determined from the rate at which  $\delta_{\Sigma\pi}$  goes through  $\pi/2$ , or from the FWHM; this is a matter of convention.

This determination of  $M$  and  $\Gamma$  from the data suffers from the following defects:

(i) The determination of  $\sin^2\delta_{\Sigma\pi}$  requires that  $R(\Sigma\pi)$  be scaled to give  $\sin^2\delta_{\Sigma\pi} = 1$  at the peak for the best fit to the data; *i.e.*, the bump must be *assumed* to arise from a resonance.

However, this assumption is supported by the analysis of the low-energy  $N\bar{K}$  data and its extrapolation below threshold.

(ii) Owing to the nearby  $N\bar{K}$  threshold, the shape of the best fit to the  $M(\Sigma\pi)$  bump is uncertain. For energies below this threshold at  $E_{N\bar{K}}$ , the general form for  $\delta_{\Sigma\pi}$  is

$$q \cot \delta_{\Sigma\pi} = \frac{1 + \kappa\alpha}{\gamma + \kappa(\alpha\gamma - \beta^2)}. \quad (1)$$

Here  $\alpha, \beta$ , and  $\gamma$  are the (generally energy-dependent)  $NN$ ,  $N\Sigma$ , and  $\Sigma\Sigma$  elements of the  $I = 0$   $S$ -wave K-matrix for the  $(\Sigma\pi, N\bar{K})$  system, and  $\kappa$  is the magnitude of the (imaginary) c.m. momentum  $k_K$  for the  $N\bar{K}$  system below threshold. The elements  $\alpha, \beta, \gamma$  are real functions of  $E$ ; they have no branch cuts at the  $\Sigma\pi$  and  $N\bar{K}$  thresholds, but they are permitted to have poles in  $E$  along the real  $E$  axis. The resonance asymmetry arises from the effect of  $\kappa$  on  $\delta_{\Sigma\pi}$ . We note that  $\delta_{\Sigma\pi} = \pi/2$  when  $\kappa = -1/\alpha$ .

Accepting this close connection of  $\delta_{\Sigma\pi}$  with the low-energy  $N\bar{K}$  data, it is natural to analyze the two sets of data together (*e.g.*, MARTIN 81), and there is now a large body of accurate  $N\bar{K}$  data for laboratory momenta between 100 and 300 MeV/ $c$  (see MILLER 84). The two sets of data span c.m. energies from 1370 MeV to 1490 MeV, and the K-matrix elements will not be energy independent over such a broad range. For the  $I = 0$  channels, a linear energy dependence for  $K^{-1}$  has been adopted routinely ever since the work of KIM 67, and it is essential when fitting the  $qR(\Sigma\pi)$  and  $N\bar{K}$  data together. However,  $qR(\Sigma\pi)$  is not always well fitted in this procedure; the value obtained for the  $\Lambda(1405)$  mass  $M$  varies a good deal with the type of fit, not a surprising result when the  $\Sigma\pi$  mass spectrum below the  $pK^-$  threshold contributes only nine data points in a total of about 200. The value of  $M$  obtained from an overall fit

is not necessarily much better than from one using only the  $qR(\Sigma\pi)$  data; and  $M$  may be a function of the representation—K-matrix,  $K^{-1}$ -matrix, relativistic-separable or nonseparable potentials, *etc.*— used in fitting over the full energy range. DALITZ 91 fitted the  $qR(\Sigma^+\pi^-)$  Hemingway data with each of the first three representations just mentioned, constrained to the  $I = 0$   $N\bar{K}$  threshold scattering length from low-energy  $N\bar{K}$  data. The (nonseparable) meson-exchange potentials of MÜLLER-GROELING 90, fitted to the low-energy  $N\bar{K}$  (and  $NK$ ) data, predicted an unstable  $N\bar{K}$  bound state with mass and width compatible with the  $\Lambda(1405)$ .

From the measurement of  $2p \rightarrow 1s$  x rays from kaonic-hydrogen, the energy-level shift  $\Delta E$  and width  $\Gamma$  of its  $1s$  state can give us two further constraints on the  $(\Sigma\pi, N\bar{K})$  system, at an energy roughly midway between those from the low-energy hydrogen bubble chamber studies and those from  $qR(\Sigma\pi)$  observations below the  $pK^-$  threshold. IWASAKI 97 have reported the first convincing observation of this x ray, with a good initial estimate:

$$\Delta E - i\Gamma/2 = (-323 \pm 63 \pm 11) - i(204 \pm 104 \pm 50) \text{ eV} . \quad (2)$$

The errors here encompass about half of the predictions made following the various analyses and/or models for the in-flight  $K^-p$  and sub-threshold  $qR(\Sigma\pi)$  data. Better measurements will be needed to discriminate between the analyses and predictions. Now that  $\Delta E$  is known with some certainty, we can anticipate much-improved data on kaonic-hydrogen, perhaps from the DAΦNE storage ring at Frascati, information vital for our quantitative understanding of the  $(\Sigma\pi, N\bar{K})$  system in this region. This will lead to better knowledge of kaonic coupling

strengths and to more reliable dispersion-theoretic arguments concerning strange-particle processes.

The present status of the  $\Lambda(1405)$  thus depends heavily on theoretical arguments, a somewhat unsatisfactory basis for a four-star rating. Nevertheless, there is no known reason to doubt its existence or quantum numbers. The 3-quark model for baryons has been broadly successful in accounting for all of the  $L^P = 1^-$  excited baryonic states (CAPSTICK 89), apart from the relatively large mass separation between the  $\Lambda(1405)$  and  $\Lambda(1520)$ . Quark model builders have no reservations about accepting the  $\Lambda(1405)$  as a 3-quark state. However, calculations with broken-chiral-symmetric models, which combine internal 3-quark configurations with external meson-baryon states (*e.g.*, VEIT 85, KAISER 95) end up with descriptions of the  $\Lambda(1405)$  dominated by the meson-baryon terms in the wavefunctions. Models using meson-baryon potentials readily fit its mass, and give  $\Delta E$  negative, as is found empirically. The problem is not so much one of “either (a) or (b),” but rather how to achieve “both (a) and (b).” Theoreticians have not yet been able to deal with the full coupled-channels system, with  $qqq$  and  $qqq\bar{q}$  configurations (at the least) being treated on the same footing. On the experimental side, better statistics are needed, both above and below the  $pK^-$  threshold. To disentangle the physics, the  $I = 1$  channels also need more attention. For example, low-energy  $pK_L^0$  interactions have not been studied at all in the last 25 years.

---

**$\Lambda(1405)$  MASS****PRODUCTION EXPERIMENTS**

| <u>VALUE (MeV)</u>  | <u>EVTS</u> | <u>DOCUMENT ID</u>     | <u>TECN</u> | <u>COMMENT</u>                |
|---|-------------|------------------------|-------------|-------------------------------|
| <b>1406.5 ± 4.0</b>   |             | <sup>1</sup> DALITZ    | 91          | M-matrix fit                  |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |             |                        |             |                               |
| 1391 ± 1  | 700         | <sup>1</sup> HEMINGWAY | 85 HBC      | $K^- p$ 4.2 GeV/c             |
| ~ 1405  | 400         | <sup>2</sup> THOMAS    | 73 HBC      | $\pi^- p$ 1.69 GeV/c          |
| 1405  | 120         | BARBARO-...            | 68B DBC     | $K^- d$ 2.1–2.7 GeV/c         |
| 1400 ± 5  | 67          | BIRMINGHAM             | 66 HBC      | $K^- p$ 3.5 GeV/c             |
| 1382 ± 8  |             | ENGLER                 | 65 HDBC     | $\pi^- p, \pi^+ d$ 1.68 GeV/c |
| 1400 ± 24   |             | MUSGRAVE               | 65 HBC      | $\bar{p} p$ 3–4 GeV/c         |
| 1410  |             | ALEXANDER              | 62 HBC      | $\pi^- p$ 2.1 GeV/c           |
| 1405  |             | ALSTON                 | 62 HBC      | $K^- p$ 1.2–0.5 GeV/c         |
| 1405  |             | ALSTON                 | 61B HBC     | $K^- p$ 1.15 GeV/c            |

**EXTRAPOLATIONS BELOW  $N\bar{K}$  THRESHOLD**

| <u>VALUE (MeV)</u>  | <u>DOCUMENT ID</u>  | <u>TECN</u> | <u>COMMENT</u>        |
|---|---------------------|-------------|-----------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                     |             |                       |
| 1411  | <sup>3</sup> MARTIN | 81          | K-matrix fit          |
| 1406  | <sup>4</sup> CHAO   | 73 DPWA     | 0-range fit (sol. B)  |
| 1421  | MARTIN              | 70 RVUE     | Constant K-matrix     |
| 1416 ± 4  | MARTIN              | 69 HBC      | Constant K-matrix     |
| 1403 ± 3  | KIM                 | 67 HBC      | K-matrix fit          |
| 1407.5 ± 1.2  | <sup>5</sup> KITTEL | 66 HBC      | 0-effective-range fit |
| 1410.7 ± 1.0  | KIM                 | 65 HBC      | 0-effective-range fit |
| 1409.6 ± 1.7  | <sup>5</sup> SAKITT | 65 HBC      | 0-effective-range fit |

 **$\Lambda(1405)$  WIDTH****PRODUCTION EXPERIMENTS**

| <u>VALUE (MeV)</u>  | <u>EVTS</u> | <u>DOCUMENT ID</u>     | <u>TECN</u> | <u>COMMENT</u>        |
|---|-------------|------------------------|-------------|-----------------------|
| <b>50 ± 2</b>   |             | <sup>1</sup> DALITZ    | 91          | M-matrix fit          |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |             |                        |             |                       |
| 32 ± 1  | 700         | <sup>1</sup> HEMINGWAY | 85 HBC      | $K^- p$ 4.2 GeV/c     |
| 45 to 55  | 400         | <sup>2</sup> THOMAS    | 73 HBC      | $\pi^- p$ 1.69 GeV/c  |
| 35  | 120         | BARBARO-...            | 68B DBC     | $K^- d$ 2.1–2.7 GeV/c |
| 50 ± 10   | 67          | BIRMINGHAM             | 66 HBC      | $K^- p$ 3.5 GeV/c     |
| 89 ± 20   |             | ENGLER                 | 65 HDBC     |                       |
| 60 ± 20   |             | MUSGRAVE               | 65 HBC      |                       |
| 35 ± 5  |             | ALEXANDER              | 62 HBC      |                       |
| 50  |             | ALSTON                 | 62 HBC      |                       |
| 20  |             | ALSTON                 | 61B HBC     |                       |

**EXTRAPOLATIONS BELOW  $N\bar{K}$  THRESHOLD**

| <u>VALUE (MeV)</u>  | <u>DOCUMENT ID</u>  | <u>TECN</u> | <u>COMMENT</u>       |
|---|---------------------|-------------|----------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                     |             |                      |
| 30  | <sup>3</sup> MARTIN | 81          | K-matrix fit         |
| 55  | <sup>4,6</sup> CHAO | 73 DPWA     | 0-range fit (sol. B) |
| 20  | MARTIN              | 70 RVUE     | Constant K-matrix    |
| 29 ± 6  | MARTIN              | 69 HBC      | Constant K-matrix    |
| 50 ± 5  | KIM                 | 67 HBC      | K-matrix fit         |
| 34.1 ± 4.1  | <sup>5</sup> KITTEL | 66 HBC      |                      |
| 37.0 ± 3.2  | KIM                 | 65 HBC      |                      |
| 28.2 ± 4.1  | <sup>5</sup> SAKITT | 65 HBC      |                      |

 **$\Lambda(1405)$  DECAY MODES**

| Mode                        | Fraction ( $\Gamma_i/\Gamma$ ) |
|-----------------------------|--------------------------------|
| $\Gamma_1$ $\Sigma\pi$      | 100 %                          |
| $\Gamma_2$ $\Lambda\gamma$  |                                |
| $\Gamma_3$ $\Sigma^0\gamma$ |                                |
| $\Gamma_4$ $N\bar{K}$       |                                |

 **$\Lambda(1405)$  PARTIAL WIDTHS**
 **$\Gamma(\Lambda\gamma)$   $\Gamma_2$** 

| <u>VALUE (keV)</u>  | <u>DOCUMENT ID</u> | <u>COMMENT</u>   |
|---|--------------------|------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                    |                  |
| 27 ± 8  | BURKHARDT 91       | Isobar model fit |

 **$\Gamma(\Sigma^0\gamma)$   $\Gamma_3$** 

| <u>VALUE (keV)</u>  | <u>DOCUMENT ID</u> | <u>COMMENT</u>   |
|---|--------------------|------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                    |                  |
| 10 ± 4 or 23 ± 7  | BURKHARDT 91       | Isobar model fit |

 **$\Lambda(1405)$  BRANCHING RATIOS**
 **$\Gamma(N\bar{K})/\Gamma(\Sigma\pi)$   $\Gamma_4/\Gamma_1$** 

| <u>VALUE</u>  | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>    |
|---|------------|--------------------|-------------|-------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |            |                    |             |                   |
| <3  | 95         | HEMINGWAY 85       | HBC         | $K^- p$ 4.2 GeV/c |

 **$\Lambda(1405)$  FOOTNOTES**

<sup>1</sup> DALITZ 91 fits the HEMINGWAY 85 data.

<sup>2</sup> THOMAS 73 data is fit by CHAO 73 (see next section).

<sup>3</sup> The MARTIN 81 fit includes the  $K^\pm p$  forward scattering amplitudes and the dispersion relations they must satisfy.

<sup>4</sup> See also the accompanying paper of THOMAS 73.

<sup>5</sup> Data of SAKITT 65 are used in the fit by KITTEL 66.

<sup>6</sup> An asymmetric shape, with  $\Gamma/2 = 41$  MeV below resonance, 14 MeV above.



**A(1405) REFERENCES**

|             |     |                |                                    |                                |
|-------------|-----|----------------|------------------------------------|--------------------------------|
| BURKHARDT   | 91  | PR C44 607     | +Lowe                              | (NOTT, UNM, BIRM)              |
| DALITZ      | 91  | JPG 17 289     | +Deloff                            | (OXFTP, WINR)                  |
| HEMINGWAY   | 85  | NP B253 742    |                                    | (CERN) J                       |
| MARTIN      | 81  | NP B179 33     |                                    | (DURH)                         |
| CHAO        | 73  | NP B56 46      | +Kraemer, Thomas, Martin           | (RHEL, CMU, LOUC)              |
| THOMAS      | 73  | NP B56 15      | +Engler, Fisk, Kraemer             | (CMU) J                        |
| MARTIN      | 70  | NP B16 479     | +Ross                              | (DURH)                         |
| MARTIN      | 69  | PR 183 1352    | +Sakitt                            | (LOUC, BNL)                    |
| Also        | 69B | PR 183 1345    | Martin, Sakitt                     | (LOUC, BNL)                    |
| BARBARO-... | 68B | PRL 21 573     | Barbaro-Galtieri, Chadwick+        | (LRL, SLAC)                    |
| KIM         | 67  | PRL 19 1074    |                                    | (YALE)                         |
| BIRMINGHAM  | 66  | PR 152 1148    |                                    | (BIRM, GLAS, LOIC, OXF, RHEL)  |
| KITTEL      | 66  | PL 21 349      | +Otter, Wacek                      | (VIEN)                         |
| ENGLER      | 65  | PRL 15 224     | +Fisk, Kraemer, Meltzer, Westgard+ | (CMU, BNL) J                   |
| KIM         | 65  | PRL 14 29      |                                    | (COLU)                         |
| MUSGRAVE    | 65  | NC 35 735      | +Petmezas+                         | (BIRM, CERN, EPOL, LOIC, SACL) |
| SAKITT      | 65  | PR 139B 719    | +Day, Glasser, Seeman, Friedman+   | (UMD, LRL)                     |
| ALEXANDER   | 62  | PRL 8 447      | +Kalbfleisch, Miller, Smith        | (LRL) I                        |
| ALSTON      | 62  | CERN Conf. 311 | +Alvarez, Ferro-Luzzi+             | (LRL) I                        |
| ALSTON      | 61B | PRL 6 698      | +Alvarez, Eberhard, Good+          | (LRL) I                        |

**OTHER RELATED PAPERS**

|  |     |   |   |                    |
|--|-----|---|---|--------------------|
| IWASAKI  | 97  | PRL 78 3067   | +Hayano, Ito, Nakamura+                     | (KEK-228 Collab.)  |
| FINK   | 90  | PR C41 2720   | +He, Landau, Schnick                        | (IBMY, ORST, ANSM) |
| LEINWEBER  | 90  | ANP 198 203   |   | (MCMS)             |
| MUELLER-GR...  | 90  | NP A513 557   | Mueller-Groeling, Holinde, Speth            | (JULI)             |
| BARRETT  | 89  | NC 102A 179   |   | (SURR)             |
| BATTY  | 89  | NC 102A 255   | +Gal  | (RAL, HEBR)        |
| CAPSTICK   | 89  | Excited Baryons '88, p. 32                            |   | (GUEL)             |
| LOWE   | 89  | NC 102A 167   |   | (BIRM)             |
| WHITEHOUSE   | 89  | PRL 63 1352   | + (BIRM, BOST, BRCO, BNL, CASE, BUDA, TRIU) |                    |
| SIEGEL   | 88  | PR C38 2221   | +Weise                                      | (REGE)             |
| WORKMAN  | 88  | PR D37 3117   | +Fearing                                    | (TRIU)             |
| SCHNICK  | 87  | PRL 58 1719   | +Landau                                     | (ORST)             |
| CAPSTICK   | 86  | PR D34 2809   | +Isgur                                      | (TNTO)             |
| JENNINGS   | 86  | PL B176 229   |   | (TRIU)             |
| MALTMAN  | 86  | PR D34 1372   | +Isgur                                      | (LANL, TNTO)       |
| ZHONG  | 86  | PL B171 471   | +Thomas, Jennings, Barrett                  | (ADLD, TRIU, SURR) |
| BURKHARDT  | 85  | NP A440 653   | +Lowe, Rosenthal                            | (NOTT, BIRM, WMIU) |
| DAREWYCH   | 85  | PR D32 1765   | +Koniuk, Isgur                              | (YORKC, TNTO)      |
| VEIT   | 85  | PR D31 1033   | +Jennings, Thomas, Barrett                  | (TRIU, ADLD, SURR) |
| KIANG  | 84  | PR C30 1638   | +Kumar, Nogami, VanDijk                     | (DALH, MCMS)       |
| MILLER   | 84  |   |   | (LOUC)             |
| Conf. Intersections between Particle and Nuclear Physics, p. 783 |     |   |   |                    |
| VANDIJK  | 84  | PR D30 937  |   | (MCMS)             |
| VEIT   | 84  | PL 137B 415   | +Jennings, Barrett, Thomas                  | (TRIU, SURR, CERN) |
| DALITZ   | 82  |   | +McGinley, Belyea, Anthony                  | (OXFTP)            |
| Heidelberg Conf., p. 201   |     |   |   |                    |
| DALITZ   | 81  |   | +McGinley                                   | (OXFTP)            |
| Low and Intermediate Energy Kaon-Nucleon Physics, p.381          |     |   |   |                    |
| MARTIN   | 81B | Low and Intermediate Energy Kaon-Nucleon Phys., p. 97 |   | (DURH)             |
| OADES  | 77  | NC 42A 462  | +Rasche                                     | (AARH, ZURI)       |
| SHAW   | 73  | Purdue Conf. 417                                      |   | (UCI)              |
| BARBARO-...  | 72  | LBL-555   | Barbaro-Galtieri                            | (LBL)              |
| DOBSON   | 72  | PR D6 3256  | +McElhaney                                  | (HAWA)             |
| RAJASEKA...  | 72  | PR D5 610   | Rajasekaran                                 | (TATA)             |
| Earlier papers also cited in RAJASEKARAN 72.                     |     |   |   |                    |
| CLINE  | 71  | PRL 26 1194   | +Laumann, Mapp                              | (WISC)             |
| MARTIN   | 71  | PL 35B 62   | +Martin, Ross                               | (DURH, LOUC, RHEL) |
| DALITZ   | 67  | PR 153 1617   | +Wong, Rajasekaran                          | (OXFTP, BOMB)      |
| DONALD   | 66  | PL 22 711   | +Edwards, Lys, Nisar, Moore                 | (LIVP)             |
| KADYK  | 66  | PRL 17 599  | +Oren, Goldhaber, Goldhaber, Trilling       | (LRL)              |
| ABRAMS   | 65  | PR 139B 454   | +Sechi-Zorn                                 | (UMD)              |