

$N(1520) D_{13}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-) \text{ Status: } ****$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

 $N(1520)$ BREIT-WIGNER MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1515 to 1530 (≈ 1520) OUR ESTIMATE			
1524 \pm 4	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
1525 \pm 10	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
1519 \pm 4	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1516 \pm 10	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
1515	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
1526 \pm 18	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
1510	LI	93	IPWA $\gamma N \rightarrow \pi N$
1504	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
1503	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
1510	BERENDS	77	IPWA $\gamma N \rightarrow \pi N$
1510	¹ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
1520	² LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

 $N(1520)$ BREIT-WIGNER WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
110 to 135 (≈ 120) OUR ESTIMATE			
124 \pm 8	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
120 \pm 15	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
114 \pm 7	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
106 \pm 4	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
106	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
143 \pm 32	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
120	LI	93	IPWA $\gamma N \rightarrow \pi N$
124	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
183	BAKER	79	DPWA $\pi^- p \rightarrow n\eta$
135	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
105	BERENDS	77	IPWA $\gamma N \rightarrow \pi N$
110	¹ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
150	² LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

$N(1520)$ POLE POSITION**REAL PART**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1505 to 1515 (\approx 1510) OUR ESTIMATE			
1515	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
1510	³ HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
1510 \pm 5	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1511	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
1514 or 1511	⁴ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
1508 or 1505	¹ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

–2×IMAGINARY PART

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
110 to 120 (\approx 115) OUR ESTIMATE			
110	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
120	³ HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
114 \pm 10	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
108	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
146 or 137	⁴ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
109 or 107	¹ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

 $N(1520)$ ELASTIC POLE RESIDUE**MODULUS $|r|$**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
34	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
32	HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
35 \pm 2	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
33	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90

PHASE θ

<u>VALUE ($^\circ$)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
– 8	HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
–12 \pm 5	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
–10	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90

N(1520) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	50–60 %
Γ_2 $N\eta$	
Γ_3 $N\pi\pi$	40–50 %
Γ_4 $\Delta\pi$	15–25 %
Γ_5 $\Delta(1232)\pi$, <i>S</i> -wave	5–12 %
Γ_6 $\Delta(1232)\pi$, <i>D</i> -wave	10–14 %
Γ_7 $N\rho$	15–25 %
Γ_8 $N\rho$, $S=1/2$, <i>D</i> -wave	
Γ_9 $N\rho$, $S=3/2$, <i>S</i> -wave	
Γ_{10} $N\rho$, $S=3/2$, <i>D</i> -wave	
Γ_{11} $N(\pi\pi)_{S\text{-wave}}^{I=0}$	<8 %
Γ_{12} $p\gamma$	0.46–0.56 %
Γ_{13} $p\gamma$, helicity=1/2	0.001–0.034 %
Γ_{14} $p\gamma$, helicity=3/2	0.44–0.53 %
Γ_{15} $n\gamma$	0.30–0.53 %
Γ_{16} $n\gamma$, helicity=1/2	0.04–0.10 %
Γ_{17} $n\gamma$, helicity=3/2	0.25–0.45 %

N(1520) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$					Γ_1/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.5 to 0.6 OUR ESTIMATE					
0.59±0.03	MANLEY	92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.58±0.03	CUTKOSKY	80	IPWA	$\pi N \rightarrow \pi N$	
0.54±0.03	HOEHLER	79	IPWA	$\pi N \rightarrow \pi N$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.61	ARNDT	95	DPWA	$\pi N \rightarrow N\pi$	
0.46±0.06	BATINIC	95	DPWA	$\pi N \rightarrow N\pi, N\eta$	

$\Gamma(N\eta)/\Gamma_{\text{total}}$					Γ_2/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.001±0.002	BATINIC	95	DPWA	$\pi N \rightarrow N\pi, N\eta$	

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1520) \rightarrow N\eta$	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.02	BAKER	79	DPWA $\pi^- p \rightarrow n\eta$
+0.011	FELTESSE	75	DPWA Soln A; see BAKER 79

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1520) \rightarrow \Delta(1232)\pi$, S-wave	$(\Gamma_1 \Gamma_5)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.26 to -0.20 OUR ESTIMATE			
-0.18 ± 0.05	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
-0.26	^{1,5} LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
-0.24	² LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1520) \rightarrow \Delta(1232)\pi$, D-wave	$(\Gamma_1 \Gamma_6)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.28 to -0.24 OUR ESTIMATE			
-0.29 ± 0.03	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
-0.21	^{1,5} LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
-0.30	² LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1520) \rightarrow N\rho$, S=3/2, S-wave	$(\Gamma_1 \Gamma_9)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.35 to -0.31 OUR ESTIMATE			
-0.35 ± 0.03	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
-0.35	^{1,5} LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
-0.24	² LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1520) \rightarrow N(\pi\pi)_{S\text{-wave}}^{I=0}$	$(\Gamma_1 \Gamma_{11})^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.22 to -0.06 OUR ESTIMATE			
-0.13	^{1,5} LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
-0.17	² LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

N(1520) PHOTON DECAY AMPLITUDES

$N(1520) \rightarrow p\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.024 ± 0.009 OUR ESTIMATE			
-0.020 ± 0.007	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
-0.028 ± 0.014	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
-0.007 ± 0.004	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.032 ± 0.005	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
-0.032 ± 0.004	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
-0.031 ± 0.009	BRATASHEV...	80	DPWA $\gamma N \rightarrow \pi N$
-0.019 ± 0.007	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
-0.0430 ± 0.0063	ISHII	80	DPWA Compton scattering

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.020 ±0.002	LI	93	IPWA	$\gamma N \rightarrow \pi N$
-0.012	WADA	84	DPWA	Compton scattering
-0.016 ±0.008	BARBOUR	78	DPWA	$\gamma N \rightarrow \pi N$
-0.008	⁶ NOELLE	78		$\gamma N \rightarrow \pi N$
-0.021	BERENDS	77	IPWA	$\gamma N \rightarrow \pi N$
-0.005 ±0.005	FELLER	76	DPWA	$\gamma N \rightarrow \pi N$

$N(1520) \rightarrow p\gamma$, helicity-3/2 amplitude $A_{3/2}$

<u>VALUE (GeV^{-1/2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
+0.166 ±0.005 OUR ESTIMATE			
0.167 ±0.005	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
0.156 ±0.022	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
0.168 ±0.013	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
0.178 ±0.003	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
0.162 ±0.003	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
0.166 ±0.005	BRATASHEV...	80	DPWA $\gamma N \rightarrow \pi N$
0.167 ±0.010	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
0.1695 ±0.0014	ISHII	80	DPWA Compton scattering

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.167 ±0.002	LI	93	IPWA	$\gamma N \rightarrow \pi N$
0.168	WADA	84	DPWA	Compton scattering
+0.157 ±0.007	BARBOUR	78	DPWA	$\gamma N \rightarrow \pi N$
0.206	⁶ NOELLE	78		$\gamma N \rightarrow \pi N$
+0.075	BERENDS	77	IPWA	$\gamma N \rightarrow \pi N$
+0.164 ±0.008	FELLER	76	DPWA	$\gamma N \rightarrow \pi N$

$N(1520) \rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

<u>VALUE (GeV^{-1/2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.059 ±0.009 OUR ESTIMATE			
-0.048 ±0.008	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
-0.066 ±0.013	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.067 ±0.004	FUJII	81	DPWA $\gamma N \rightarrow \pi N$
-0.076 ±0.006	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
-0.071 ±0.011	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
-0.056 ±0.011	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
-0.050 ±0.014	TAKEDA	80	DPWA $\gamma N \rightarrow \pi N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.058 ±0.003	LI	93	IPWA	$\gamma N \rightarrow \pi N$
-0.055 ±0.014	BARBOUR	78	DPWA	$\gamma N \rightarrow \pi N$
-0.060	⁶ NOELLE	78		$\gamma N \rightarrow \pi N$

$N(1520) \rightarrow n\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE ($\text{GeV}^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
-0.139±0.011 OUR ESTIMATE			
-0.140±0.010	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
-0.124±0.009	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.158±0.003	FUJII	81	DPWA $\gamma N \rightarrow \pi N$
-0.147±0.008	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
-0.148±0.009	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
-0.144±0.015	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
-0.118±0.011	TAKEDA	80	DPWA $\gamma N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
-0.131±0.003	LI	93	IPWA $\gamma N \rightarrow \pi N$
-0.141±0.015	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
-0.127	⁶ NOELLE	78	$\gamma N \rightarrow \pi N$

 $N(1520)$ FOOTNOTES

- ¹ LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- ² From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- ³ See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- ⁴ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- ⁵ LONGACRE 77 considers this coupling to be well determined.
- ⁶ Converted to our conventions using $M = 1528$ MeV, $\Gamma = 187$ MeV from NOELLE 78.

 $N(1520)$ REFERENCES

For early references, see Physics Letters **111B** 70 (1982). For very early references, see Reviews of Modern Physics **37** 633 (1965).

ARNDT	96	PR C53 430	+Strakovsky, Workman	(VPI)
ARNDT	95	PR C52 2120	+Strakovsky, Workman, Pavan	(VPI, BRCO)
BATINIC	95	PR C51 2310	+Slaus, Svarc, Nefkens	(BOSK, UCLA)
HOEHLER	93	πN Newsletter 9 1		(KARL)
LI	93	PR C47 2759	+Arndt, Roper, Workman	(VPI)
MANLEY	92	PR D45 4002	+Saleski	(KENT) IJP
Also	84	PR D30 904	Manley, Arndt, Goradia, Teplitz	(VPI)
ARNDT	91	PR D43 2131	+Li, Roper, Workman, Ford	(VPI, TELE) IJP
WADA	84	NP B247 313	+Egawa, Imanishi, Ishii, Kato, Ukai+	(INUS)
CRAWFORD	83	NP B211 1	+Morton	(GLAS)
PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
AWAJI	81	Bonn Conf. 352	+Kajikawa	(NAGO)
Also	82	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+	(NAGO)
FUJII	81	NP B187 53	+Hayashii, Iwata, Kajikawa+	(NAGO, OSAK)
ARAI	80	Toronto Conf. 93		(INUS)
Also	82	NP B194 251	Arai, Fujii	(INUS)
BRATASHEV...	80	NP B166 525	Bratashevskij, Gorbenko, Derebchinskij+	(KFTI)

CRAWFORD	80	Toronto Conf. 107		(GLAS)
CUTKOSKY	80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP
ISHII	80	NP B165 189	+Egawa, Kato, Miyachi+	(KYOT, INUS)
TAKEDA	80	NP B168 17	+Arai, Fujii, Ikeda, Iwasaki+	(TOKY, INUS)
BAKER	79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+	(RHEL) IJP
HOEHLER	79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80	Toronto Conf. 3	Koch	(KARLT) IJP
BARBOUR	78	NP B141 253	+Crawford, Parsons	(GLAS)
LONGACRE	78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+	(LBL, SLAC)
NOELLE	78	PTP 60 778		(NAGO)
BERENDS	77	NP B136 317	+Donnachie	(LEID, MCHS) IJP
LONGACRE	77	NP B122 493	+Dolbeau	(SACL) IJP
Also	76	NP B108 365	Dolbeau, Triantis, Neveu, Cadiet	(SACL) IJP
FELLER	76	NP B104 219	+Fukushima, Horikawa, Kajikawa+	(NAGO, OSAK) IJP
FELTESSE	75	NP B93 242	+Ayed, Bareyre, Borgeaud, David+	(SACL) IJP
LONGACRE	75	PL 55B 415	+Rosenfeld, Lasinski, Smadja+	(LBL, SLAC) IJP
