

$N(1675) D_{15}$

$$I(J^P) = \frac{1}{2}(\frac{5}{2}^-) \text{ Status: } ****$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

 $N(1675)$ BREIT-WIGNER MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1670 to 1685 (\approx 1675) OUR ESTIMATE			
1676 \pm 2	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
1675 \pm 10	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
1679 \pm 8	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1673 \pm 5	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
1673	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
1683 \pm 19	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
1666	LI	93	IPWA $\gamma N \rightarrow \pi N$
1685	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
1670	SAXON	80	DPWA $\pi^- p \rightarrow \Lambda K^0$
1680	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
1650	¹ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
1660	² LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

 $N(1675)$ BREIT-WIGNER WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
140 to 180 (\approx 150) OUR ESTIMATE			
159 \pm 7	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
160 \pm 20	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
120 \pm 15	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
154 \pm 7	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
154	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
142 \pm 23	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
136	LI	93	IPWA $\gamma N \rightarrow \pi N$
191	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
40	SAXON	80	DPWA $\pi^- p \rightarrow \Lambda K^0$
88	BAKER	79	DPWA $\pi^- p \rightarrow n\eta$
192	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
130	¹ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
150	² LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

$N(1675)$ POLE POSITION**REAL PART**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1655 to 1665 (≈ 1660) OUR ESTIMATE			
1663	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
1656	³ HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
1660 \pm 10	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1655	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
1663 or 1668	⁴ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
1649 or 1650	¹ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

-2xIMAGINARY PART

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
125 to 155 (≈ 140) OUR ESTIMATE			
152	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
126	³ HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
140 \pm 10	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
124	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
146 or 171	⁴ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
127 or 127	¹ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

 $N(1675)$ ELASTIC POLE RESIDUE**MODULUS $|r|$**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
29	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
23	HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
31 \pm 5	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
28	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90

PHASE θ

<u>VALUE ($^\circ$)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
- 6	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
-22	HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
-30 \pm 10	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
-17	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90

N(1675) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	40–50 %
Γ_2 $N\eta$	
Γ_3 ΛK	<1 %
Γ_4 ΣK	
Γ_5 $N\pi\pi$	50–60 %
Γ_6 $\Delta\pi$	50–60 %
Γ_7 $\Delta(1232)\pi$, <i>D</i> -wave	
Γ_8 $\Delta(1232)\pi$, <i>G</i> -wave	
Γ_9 $N\rho$	< 1–3 %
Γ_{10} $N\rho$, $S=1/2$, <i>D</i> -wave	
Γ_{11} $N\rho$, $S=3/2$, <i>D</i> -wave	
Γ_{12} $N\rho$, $S=3/2$, <i>G</i> -wave	
Γ_{13} $N(\pi\pi)_{S\text{-wave}}^{I=0}$	
Γ_{14} $p\gamma$	0.004–0.023 %
Γ_{15} $p\gamma$, helicity=1/2	0.0–0.015 %
Γ_{16} $p\gamma$, helicity=3/2	0.0–0.011 %
Γ_{17} $n\gamma$	0.02–0.12 %
Γ_{18} $n\gamma$, helicity=1/2	0.006–0.046 %
Γ_{19} $n\gamma$, helicity=3/2	0.01–0.08 %

N(1675) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.4 to 0.5 OUR ESTIMATE	
0.47±0.02	MANLEY 92 IPWA $\pi N \rightarrow \pi N \ \& \ N\pi\pi$
0.38±0.05	CUTKOSKY 80 IPWA $\pi N \rightarrow \pi N$
0.38±0.03	HOEHLER 79 IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●	
0.38	ARNDT 95 DPWA $\pi N \rightarrow N\pi$
0.31±0.06	BATINIC 95 DPWA $\pi N \rightarrow N\pi, N\eta$
$\Gamma(N\eta)/\Gamma_{\text{total}}$	Γ_2/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●	
0.001±0.001	BATINIC 95 DPWA $\pi N \rightarrow N\pi, N\eta$
$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1675) \rightarrow N\eta$	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●	
−0.07	BAKER 79 DPWA $\pi^- p \rightarrow n\eta$
+0.009	FELTESSE 75 DPWA Soln A; see BAKER 79

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1675) \rightarrow \Lambda K$ $(\Gamma_1 \Gamma_3)^{1/2} / \Gamma$
VALUE DOCUMENT ID TECN COMMENT
 ± 0.04 to ± 0.08 OUR ESTIMATE

-0.01	BELL	83	DPWA	$\pi^- p \rightarrow \Lambda K^0$
+0.036	⁵ SAXON	80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.034 ± 0.006	DEVENISH	74B		Fixed- t dispersion rel.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1675) \rightarrow \Sigma K$ $(\Gamma_1 \Gamma_4)^{1/2} / \Gamma$
VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.003	⁶ DEANS	75	DPWA	$\pi N \rightarrow \Sigma K$

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1675) \rightarrow \Delta(1232)\pi, D\text{-wave}$ $(\Gamma_1 \Gamma_7)^{1/2} / \Gamma$
VALUE DOCUMENT ID TECN COMMENT

+0.46 to +0.50 OUR ESTIMATE				
$+0.496 \pm 0.003$	MANLEY	92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
+0.46	^{1,7} LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.50	² LONGACRE	75	IPWA	$\pi N \rightarrow N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.5	⁸ NOVOSELLER	78	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1675) \rightarrow N\rho, S=1/2, D\text{-wave}$ $(\Gamma_1 \Gamma_{10})^{1/2} / \Gamma$
VALUE DOCUMENT ID TECN COMMENT

$+0.04 \pm 0.02$	MANLEY	92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
------------------	--------	----	------	--------------------------------------

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1675) \rightarrow N\rho, S=3/2, D\text{-wave}$ $(\Gamma_1 \Gamma_{11})^{1/2} / \Gamma$
VALUE DOCUMENT ID TECN COMMENT

-0.12 to -0.06 OUR ESTIMATE				
-0.03 ± 0.02	MANLEY	92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
-0.15	^{1,7} LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1675) \rightarrow N(\pi\pi)_{S\text{-wave}}^{I=0}$ $(\Gamma_1 \Gamma_{13})^{1/2} / \Gamma$
VALUE DOCUMENT ID TECN COMMENT

+0.03	^{1,7} LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
-------	-------------------------	----	------	-----------------------------

$N(1675)$ PHOTON DECAY AMPLITUDES **$N(1675) \rightarrow p\gamma$, helicity-1/2 amplitude $A_{1/2}$**

<u>VALUE (GeV^{-1/2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
+0.019±0.008 OUR ESTIMATE			
0.015±0.010	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
0.021±0.011	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
0.034±0.005	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
0.006±0.005	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
0.006±0.004	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
0.023±0.015	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.012±0.002	LI	93	IPWA $\gamma N \rightarrow \pi N$
+0.022±0.010	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
+0.034±0.004	FELLER	76	DPWA $\gamma N \rightarrow \pi N$

 $N(1675) \rightarrow p\gamma$, helicity-3/2 amplitude $A_{3/2}$

<u>VALUE (GeV^{-1/2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
+0.015±0.009 OUR ESTIMATE			
0.010±0.007	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
0.015±0.009	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
0.024±0.008	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
0.030±0.004	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
0.029±0.004	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
0.003±0.012	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.021±0.002	LI	93	IPWA $\gamma N \rightarrow \pi N$
+0.015±0.006	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
+0.019±0.009	FELLER	76	DPWA $\gamma N \rightarrow \pi N$

 $N(1675) \rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

<u>VALUE (GeV^{-1/2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.043±0.012 OUR ESTIMATE			
-0.049±0.010	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
-0.057±0.024	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.033±0.004	FUJII	81	DPWA $\gamma N \rightarrow \pi N$
-0.039±0.017	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
-0.025±0.027	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
-0.059±0.015	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
-0.021±0.011	TAKEDA	80	DPWA $\gamma N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
-0.060±0.003	LI	93	IPWA $\gamma N \rightarrow \pi N$
-0.066±0.020	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$

$N(1675) \rightarrow n\gamma$, helicity-3/2 amplitude $A_{3/2}$

<u>VALUE (GeV^{-1/2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.058±0.013 OUR ESTIMATE			
-0.051±0.010	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
-0.077±0.018	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.069±0.004	FUJII	81	DPWA $\gamma N \rightarrow \pi N$
-0.066±0.026	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
-0.071±0.022	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
-0.059±0.020	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
-0.030±0.012	TAKEDA	80	DPWA $\gamma N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
-0.074±0.003	LI	93	IPWA $\gamma N \rightarrow \pi N$
-0.073±0.014	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$

 $N(1675)$ FOOTNOTES

- ¹ LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- ² From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- ³ See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- ⁴ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- ⁵ SAXON 80 finds the coupling phase is near 90°.
- ⁶ The range given is from the four best solutions. DEANS 75 disagrees with $\pi^+ p \rightarrow \Sigma^+ K^+$ data of WINNIK 77 around 1920 MeV.
- ⁷ LONGACRE 77 considers this coupling to be well determined.
- ⁸ A Breit-Wigner fit to the HERNDON 75 IPWA.

 $N(1675)$ REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

ARNDT	96	PR C53 430	+Strakovsky, Workman	(VPI)
ARNDT	95	PR C52 2120	+Strakovsky, Workman, Pavan	(VPI, BRCO)
BATINIC	95	PR C51 2310	+Slaus, Svarc, Nefkens	(BOSK, UCLA)
HOEHLER	93	πN Newsletter 9 1		(KARL)
LI	93	PR C47 2759	+Arndt, Roper, Workman	(VPI)
MANLEY	92	PR D45 4002	+Saleski	(KENT) IJP
Also	84	PR D30 904	Manley, Arndt, Goradia, Teplitz	(VPI)
ARNDT	91	PR D43 2131	+Li, Roper, Workman, Ford	(VPI, TELE) IJP
BELL	83	NP B222 389	+Blissett, Broome, Daley, Hart, Lintern+	(RL) IJP
CRAWFORD	83	NP B211 1	+Morton	(GLAS)
PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
AWAJI	81	Bonn Conf. 352	+Kajikawa	(NAGO)
Also	82	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+	(NAGO)
FUJII	81	NP B187 53	+Hayashii, Iwata, Kajikawa+	(NAGO, OSAK)
ARAI	80	Toronto Conf. 93		(INUS)
Also	82	NP B194 251	Arai, Fujii	(INUS)
CRAWFORD	80	Toronto Conf. 107		(GLAS)
CUTKOSKY	80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP

SAXON	80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+	(RHEL, BRIS) IJP
TAKEDA	80	NP B168 17	+Arai, Fujii, Ikeda, Iwasaki+	(TOKY, INUS)
BAKER	79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+	(RHEL) IJP
HOEHLER	79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80	Toronto Conf. 3	Koch	(KARLT) IJP
BARBOUR	78	NP B141 253	+Crawford, Parsons	(GLAS)
LONGACRE	78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+	(LBL, SLAC)
NOVOSELLER	78	NP B137 509		(CIT) IJP
Also	78B	NP B137 445	Novoseller	(CIT) IJP
LONGACRE	77	NP B122 493	+Dolbeau	(SACL) IJP
Also	76	NP B108 365	Dolbeau, Triantis, Neveu, Cadiet	(SACL) IJP
WINNIK	77	NP B128 66	+Toaff, Revel, Goldberg, Berny	(HAIF) I
FELLER	76	NP B104 219	+Fukushima, Horikawa, Kajikawa+	(NAGO, OSAK) IJP
DEANS	75	NP B96 90	+Mitchell, Montgomery+	(SFLA, ALAH) IJP
FELTESSE	75	NP B93 242	+Ayed, Bareyre, Borgeaud, David+	(SACL) IJP
HERNDON	75	PR D11 3183	+Longacre, Miller, Rosenfeld+	(LBL, SLAC)
LONGACRE	75	PL 55B 415	+Rosenfeld, Lasinski, Smadja+	(LBL, SLAC) IJP
DEVENISH	74B	NP B81 330	+Froggatt, Martin	(DESY, NORD, LOUC)
