


$\Upsilon(1S)$

$$J^{PC} = 0^{-}(1^{- -})$$

$\Upsilon(1S)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9460.37 ± 0.21 OUR AVERAGE	Error includes scale factor of 2.7. See the ideogram below.		
9460.60 ± 0.09 ± 0.05	¹ BARU	92B REDE	$e^+ e^- \rightarrow$ hadrons
9460.6 ± 0.4	² ARTAMONOV	84 REDE	$e^+ e^- \rightarrow$ hadrons
9459.97 ± 0.11 ± 0.07	MACKAY	84 REDE	$e^+ e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
9460.59 ± 0.12	BARU	86 REDE	$e^+ e^- \rightarrow$ hadrons
¹ Superseding BARU 86.			
² Value includes data of ARTAMONOV 82.			



$\Upsilon(1S)$ mass (MeV)

$\Upsilon(1S)$ WIDTH

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>
52.5 ± 1.8 OUR EVALUATION	See the Note on Width Determinations of the Υ states

$\Upsilon(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\tau^+ \tau^-$	$(2.67^{+0.14}_{-0.16}) \%$	
Γ_2 $e^+ e^-$	$(2.52 \pm 0.17) \%$	
Γ_3 $\mu^+ \mu^-$	$(2.48 \pm 0.07) \%$	S=1.1
Hadronic decays		
Γ_4 $J/\psi(1S)$ anything	$(1.1 \pm 0.4) \times 10^{-3}$	
Γ_5 $\rho\pi$	$< 2 \times 10^{-4}$	CL=90%
Γ_6 $\pi^+ \pi^-$	$< 5 \times 10^{-4}$	CL=90%
Γ_7 $K^+ K^-$	$< 5 \times 10^{-4}$	CL=90%
Γ_8 $p\bar{p}$	$< 5 \times 10^{-4}$	CL=90%
Γ_9 $D^*(2010)^\pm$ anything		
Radiative decays		
Γ_{10} $\gamma 2h^+ 2h^-$	$(7.0 \pm 1.5) \times 10^{-4}$	
Γ_{11} $\gamma 3h^+ 3h^-$	$(5.4 \pm 2.0) \times 10^{-4}$	
Γ_{12} $\gamma 4h^+ 4h^-$	$(7.4 \pm 3.5) \times 10^{-4}$	
Γ_{13} $\gamma \pi^+ \pi^- K^+ K^-$	$(2.9 \pm 0.9) \times 10^{-4}$	
Γ_{14} $\gamma 2\pi^+ 2\pi^-$	$(2.5 \pm 0.9) \times 10^{-4}$	
Γ_{15} $\gamma 3\pi^+ 3\pi^-$	$(2.5 \pm 1.2) \times 10^{-4}$	
Γ_{16} $\gamma 2\pi^+ 2\pi^- K^+ K^-$	$(2.4 \pm 1.2) \times 10^{-4}$	
Γ_{17} $\gamma \pi^+ \pi^- p\bar{p}$	$(1.5 \pm 0.6) \times 10^{-4}$	
Γ_{18} $\gamma 2\pi^+ 2\pi^- p\bar{p}$	$(4 \pm 6) \times 10^{-5}$	
Γ_{19} $\gamma 2K^+ 2K^-$	$(2.0 \pm 2.0) \times 10^{-5}$	
Γ_{20} $\gamma \eta'(958)$	$< 1.3 \times 10^{-3}$	CL=90%
Γ_{21} $\gamma \eta$	$< 3.5 \times 10^{-4}$	CL=90%
Γ_{22} $\gamma f_2'(1525)$	$< 1.4 \times 10^{-4}$	CL=90%
Γ_{23} $\gamma f_2(1270)$	$< 1.3 \times 10^{-4}$	CL=90%
Γ_{24} $\gamma \eta(1440)$	$< 8.2 \times 10^{-5}$	CL=90%
Γ_{25} $\gamma f_J(1710) \rightarrow \gamma K\bar{K}$	$< 2.6 \times 10^{-4}$	CL=90%
Γ_{26} $\gamma f_0(2200) \rightarrow \gamma K^+ K^-$	$< 2 \times 10^{-4}$	CL=90%
Γ_{27} $\gamma f_J(2220) \rightarrow \gamma K^+ K^-$	$< 1.5 \times 10^{-5}$	CL=90%
Γ_{28} $\gamma \eta(2225) \rightarrow \gamma \phi\phi$	$< 3 \times 10^{-3}$	CL=90%
Γ_{29} γX $X = \text{pseudoscalar with } m < 7.2 \text{ GeV}$	$< 3 \times 10^{-5}$	CL=90%
Γ_{30} $\gamma X\bar{X}$ $X\bar{X} = \text{vectors with } m < 3.1 \text{ GeV}$	$< 1 \times 10^{-3}$	CL=90%

 $\Upsilon(1S) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(e^+ e^-) \times \Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_3/\Gamma$
VALUE (eV)	DOCUMENT ID TECN COMMENT
$31.2 \pm 1.6 \pm 1.7$	KOBEL 92 CBAL $e^+ e^- \rightarrow \mu^+ \mu^-$

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_0\Gamma_2/\Gamma$

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.216±0.027 OUR AVERAGE			
1.187±0.023±0.031	³ BARU	92B MD1	$e^+e^- \rightarrow \text{hadrons}$
1.23 ±0.02 ±0.05	³ JAKUBOWSKI	88 CBAL	$e^+e^- \rightarrow \text{hadrons}$
1.37 ±0.06 ±0.09	⁴ GILES	84B CLEO	$e^+e^- \rightarrow \text{hadrons}$
1.23 ±0.08 ±0.04	⁴ ALBRECHT	82 DASP	$e^+e^- \rightarrow \text{hadrons}$
1.13 ±0.07 ±0.11	⁴ NICZYPORUK	82 LENA	$e^+e^- \rightarrow \text{hadrons}$
1.09 ±0.25	⁴ BOCK	80 CNTR	$e^+e^- \rightarrow \text{hadrons}$
1.35 ±0.14	⁵ BERGER	79 PLUT	$e^+e^- \rightarrow \text{hadrons}$

³ Radiative corrections evaluated following KURAEV 85.

⁴ Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

⁵ Radiative corrections reevaluated by ALEXANDER 89 using $B(\mu\mu) = 0.026$.

$\Upsilon(1S)$ PARTIAL WIDTHS

 $\Gamma(e^+e^-) \qquad \Gamma_2$

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.32±0.04±0.03	⁶ ALBRECHT	95E ARG	$e^+e^- \rightarrow \text{hadrons}$

⁶ Applying the formula of Kuraev and Fadin.

$\Upsilon(1S)$ BRANCHING RATIOS

 $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}} \qquad \Gamma_1/\Gamma$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0267^{+0.0014}_{-0.0016} OUR AVERAGE				
0.0261±0.0012 ^{+0.0009} _{-0.0013}	25k	CINABRO	94B CLE2	$e^+e^- \rightarrow \tau^+\tau^-$
0.027 ±0.004 ±0.002		⁷ ALBRECHT	85C ARG	$\Upsilon(2S) \rightarrow$
0.034 ±0.004 ±0.004		GILES	83 CLEO	$e^+e^- \rightarrow \tau^+\tau^-$ $\pi^+\pi^-\tau^+\tau^-$

⁷ Using $B(\Upsilon(1S) \rightarrow ee) = B(\Upsilon(1S) \rightarrow \mu\mu) = 0.0256$; not used for width evaluations.

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}} \qquad \Gamma_3/\Gamma$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0248±0.0007 OUR AVERAGE	Error includes scale factor of 1.1.			
0.0212±0.0020±0.0010		⁸ BARU	92 MD1	$e^+e^- \rightarrow$ $\mu^+\mu^-$
0.0231±0.0012±0.0010		⁸ KOBEL	92 CBAL	$e^+e^- \rightarrow$ $\mu^+\mu^-$
0.0252±0.0007±0.0007		CHEN	89B CLEO	$e^+e^- \rightarrow$ $\mu^+\mu^-$
0.0261±0.0009±0.0011		KAARSBERG	89 CSB2	$e^+e^- \rightarrow$ $\mu^+\mu^-$
0.0230±0.0025±0.0013	86	ALBRECHT	87 ARG	$\Upsilon(2S) \rightarrow$ $\mu^+\mu^-$
0.029 ±0.003 ±0.002	864	BESSON	84 CLEO	$\Upsilon(2S) \rightarrow$ $\pi^+\pi^-\mu^+\mu^-$ $\pi^+\pi^-\mu^+\mu^-$

0.027 ± 0.003 ± 0.003	ANDREWS	83	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.032 ± 0.013 ± 0.003	ALBRECHT	82	DASP	$e^+e^- \rightarrow \mu^+\mu^-$
0.038 ± 0.015 ± 0.002	NICZYPORUK	82	LENA	$e^+e^- \rightarrow \mu^+\mu^-$
0.014 ^{+0.034} _{-0.014}	BOCK	80	CNTR	$e^+e^- \rightarrow \mu^+\mu^-$
0.022 ± 0.020	BERGER	79	PLUT	$e^+e^- \rightarrow \mu^+\mu^-$

⁸ Taking into account interference between the resonance and continuum.

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$		Γ_2/Γ			
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.0252 ± 0.0017 OUR AVERAGE					
0.0242 ± 0.0014 ± 0.0014	307	ALBRECHT	87 ARG	$\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$	
0.028 ± 0.003 ± 0.002	826	BESSON	84 CLEO	$\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$	
0.051 ± 0.030		BERGER	80c PLUT	$e^+e^- \rightarrow e^+e^-$	

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$		Γ_4/Γ			
<u>VALUE (units 10⁻³)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 0.68	90	ALBRECHT	92J ARG	$e^+e^- \rightarrow e^+e^-X,$ $e^+e^- \rightarrow \mu^+\mu^-X$	
1.1 ± 0.4 ± 0.2		⁹ FULTON	89 CLEO	$e^+e^- \rightarrow \mu^+\mu^-X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.7	90	MASCHMANN	90 CBAL	$e^+e^- \rightarrow \text{hadrons}$	
< 20	90	NICZYPORUK	83 LENA		

⁹ Using $B((J/\psi) \rightarrow \mu^+\mu^-) = (6.9 \pm 0.9)\%$.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$		Γ_6/Γ			
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 5	90	BARU	92 MD1	$\Upsilon(1S) \rightarrow \pi^+\pi^-$	

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$		Γ_7/Γ			
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 5	90	BARU	92 MD1	$\Upsilon(1S) \rightarrow K^+K^-$	

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$		Γ_8/Γ			
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 5	90	¹⁰ BARU	96 MD1	$\Upsilon(1S) \rightarrow p\bar{p}$	

¹⁰ Supersedes BARU 92 in this node.

$\Gamma(\gamma X)/\Gamma_{\text{total}}$		Γ_{29}/Γ			
<i>(X = pseudoscalar with $m < 7.2$ GeV)</i>					
<u>VALUE (units 10⁻⁵)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 3	90	¹¹ BALEST	95 CLEO	$e^+e^- \rightarrow \gamma + X$	

¹¹ For a noninteracting pseudoscalar X with mass < 7.2 GeV.

$$\Gamma(\gamma X \bar{X})/\Gamma_{\text{total}} \qquad \Gamma_{30}/\Gamma$$

($X \bar{X}$ = vectors with $m < 3.1$ GeV)

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1	90	¹² BALEST	95 CLEO	$e^+ e^- \rightarrow \gamma + X \bar{X}$

¹²For a noninteracting vector X with mass < 3.1 GeV.

$$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \qquad \Gamma_{14}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.5 \pm 0.7 \pm 0.5$	26 ± 7	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$$\Gamma(\gamma \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}} \qquad \Gamma_{13}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.9 \pm 0.7 \pm 0.6$	29 ± 8	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$$\Gamma(\gamma \pi^+ \pi^- \rho \bar{\rho})/\Gamma_{\text{total}} \qquad \Gamma_{17}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.5 \pm 0.5 \pm 0.3$	22 ± 6	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$$\Gamma(\gamma 2K^+ 2K^-)/\Gamma_{\text{total}} \qquad \Gamma_{19}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.2 ± 0.2	2 ± 2	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$$\Gamma(\gamma 3\pi^+ 3\pi^-)/\Gamma_{\text{total}} \qquad \Gamma_{15}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.5 \pm 0.9 \pm 0.8$	17 ± 5	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$$\Gamma(\gamma 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}} \qquad \Gamma_{16}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.4 \pm 0.9 \pm 0.8$	18 ± 7	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$$\Gamma(\gamma 2\pi^+ 2\pi^- \rho \bar{\rho})/\Gamma_{\text{total}} \qquad \Gamma_{18}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.4 \pm 0.4 \pm 0.4$	7 ± 6	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$$\Gamma(\gamma 2h^+ 2h^-)/\Gamma_{\text{total}} \qquad \Gamma_{10}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$7.0 \pm 1.1 \pm 1.0$	80 ± 12	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$$\Gamma(\gamma 3h^+ 3h^-)/\Gamma_{\text{total}} \qquad \Gamma_{11}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.4 \pm 1.5 \pm 1.3$	39 ± 11	FULTON	90B CLEO	$e^+ e^- \rightarrow$ hadrons

$\Gamma(\gamma 4h^+ 4h^-)/\Gamma_{\text{total}}$			Γ_{12}/Γ		
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
$7.4 \pm 2.5 \pm 2.5$	36 ± 12	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$	

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$			Γ_5/Γ		
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
< 2	90	FULTON	90B	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 10	90	BLINOV	90 MD1	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$	
< 21	90	NICZYPORUK	83 LENA	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$	

$\Gamma(D^*(2010)^\pm \text{ anything})/\Gamma_{\text{total}}$			Γ_9/Γ		
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
< 19	90	¹³ ALBRECHT	92J ARG	$e^+ e^- \rightarrow D^0 \pi^\pm X$	
¹³ For $x_p > 0.2$.					

$\Gamma(\gamma\eta(1440))/\Gamma_{\text{total}}$			Γ_{24}/Γ		
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
< 8.2	90	¹⁴ FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ \pi^\mp K_S^0$	
¹⁴ Includes unknown branching ratio of $\eta(1440) \rightarrow K^\pm \pi^\mp K_S^0$.					

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$			Γ_{20}/Γ		
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
< 1.3	90	SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$	

$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$			Γ_{21}/Γ		
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
< 3.5	90	SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$	

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$			Γ_{22}/Γ		
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
< 14	90	¹⁵ FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 19.4	90	¹⁵ ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	
¹⁵ Assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$.					

$\Gamma(\gamma f_J(1710) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$			Γ_{25}/Γ		
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
< 2.6	90	¹⁶ ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.3	90	16 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 19	90	16 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K_S^0 K_S^0$
< 8	90	17 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
< 24	90	18 SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$

¹⁶ Assuming $B(f_J(1710) \rightarrow K\bar{K}) = 0.38$.

¹⁷ Assuming $B(f_J(1710) \rightarrow \pi\pi) = 0.04$.

¹⁸ Assuming $B(f_J(1710) \rightarrow \eta\eta) = 0.18$.

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 13	90	19 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 21	90	19 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
< 81	90	SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$

¹⁹ Using $B(f_2(1270) \rightarrow \pi\pi) = 0.84$.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.5	90	20 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.9	90	20 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 20	90	20 BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

²⁰ Including unknown branching ratio of $f_J(2220) \rightarrow K^+ K^-$.

$\Gamma(\gamma \eta(2225) \rightarrow \gamma \phi \phi)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.003	90	21 BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^- K^+ K^-$

²¹ Assuming that the $\eta(2225)$ decays only into $\phi\phi$.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.0002	90	22 BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

²² Assuming that the $f_0(2200)$ decays only into $K^+ K^-$.

$\tau(1S)$ REFERENCES

BARU	96	PRPL 267 71	+Blinov, Blinov, Bondar+	(NOVO)
ALBRECHT	95E	ZPHY C65 619	+Hamacher+	(ARGUS Collab.)
BALEST	95	PR D51 2053	+Cho, Ford, Johnson+	(CLEO Collab.)
CINABRO	94B	PL B340 129	+Liu, Saulnier, Wilson+	(CLEO Collab.)
ALBRECHT	92J	ZPHY C55 25	+Ehrlichmann, Hamacher+	(ARGUS Collab.)
BARU	92	ZPHY C54 229	+Beilin, Blinov+	(NOVO)
BARU	92B	ZPHY C56 547	+Blinov, Blinov, Bondar+	(NOVO)
KOBEL	92	ZPHY C53 193	+Antreasyan, Bartels, Besset+	(Crystal Ball Collab.)
BLINOV	90	PL B245 311	+Bondar+	(NOVO)
FULTON	90B	PR D41 1401	+Hempstead+	(CLEO Collab.)
MASCHMANN	90	ZPHY C46 555	+Antreasyan, Bartels, Besset+	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	+Boeckmann, Glaeser, Harder+	(ARGUS Collab.)
ALEXANDER	89	NP B320 45	+Bonvicini, Drell, Frey, Luth	(LBL, MICH, SLAC)
BARU	89	ZPHY C42 505	+Beilin, Blinov, Blinov+	(NOVO)
CHEN	89B	PR D39 3528	+Mcllwain, Miller+	(CLEO Collab.)
FULTON	89	PL B224 445	+Haas, Hempstead+	(CLEO Collab.)
KAARSBERG	89	PRL 62 2077	+Heintz+	(CUSB Collab.)
BUCHMUEL...	88	HE e^+e^- Physics 412	Buchmueller, Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI	88	ZPHY C40 49	+Antreasyan, Bartels+	(Crystal Ball Collab.) IGJPC
SCHMITT	88	ZPHY C40 199	+Antreasyan+	(Crystal Ball Collab.)
ALBRECHT	87	ZPHY C35 283	+Binder, Boeckmann, Glaeser+	(ARGUS Collab.)
BARU	86	ZPHY C30 551	+Blinov, Bondar, Bukin+	(NOVO)
ALBRECHT	85C	PL 154B 452	+Drescher, Heller+	(ARGUS Collab.)
KURAEV	85	SJNP 41 466	+Fadin	(NOVO)
Translated from YAF 41 733.				
ARTAMONOV	84	PL 137B 272	+Baru, Blinov, Bondar+	(NOVO)
BESSON	84	PR D30 1433	+Green, Hicks, Namjoshi, Sannes+	(CLEO Collab.)
GILES	84B	PR D29 1285	+Hassard, Hempstead, Kinoshita+	(CLEO Collab.)
MACKAY	84	PR D29 2483	+Hasard, Giles, Hempstead+	(CUSB Collab.)
ANDREWS	83	PRL 50 807	+Avery, Berkelman, Cassel+	(CLEO Collab.)
GILES	83	PRL 50 877	+ (HARV, OSU, ROCH, RUTG, SYRA, VAND+)	
NICZYPORUK	83	ZPHY C17 197	+Jakubowski, Zeludziejewicz+	(LENA Collab.)
ALBRECHT	82	PL 116B 383	+Hofmann+ (DESY, DORT, HEIDH, LUND, ITEP)	
ARTAMONOV	82	PL 118B 225	+Baru, Blinov, Bondar, Bukin, Groshev+	(NOVO)
NICZYPORUK	82	ZPHY C15 299	+Folger, Bienlein+	(LENA Collab.)
BERGER	80C	PL 93B 497	+Lackas, Raupach+	(PLUTO Collab.)
BOCK	80	ZPHY C6 125	+Blonar, Blum+ (HEIDP, MPIM, DESY, HAMB)	
BERGER	79	ZPHY C1 343	+Alexander+ (PLUTO Collab.)	

OTHER RELATED PAPERS

KOENIGS...	86	DESY 86/136	Koenigsmann	(DESY)
ALBRECHT	84	PL 134B 137	+Drescher, Heller+	(ARGUS Collab.)
ARTAMONOV	84	PL 137B 272	+Baru, Blinov, Bondar+	(NOVO)
ARTAMONOV	82	PL 118B 225	+Baru, Blinov, Bondar, Bukin, Groshev+	(NOVO)
BERGER	78	PL 76B 243	+Alexander, Daum+	(PLUTO Collab.)
BIENLEIN	78	PL 78B 360	+Glawe, Bock, Blonar+ (DESY, HAMB, HEIDP, MPIM)	
DARDEN	78	PL 76B 246	+Hofmann, Schubert+ (DESY, DORT, HEIDH, LUND)	
GARELICK	78	PR D18 945	+Gauthier, Hicks, Oliver+ (NEAS, WASH, TUFTS)	
KAPLAN	78	PRL 40 435	+Appel, Herb, Hom+ (STON, FNAL, COLU)	
YOH	78	PRL 41 684	+Herb, Hom, Lederman+ (COLU, FNAL, STON)	
COBB	77	PL 72B 273	+Iwata, Fabjan+ (BNL, CERN, SYRA, YALE)	
HERB	77	PRL 39 252	+Hom, Lederman, Appel, Ito+ (COLU, FNAL, STON)	
INNES	77	PRL 39 1240	+Appel, Brown, Herb, Hom+ (COLU, FNAL, STON)	