b-baryon ADMIXTURE (Λ_b , Ξ_b , Σ_b , Ω_b)

b-baryon ADMIXTURE MEAN LIFE

Each measurement of the *b*-baryon mean life is an average over an admixture of various *b* baryons which decay weakly. Different techniques emphasize different admixtures of produced particles, which could result in a different *b*-baryon mean life.

"OUR EVALUATION" is an average of the data listed below performed by the LEP B Lifetimes Working Group as described in our review "Production and Decay of b-flavored Hadrons" in the B^\pm Section of these Listings. The averaging procedure takes into account correlations between the measurements and asymmetric lifetime errors.

$VALUE (10^{-12} \text{ s})$	EVTS	DOCUMENT ID		TECN	COMMENT	
1.20±0.07 OUR EVALUATION						
$1.20\!\pm\!0.08\!\pm\!0.06$		¹ BARATE	98 D	ALEP	$e^+e^- \rightarrow Z$	
$1.46 ^{+ 0.22 + 0.07}_{- 0.21 - 0.09}$		ABREU	96 D	DLPH	Excess $\Lambda \ell^- \pi^+$, decay lengths	
$1.10^{igoplus 0.19}_{-0.17}\!\pm\!0.09$		ABREU	96 D	DLPH	Excess $\Lambda \mu^-$ impact parameters	
$1.16 \pm 0.11 \pm 0.06$		AKERS	96	OPAL	Excess $\Lambda \ell^-$, decay lengths and impact parameters	
$1.27^{igoplus 0.35}_{-0.29} \pm 0.09$		ABREU	95 S	DLPH	Excess $p\mu^-$, decay lengths	
\bullet \bullet We do not use	the following	g data for averages	s, fits,	limits,	etc. • • •	
$1.25\!\pm\!0.11\!\pm\!0.05$		² ABREU	96 D	DLPH	Combined result	
$1.05 {+0.12\atop -0.11} \pm 0.09$	290	BUSKULIC	95L	ALEP	Repl. by BARATE 98D	
$1.04^{+0.48}_{-0.38}{\pm}0.10$	11	³ ABREU	93F	DLPH	Excess $\Lambda\mu^-$, decay lengths	
$1.05^{+0.23}_{-0.20}\pm0.08$	157	⁴ AKERS	93	OPAL	Excess $\Lambda\ell^-$, decay lengths	
$1.12^{\ +\ 0.32}_{\ -\ 0.29}\!\pm\!0.16$	101	⁵ BUSKULIC	921	ALEP	Excess $\Lambda \ell^-$, impact parameters	

 $^{^{1}}$ Measured using the excess of $\Lambda\ell^{-}$, lepton impact parameter.

² Combined result of the three ABREU 96D methods and ABREU 95S.

³ ABREU 93F superseded by ABREU 96D.

⁴ AKERS 93 superseded by AKERS 96.

⁵ BUSKULIC 921 superseded by BUSKULIC 95L.

b-baryon ADMIXTURE $(\Lambda_b, \Xi_b, \Sigma_b, \Omega_b)$

These branching fractions are actually an average over weakly decaying b-baryons weighted by their production rates in Z decay (or high-energy $p\overline{p})$, branching ratios, and detection efficiencies. They scale with the LEP Λ_b production fraction B(b \rightarrow $\Lambda_b)$ and are evaluated for our value B(b \rightarrow $\Lambda_b) = (10.1^{+3.9}_{-3.1})\%$.

The branching fractions B(b-baryon $\to \Lambda \ell^- \overline{\nu}_\ell$ anything) and B($\Lambda_b^0 \to \Lambda_c^+ \ell^- \overline{\nu}_\ell$ anything) are not pure measurements because the underlying measured products of these with B($b \to \Lambda_b$) were used to determine B($b \to \Lambda_b$), as described in the note "Production and Decay of b-Flavored Hadrons."

	Mode	Fraction (Γ_i/Γ)
$\overline{\Gamma_1}$	$ ho\mu^-\overline{ u}$ anything	(4.9± 2.4) %
Γ_2	$arLambda \ell^- \overline{ u}_\ell$ anything	$(\ 3.1 + \ 1.0 \ 1.2) \%$
Γ_4	$\Lambda\ell^+ u_\ell$ anything $\Lambda_c^+\ell^-\overline{ u}_\ell$ anything	
Γ ₆	$\Lambda/\overline{\Lambda}$ anything	$(35 \begin{array}{cc} +12 \\ -14 \end{array})\%$
Γ ₇	$ar{arXi}^-\ell^-\overline{ u}_\ell$ anything	$(5.5^{+}_{-2.4}) \times 10^{-3}$

b-baryon ADMIXTURE $(\Lambda_b, \Xi_b, \Sigma_b, \Omega_b)$ BRANCHING RATIOS

$\Gamma(p\mu^-\overline{ u})$ anything $\Gamma(p\mu^-\overline{ u})$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.049 + 0.018 + 0.015 -0.015 - 0.019	125	⁶ ABREU	95s DLPH	$e^+e^- ightarrow Z$	

⁶ ABREU 95S reports [B(*b*-baryon $\to p\mu^-\overline{\nu}$ anything) \times B($\overline{b}\to \Lambda_b$)] = 0.0049 \pm 0.0011 $^{+0.0015}_{-0.0011}$. We divide by our best value B($\overline{b}\to \Lambda_b$) = $(10.1^{+3.9}_{-3.1})\times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda \ell^- \overline{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}$

The values and averages in this section serve only to show what values result if one assumes our $B(b \to \Lambda_h)$. They cannot be thought of as measurements since the

underlying product branching fractions were also used to determinine $B(b \to \Lambda_b)$ as described in the note on "Production and Decay of *b*-Flavored Hadrons."

described in the r	ote on " <i>EVTS</i>	Production and D <u>DOCUMENT I</u> E			ored Hadrons." <i>COMMENT</i>	~
0.031 + 0.010 OUR AVE						
-0.012 $0.032 \pm 0.004 + 0.010$ -0.012		⁷ BARATE	98 D	ALEP	$e^+e^- \rightarrow Z$	
$0.029 \pm 0.003 ^{+0.009}_{-0.011}$		⁸ AKERS	96	OPAL	Excess of $\Lambda\ell$	$^-$ over $\hbar\ell^+$
$0.030 \pm 0.007 {+0.009 \atop -0.011}$	262	⁹ ABREU	95 S	DLPH	Excess of $\Lambda\ell$	$^-$ over $\it \Lambda \ell^+$
$0.060 \pm 0.012 ^{+ 0.019}_{- 0.023}$	290	¹⁰ BUSKULIC	95L	ALEP	Excess of $\Lambda\ell$	$^-$ over $\hbar\ell^+$
• • • We do not use th	e followi	ng data for averag	ges, fits	, limits,	etc. • • •	
seen	157	¹¹ AKERS	93	OPAL	Excess of $\Lambda\ell$	$^-$ over $\varLambda\ell^+$
$0.069 \pm 0.020 {}^{+ 0.021}_{- 0.027}$	101	¹² BUSKULIC	921	ALEP	Excess of $\Lambda\ell$	$^-$ over $\it \Lambda \ell^+$
from using our best 8 AKERS 96 reports 0.00023 ± 0.00025. Our first error is the from using our best 9 ABREU 95S reports 0.0006 ± 0.0004. We first error is their exusing our best value 10 BUSKULIC 95L reports 0.0006 ± 0.0010. We first error is their exusing our best value 11 AKERS 93 supersed 12 BUSKULIC 92I reports 0.0010 ± 0.0018. We first error is their exusing our best value 11 AKERS 93 supersed 12 BUSKULIC 92I reports 0.0010 ± 0.0018. We first error is their exusing our best value 11 AKERS 93 supersed 12 BUSKULIC 92I reports 0.0010 ± 0.0018. We first error is their exusing our best value 11 AKERS 93 supersed 12 BUSKULIC 92I reports 0.0010 ± 0.0018. We first error is their exusing our best value 11 AKERS 93 supersed 12 BUSKULIC 92I reports 0.0010 ± 0.0018.	[B(b-bar) We divide ir experiments. In the divide experiments.	yon $\rightarrow \Lambda \ell^- \overline{\nu}_{\ell}$ de by our best value tryon $\rightarrow \Lambda \ell^- \overline{\nu}_{\ell}$ by our best value tryon $\rightarrow \Lambda \ell^-$ by our best value tryon by our best value tryon by our best value tryon by BUSKUL	anythin alue $\mathrm{B}(\overline{b})$ our set $\mathrm{B}(\overline{b})$ anythin $\mathrm{B}(\overline{b})$ second $\mathrm{B}(\overline{b})$ anythin $\mathrm{B}(\overline{b})$ second $\mathrm{B}(\overline{b})$ second $\mathrm{B}(\overline{b})$ anythin $\mathrm{B}(\overline{b})$ second	$(a,b) \times (b,b) \times (b,b) \times (a,b) = (a,b) \times (a,b$	$(\overline{b} ightarrow \Lambda_b)] = b = (10.1^{+3}_{-3})$ for is the systemation of the systematic	$0.00291 \pm 0.00291 \pm 0.00291 \pm 0.00291 \pm 0.0030 \pm 0.0030 \pm 0.0061 \pm 0.0061 \pm 0.0061 \pm 0.0070 \pm 0.0070$
$\Gamma(\Lambda \ell^+ \nu_\ell \text{ anything})/\nu_{ALUE}$	Γ(Λany	•)	TECN	<u>COMMENT</u>	Γ_3/Γ_4
$0.070 \pm 0.012 \pm 0.007$	_				$e^+e^- \rightarrow Z$	
$\Gamma(\Lambda/\overline{\Lambda})$ anything Γ_{t}	otal	<u>DOCUMENT II</u>)	<u>TECN</u>	<u>COMMENT</u>	Γ ₆ /Γ
$0.35^{+0.12}_{-0.14}$ OUR AVER	AGE					
$a_{00} + a_{00} + 0.12$		13	^-	0544		

HTTP://PDG.LBL.GOV

Page 3

- ¹³ ACKERSTAFF 97N reports $[B(b\text{-baryon} \to \Lambda/\overline{\Lambda}\text{anything}) \times B(\overline{b} \to \Lambda_b)] = 0.0393 \pm 0.0046 \pm 0.0037$. We divide by our best value $B(\overline{b} \to \Lambda_b) = (10.1^{+3.9}_{-3.1}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ¹⁴ ABREU 95C reports $0.28^{+0.17}_{-0.12}$ for B($\overline{b} \rightarrow \Lambda_b$) = 0.08 ± 0.02 . We rescale to our best value B($\overline{b} \rightarrow \Lambda_b$) = $(10.1^{+3.9}_{-3.1}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Xi^-\ell^-\overline{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}$

 Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
	·		

$0.0055^{+0.0020}_{-0.0024}$ OUR AVERAGE

$0.0053\!\pm\!0.0013\!+\!0.0016\\-0.0021$	¹⁵ BUSKULIC	96⊤ ALEP	Excess $\Xi^-\ell^-$ over $\Xi^-\ell^+$
$0.0058\!\pm\!0.0023\!+\!0.0018\\-0.0023$	¹⁶ ABREU	95∨ DLPH	Excess $\Xi^-\ell^-$ over $\Xi^-\ell^+$

- ¹⁵ BUSKULIC 96T reports [B(b-baryon $\to \bar{\Xi}^-\ell^-\overline{\nu}_\ell$ anything) \times B($\bar{b} \to \Lambda_b$)] = 0.00054 \pm 0.00011 \pm 0.00008. We divide by our best value B($\bar{b} \to \Lambda_b$) = $(10.1^{+3.9}_{-3.1}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ¹⁶ ABREU 95V reports $[B(b\text{-baryon} \to \overline{\Xi}^-\ell^-\overline{\nu}_\ell \text{ anything}) \times B(\overline{b} \to \Lambda_b)] = 0.00059 \pm 0.00021 \pm 0.0001$. We divide by our best value $B(\overline{b} \to \Lambda_b) = (10.1^{+3.9}_{-3.1}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

b-baryon ADMIXTURE (Λ_b , Ξ_b , Σ_b , Ω_b) REFERENCES

BARATE	98D	EPJ C2 197	R. Barate+	(ALEPH Collab.)
ACKERSTAFF	97N	ZPHY C74 423	K. Ackerstaff+	(OPAL Collab.)
ABREU	96D	ZPHY C71 199	+Adam, Adye, Agasi $+$	(DELPHI Collab.)
AKERS	96	ZPHY C69 195	+Alexander, Allison, Altekamp+	(OPAL Collab.)
BUSKULIC	96T	PL B384 449	+De Bonis, Decamp, Ghez+	(ALEPH Collab.)
ABREU	95C	PL B347 447	+Adam, Adye, Agasi $+$	(DELPHI Collab.)
ABREU	95S	ZPHY C68 375	+Adam, Adye, Agasi $+$	(DELPHI Collab.)
ABREU	95V	ZPHY C68 541	+Adam, Adye, Agasi $+$	(DELPHI Collab.)
BUSKULIC	95L	PL B357 685	+Casper, De Bonis, Decamp+	(ALEPH Collab.)
ABREU	93F	PL B311 379	+Adam, Adye, Agasi+	(DELPHI Collab.)
AKERS	93	PL B316 435	+Alexander, Allison, Anderson+	(OPAL Collab.)
BUSKULIC	92I	PL B297 449	+Decamp, Goy, Lees $+$	(ALEPH Collab.)