

$\Lambda(1670) S_{01}$

$I(J^P) = 0(\frac{1}{2}^-)$ Status: ****

The measurements of the mass, width, and elasticity published before 1974 are now obsolete and have been omitted. They were last listed in our 1982 edition Physics Letters **111B** (1982).

$\Lambda(1670)$ MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|------|--|
| 1660 to 1680 (≈ 1670) OUR ESTIMATE | | | |
| 1670.8 \pm 1.7 | KOISO | 85 | DPWA $K^- p \rightarrow \Sigma \pi$ |
| 1667 \pm 5 | GOPAL | 80 | DPWA $\bar{K} N \rightarrow \bar{K} N$ |
| 1671 \pm 3 | ALSTON-... | 78 | DPWA $\bar{K} N \rightarrow \bar{K} N$ |
| 1670 \pm 5 | GOPAL | 77 | DPWA $\bar{K} N$ multichannel |
| 1675 \pm 2 | HEPP | 76B | DPWA $K^- N \rightarrow \Sigma \pi$ |
| 1679 \pm 1 | KANE | 74 | DPWA $K^- p \rightarrow \Sigma \pi$ |
| 1665 \pm 5 | PREVOST | 74 | DPWA $K^- N \rightarrow \Sigma(1385)\pi$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 1669 \pm 2 | ABAEV | 96 | DPWA $\pi^- p \rightarrow \eta n$ |
| 1664 | ¹ MARTIN | 77 | DPWA $\bar{K} N$ multichannel |

$\Lambda(1670)$ WIDTH

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|------|--|
| 25 to 50 (≈ 35) OUR ESTIMATE | | | |
| 34.1 \pm 3.7 | KOISO | 85 | DPWA $K^- p \rightarrow \Sigma \pi$ |
| 29 \pm 5 | GOPAL | 80 | DPWA $\bar{K} N \rightarrow \bar{K} N$ |
| 29 \pm 5 | ALSTON-... | 78 | DPWA $\bar{K} N \rightarrow \bar{K} N$ |
| 45 \pm 10 | GOPAL | 77 | DPWA $\bar{K} N$ multichannel |
| 46 \pm 5 | HEPP | 76B | DPWA $K^- N \rightarrow \Sigma \pi$ |
| 40 \pm 3 | KANE | 74 | DPWA $K^- p \rightarrow \Sigma \pi$ |
| 19 \pm 5 | PREVOST | 74 | DPWA $K^- N \rightarrow \Sigma(1385)\pi$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 21 \pm 4 | ABAEV | 96 | DPWA $\pi^- p \rightarrow \eta n$ |
| 12 | ¹ MARTIN | 77 | DPWA $\bar{K} N$ multichannel |

$\Lambda(1670)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|------------------------------|--------------------------------|
| Γ_1 $N\bar{K}$ | 15–25 % |
| Γ_2 $\Sigma \pi$ | 20–60 % |
| Γ_3 $\Lambda \eta$ | 15–35 % |
| Γ_4 $\Sigma(1385)\pi$ | |

The above branching fractions are our estimates, not fits or averages.

$\Lambda(1670)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

| $\Gamma(N\bar{K})/\Gamma_{\text{total}}$ | Γ_1/Γ |
|--|---|
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |

0.15 to 0.25 OUR ESTIMATE

| | | | | |
|---|---------------------|----|------|---------------------------------|
| 0.18±0.03 | GOPAL | 80 | DPWA | $\bar{K}N \rightarrow \bar{K}N$ |
| 0.17±0.03 | ALSTON-... | 78 | DPWA | $\bar{K}N \rightarrow \bar{K}N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 0.20±0.03 | GOPAL | 77 | DPWA | See GOPAL 80 |
| 0.15 | ¹ MARTIN | 77 | DPWA | $\bar{K}N$ multichannel |

| $(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Sigma\pi$ | $(\Gamma_1\Gamma_2)^{1/2}/\Gamma$ |
|--|---|
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |

| | | | | |
|---|---------------------|-----|------|----------------------------------|
| −0.26±0.02 | KOISO | 85 | DPWA | $K^-p \rightarrow \Sigma\pi$ |
| −0.31±0.03 | GOPAL | 77 | DPWA | $\bar{K}N$ multichannel |
| −0.29±0.03 | HEPP | 76B | DPWA | $K^-N \rightarrow \Sigma\pi$ |
| −0.23±0.03 | LONDON | 75 | HLBC | $K^-p \rightarrow \Sigma^0\pi^0$ |
| −0.27±0.02 | KANE | 74 | DPWA | $K^-p \rightarrow \Sigma\pi$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| −0.13 | ¹ MARTIN | 77 | DPWA | $\bar{K}N$ multichannel |

| $(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Lambda\eta$ | $(\Gamma_1\Gamma_3)^{1/2}/\Gamma$ |
|--|---|
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |

| | | | | |
|---|---------------|-----|------|-----------------------------|
| +0.20±0.05 | BAXTER | 73 | DPWA | $K^-p \rightarrow$ neutrals |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 0.06 | ABAEV | 96 | DPWA | $\pi^-p \rightarrow \eta n$ |
| 0.24 | KIM | 71 | DPWA | K-matrix analysis |
| 0.26 | ARMENTEROS69C | HBC | | |
| 0.20 or 0.23 | BERLEY | 65 | HBC | |

| $(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Sigma(1385)\pi$ | $(\Gamma_1\Gamma_4)^{1/2}/\Gamma$ |
|--|---|
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |

| | | | | |
|------------|---------|----|------|------------------------------------|
| −0.18±0.05 | PREVOST | 74 | DPWA | $K^-N \rightarrow \Sigma(1385)\pi$ |
|------------|---------|----|------|------------------------------------|

$\Lambda(1670)$ FOOTNOTES

¹MARTIN 77 obtains identical resonance parameters from a T-matrix pole and from a Breit-Wigner fit.

$\Lambda(1670)$ REFERENCES

| | | | | |
|------------|-----|-------------------------------------|-----------------------------------|--------------------------------|
| ABAEV | 96 | PR C53 385 | +Nefkens | (UCLA) |
| KOISO | 85 | NP A433 619 | +Sai, Yamamoto, Kofler | (TOKY, MASA) |
| PDG | 82 | PL 111B | Roos, Porter, Aguilar-Benitez+ | (HELSE, CIT, CERN) |
| GOPAL | 80 | Toronto Conf. 159 | | (RHEL) IJP |
| ALSTON-... | 78 | PR D18 182 | Alston-Garnjost, Kenney+ | (LBL, MTHO, CERN) IJP |
| Also | 77 | PRL 38 1007 | Alston-Garnjost, Kenney+ | (LBL, MTHO, CERN) IJP |
| GOPAL | 77 | NP B119 362 | +Ross, VanHorn, McPherson+ | (LOIC, RHEL) IJP |
| MARTIN | 77 | NP B127 349 | +Pidcock, Moorhouse | (LOUC, GLAS) IJP |
| Also | 77B | NP B126 266 | Martin, Pidcock | (LOUC) |
| Also | 77C | NP B126 285 | Martin, Pidcock | (LOUC) IJP |
| HEPP | 76B | PL 65B 487 | +Braun, Grimm, Strobele+ | (CERN, HEIDH, MPIM) IJP |
| LONDON | 75 | NP B85 289 | +Yu, Boyd+ | (BNL, CERN, EPOL, ORSAY, TORI) |
| KANE | 74 | LBL-2452 | | (LBL) IJP |
| PREVOST | 74 | NP B69 246 | +Barloutaud+ | (SACL, CERN, HEID) |
| BAXTER | 73 | NP B67 125 | +Buckingham, Corbett, Dunn+ | (OXF) IJP |
| KIM | 71 | PRL 27 356 | | (HARV) IJP |
| Also | 70 | Duke Conf. 161 | Kim | (HARV) IJP |
| ARMENTEROS | 69C | Lund Paper 229 | +Baillon+ | (CERN, HEID, SACL) IJP |
| | | Values are quoted in LEVI-SETTI 69. | | |
| BERLEY | 65 | PRL 15 641 | +Connolly, Hart, Rahm, Stonehill+ | (BNL) IJP |