

# $\Lambda(1830) D_{05}$

$$I(J^P) = 0(\frac{5}{2}^-) \text{ Status: } ****$$

For results published before 1973 (they are now obsolete), see our 1982 edition Physics Letters **111B** (1982).

The best evidence for this resonance is in the  $\Sigma\pi$  channel.

## $\Lambda(1830)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1810 to 1830 (<math>\approx 1830</math>) OUR ESTIMATE</b>			
1831 $\pm$ 10	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
1825 $\pm$ 10	GOPAL	77	DPWA $\bar{K}N$ multichannel
1825 $\pm$ 1	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1817 or 1818	<sup>1</sup> MARTIN	77	DPWA $\bar{K}N$ multichannel

## $\Lambda(1830)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>60 to 110 (<math>\approx 95</math>) OUR ESTIMATE</b>			
100 $\pm$ 10	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
94 $\pm$ 10	GOPAL	77	DPWA $\bar{K}N$ multichannel
119 $\pm$ 3	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
56 or 56	<sup>1</sup> MARTIN	77	DPWA $\bar{K}N$ multichannel

## $\Lambda(1830)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $N\bar{K}$	3–10 %
$\Gamma_2$ $\Sigma\pi$	35–75 %
$\Gamma_3$ $\Sigma(1385)\pi$	>15 %
$\Gamma_4$ $\Sigma(1385)\pi, D\text{-wave}$	
$\Gamma_5$ $\Lambda\eta$	

The above branching fractions are our estimates, not fits or averages.

## $\Lambda(1830)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on  $\Lambda$  and  $\Sigma$  Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
<b>0.03 to 0.10 OUR ESTIMATE</b>				
0.08 $\pm$ 0.03	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$	
0.02 $\pm$ 0.02	ALSTON-...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				

0.04±0.03	GOPAL	77	DPWA	See GOPAL 80
0.04 or 0.04	<sup>1</sup> MARTIN	77	DPWA	$\bar{K}N$ multichannel

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$  in  $N\bar{K} \rightarrow \Lambda(1830) \rightarrow \Sigma \pi$   $(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.17±0.03	GOPAL	77	DPWA $\bar{K}N$ multichannel
-0.15±0.01	KANE	74	DPWA $K^- p \rightarrow \Sigma \pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.17 or -0.17	<sup>1</sup> MARTIN	77	DPWA	$\bar{K}N$ multichannel
----------------	---------------------	----	------	-------------------------

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$  in  $N\bar{K} \rightarrow \Lambda(1830) \rightarrow \Lambda \eta$   $(\Gamma_1 \Gamma_5)^{1/2} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.044±0.020	RADER	73	MPWA

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$  in  $N\bar{K} \rightarrow \Lambda(1830) \rightarrow \Sigma(1385) \pi$   $(\Gamma_1 \Gamma_3)^{1/2} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.141±0.014	<sup>2</sup> CAMERON	78	DPWA $K^- p \rightarrow \Sigma(1385) \pi$
+0.13 ±0.03	PREVOST	74	DPWA $K^- N \rightarrow \Sigma(1385) \pi$

### $\Lambda(1830)$ FOOTNOTES

<sup>1</sup> The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.

<sup>2</sup> The CAMERON 78 upper limit on G-wave decay is 0.03. The published sign has been changed to be in accord with the baryon-first convention.

### $\Lambda(1830)$ REFERENCES

PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159		(RHEL) IJP
ALSTON...	78	PR D18 182	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
Also	77	PRL 38 1007	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
CAMERON	78	NP B143 189	+Franeek, Gopal, Bacon, Butterworth+	(RHEL, LOIC) IJP
GOPAL	77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
MARTIN	77	NP B127 349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
Also	77B	NP B126 266	Martin, Pidcock	(LOUC)
Also	77C	NP B126 285	Martin, Pidcock	(LOUC) IJP
KANE	74	LBL-2452		(LBL) IJP
PREVOST	74	NP B69 246	+Barloutaud+	(SACL, CERN, HEID)
RADER	73	NC 16A 178	+Barloutaud+	(SACL, HEID, CERN, RHEL, CDEF)