

# $\Sigma(1880) P_{11}$

$$I(J^P) = 1(\frac{1}{2}^+) \quad \text{Status: } **$$

## OMITTED FROM SUMMARY TABLE

A  $P_{11}$  resonance is suggested by several partial-wave analyses, but with wide variations in the mass and other parameters. We list here all claims which lie well above the  $P_{11} \Sigma(1770)$ .

### $\Sigma(1880)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>\approx 1880</math> OUR ESTIMATE</b>			
1826 $\pm$ 20	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
1870 $\pm$ 10	CAMERON	78B	DPWA $K^- p \rightarrow N\bar{K}^*$
1847 or 1863	<sup>1</sup> MARTIN	77	DPWA $\bar{K}N$ multichannel
1960 $\pm$ 30	<sup>2</sup> BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
1985 $\pm$ 50	VANHORN	75	DPWA $K^- p \rightarrow \Lambda\pi^0$
1898	<sup>3</sup> LEA	73	DPWA Multichannel K-matrix
$\sim 1850$	ARMENTEROS70	70	IPWA $\bar{K}N \rightarrow \bar{K}N$
1950 $\pm$ 50	BARBARO-...	70	DPWA $K^- N \rightarrow \Lambda\pi$
1920 $\pm$ 30	LITCHFIELD	70	DPWA $K^- N \rightarrow \Lambda\pi$
1850	BAILEY	69	DPWA $\bar{K}N \rightarrow \bar{K}N$
1882 $\pm$ 40	SMART	68	DPWA $K^- N \rightarrow \Lambda\pi$

### $\Sigma(1880)$ WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
86 $\pm$ 15	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
80 $\pm$ 10	CAMERON	78B	DPWA $K^- p \rightarrow N\bar{K}^*$
216 or 220	<sup>1</sup> MARTIN	77	DPWA $\bar{K}N$ multichannel
260 $\pm$ 40	<sup>2</sup> BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
220 $\pm$ 140	VANHORN	75	DPWA $K^- p \rightarrow \Lambda\pi^0$
222	<sup>3</sup> LEA	73	DPWA Multichannel K-matrix
$\sim 30$	ARMENTEROS70	70	IPWA $\bar{K}N \rightarrow \bar{K}N$
200 $\pm$ 50	BARBARO-...	70	DPWA $K^- N \rightarrow \Lambda\pi$
170 $\pm$ 40	LITCHFIELD	70	DPWA $K^- N \rightarrow \Lambda\pi$
200	BAILEY	69	DPWA $\bar{K}N \rightarrow \bar{K}N$
222 $\pm$ 150	SMART	68	DPWA $K^- N \rightarrow \Lambda\pi$

### $\Sigma(1880)$ DECAY MODES

Mode
$\Gamma_1$ $N\bar{K}$
$\Gamma_2$ $\Lambda\pi$
$\Gamma_3$ $\Sigma\pi$
$\Gamma_4$ $N\bar{K}^*(892)$ , $S=1/2$ , $P$ -wave
$\Gamma_5$ $N\bar{K}^*(892)$ , $S=3/2$ , $P$ -wave

## $\Sigma(1880)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on  $\Lambda$  and  $\Sigma$  Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.06±0.02	GOPAL    80    DPWA $\bar{K}N \rightarrow \bar{K}N$
0.27 or 0.27	<sup>1</sup> MARTIN    77    DPWA $\bar{K}N$ multichannel
0.31	<sup>3</sup> LEA    73    DPWA    Multichannel K-matrix
0.20	ARMENTEROS70    IPWA $\bar{K}N \rightarrow \bar{K}N$
0.22	BAILEY    69    DPWA $\bar{K}N \rightarrow \bar{K}N$

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1880) \rightarrow \Lambda\pi$	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
−0.24 or −0.24	<sup>1</sup> MARTIN    77    DPWA $\bar{K}N$ multichannel
−0.12 ±0.02	<sup>2</sup> BAILLON    75    IPWA $\bar{K}N \rightarrow \Lambda\pi$
+0.05 <sup>+0.07</sup> −0.02	VANHORN    75    DPWA $K^- p \rightarrow \Lambda\pi^0$
−0.169±0.119	DEVENISH    74B    Fixed- <i>t</i> dispersion rel.
−0.30	<sup>3</sup> LEA    73    DPWA    Multichannel K-matrix
−0.09 ±0.04	BARBARO-...    70    DPWA $K^- N \rightarrow \Lambda\pi$
−0.14 ±0.03	LITCHFIELD    70    DPWA $K^- N \rightarrow \Lambda\pi$
−0.11 ±0.03	SMART    68    DPWA $K^- N \rightarrow \Lambda\pi$

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1880) \rightarrow \Sigma\pi$	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
+0.30 or +0.29	<sup>1</sup> MARTIN    77    DPWA $\bar{K}N$ multichannel
not seen	<sup>3</sup> LEA    73    DPWA    Multichannel K-matrix

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1880) \rightarrow N\bar{K}^*(892), S=1/2, P\text{-wave}$	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
−0.05±0.03	<sup>4</sup> CAMERON    78B    DPWA $K^- p \rightarrow N\bar{K}^*$

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1880) \rightarrow N\bar{K}^*(892), S=3/2, P\text{-wave}$	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
+0.11±0.03	CAMERON    78B    DPWA $K^- p \rightarrow N\bar{K}^*$

### $\Sigma(1880)$ FOOTNOTES

- <sup>1</sup> The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
- <sup>2</sup> From solution 1 of BAILLON 75; not present in solution 2.
- <sup>3</sup> Only unconstrained states from table 1 of LEA 73 are listed.
- <sup>4</sup> The published sign has been changed to be in accord with the baryon-first convention.

## Σ(1880) REFERENCES

GOPAL	80	Toronto Conf.	159		(RHEL) IJP
CAMERON	78B	NP B146	327	+FraneK, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
MARTIN	77	NP B127	349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
Also	77B	NP B126	266	Martin, Pidcock	(LOUC)
Also	77C	NP B126	285	Martin, Pidcock	(LOUC) IJP
BAILLON	75	NP B94	39	+Litchfield	(CERN, RHEL) IJP
VANHORN	75	NP B87	145		(LBL) IJP
Also	75B	NP B87	157	VanHorn	(LBL) IJP
DEVENISH	74B	NP B81	330	+Froggatt, Martin	(DESY, NORD, LOUC)
LEA	73	NP B56	77	+Martin, Moorhouse+	(RHEL, LOUC, GLAS, AARH) IJP
ARMENTEROS	70	Duke Conf.	123	+Baillon+	(CERN, HEID, SACL) IJP
BARBARO-...	70	Duke Conf.	173	Barbaro-Galtieri	(LRL) IJP
LITCHFIELD	70	NP B22	269		(RHEL) IJP
BAILEY	69	Thesis UCRL	50617		(LLL) IJP
SMART	68	PR 169	1330		(LRL) IJP

---