

$$I(J^P) = \frac{1}{2}(0^-)$$

D^\pm MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , and $D_s^{*\pm}$ mass and mass difference measurements.

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1869.3 ± 0.5 OUR FIT	Error includes scale factor of 1.1.			
1869.4 ± 0.5 OUR AVERAGE				
1870.0 ± 0.5 ± 1.0	317	BARLAG	90C ACCM	π^- Cu 230 GeV
1863 ± 4		DERRICK	84 HRS	$e^+ e^-$ 29 GeV
1869.4 ± 0.6		¹ TRILLING	81 RVUE	$e^+ e^-$ 3.77 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1875 ± 10	9	ADAMOVICH	87 EMUL	Photoproduction
1860 ± 16	6	ADAMOVICH	84 EMUL	Photoproduction
1868.4 ± 0.5		¹ SCHINDLER	81 MRK2	$e^+ e^-$ 3.77 GeV
1874 ± 5		GOLDHABER	77 MRK1	D^0 , D^+ recoil spectra
1868.3 ± 0.9		¹ PERUZZI	77 MRK1	$e^+ e^-$ 3.77 GeV
1874 ± 11		PICCOLO	77 MRK1	$e^+ e^-$ 4.03, 4.41 GeV
1876 ± 15	50	PERUZZI	76 MRK1	$K^\mp \pi^\pm \pi^\pm$

¹ PERUZZI 77 and SCHINDLER 81 errors do not include the 0.13% uncertainty in the absolute SPEAR energy calibration. TRILLING 81 uses the high precision $J/\psi(1S)$ and $\psi(2S)$ measurements of ZHOLENTZ 80 to determine this uncertainty and combines the PERUZZI 77 and SCHINDLER 81 results to obtain the value quoted.

D^\pm MEAN LIFE

Measurements with an error $> 0.1 \times 10^{-12}$ s are omitted from the average, and those with an error $> 0.2 \times 10^{-12}$ s have been omitted from the Listings.

<u>VALUE (10^{-12} s)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.057 ± 0.015 OUR AVERAGE				
1.048 ± 0.015 ± 0.011	9k	FRABETTI	94D E687	$D^+ \rightarrow K^- \pi^+ \pi^+$
1.075 ± 0.040 ± 0.018	2455	FRABETTI	91 E687	γ Be, $D^+ \rightarrow K^- \pi^+ \pi^+$
1.03 ± 0.08 ± 0.06	200	ALVAREZ	90 NA14	γ , $D^+ \rightarrow K^- \pi^+ \pi^+$
1.05 ^{+0.077} _{-0.072}	317	² BARLAG	90C ACCM	π^- Cu 230 GeV
1.05 ± 0.08 ± 0.07	363	ALBRECHT	88I ARG	$e^+ e^-$ 10 GeV
1.090 ± 0.030 ± 0.025	2992	RAAB	88 E691	Photoproduction

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.12	$\begin{matrix} +0.14 \\ -0.11 \end{matrix}$	149	AGUILAR-...	87D HYBR	$\pi^- p$ and pp
1.09	$\begin{matrix} +0.19 \\ -0.15 \end{matrix}$	59	BARLAG	87B ACCM	K^- and π^- 200 GeV
1.14	± 0.16	247	CSORNA	87 CLEO	$e^+ e^-$ 10 GeV
1.09	± 0.14	74	³ PALKA	87B SILI	π Be 200 GeV
0.86	$\begin{matrix} \pm 0.13 \\ -0.03 \end{matrix}$	48	ABE	86 HYBR	γp 20 GeV

² BARLAG 90C estimates the systematic error to be negligible.

³ PALKA 87B observes this in $D^+ \rightarrow \bar{K}^*(892) e \nu$.

D^+ DECAY MODES

D^- modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Inclusive modes		
Γ_1 e^+ anything	$(17.2 \pm 1.9) \%$	
Γ_2 K^- anything	$(24.2 \pm 2.8) \%$	S=1.4
Γ_3 \bar{K}^0 anything + K^0 anything	$(59 \pm 7) \%$	
Γ_4 K^+ anything	$(5.8 \pm 1.4) \%$	
Γ_5 η anything	[a] < 13 %	CL=90%
Γ_6 μ^+ anything		
Leptonic and semileptonic modes		
Γ_7 $\mu^+ \nu_\mu$	$(8 \begin{matrix} +17 \\ -5 \end{matrix}) \times 10^{-4}$	
Γ_8 $\bar{K}^0 \ell^+ \nu_\ell$	[b] $(6.8 \pm 0.8) \%$	
Γ_9 $\bar{K}^0 e^+ \nu_e$	$(6.7 \pm 0.9) \%$	
Γ_{10} $\bar{K}^0 \mu^+ \nu_\mu$	$(7.0 \begin{matrix} +3.0 \\ -2.0 \end{matrix}) \%$	
Γ_{11} $K^- \pi^+ e^+ \nu_e$	$(4.1 \begin{matrix} +0.9 \\ -0.7 \end{matrix}) \%$	
Γ_{12} $\bar{K}^*(892)^0 e^+ \nu_e$ $\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	$(3.2 \pm 0.33) \%$	
Γ_{13} $K^- \pi^+ e^+ \nu_e$ nonresonant	< 7 $\times 10^{-3}$	CL=90%
Γ_{14} $K^- \pi^+ \mu^+ \nu_\mu$	$(3.2 \pm 0.4) \%$	S=1.1
In the fit as $\frac{2}{3}\Gamma_{26} + \Gamma_{16}$, where $\frac{2}{3}\Gamma_{26} = \Gamma_{15}$.		
Γ_{15} $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ $\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	$(2.9 \pm 0.4) \%$	

Γ_{16}	$K^- \pi^+ \mu^+ \nu_\mu$ nonresonant	$(2.7 \pm 1.1) \times 10^{-3}$	
Γ_{17}	$\bar{K}^0 \pi^+ \pi^- e^+ \nu_e$		
Γ_{18}	$K^- \pi^+ \pi^0 e^+ \nu_e$		
Γ_{19}	$(\bar{K}^*(892) \pi)^0 e^+ \nu_e$	< 1.2	% CL=90%
Γ_{20}	$(\bar{K} \pi \pi)^0 e^+ \nu_e$ non- $\bar{K}^*(892)$	< 9	$\times 10^{-3}$ CL=90%
Γ_{21}	$K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	< 1.4	$\times 10^{-3}$ CL=90%
Γ_{22}	$\pi^0 \ell^+ \nu_\ell$	[c] $(3.1 \pm 1.5) \times 10^{-3}$	
Γ_{23}	$\pi^+ \pi^- e^+ \nu_e$		

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

Γ_{24}	$\bar{K}^*(892)^0 \ell^+ \nu_\ell$	[b] $(4.7 \pm 0.4) \%$	
Γ_{25}	$\bar{K}^*(892)^0 e^+ \nu_e$	$(4.8 \pm 0.5) \%$	
Γ_{26}	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$	$(4.4 \pm 0.6) \%$	S=1.1
Γ_{27}	$\rho^0 e^+ \nu_e$	$(2.2 \pm 0.8) \times 10^{-3}$	
Γ_{28}	$\rho^0 \mu^+ \nu_\mu$	$(2.7 \pm 0.7) \times 10^{-3}$	
Γ_{29}	$\phi e^+ \nu_e$	< 2.09	% CL=90%
Γ_{30}	$\phi \mu^+ \nu_\mu$	< 3.72	% CL=90%
Γ_{31}	$\eta \ell^+ \nu_\ell$	< 5	$\times 10^{-3}$ CL=90%
Γ_{32}	$\eta'(958) \mu^+ \nu_\mu$	< 9	$\times 10^{-3}$ CL=90%

Hadronic modes with a \bar{K} or $\bar{K}K\bar{K}$

Γ_{33}	$\bar{K}^0 \pi^+$	$(2.89 \pm 0.26) \%$	S=1.1
Γ_{34}	$K^- \pi^+ \pi^+$	[d] $(9.0 \pm 0.6) \%$	
Γ_{35}	$\bar{K}^*(892)^0 \pi^+$ $\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	$(1.27 \pm 0.13) \%$	
Γ_{36}	$\bar{K}_0^*(1430)^0 \pi^+$ $\times B(\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+)$	$(2.3 \pm 0.3) \%$	
Γ_{37}	$\bar{K}^*(1680)^0 \pi^+$ $\times B(\bar{K}^*(1680)^0 \rightarrow K^- \pi^+)$	$(3.7 \pm 0.8) \times 10^{-3}$	
Γ_{38}	$K^- \pi^+ \pi^+$ nonresonant	$(8.5 \pm 0.8) \%$	
Γ_{39}	$\bar{K}^0 \pi^+ \pi^0$	[d] $(9.7 \pm 3.0) \%$	S=1.1
Γ_{40}	$\bar{K}^0 \rho^+$	$(6.6 \pm 2.5) \%$	
Γ_{41}	$\bar{K}^*(892)^0 \pi^+$ $\times B(\bar{K}^{*0} \rightarrow \bar{K}^0 \pi^0)$	$(6.3 \pm 0.4) \times 10^{-3}$	
Γ_{42}	$\bar{K}^0 \pi^+ \pi^0$ nonresonant	$(1.3 \pm 1.1) \%$	
Γ_{43}	$K^- \pi^+ \pi^+ \pi^0$	[d] $(6.4 \pm 1.1) \%$	
Γ_{44}	$\bar{K}^*(892)^0 \rho^+$ total $\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	$(1.4 \pm 0.9) \%$	
Γ_{45}	$\bar{K}_1(1400)^0 \pi^+$ $\times B(\bar{K}_1(1400)^0 \rightarrow K^- \pi^+ \pi^0)$	$(2.2 \pm 0.6) \%$	
Γ_{46}	$K^- \rho^+ \pi^+$ total	$(3.1 \pm 1.1) \%$	
Γ_{47}	$K^- \rho^+ \pi^+$ 3-body	$(1.1 \pm 0.4) \%$	

Γ ₄₈	$\bar{K}^*(892)^0 \pi^+ \pi^0$ total	(4.5 ± 0.9) %	
	× B($\bar{K}^{*0} \rightarrow K^- \pi^+$)		
Γ ₄₉	$\bar{K}^*(892)^0 \pi^+ \pi^0$ 3-body	(2.8 ± 0.9) %	
	× B($\bar{K}^{*0} \rightarrow K^- \pi^+$)		
Γ ₅₀	$K^*(892)^- \pi^+ \pi^+$ 3-body	(7 ± 3) × 10 ⁻³	
	× B($K^{*-} \rightarrow K^- \pi^0$)		
Γ ₅₁	$K^- \pi^+ \pi^+ \pi^0$ nonresonant	[e] (1.2 ± 0.6) %	
Γ ₅₂	$\bar{K}^0 \pi^+ \pi^+ \pi^-$	[d] (7.0 ± 0.9) %	
Γ ₅₃	$\bar{K}^0 a_1(1260)^+$	(4.0 ± 0.9) %	
	× B($a_1(1260)^+ \rightarrow \pi^+ \pi^+ \pi^-$)		
Γ ₅₄	$\bar{K}_1(1400)^0 \pi^+$	(2.2 ± 0.6) %	
	× B($\bar{K}_1(1400)^0 \rightarrow \bar{K}^0 \pi^+ \pi^-$)		
Γ ₅₅	$K^*(892)^- \pi^+ \pi^+$ 3-body	(1.4 ± 0.6) %	
	× B($K^{*-} \rightarrow \bar{K}^0 \pi^-$)		
Γ ₅₆	$\bar{K}^0 \rho^0 \pi^+$ total	(4.2 ± 0.9) %	
Γ ₅₇	$\bar{K}^0 \rho^0 \pi^+$ 3-body	(5 ± 5) × 10 ⁻³	
Γ ₅₈	$\bar{K}^0 \pi^+ \pi^+ \pi^-$ nonresonant	(8 ± 4) × 10 ⁻³	
Γ ₅₉	$K^- \pi^+ \pi^+ \pi^+ \pi^-$	[d] (7.2 ± 1.0) × 10 ⁻³	
Γ ₆₀	$\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^-$	(5.4 ± 2.3) × 10 ⁻³	
	× B($\bar{K}^{*0} \rightarrow K^- \pi^+$)		
Γ ₆₁	$\bar{K}^*(892)^0 \rho^0 \pi^+$	(1.9 ± 1.1) × 10 ⁻³	
	× B($\bar{K}^{*0} \rightarrow K^- \pi^+$)		
Γ ₆₂	$\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^-$ no-ρ	(2.9 ± 1.1) × 10 ⁻³	
	× B($\bar{K}^{*0} \rightarrow K^- \pi^+$)		
Γ ₆₃	$K^- \rho^0 \pi^+ \pi^+$	(3.1 ± 0.9) × 10 ⁻³	
Γ ₆₄	$K^- \pi^+ \pi^+ \pi^+ \pi^-$ nonresonant	< 2.3 × 10 ⁻³	CL=90%
Γ ₆₅	$K^- \pi^+ \pi^+ \pi^0 \pi^0$	(2.2 ± 5.0) %	
		- 0.9) %	
Γ ₆₆	$\bar{K}^0 \pi^+ \pi^+ \pi^- \pi^0$	(5.4 ± 3.0) %	
		- 1.4) %	
Γ ₆₇	$\bar{K}^0 \pi^+ \pi^+ \pi^+ \pi^- \pi^-$	(8 ± 7) × 10 ⁻⁴	
Γ ₆₈	$K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^0$	(2.0 ± 1.8) × 10 ⁻³	
Γ ₆₉	$\bar{K}^0 \bar{K}^0 K^+$	(1.8 ± 0.8) %	

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

Γ ₇₀	$\bar{K}^0 \rho^+$	(6.6 ± 2.5) %	
Γ ₇₁	$\bar{K}^0 a_1(1260)^+$	(8.0 ± 1.7) %	
Γ ₇₂	$\bar{K}^0 a_2(1320)^+$	< 3 × 10 ⁻³	CL=90%
Γ ₇₃	$\bar{K}^*(892)^0 \pi^+$	(1.90 ± 0.19) %	
Γ ₇₄	$\bar{K}^*(892)^0 \rho^+$ total	[e] (2.1 ± 1.3) %	
Γ ₇₅	$\bar{K}^*(892)^0 \rho^+$ S-wave	[e] (1.6 ± 1.6) %	
Γ ₇₆	$\bar{K}^*(892)^0 \rho^+$ P-wave	< 1 × 10 ⁻³	CL=90%
Γ ₇₇	$\bar{K}^*(892)^0 \rho^+$ D-wave	(10 ± 7) × 10 ⁻³	

Γ_{78}	$\bar{K}^*(892)^0 \rho^+ D\text{-wave longitudinal}$	< 7	$\times 10^{-3}$	CL=90%
Γ_{79}	$\bar{K}_1(1270)^0 \pi^+$	< 7	$\times 10^{-3}$	CL=90%
Γ_{80}	$\bar{K}_1(1400)^0 \pi^+$	(4.9 ± 1.2)	%	
Γ_{81}	$\bar{K}^*(1410)^0 \pi^+$	< 7	$\times 10^{-3}$	CL=90%
Γ_{82}	$\bar{K}_0^*(1430)^0 \pi^+$	(3.7 ± 0.4)	%	
Γ_{83}	$\bar{K}^*(1680)^0 \pi^+$	(1.43 ± 0.30)	%	
Γ_{84}	$\bar{K}^*(892)^0 \pi^+ \pi^0$ total	(6.7 ± 1.4)	%	
Γ_{85}	$\bar{K}^*(892)^0 \pi^+ \pi^0$ 3-body	[e] (4.2 ± 1.4)	%	
Γ_{86}	$K^*(892)^- \pi^+ \pi^+$ total			
Γ_{87}	$K^*(892)^- \pi^+ \pi^+$ 3-body	(2.0 ± 0.9)	%	
Γ_{88}	$K^- \rho^+ \pi^+$ total	(3.1 ± 1.1)	%	
Γ_{89}	$K^- \rho^+ \pi^+$ 3-body	(1.1 ± 0.4)	%	
Γ_{90}	$\bar{K}^0 \rho^0 \pi^+$ total	(4.2 ± 0.9)	%	CL=90%
Γ_{91}	$\bar{K}^0 \rho^0 \pi^+$ 3-body	(5 ± 5)	$\times 10^{-3}$	
Γ_{92}	$\bar{K}^0 f_0(980) \pi^+$	< 5	$\times 10^{-3}$	CL=90%
Γ_{93}	$\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^-$	(8.1 ± 3.4)	$\times 10^{-3}$	S=1.7
Γ_{94}	$\bar{K}^*(892)^0 \rho^0 \pi^+$	$(2.9 \begin{smallmatrix} +1.7 \\ -1.5 \end{smallmatrix})$	$\times 10^{-3}$	S=1.8
Γ_{95}	$\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^-$ no- ρ	(4.3 ± 1.7)	$\times 10^{-3}$	
Γ_{96}	$K^- \rho^0 \pi^+ \pi^+$	(3.1 ± 0.9)	$\times 10^{-3}$	

Pionic modes

Γ_{97}	$\pi^+ \pi^0$	(2.5 ± 0.7)	$\times 10^{-3}$	
Γ_{98}	$\pi^+ \pi^+ \pi^-$	(3.6 ± 0.4)	$\times 10^{-3}$	
Γ_{99}	$\rho^0 \pi^+$	(1.05 ± 0.31)	$\times 10^{-3}$	
Γ_{100}	$\pi^+ \pi^+ \pi^-$ nonresonant	(2.2 ± 0.4)	$\times 10^{-3}$	
Γ_{101}	$\pi^+ \pi^+ \pi^- \pi^0$	$(1.9 \begin{smallmatrix} +1.5 \\ -1.2 \end{smallmatrix})$	%	
Γ_{102}	$\eta \pi^+ \times B(\eta \rightarrow \pi^+ \pi^- \pi^0)$	(1.7 ± 0.6)	$\times 10^{-3}$	
Γ_{103}	$\omega \pi^+ \times B(\omega \rightarrow \pi^+ \pi^- \pi^0)$	< 6	$\times 10^{-3}$	CL=90%
Γ_{104}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^-$	(2.1 ± 0.4)	$\times 10^{-3}$	
Γ_{105}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0$	$(2.9 \begin{smallmatrix} +2.9 \\ -2.0 \end{smallmatrix})$	$\times 10^{-3}$	

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

Γ_{106}	$\eta \pi^+$	(3.0 ± 0.6)	$\times 10^{-3}$	
Γ_{107}	$\rho^0 \pi^+$	(1.05 ± 0.31)	$\times 10^{-3}$	
Γ_{108}	$\omega \pi^+$	< 7	$\times 10^{-3}$	CL=90%
Γ_{109}	$\eta \rho^+$	< 7	$\times 10^{-3}$	CL=90%
Γ_{110}	$\eta'(958) \pi^+$	(5.0 ± 1.0)	$\times 10^{-3}$	
Γ_{111}	$\eta'(958) \rho^+$	< 5	$\times 10^{-3}$	CL=90%

Hadronic modes with a $K\bar{K}$ pair

Γ_{112}	$K^+\bar{K}^0$		$(7.4 \pm 1.0) \times 10^{-3}$	
Γ_{113}	$K^+K^-\pi^+$	[d]	$(8.8 \pm 0.8) \times 10^{-3}$	
Γ_{114}	$\phi\pi^+ \times B(\phi \rightarrow K^+K^-)$		$(3.0 \pm 0.3) \times 10^{-3}$	
Γ_{115}	$K^+\bar{K}^*(892)^0$ $\times B(\bar{K}^{*0} \rightarrow K^-\pi^+)$		$(2.8 \pm 0.4) \times 10^{-3}$	
Γ_{116}	$K^+K^-\pi^+$ nonresonant		$(4.5 \pm 0.9) \times 10^{-3}$	
Γ_{117}	$K^0\bar{K}^0\pi^+$		—	
Γ_{118}	$K^*(892)^+\bar{K}^0$ $\times B(K^{*+} \rightarrow K^0\pi^+)$		$(2.1 \pm 1.0) \%$	
Γ_{119}	$K^+K^-\pi^+\pi^0$		—	
Γ_{120}	$\phi\pi^+\pi^0 \times B(\phi \rightarrow K^+K^-)$		$(1.1 \pm 0.5) \%$	
Γ_{121}	$\phi\rho^+ \times B(\phi \rightarrow K^+K^-)$		$< 7 \times 10^{-3}$	CL=90%
Γ_{122}	$K^+K^-\pi^+\pi^0$ non- ϕ		$(1.5 \pm_{-0.6}^{+0.7}) \%$	
Γ_{123}	$K^+\bar{K}^0\pi^+\pi^-$		$< 2 \%$	CL=90%
Γ_{124}	$K^0K^-\pi^+\pi^+$		$(1.0 \pm 0.6) \%$	
Γ_{125}	$K^*(892)^+\bar{K}^*(892)^0$ $\times B^2(K^{*+} \rightarrow K^0\pi^+)$		$(1.2 \pm 0.5) \%$	
Γ_{126}	$K^0K^-\pi^+\pi^+$ non- $K^{*+}\bar{K}^{*0}$		$< 7.9 \times 10^{-3}$	CL=90%
Γ_{127}	$K^+K^-\pi^+\pi^+\pi^-$		—	
Γ_{128}	$\phi\pi^+\pi^+\pi^-$ $\times B(\phi \rightarrow K^+K^-)$		$< 1 \times 10^{-3}$	CL=90%
Γ_{129}	$K^+K^-\pi^+\pi^+\pi^-$ nonresonant		$< 3 \%$	CL=90%

Fractions of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

Γ_{130}	$\phi\pi^+$		$(6.1 \pm 0.6) \times 10^{-3}$	
Γ_{131}	$\phi\pi^+\pi^0$		$(2.3 \pm 1.0) \%$	
Γ_{132}	$\phi\rho^+$		$< 1.4 \%$	CL=90%
Γ_{133}	$\phi\pi^+\pi^+\pi^-$		$< 2 \times 10^{-3}$	CL=90%
Γ_{134}	$K^+\bar{K}^*(892)^0$		$(4.2 \pm 0.5) \times 10^{-3}$	
Γ_{135}	$K^*(892)^+\bar{K}^0$		$(3.2 \pm 1.5) \%$	
Γ_{136}	$K^*(892)^+\bar{K}^*(892)^0$		$(2.6 \pm 1.1) \%$	

Doubly Cabibbo suppressed (DC) modes, $\Delta C = 1$ weak neutral current (C1) modes, or Lepton Family number (LF) or Lepton number (L) violating modes

Γ_{137}	$K^+\pi^+\pi^-$	DC	$(6.8 \pm 1.5) \times 10^{-4}$	
Γ_{138}	$K^+\rho^0$	DC	$(2.5 \pm 1.2) \times 10^{-4}$	
Γ_{139}	$K^*(892)^0\pi^+$	DC	$(3.6 \pm 1.6) \times 10^{-4}$	
Γ_{140}	$K^+\pi^+\pi^-$ nonresonant	DC	$(2.4 \pm 1.2) \times 10^{-4}$	
Γ_{141}	$K^+K^+K^-$	DC	$< 1.4 \times 10^{-4}$	CL=90%
Γ_{142}	ϕK^+	DC	$< 1.3 \times 10^{-4}$	CL=90%
Γ_{143}	$\pi^+e^+e^-$	C1	$< 6.6 \times 10^{-5}$	CL=90%

Γ_{144}	$\pi^+ \mu^+ \mu^-$	<i>CI</i>	< 1.8	$\times 10^{-5}$	CL=90%
Γ_{145}	$\rho^+ \mu^+ \mu^-$	<i>CI</i>	< 5.6	$\times 10^{-4}$	CL=90%
Γ_{146}	$K^+ e^+ e^-$		[f] < 2.0	$\times 10^{-4}$	CL=90%
Γ_{147}	$K^+ \mu^+ \mu^-$		[f] < 9.7	$\times 10^{-5}$	CL=90%
Γ_{148}	$\pi^+ e^+ \mu^-$	<i>LF</i>	< 1.1	$\times 10^{-4}$	CL=90%
Γ_{149}	$\pi^+ e^- \mu^+$	<i>LF</i>	< 1.3	$\times 10^{-4}$	CL=90%
Γ_{150}	$K^+ e^+ \mu^-$	<i>LF</i>	< 1.3	$\times 10^{-4}$	CL=90%
Γ_{151}	$K^+ e^- \mu^+$	<i>LF</i>	< 1.2	$\times 10^{-4}$	CL=90%
Γ_{152}	$\pi^- e^+ e^+$	<i>L</i>	< 1.1	$\times 10^{-4}$	CL=90%
Γ_{153}	$\pi^- \mu^+ \mu^+$	<i>L</i>	< 8.7	$\times 10^{-5}$	CL=90%
Γ_{154}	$\pi^- e^+ \mu^+$	<i>L</i>	< 1.1	$\times 10^{-4}$	CL=90%
Γ_{155}	$\rho^- \mu^+ \mu^+$	<i>L</i>	< 5.6	$\times 10^{-4}$	CL=90%
Γ_{156}	$K^- e^+ e^+$	<i>L</i>	< 1.2	$\times 10^{-4}$	CL=90%
Γ_{157}	$K^- \mu^+ \mu^+$	<i>L</i>	< 1.2	$\times 10^{-4}$	CL=90%
Γ_{158}	$K^- e^+ \mu^+$	<i>L</i>	< 1.3	$\times 10^{-4}$	CL=90%
Γ_{159}	$K^*(892)^- \mu^+ \mu^+$	<i>L</i>	< 8.5	$\times 10^{-4}$	CL=90%

Γ_{160} A dummy mode used by the fit. $(33 \pm 5) \%$

- [a] This is a weighted average of D^\pm (44%) and D^0 (56%) branching fractions. See " D^+ and $D^0 \rightarrow (\eta \text{ anything}) / (\text{total } D^+ \text{ and } D^0)$ " under " D^+ Branching Ratios" in these Particle Listings.
- [b] This value averages the e^+ and μ^+ branching fractions, after making a small phase-space adjustment to the μ^+ fraction to be able to use it as an e^+ fraction; hence our ℓ^+ here is really an e^+ .
- [c] An ℓ indicates an e or a μ mode, not a sum over these modes.
- [d] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers.
- [e] The two experiments measuring this fraction are in serious disagreement. See the Particle Listings.
- [f] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.

CONSTRAINED FIT INFORMATION

An overall fit to 32 branching ratios uses 54 measurements and one constraint to determine 20 parameters. The overall fit has a $\chi^2 = 20.8$ for 35 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_{11}	5										
x_{16}	4	2									
x_{25}	18	29	8								
x_{26}	14	7	31	25							
x_{33}	38	9	8	31	25						
x_{34}	32	16	14	56	45	55					
x_{39}	0	0	0	0	0	0	0				
x_{43}	7	4	3	13	10	12	23	0			
x_{52}	9	5	4	17	14	16	30	0	18		
x_{59}	15	8	7	28	22	27	49	0	11	15	
x_{73}	21	11	9	37	29	36	65	0	15	20	
x_{80}	5	3	2	9	7	8	16	0	31	37	
x_{87}	3	1	1	5	4	5	9	0	29	13	
x_{93}	5	2	2	9	7	8	15	0	3	5	
x_{94}	3	2	1	6	5	6	11	0	2	3	
x_{98}	19	10	9	35	28	33	61	0	14	18	
x_{100}	11	5	5	19	15	18	34	0	8	10	
x_{112}	22	7	6	23	18	53	41	0	9	12	
x_{160}	-35	-26	-12	-41	-34	-38	-55	-58	-46	-45	
	x_9	x_{11}	x_{16}	x_{25}	x_{26}	x_{33}	x_{34}	x_{39}	x_{43}	x_{52}	
x_{73}	32										
x_{80}	8	10									
x_{87}	4	6	12								
x_{93}	29	10	2	1							
x_{94}	8	7	2	1	15						
x_{98}	30	40	10	5	9	7					
x_{100}	16	22	5	3	5	4	43				
x_{112}	20	26	6	4	6	4	25	14			
x_{160}	-30	-38	-46	-32	-16	-10	-35	-19	-27		
	x_{59}	x_{73}	x_{80}	x_{87}	x_{93}	x_{94}	x_{98}	x_{100}	x_{112}		

D^+ BRANCHING RATIOS

See the "Note on D Mesons" above. Some now-obsolete measurements have been omitted from these Listings.

———— Inclusive modes ————

$\Gamma(e^+ \text{ anything})/\Gamma_{\text{total}}$				Γ_1/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.172±0.019 OUR AVERAGE				
0.20 ^{+0.09} _{-0.07}		AGUILAR-...	87E HYBR	$\pi p, p p$ 360, 400 GeV
0.170±0.019±0.007	158	BALTRUSAIT..85B	MRK3	$e^+ e^-$ 3.77 GeV
0.168±0.064	23	SCHINDLER	81 MRK2	$e^+ e^-$ 3.771 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.220 ^{+0.044} _{-0.022}		BACINO	80 DLCO	$e^+ e^-$ 3.77 GeV

D^+ and $D^0 \rightarrow (e^+ \text{ anything}) / (\text{total } D^+ \text{ and } D^0)$

If measured at the $\psi(3770)$, this quantity is a weighted average of D^+ (44%) and D^0 (56%) branching fractions. Only experiments at $E_{\text{cm}} = 3.77$ GeV are included in the average here. We don't put this result in the Meson Summary Table.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.110±0.011 OUR AVERAGE Error includes scale factor of 1.1.				
0.117±0.011	295	BALTRUSAIT..85B	MRK3	$e^+ e^-$ 3.77 GeV
0.10 ±0.032		⁴ SCHINDLER	81 MRK2	$e^+ e^-$ 3.771 GeV
0.072±0.028		FELLER	78 MRK1	$e^+ e^-$ 3.772 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.096±0.004±0.011	2207	⁵ ALBRECHT	96C ARG	$e^+ e^- \approx 10$ GeV
0.134±0.015±0.010		⁶ ABE	93E VNS	$e^+ e^-$ 58 GeV
0.098±0.009 ^{+0.006} _{-0.005}	240	⁷ ALBRECHT	92F ARG	$e^+ e^- \approx 10$ GeV
0.096±0.007±0.015		⁸ ONG	88 MRK2	$e^+ e^-$ 29 GeV
0.116 ^{+0.011} _{-0.009}		⁸ PAL	86 DLCO	$e^+ e^-$ 29 GeV
0.091±0.009±0.013		⁸ AIHARA	85 TPC	$e^+ e^-$ 29 GeV
0.092±0.022±0.040		⁸ ALTHOFF	84J TASS	$e^+ e^-$ 34.6 GeV
0.091±0.013		⁸ KOOP	84 DLCO	See PAL 86
0.08 ±0.015		⁹ BACINO	79 DLCO	$e^+ e^-$ 3.772 GeV

⁴ Isolates D^+ and $D^0 \rightarrow e^+ X$ and weights for relative production (44%–56%).

⁵ ALBRECHT 96C uses e^- in the hemisphere opposite to $D^{*+} \rightarrow D^0 \pi^+$ events.

⁶ ABE 93E also measures forward-backward asymmetries and fragmentation functions for c and b quarks.

⁷ ALBRECHT 92F uses the excess of right-sign over wrong-sign leptons in a sample of events tagged by fully reconstructed $D^*(2010)^+ \rightarrow D^0 \pi^+$ decays.

⁸ Average BR for charm $\rightarrow e^+ X$. Unlike at $E_{\text{cm}} = 3.77$ GeV, the admixture of charmed mesons is unknown.


⁹ Not independent of BACINO 80 measurements of $\Gamma(e^+ \text{ anything})/\Gamma_{\text{total}}$ for the D^+ and D^0 separately.

$\Gamma(K^- \text{ anything})/\Gamma_{\text{total}}$

Γ_2/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.242±0.028 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
0.278 ^{+0.036} -0.031		¹⁰ BARLAG	92C ACCM	π^- Cu 230 GeV
0.271±0.023±0.024		COFFMAN	91 MRK3	e^+e^- 3.77 GeV
0.17 ±0.07		AGUILAR-...	87E HYBR	$\pi p, pp$ 360, 400 GeV
0.19 ±0.05	26	SCHINDLER	81 MRK2	e^+e^- 3.771 GeV
0.10 ±0.07	3	VUILLEMIN	78 MRK1	e^+e^- 3.772 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.16 ^{+0.08} -0.07		AGUILAR-...	86B HYBR	See AGUILAR-BENITEZ 87E

¹⁰ BARLAG 92C computes the branching fraction using topological normalization.



$\Gamma(K^- \text{ anything})/\Gamma_{\text{total}}$

$[\Gamma(\bar{K}^0 \text{ anything}) + \Gamma(K^0 \text{ anything})]/\Gamma_{\text{total}}$

Γ_3/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.59 ±0.07 OUR AVERAGE				
0.612±0.065±0.043		COFFMAN	91 MRK3	e^+e^- 3.77 GeV
0.52 ±0.18	15	SCHINDLER	81 MRK2	e^+e^- 3.771 GeV
0.39 ±0.29	3	VUILLEMIN	78 MRK1	e^+e^- 3.772 GeV

$\Gamma(K^+ \text{ anything})/\Gamma_{\text{total}}$

Γ_4/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.058 ± 0.014 OUR AVERAGE				
0.055 ± 0.013 ± 0.009		COFFMAN	91 MRK3	$e^+ e^-$ 3.77 GeV
0.08 $\begin{smallmatrix} +0.06 \\ -0.05 \end{smallmatrix}$		AGUILAR-...	87E HYBR	$\pi p, p p$ 360, 400 GeV
0.06 ± 0.04	12	SCHINDLER	81 MRK2	$e^+ e^-$ 3.771 GeV
0.06 ± 0.06	2	VUILLEMIN	78 MRK1	$e^+ e^-$ 3.772 GeV

$D^+ \text{ and } D^0 \rightarrow (\eta \text{ anything}) / (\text{total } D^+ \text{ and } D^0)$

If measured at the $\psi(3770)$, this quantity is a weighted average of D^+ (44%) and D^0 (56%) branching fractions. Only the experiment at $E_{\text{cm}} = 3.77$ GeV is used.

VALUE	DOCUMENT ID	TECN	COMMENT
< 0.13	PARTRIDGE	81 CBAL	$e^+ e^-$ 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
< 0.02	¹¹ BRANDELIK	79 DASP	$e^+ e^-$ 4.03 GeV
¹¹ The BRANDELIK 79 result is based on the absence of an η signal at $E_{\text{cm}} = 4.03$ GeV. PARTRIDGE 81 observes a substantially higher η cross section at 4.03 GeV.			

$\Gamma(c/\bar{c} \rightarrow \mu^+ \text{ anything})/\Gamma(c/\bar{c} \rightarrow \text{anything})$

This is the average branching ratio for charm $\rightarrow \mu^+ X$. The mixture of charmed particles is unknown and may actually contain states other than D mesons. We don't put this result in the Meson Summary Table.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.081 $\begin{smallmatrix} +0.010 \\ -0.009 \end{smallmatrix}$ OUR AVERAGE				
0.086 ± 0.017 $\begin{smallmatrix} +0.008 \\ -0.007 \end{smallmatrix}$	69	¹² ALBRECHT	92F ARG	$e^+ e^- \approx 10$ GeV
0.078 ± 0.009 ± 0.012		ONG	88 MRK2	$e^+ e^-$ 29 GeV
0.078 ± 0.015 ± 0.02		BARTEL	87 JADE	$e^+ e^-$ 34.6 GeV
0.082 ± 0.012 $\begin{smallmatrix} +0.02 \\ -0.01 \end{smallmatrix}$		ALTHOFF	84G TASS	$e^+ e^-$ 34.5 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.089 ± 0.018 ± 0.025		BARTEL	85J JADE	See BARTEL 87
¹² ALBRECHT 92F uses the excess of right-sign over wrong-sign leptons in a sample of events tagged by fully reconstructed $D^*(2010)^+ \rightarrow D^0 \pi^+$ decays.				

Leptonic and semileptonic modes

$\Gamma(\mu^+ \nu_\mu)/\Gamma_{\text{total}}$

Γ_7/Γ

See the "Note on Pseudoscalar-Meson Decay Constants" in the Listings for the π^\pm .

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.0008 $\begin{smallmatrix} +0.0016 + 0.0005 \\ -0.0005 - 0.0002 \end{smallmatrix}$		1	¹³ BAI	98B BES	$e^+ e^- \rightarrow D^{*+} D^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.00072	90		ADLER	88B MRK3	$e^+ e^-$ 3.77 GeV
< 0.02	90	0	¹⁴ AUBERT	83 SPEC	$\mu^+ \text{ Fe}$, 250 GeV
¹³ BAI 98B obtains $f_D = (300 \begin{smallmatrix} +180 + 80 \\ -150 - 40 \end{smallmatrix})$ MeV from this measurement.					
¹⁴ AUBERT 83 obtains an upper limit 0.014 assuming the final state contains equal amounts of (D^+, D^-) , (D^+, \bar{D}^0) , (D^-, D^0) , and (D^0, \bar{D}^0) . We quote the limit they get under more general assumptions.					

$\Gamma(\bar{K}^0 \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_8/Γ

We average our $\bar{K}^0 e^+ \nu_e$ and $\bar{K}^0 \mu^+ \nu_\mu$ branching fractions, after multiplying the latter by a phase-space factor of 1.03 to be able to use it with the $\bar{K}^0 e^+ \nu_e$ fraction. Hence our ℓ^+ here is really an e^+ .

VALUE	DOCUMENT ID	COMMENT
0.068 ± 0.008 OUR AVERAGE		
0.067 ± 0.009	PDG 98	Our $\Gamma(\bar{K}^0 e^+ \nu_e)/\Gamma_{\text{total}}$
0.072 ^{+0.031} _{-0.020}	PDG 98	1.03 × our $\Gamma(\bar{K}^0 \mu^+ \nu_\mu)/\Gamma_{\text{total}}$

$\Gamma(\bar{K}^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.067 ± 0.009 OUR FIT				
0.06 ^{+0.022} _{-0.013} ± 0.007	13	BAI 91	MRK3	$e^+ e^- \approx 3.77$ GeV

$\Gamma(\bar{K}^0 e^+ \nu_e)/\Gamma(\bar{K}^0 \pi^+)$ Γ_9/Γ_{33}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
2.32 ± 0.31 OUR FIT				
2.60 ± 0.35 ± 0.26	186	¹⁵ BEAN 93C	CLE2	$e^+ e^- \approx \Upsilon(4S)$

¹⁵ BEAN 93C uses $\bar{K}^0 \mu^+ \nu_\mu$ as well as $\bar{K}^0 e^+ \nu_e$ events and makes a small phase-space adjustment to the number of the μ^+ events to use them as e^+ events.

$\Gamma(\bar{K}^0 e^+ \nu_e)/\Gamma(K^- \pi^+ \pi^+)$ Γ_9/Γ_{34}

VALUE	DOCUMENT ID	TECN	COMMENT
0.74 ± 0.10 OUR FIT			
0.66 ± 0.09 ± 0.14	ANJOS 91C	E691	γ Be 80–240 GeV

$\Gamma(\bar{K}^0 \mu^+ \nu_\mu)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.07 ^{+0.028} _{-0.016} ± 0.012	14	BAI 91	MRK3	$e^+ e^- \approx 3.77$ GeV

$\Gamma(\bar{K}^0 \mu^+ \nu_\mu)/\Gamma(\mu^+ \text{ anything})$ Γ_{10}/Γ_6

VALUE	EVTS	DOCUMENT ID	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.76 ± 0.06	84	¹⁶ AOKI 88	π^- emulsion

¹⁶ From topological branching ratios in emulsion with an identified muon.

$\Gamma(K^- \pi^+ e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.041^{+0.009}_{-0.007} OUR FIT					
0.035 ^{+0.012} _{-0.007} ± 0.004	14	¹⁷ BAI 91	MRK3	$e^+ e^- \approx 3.77$ GeV	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.057	90	¹⁸ AGUILAR-...	87F HYBR	$\pi p, p p$ 360, 400 GeV	

¹⁷ BAI 91 finds that a fraction $0.79^{+0.15+0.09}_{-0.17-0.03}$ of combined D^+ and D^0 decays to $\bar{K} \pi e^+ \nu_e$ (24 events) are $\bar{K}^*(892) e^+ \nu_e$.

¹⁸ AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization.

$\Gamma(\bar{K}^*(892)^0 \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_{24} / Γ

We average our $\bar{K}^{*0} e^+ \nu_e$ and $\bar{K}^{*0} \mu^+ \nu_\mu$ branching fractions, after multiplying the latter by a phase-space factor of 1.05 to be able to use it with the $\bar{K}^{*0} e^+ \nu_e$ fraction. Hence our ℓ^+ here is really an e^+ .

VALUE	DOCUMENT ID	COMMENT
0.047 ± 0.004 OUR AVERAGE		
0.048 ± 0.005	PDG	98 Our $\Gamma(\bar{K}^{*0} e^+ \nu_e) / \Gamma_{\text{total}}$
0.046 ± 0.006	PDG	98 $1.05 \times$ our $\Gamma(\bar{K}^{*0} \mu^+ \nu_\mu) / \Gamma_{\text{total}}$

$\Gamma(\bar{K}^*(892)^0 e^+ \nu_e) / \Gamma(K^- \pi^+ e^+ \nu_e)$ $\Gamma_{25} / \Gamma_{11}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.16^{+0.21}_{-0.24} OUR FIT				
1.0 ± 0.3	35	ADAMOVICH	91	OMEG π^- 340 GeV

$\Gamma(\bar{K}^*(892)^0 e^+ \nu_e) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{25} / \Gamma_{34}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.53 ± 0.05 OUR FIT				
0.54 ± 0.05 OUR AVERAGE				
0.67 ± 0.09 ± 0.07	710	¹⁹ BEAN	93C	CLE2 $e^+ e^- \approx \Upsilon(4S)$
0.62 ± 0.15 ± 0.09	35	ADAMOVICH	91	OMEG π^- 340 GeV
0.55 ± 0.08 ± 0.10	880	ALBRECHT	91	ARG $e^+ e^- \approx 10.4$ GeV
0.49 ± 0.04 ± 0.05		ANJOS	89B	E691 Photoproduction

¹⁹ BEAN 93C uses $\bar{K}^{*0} \mu^+ \nu_\mu$ as well as $\bar{K}^{*0} e^+ \nu_e$ events and makes a small phase-space adjustment to the number of the μ^+ events to use them as e^+ events.

$\Gamma(K^- \pi^+ e^+ \nu_e \text{ nonresonant}) / \Gamma_{\text{total}}$ Γ_{13} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.007	90	²⁰ ANJOS	89B	E691 Photoproduction

²⁰ ANJOS 89B assumes a $\Gamma(D^+ \rightarrow K^- \pi^+ \pi^+) / \Gamma_{\text{total}} = 9.1 \pm 1.3 \pm 0.4\%$.

$\Gamma(K^- \pi^+ \mu^+ \nu_\mu) / \Gamma_{\text{total}}$ $\Gamma_{14} / \Gamma = (\Gamma_{16} + \frac{2}{3} \Gamma_{26}) / \Gamma$

VALUE	DOCUMENT ID
0.032 ± 0.004 OUR FIT	Error includes scale factor of 1.1.

$\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu) / \Gamma_{\text{total}}$ Γ_{26} / Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.044 ± 0.006 OUR FIT				Error includes scale factor of 1.1.
0.0325 ± 0.0071 ± 0.0075	224	²¹ KODAMA	92C	E653 π^- emulsion 600 GeV

²¹ KODAMA 92C measures $\Gamma(D^+ \rightarrow \bar{K}^{*0} \mu^+ \nu_\mu) / \Gamma(D^0 \rightarrow K^- \mu^+ \nu_\mu) = 0.43 \pm 0.09 \pm 0.09$ and then uses $\Gamma(D^0 \rightarrow K^- \mu^+ \nu_\mu) = (7.0 \pm 0.7) \times 10^{10} \text{ s}^{-1}$ to get the quoted branching fraction. See also the footnote to KODAMA 92C in the next data block.

$\Gamma(\overline{K}^*(892)^0 \mu^+ \nu_\mu) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{26} / \Gamma_{34}$

Unseen decay modes of the $\overline{K}^*(892)^0$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.49 ± 0.06 OUR FIT

0.53 ± 0.06 OUR AVERAGE

0.56 ± 0.04 ± 0.06	875	FRABETTI	93E E687	γ Be $\overline{E}_\gamma \approx 200$ GeV
--------------------	-----	----------	----------	---

0.46 ± 0.07 ± 0.08	224	²² KODAMA	92C E653	π^- emulsion 600 GeV
--------------------	-----	----------------------	----------	--------------------------

²² KODAMA 92C uses the same $\overline{K}^{*0} \mu^+ \nu_\mu$ events normalizing instead with $D^0 \rightarrow K^- \mu^+ \nu_\mu$ events, as reported in the preceding data block.

$\Gamma(K^- \pi^+ \mu^+ \nu_\mu \text{ nonresonant}) / \Gamma(K^- \pi^+ \mu^+ \nu_\mu)$ $\Gamma_{16} / \Gamma_{14} = \Gamma_{16} / (\Gamma_{16} + \frac{2}{3} \Gamma_{26})$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.083 ± 0.029 OUR FIT

0.083 ± 0.029

FRABETTI	93E E687	< 0.12 (90% CL)
----------	----------	-----------------

$\Gamma(\overline{K}^0 \pi^+ \pi^- e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{17} / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.022 $^{+0.047}_{-0.006} \pm 0.004$	1	²³ AGUILAR-...	87F HYBR	$\pi p, pp$ 360, 400 GeV
--------------------------------------	---	---------------------------	----------	--------------------------

²³ AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization.

$\Gamma(K^- \pi^+ \pi^0 e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{18} / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.044 $^{+0.052}_{-0.013} \pm 0.007$	2	²⁴ AGUILAR-...	87F HYBR	$\pi p, pp$ 360, 400 GeV
--------------------------------------	---	---------------------------	----------	--------------------------

²⁴ AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization.

$\Gamma((\overline{K}^*(892) \pi)^0 e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{19} / Γ

Unseen decay modes of the $\overline{K}^*(892)$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 0.012	90	ANJOS	92 E691	Photoproduction
---------	----	-------	---------	-----------------

$\Gamma((\overline{K} \pi \pi)^0 e^+ \nu_e \text{ non-}\overline{K}^*(892)) / \Gamma_{\text{total}}$ Γ_{20} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 0.009	90	ANJOS	92 E691	Photoproduction
---------	----	-------	---------	-----------------

$\Gamma(K^- \pi^+ \pi^0 \mu^+ \nu_\mu) / \Gamma(K^- \pi^+ \mu^+ \nu_\mu)$ $\Gamma_{21} / \Gamma_{14} = \Gamma_{21} / (\Gamma_{16} + \frac{2}{3} \Gamma_{26})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 0.042	90	FRABETTI	93E E687	γ Be $\overline{E}_\gamma \approx 200$ GeV
---------	----	----------	----------	---

$\Gamma(\pi^0 \ell^+ \nu_\ell) / \Gamma(\bar{K}^0 \ell^+ \nu_\ell)$ Γ_{22} / Γ_8

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

$0.046 \pm 0.014 \pm 0.017$	100	²⁵ BARTELT	97 CLE2	$e^+ e^- \approx \gamma(4S)$
-----------------------------	-----	-----------------------	---------	------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.085 \pm 0.027 \pm 0.014$	53	²⁶ ALAM	93 CLE2	See BARTELT 97
-----------------------------	----	--------------------	---------	----------------

²⁵ BARTELT 97 thus directly measures the product of ratios squared of CKM matrix elements and form factors at $q^2=0$: $|V_{cd}/V_{cs}|^2 \cdot |f_+^\pi(0)/f_+^K(0)|^2 = 0.046 \pm 0.014 \pm 0.017$.

²⁶ ALAM 93 thus directly measures the product of ratios squared of CKM matrix elements and form factors at $q^2=0$: $|V_{cd}/V_{cs}|^2 \cdot |f_+^\pi(0)/f_+^K(0)|^2 = 0.085 \pm 0.027 \pm 0.014$.

$\Gamma(\pi^+ \pi^- e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{23} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.057	90	²⁷ AGUILAR-...	87F HYBR	$\pi p, p p$ 360, 400 GeV
----------	----	---------------------------	----------	---------------------------

²⁷ AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization.

$\Gamma(\rho^0 e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{27} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0037	90	BAI	91 MRK3	$e^+ e^- \approx 3.77$ GeV
-----------	----	-----	---------	----------------------------

$\Gamma(\rho^0 e^+ \nu_e) / \Gamma(\bar{K}^*(892)^0 e^+ \nu_e)$ $\Gamma_{27} / \Gamma_{25}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

$0.045 \pm 0.014 \pm 0.009$	49	²⁸ AITALA	97 E791	π^- nucleus, 500 GeV
-----------------------------	----	----------------------	---------	--------------------------

²⁸ AITALA 97 explicitly subtracts $D^+ \rightarrow \eta' e^+ \nu_e$ and other backgrounds to get this result.

$\Gamma(\rho^0 \mu^+ \nu_\mu) / \Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)$ $\Gamma_{28} / \Gamma_{26}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.061 ± 0.014 OUR AVERAGE

$0.051 \pm 0.015 \pm 0.009$	54	²⁹ AITALA	97 E791	π^- nucleus, 500 GeV
-----------------------------	----	----------------------	---------	--------------------------

$0.079 \pm 0.019 \pm 0.013$	39	³⁰ FRABETTI	97 E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV
-----------------------------	----	------------------------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.044^{+0.031}_{-0.025} \pm 0.014$	4	³¹ KODAMA	93C E653	π^- emulsion 600 GeV
-------------------------------------	---	----------------------	----------	--------------------------

²⁹ AITALA 97 explicitly subtracts $D^+ \rightarrow \eta' \mu^+ \nu_\mu$ and other backgrounds to get this result.

³⁰ Because the reconstruction efficiency for photons is low, this FRABETTI 97 result also includes any $D^+ \rightarrow \eta' \mu^+ \nu_\mu \rightarrow \gamma \rho^0 \mu^+ \nu_\mu$ events in the numerator.

³¹ This KODAMA 93C result is based on a final signal of $4.0^{+2.8}_{-2.3} \pm 1.3$ events; the estimates of backgrounds that affect this number are somewhat model dependent.

$\Gamma(\phi e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{29} / Γ

Decay modes of the ϕ not included in the search are corrected for.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.0209	90	BAI	91 MRK3	$e^+ e^- \approx 3.77$ GeV
--------------------------------	----	-----	---------	----------------------------

$\Gamma(\phi\mu^+\nu_\mu)/\Gamma_{\text{total}}$ Γ_{30}/Γ

Decay modes of the ϕ not included in the search are corrected for.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0372	90	BAI	91 MRK3	$e^+e^- \approx 3.77$ GeV

$\Gamma(\eta\ell^+\nu_\ell)/\Gamma(\pi^0\ell^+\nu_\ell)$ Γ_{31}/Γ_{22}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	BARTELT	97 CLE2	$e^+e^- \approx \Upsilon(4S)$

$\Gamma(\eta'(958)\mu^+\nu_\mu)/\Gamma(\bar{K}^*(892)^0\mu^+\nu_\mu)$ Γ_{32}/Γ_{26}

Decay modes of the $\eta'(958)$ not included in the search are corrected for.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.20	90	KODAMA	93B E653	π^- emulsion 600 GeV

———— Hadronic modes with a \bar{K} or $\bar{K}K\bar{K}$ ————

$\Gamma(\bar{K}^0\pi^+)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.0289 ± 0.0026 OUR FIT Error includes scale factor of 1.1.

0.032 ± 0.004 OUR AVERAGE

0.032 ± 0.005 ± 0.002	161	ADLER	88C MRK3	e^+e^- 3.77 GeV
0.033 ± 0.009	36	³² SCHINDLER	81 MRK2	e^+e^- 3.771 GeV
0.033 ± 0.013	17	³³ PERUZZI	77 MRK1	e^+e^- 3.77 GeV

³²SCHINDLER 81 (MARK-2) measures $\sigma(e^+e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.14 ± 0.03 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

³³PERUZZI 77 (MARK-1) measures $\sigma(e^+e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.14 ± 0.05 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

$\Gamma(\bar{K}^0\pi^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{33}/Γ_{34}

It is generally assumed for modes such as $D^+ \rightarrow \bar{K}^0\pi^+$ that

$$\Gamma(D^+ \rightarrow \bar{K}^0\pi^+) = 2\Gamma(D^+ \rightarrow K_S^0\pi^+);$$

it is the latter Γ that is actually measured. BIGI 95 points out that interference between Cabibbo-allowed and doubly Cabibbo-suppressed amplitudes, where both occur, could invalidate this assumption by a few percent.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.321 ± 0.025 OUR FIT Error includes scale factor of 1.1.

0.32 ± 0.04 OUR AVERAGE Error includes scale factor of 1.4.

0.348 ± 0.024 ± 0.022	473	³⁴ BISHAI	97 CLE2	$e^+e^- \approx \Upsilon(4S)$
0.274 ± 0.030 ± 0.031	264	ANJOS	90C E691	Photoproduction

³⁴See BISHAI 97 for an isospin analysis of $D^+ \rightarrow \bar{K}\pi$ amplitudes.

$\Gamma(K^-\pi^+\pi^+)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.090 ± 0.006 OUR FIT

0.091 ± 0.007 OUR AVERAGE

0.093 ± 0.006 ± 0.008	1502	³⁵ BALEST	94 CLE2	$e^+e^- \approx \Upsilon(4S)$
0.091 ± 0.013 ± 0.004	1164	ADLER	88C MRK3	e^+e^- 3.77 GeV
0.091 ± 0.019	239	³⁶ SCHINDLER	81 MRK2	e^+e^- 3.771 GeV
0.086 ± 0.020	85	³⁷ PERUZZI	77 MRK1	e^+e^- 3.77 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.064^{+0.015}_{-0.014}$		³⁸ BARLAG	92C ACCM	π^- Cu 230 GeV
$0.063^{+0.028}_{-0.014} \pm 0.011$	8	³⁸ AGUILAR-...	87F HYBR	$\pi p, pp$ 360, 400 GeV

³⁵ BALEST 94 measures the ratio of $D^+ \rightarrow K^- \pi^+ \pi^+$ and $D^0 \rightarrow K^- \pi^+$ branching fractions to be $2.35 \pm 0.16 \pm 0.16$ and uses their absolute measurement of the $D^0 \rightarrow K^- \pi^+$ fraction (AKERIB 93).

³⁶ SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.38 ± 0.05 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

³⁷ PERUZZI 77 (MARK-1) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.36 ± 0.06 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

³⁸ AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.

$\Gamma(\bar{K}^*(892)^0 \pi^+) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{73} / \Gamma_{34}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.212 ± 0.016 OUR FIT				
0.210 ± 0.015 OUR AVERAGE				
$0.206 \pm 0.009 \pm 0.014$		FRABETTI	94G E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV
$0.255 \pm 0.014 \pm 0.050$		ANJOS	93 E691	γ Be 90–260 GeV
$0.21 \pm 0.06 \pm 0.06$		ALVAREZ	91B NA14	Photoproduction
$0.20 \pm 0.02 \pm 0.11$		ADLER	87 MRK3	$e^+ e^-$ 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.053	90	SCHINDLER	81 MRK2	$e^+ e^-$ 3.771 GeV

$\Gamma(\bar{K}_0^*(1430)^0 \pi^+) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{82} / \Gamma_{34}$

Unseen decay modes of the $\bar{K}_0^*(1430)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.41 ± 0.04 OUR AVERAGE			
$0.458 \pm 0.035 \pm 0.094$	FRABETTI	94G E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV
$0.400 \pm 0.031 \pm 0.027$	ANJOS	93 E691	γ Be 90–260 GeV

$\Gamma(\bar{K}^*(1680)^0 \pi^+) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{83} / \Gamma_{34}$

Unseen decay modes of the $\bar{K}^*(1680)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.160 ± 0.032 OUR AVERAGE Error includes scale factor of 1.1.			
$0.182 \pm 0.023 \pm 0.028$	FRABETTI	94G E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV
$0.113 \pm 0.015 \pm 0.050$	ANJOS	93 E691	γ Be 90–260 GeV

$\Gamma(K^- \pi^+ \pi^+ \text{ nonresonant}) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{38} / \Gamma_{34}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.95 ± 0.07 OUR AVERAGE			
$0.998 \pm 0.037 \pm 0.072$	FRABETTI	94G E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV
$0.838 \pm 0.088 \pm 0.275$	ANJOS	93 E691	γ Be 90–260 GeV
$0.79 \pm 0.07 \pm 0.15$	ADLER	87 MRK3	$e^+ e^-$ 3.77 GeV

$\Gamma(\bar{K}^0 \pi^+ \pi^0) / \Gamma_{\text{total}}$ Γ_{39} / Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.097 ± 0.030 OUR FIT Error includes scale factor of 1.1.

0.107 ± 0.029 OUR AVERAGE

0.102 ± 0.025 ± 0.016	159	ADLER	88C MRK3	$e^+ e^-$ 3.77 GeV
0.19 ± 0.12	10	³⁹ SCHINDLER	81 MRK2	$e^+ e^-$ 3.771 GeV

³⁹SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.78 ± 0.48 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

$\Gamma(\bar{K}^0 \rho^+) / \Gamma(\bar{K}^0 \pi^+ \pi^0)$ $\Gamma_{40} / \Gamma_{39}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.68 ± 0.08 ± 0.12 ADLER 87 MRK3 $e^+ e^-$ 3.77 GeV

$\Gamma(\bar{K}^*(892)^0 \pi^+) / \Gamma(\bar{K}^0 \pi^+ \pi^0)$ $\Gamma_{73} / \Gamma_{39}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.20 ± 0.06 OUR FIT

0.57 ± 0.18 ± 0.18 ADLER 87 MRK3 $e^+ e^-$ 3.77 GeV

$\Gamma(\bar{K}^0 \pi^+ \pi^0 \text{ nonresonant}) / \Gamma(\bar{K}^0 \pi^+ \pi^0)$ $\Gamma_{42} / \Gamma_{39}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.13 ± 0.07 ± 0.08 ADLER 87 MRK3 $e^+ e^-$ 3.77 GeV

$\Gamma(K^- \pi^+ \pi^+ \pi^0) / \Gamma_{\text{total}}$ Γ_{43} / Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.064 ± 0.011 OUR FIT

0.058 ± 0.012 ± 0.012 142 COFFMAN 92B MRK3 $e^+ e^-$ 3.77 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.034 $^{+0.056}_{-0.070}$ 40 BARLAG 92C ACCM π^- Cu 230 GeV

0.022 $^{+0.047}_{-0.006} \pm 0.004$ 1 40 AGUILAR-... 87F HYBR $\pi p, p p$ 360, 400 GeV

0.063 $^{+0.014}_{-0.013} \pm 0.012$ 175 BALTRUSAIT..86E MRK3 See COFFMAN 92B

⁴⁰AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.

$\Gamma(K^- \pi^+ \pi^+ \pi^0) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{43} / \Gamma_{34}$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.71 ± 0.12 OUR FIT

0.76 ± 0.11 ± 0.12 91 ANJOS 92C E691 γ Be 90–260 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.69 ± 0.10 ± 0.16 ANJOS 89E E691 See ANJOS 92C

0.57 $^{+0.65}_{-0.17}$ 1 AGUILAR-... 83B HYBR $\pi^- p$, 360 GeV

$\Gamma(\bar{K}^*(892)^0 \rho^+ \text{ total}) / \Gamma(K^- \pi^+ \pi^+ \pi^0)$ $\Gamma_{74} / \Gamma_{43}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.33 ± 0.165 ± 0.12 ⁴¹ANJOS 92C E691 γ Be 90–260 GeV

⁴¹See, however, the next entry, where the two experiments disagree completely.

$\Gamma(\bar{K}^*(892)^0 \rho^+ S\text{-wave})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{75}/Γ_{43}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included. The two experiments here disagree completely.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.26 ± 0.25 OUR AVERAGE	Error includes scale factor of 3.1.		
0.15 ± 0.075 ± 0.045	ANJOS	92C E691	γ Be 90–260 GeV
0.833 ± 0.116 ± 0.165	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

$\Gamma(\bar{K}^*(892)^0 \rho^+ P\text{-wave})/\Gamma_{\text{total}}$ Γ_{76}/Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.001	90	ANJOS	92C E691	γ Be 90–260 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.005	90	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

$\Gamma(\bar{K}^*(892)^0 \rho^+ D\text{-wave})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{77}/Γ_{43}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.15 ± 0.09 ± 0.045	ANJOS	92C E691	γ Be 90–260 GeV

$\Gamma(\bar{K}^*(892)^0 \rho^+ D\text{-wave longitudinal})/\Gamma_{\text{total}}$ Γ_{78}/Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.007	90	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

$\Gamma(\bar{K}_1(1400)^0 \pi^+)/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{80}/Γ_{43}

Unseen decay modes of the $\bar{K}_1(1400)^0$ are included.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.77 ± 0.20 OUR FIT			
0.907 ± 0.218 ± 0.180	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

$\Gamma(K^- \rho^+ \pi^+ \text{total})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{88}/Γ_{43}

This includes $\bar{K}^*(892)^0 \rho^+$, etc. The next entry gives the specifically 3-body fraction.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.48 ± 0.13 ± 0.09	ANJOS	92C E691	γ Be 90–260 GeV

$\Gamma(K^- \rho^+ \pi^+ 3\text{-body})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{89}/Γ_{43}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.17 ± 0.06 OUR AVERAGE			
0.18 ± 0.08 ± 0.04	ANJOS	92C E691	γ Be 90–260 GeV
0.159 ± 0.065 ± 0.060	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

$\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^0 \text{total})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{84}/Γ_{43}

This includes $\bar{K}^*(892)^0 \rho^+$, etc. The next two entries give the specifically 3-body fraction. Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.05 ± 0.11 ± 0.08	ANJOS	92C E691	γ Be 90–260 GeV

$\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^0 \text{ 3-body})/\Gamma_{\text{total}}$ Γ_{85}/Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.008	90	⁴² COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV
--------	----	-----------------------	----------	--------------------

⁴² See, however, the next entry: ANJOS 92C sees a large signal in this channel.

$\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^0 \text{ 3-body})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{85}/Γ_{43}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.66 ± 0.09 ± 0.17	ANJOS	92C E691	γ Be 90–260 GeV
---------------------------	-------	----------	------------------------

$\Gamma(K^*(892)^- \pi^+ \pi^+ \text{ 3-body})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{87}/Γ_{43}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.32 ± 0.14 OUR FIT Error includes scale factor of 1.1.

0.24 ± 0.12 ± 0.09	ANJOS	92C E691	γ Be 90–260 GeV
---------------------------	-------	----------	------------------------

$\Gamma(K^- \pi^+ \pi^+ \pi^0 \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.002	90	⁴³ ANJOS	92C E691	γ Be 90–260 GeV
--------	----	---------------------	----------	------------------------

⁴³ Whereas ANJOS 92C finds no signal here, COFFMAN 92B finds a fairly large one; see the next entry.

$\Gamma(K^- \pi^+ \pi^+ \pi^0 \text{ nonresonant})/\Gamma(K^- \pi^+ \pi^+ \pi^0)$ Γ_{51}/Γ_{43}

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.184 ± 0.070 ± 0.050	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV
------------------------------	---------	----------	--------------------

$\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.070 ± 0.009 OUR FIT

0.071 ± 0.016 OUR AVERAGE

0.066 ± 0.015 ± 0.005	168	ADLER	88C MRK3	$e^+ e^-$ 3.77 GeV
-----------------------	-----	-------	----------	--------------------

0.12 ± 0.05	21	⁴⁴ SCHINDLER	81 MRK2	$e^+ e^-$ 3.771 GeV
-------------	----	-------------------------	---------	---------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.042 ^{+0.019} _{-0.017}		⁴⁵ BARLAG	92C ACCM	π^- Cu 230 GeV
---	--	----------------------	----------	--------------------

0.243 ^{+0.064} _{-0.041} ± 0.041	11	⁴⁵ AGUILAR-...	87F HYBR	$\pi p, p p$ 360, 400 GeV
---	----	---------------------------	----------	---------------------------

⁴⁴ SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.51 ± 0.08 nb. We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

⁴⁵ AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.

$\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{52}/Γ_{34}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.78 ± 0.10 OUR FIT

0.77 ± 0.07 ± 0.11	229	ANJOS	92C E691	γ Be 90–260 GeV
---------------------------	-----	-------	----------	------------------------

$\Gamma(\bar{K}^0 a_1(1260)^+)/\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)$ Γ_{71}/Γ_{52}

Unseen decay modes of the $a_1(1260)^+$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
1.15 ± 0.19 OUR AVERAGE	Error includes scale factor of 1.1.		
1.66 ± 0.28 ± 0.40	ANJOS	92C E691	γ Be 90–260 GeV
1.078 ± 0.114 ± 0.140	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

$\Gamma(\bar{K}^0 a_2(1320)^+)/\Gamma_{\text{total}}$ Γ_{72}/Γ

Unseen decay modes of the $a_2(1320)^+$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.003	90	ANJOS	92C E691	γ Be 90–260 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.008	90	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

$\Gamma(\bar{K}_1(1270)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{79}/Γ

Unseen decay modes of the $\bar{K}_1(1270)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.007	90	ANJOS	92C E691	γ Be 90–260 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.011	90	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

$\Gamma(\bar{K}_1(1400)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{80}/Γ

Unseen decay modes of the $\bar{K}_1(1400)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.009	90	⁴⁶ ANJOS	92C E691	γ Be 90–260 GeV
⁴⁶ ANJOS 92C sees no evidence for $\bar{K}_1(1400)^0 \pi^+$ in either the $\bar{K}^0 \pi^+ \pi^+ \pi^-$ or $K^- \pi^+ \pi^+ \pi^0$ channels, whereas COFFMAN 92B finds the $\bar{K}_1(1400)^0 \pi^+$ branching fraction to be large; see the next entry.				

$\Gamma(\bar{K}_1(1400)^0 \pi^+)/\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)$ Γ_{80}/Γ_{52}

Unseen decay modes of the $\bar{K}_1(1400)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.70 ± 0.17 OUR FIT			
0.623 ± 0.106 ± 0.180	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

$\Gamma(\bar{K}^*(1410)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{81}/Γ

Unseen decay modes of the $\bar{K}^*(1410)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.007	90	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

$\Gamma(K^*(892)^- \pi^+ \pi^+ \text{total})/\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)$ Γ_{86}/Γ_{52}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.41 ± 0.14	14	ALEEV	94 BIS2	nN 20–70 GeV

$\Gamma(K^*(892)^- \pi^+ \pi^+ \text{3-body})/\Gamma_{\text{total}}$ Γ_{87}/Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.020±0.009 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.013 90 COFFMAN 92B MRK3 $e^+ e^-$ 3.77 GeV

$\Gamma(K^*(892)^- \pi^+ \pi^+ \text{3-body})/\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)$ Γ_{87}/Γ_{52}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.29±0.13 OUR FIT	Error includes scale factor of 1.1.		
0.50±0.09±0.21	ANJOS	92C E691	γ Be 90–260 GeV

$\Gamma(\bar{K}^0 \rho^0 \pi^+ \text{total})/\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)$ Γ_{90}/Γ_{52}

This includes $\bar{K}^0 a_1(1260)^+$. The next two entries give the specifically 3-body reaction.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.60±0.10±0.17	90	ANJOS	92C E691	γ Be 90–260 GeV

$\Gamma(\bar{K}^0 \rho^0 \pi^+ \text{3-body})/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.004	90	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.004 90 COFFMAN 92B MRK3 $e^+ e^-$ 3.77 GeV

$\Gamma(\bar{K}^0 \rho^0 \pi^+ \text{3-body})/\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)$ Γ_{91}/Γ_{52}

VALUE	DOCUMENT ID	TECN	COMMENT
0.07±0.04±0.06	ANJOS	92C E691	γ Be 90–260 GeV

$\Gamma(\bar{K}^0 f_0(980) \pi^+)/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.005	90	ANJOS	92C E691	γ Be 90–260 GeV

$\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^- \text{nonresonant})/\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-)$ Γ_{58}/Γ_{52}

VALUE	DOCUMENT ID	TECN	COMMENT
0.12±0.06 OUR AVERAGE			
0.10±0.04 ±0.06	ANJOS	92C E691	γ Be 90–260 GeV
0.17±0.056±0.100	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

$\Gamma(K^- \pi^+ \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0037 ^{+0.0012} _{-0.0010}	⁴⁷ BARLAG	92C ACCM	π^- Cu 230 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

⁴⁷ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K^- \pi^+ \pi^+ \pi^+ \pi^-)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{59}/Γ_{34}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.080±0.009 OUR FIT				
0.083±0.009 OUR AVERAGE				
0.077±0.008±0.010	239	FRABETTI	97C E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV
0.09 ±0.01 ±0.01	113	ANJOS	90D E691	Photoproduction

$$\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^-) / \Gamma(K^- \pi^+ \pi^+ \pi^-) \quad \Gamma_{93} / \Gamma_{59}$$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
1.1 ± 0.4 OUR FIT Error includes scale factor of 1.8.			
1.25 ± 0.12 ± 0.23	ANJOS	90D E691	Photoproduction

$$\Gamma(\bar{K}^*(892)^0 \rho^0 \pi^+) / \Gamma(K^- \pi^+ \pi^+) \quad \Gamma_{94} / \Gamma_{34}$$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.032^{+0.019}_{-0.017} OUR FIT Error includes scale factor of 1.8.			
0.023 ± 0.010 ± 0.006	FRABETTI	97C E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV

$$\Gamma(\bar{K}^*(892)^0 \rho^0 \pi^+) / \Gamma(\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^-) \quad \Gamma_{94} / \Gamma_{93}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.36^{+0.24}_{-0.20} OUR FIT Error includes scale factor of 1.8.			
0.75 ± 0.17 ± 0.19	ANJOS	90D E691	Photoproduction

$$\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^- \text{ no-}\rho) / \Gamma(K^- \pi^+ \pi^+) \quad \Gamma_{95} / \Gamma_{34}$$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.048 ± 0.015 ± 0.011	FRABETTI	97C E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV

$$\Gamma(K^- \rho^0 \pi^+ \pi^+) / \Gamma(K^- \pi^+ \pi^+) \quad \Gamma_{63} / \Gamma_{34}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.034 ± 0.009 ± 0.005	FRABETTI	97C E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV

$$\Gamma(K^- \pi^+ \pi^+ \pi^+ \pi^- \text{ nonresonant}) / \Gamma(K^- \pi^+ \pi^+) \quad \Gamma_{64} / \Gamma_{34}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.026	90	FRABETTI	97C E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV

$$\Gamma(K^- \pi^+ \pi^+ \pi^0 \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{65} / \Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.022^{+0.047}_{-0.008} ± 0.004	1	⁴⁸ AGUILAR-...	87F HYBR	$\pi p, pp$ 360, 400 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.015	⁴⁸ BARLAG	92C ACCM	π^- Cu	230 GeV
---------	----------------------	----------	------------	---------

⁴⁸ AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.

$$\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{66} / \Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.054^{+0.030}_{-0.014} OUR AVERAGE				
0.099 ^{+0.036} _{-0.070}		⁴⁹ BARLAG	92C ACCM	π^- Cu
0.044 ^{+0.052} _{-0.013} ± 0.007	2	⁴⁹ AGUILAR-...	87F HYBR	$\pi p, pp$ 360, 400 GeV

⁴⁹ AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.

$$\Gamma(\overline{K}^0 \pi^+ \pi^+ \pi^- \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{67} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0008 ± 0.0007	50 BARLAG	92C ACCM	π^- Cu 230 GeV

⁵⁰ BARLAG 92C computes the branching fraction using topological normalization.

$$\Gamma(K^- \pi^+ \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{68} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0020 ± 0.0018	51 BARLAG	92C ACCM	π^- Cu 230 GeV

⁵¹ BARLAG 92C computes the branching fraction using topological normalization.

$$\Gamma(\overline{K}^0 \overline{K}^0 K^+) / \Gamma(K^- \pi^+ \pi^+) \quad \Gamma_{69} / \Gamma_{34}$$

VALUE	EVS	DOCUMENT ID	TECN	COMMENT
0.20 ± 0.09 OUR AVERAGE				Error includes scale factor of 2.4.
0.14 ± 0.04 ± 0.02	39	ALBRECHT	94i ARG	$e^+ e^- \approx 10$ GeV
0.34 ± 0.07	70	AMMAR	91 CLEO	$e^+ e^- \approx 10.5$ GeV

Pionic modes

$$\Gamma(\pi^+ \pi^0) / \Gamma(K^- \pi^+ \pi^+) \quad \Gamma_{97} / \Gamma_{34}$$

VALUE	EVS	DOCUMENT ID	TECN	COMMENT
0.028 ± 0.006 ± 0.005	34	SELEN	93 CLE2	$e^+ e^- \approx \Upsilon(4S)$

$$\Gamma(\pi^+ \pi^+ \pi^-) / \Gamma(K^- \pi^+ \pi^+) \quad \Gamma_{98} / \Gamma_{34}$$

VALUE	EVS	DOCUMENT ID	TECN	COMMENT
0.0406 ± 0.0034 OUR FIT				
0.0403 ± 0.0035 OUR AVERAGE				
0.043 ± 0.003 ± 0.003	236	FRABETTI	97D E687	γ Be ≈ 200 GeV
0.032 ± 0.011 ± 0.003	20	ADAMOVICH	93 WA82	π^- 340 GeV
0.035 ± 0.007 ± 0.003		ANJOS	89 E691	Photoproduction
0.042 ± 0.016 ± 0.010	57	BALTRUSAIT..85E	MRK3	$e^+ e^-$ 3.77 GeV

$$\Gamma(\rho^0 \pi^+) / \Gamma(\pi^+ \pi^+ \pi^-) \quad \Gamma_{99} / \Gamma_{98}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.289 ± 0.055 ± 0.058	52 FRABETTI	97D E687	γ Be ≈ 200 GeV

⁵² FRABETTI 97D also includes $f_2(1270)\pi^+$ and $f_0(980)\pi^+$ modes in the fit, but the resulting decay fractions are not statistically significant.

$$\Gamma(\rho^0 \pi^+) / \Gamma(K^- \pi^+ \pi^+) \quad \Gamma_{99} / \Gamma_{34}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.015	90	ANJOS	89 E691	Photoproduction

$$\Gamma(\pi^+ \pi^+ \pi^- \text{ nonresonant}) / \Gamma(\pi^+ \pi^+ \pi^-) \quad \Gamma_{100} / \Gamma_{98}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.62 ± 0.11 OUR FIT			
0.589 ± 0.105 ± 0.081	53 FRABETTI	97D E687	γ Be ≈ 200 GeV

⁵³ FRABETTI 97D also includes $f_2(1270)\pi^+$ and $f_0(980)\pi^+$ modes in the fit, but the resulting decay fractions are not statistically significant.

$\Gamma(\pi^+\pi^+\pi^-\text{ nonresonant})/\Gamma(K^-\pi^+\pi^+)$ Γ_{100}/Γ_{34}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.025±0.005 OUR FIT			
0.027±0.007±0.002	ANJOS	89 E691	Photoproduction

$\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{101}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.019^{+0.015}_{-0.012}	54 BARLAG	92C ACCM	π^- Cu 230 GeV

⁵⁴ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma(K^-\pi^+\pi^+)$ Γ_{101}/Γ_{34}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.4	90	ANJOS	89E E691	Photoproduction

$\Gamma(\eta\pi^+)/\Gamma(\phi\pi^+)$ $\Gamma_{106}/\Gamma_{130}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.49±0.08	275	JESSOP	98 CLE2	$e^+e^- \approx \gamma(4S)$

$\Gamma(\eta\pi^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{106}/Γ_{34}

Unseen decay modes of the η are included.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.083±0.023±0.014		99	DAOUDI	92 CLE2	See JESSOP 98
<0.12	90		ANJOS	89E E691	Photoproduction

$\Gamma(\omega\pi^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{108}/Γ_{34}

Unseen decay modes of the ω are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.08	90	ANJOS	89E E691	Photoproduction

$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}}$ Γ_{104}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.0010 ^{+0.0008} _{-0.0007}	55 BARLAG	92C ACCM	π^- Cu 230 GeV

⁵⁵ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-)/\Gamma(K^-\pi^+\pi^+)$ Γ_{104}/Γ_{34}

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.023±0.004±0.002		58	FRABETTI	97C E687	$\gamma\text{Be}, \bar{E}_\gamma \approx 200$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.019	90	ANJOS	89 E691	Photoproduction
--------	----	-------	---------	-----------------

$\Gamma(\eta\rho^+)/\Gamma(\phi\pi^+)$ $\Gamma_{109}/\Gamma_{130}$

Unseen decay modes of the η are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.11	90	JESSOP	98 CLE2	$e^+e^- \approx \gamma(4S)$

$\Gamma(\eta\rho^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{109}/Γ_{34}

Unseen decay modes of the η are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.13	90	DAOUDI	92 CLE2	See JESSOP 98
-------	----	--------	---------	---------------

$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{105}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$0.0029^{+0.0029}_{-0.0020}$	⁵⁶ BARLAG	92C ACCM	π^- Cu 230 GeV
------------------------------	----------------------	----------	--------------------

⁵⁶ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\eta'(958)\pi^+)/\Gamma(\phi\pi^+)$ $\Gamma_{110}/\Gamma_{130}$

Unseen decay modes of the $\eta'(958)$ are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.82 ± 0.14	126	JESSOP	98 CLE2	$e^+e^- \approx \gamma(4S)$
-----------------	-----	--------	---------	-----------------------------

$\Gamma(\eta'(958)\pi^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{110}/Γ_{34}

Unseen decay modes of the $\eta'(958)$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.1	90	DAOUDI	92 CLE2	See JESSOP 98
<0.1	90	ALVAREZ	91 NA14	Photoproduction
<0.13	90	ANJOS	91B E691	$\gamma\text{Be}, \bar{E}_\gamma \approx 145$ GeV

$\Gamma(\eta'(958)\rho^+)/\Gamma(\phi\pi^+)$ $\Gamma_{111}/\Gamma_{130}$

Unseen decay modes of the $\eta'(958)$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.86	90	JESSOP	98 CLE2	$e^+e^- \approx \gamma(4S)$
-------	----	--------	---------	-----------------------------

$\Gamma(\eta'(958)\rho^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{111}/Γ_{34}

Unseen decay modes of the $\eta'(958)$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.17	90	DAOUDI	92 CLE2	See JESSOP 98
-------	----	--------	---------	---------------

————— Hadronic modes with a $K\bar{K}$ pair —————

$\Gamma(K^+\bar{K}^0)/\Gamma(\bar{K}^0\pi^+)$ Γ_{112}/Γ_{33}

It is generally assumed for modes such as $D^+ \rightarrow \bar{K}^0\pi^+$ that

$$\Gamma(D^+ \rightarrow \bar{K}^0\pi^+) = 2\Gamma(D^+ \rightarrow K_S^0\pi^+);$$

it is the latter Γ that is actually measured. BIG1 95 points out that interference between Cabibbo-allowed and doubly Cabibbo-suppressed amplitudes, where both occur, could invalidate this assumption by a few percent.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.255±0.029 OUR FIT				
0.263±0.035 OUR AVERAGE				
0.25 ±0.04 ±0.02	129	FRABETTI	95 E687	γ Be $\bar{E}_\gamma \approx 200$ GeV
0.271±0.065±0.039	69	ANJOS	90C E691	γ Be
0.317±0.086±0.048	31	BALTRUSAIT..85E	MRK3	e^+e^- 3.77 GeV
0.25 ±0.15	6	SCHINDLER	81 MRK2	e^+e^- 3.771 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.222±0.041±0.029 70 ⁵⁷ BISHAI 97 CLE2 $e^+e^- \approx \Upsilon(4S)$

⁵⁷ This BISHAI 97 result is redundant with results elsewhere in the Listings.

$\Gamma(K^+\bar{K}^0)/\Gamma(K^-\pi^+\pi^+)$ Γ_{112}/Γ_{34}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.082±0.010 OUR FIT				
0.077±0.014±0.007	70	⁵⁸ BISHAI	97 CLE2	$e^+e^- \approx \Upsilon(4S)$

⁵⁸ See BISHAI 97 for an isospin analysis of $D^+ \rightarrow K\bar{K}$ amplitudes.

$\Gamma(K^+K^-\pi^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{113}/Γ_{34}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0976±0.0042±0.0046	FRABETTI	95B E687	Dalitz plot analysis

$\Gamma(\phi\pi^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{130}/Γ_{34}

Unseen decay modes of the ϕ are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.068±0.005 OUR AVERAGE				
0.058±0.006±0.006		FRABETTI	95B E687	Dalitz plot analysis
0.062±0.017±0.006	19	ADAMOVICH	93 WA82	π^- 340 GeV
0.077±0.011±0.005	128	DAOUDI	92 CLE2	$e^+e^- \approx 10.5$ GeV
0.098±0.032±0.014	12	ALVAREZ	90C NA14	Photoproduction
0.071±0.008±0.007	84	ANJOS	88 E691	Photoproduction
0.084±0.021±0.011	21	BALTRUSAIT..85E	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^+\bar{K}^*(892)^0)/\Gamma(K^-\pi^+\pi^+)$ Γ_{134}/Γ_{34}

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.047±0.005 OUR AVERAGE Error includes scale factor of 1.2.				
0.044±0.003±0.004		⁵⁹ FRABETTI	95B E687	Dalitz plot analysis
0.058±0.009±0.006	73	ANJOS	88 E691	Photoproduction
0.048±0.021±0.011	14	BALTRUSAIT..85E	MRK3	e^+e^- 3.77 GeV

⁵⁹ See FRABETTI 95B for evidence also of $\bar{K}_0^*(1430)^0 K^+$ in the $D^+ \rightarrow K^+K^-\pi^+$ Dalitz plot.

$\Gamma(K^+K^-\pi^+ \text{ nonresonant})/\Gamma(K^-\pi^+\pi^+)$ Γ_{116}/Γ_{34}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.050±0.009 OUR AVERAGE				
0.049±0.008±0.006	95	ANJOS	88 E691	Photoproduction
0.059±0.026±0.009	37	BALTRUSAIT..85E	MRK3	e^+e^- 3.77 GeV

$\Gamma(K^*(892)^+\bar{K}^0)/\Gamma(\bar{K}^0\pi^+)$ Γ_{135}/Γ_{33}

Unseen decay modes of the $K^*(892)^+$ are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
1.1±0.3±0.4	67	FRABETTI	95 E687	γ Be $\bar{E}_\gamma \approx 200$ GeV

$\Gamma(\phi\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{131}/Γ

Unseen decay modes of the ϕ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.023±0.010	⁶⁰ BARLAG	92C ACCM	π^- Cu 230 GeV

⁶⁰ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\phi\pi^+\pi^0)/\Gamma(K^-\pi^+\pi^+)$ Γ_{131}/Γ_{34}

Unseen decay modes of the ϕ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.58	90	ALVAREZ	90C NA14	Photoproduction
<0.28	90	ANJOS	89E E691	Photoproduction

$\Gamma(\phi\rho^+)/\Gamma(K^-\pi^+\pi^+)$ Γ_{132}/Γ_{34}

Unseen decay modes of the ϕ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.16	90	DAOUDI	92 CLE2	$e^+e^- \approx 10.5$ GeV

$\Gamma(K^+K^-\pi^+\pi^0 \text{ non-}\phi)/\Gamma_{\text{total}}$ Γ_{122}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.015^{+0.007}_{-0.006}	⁶¹ BARLAG	92C ACCM	π^- Cu 230 GeV

⁶¹ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K^+K^-\pi^+\pi^0 \text{ non-}\phi)/\Gamma(K^-\pi^+\pi^+)$ Γ_{122}/Γ_{34}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.25	90	ANJOS	89E E691	Photoproduction

$\Gamma(K^+\bar{K}^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{123}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.02	90	ALBRECHT	92B ARG	$e^+e^- \simeq 10.4$ GeV

$\Gamma(K^0K^-\pi^+\pi^+)/\Gamma_{\text{total}}$ Γ_{124}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.01 ±0.005±0.003	ALBRECHT	92B ARG	$e^+e^- \simeq 10.4$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.003	⁶² BARLAG	92C ACCM	π^- Cu 230 GeV
--------	----------------------	----------	--------------------

⁶² BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K^*(892)^+\bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{136}/Γ

Unseen decay modes of the $K^*(892)$'s are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.026±0.008±0.007	ALBRECHT	92B ARG	$e^+e^- \simeq 10.4$ GeV

$\Gamma(K^0 K^- \pi^+ \pi^+ \text{non-}K^{*+} \bar{K}^{*0})/\Gamma_{\text{total}}$ Γ_{126}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0079	90	ALBRECHT	92B ARG	$e^+ e^- \simeq 10.4$ GeV

$\Gamma(\phi \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{133}/Γ

Unseen decay modes of the ϕ are included.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.002	90	0	ANJOS	88 E691	Photoproduction

$\Gamma(\phi \pi^+ \pi^+ \pi^-)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{133}/Γ_{34}

Unseen decay modes of the ϕ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.031	90	ALVAREZ	90C NA14	Photoproduction
--------	----	---------	----------	-----------------

$\Gamma(\phi \pi^+ \pi^+ \pi^-)/\Gamma(\phi \pi^+)$ $\Gamma_{133}/\Gamma_{130}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.6	90	FRABETTI	92 E687	γ Be
------	----	----------	---------	-------------

$\Gamma(K^+ K^- \pi^+ \pi^+ \pi^- \text{nonresonant})/\Gamma_{\text{total}}$ Γ_{129}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.03	90	12	ANJOS	88 E691	Photoproduction

————— Rare or forbidden modes —————

$\Gamma(K^+ \pi^+ \pi^-)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{137}/Γ_{34}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.0075 ± 0.0016 OUR AVERAGE

0.0077 ± 0.0017 ± 0.0008	59	AITALA	97C E791	π^- nucleus, 500 GeV
0.0072 ± 0.0023 ± 0.0017	21	FRABETTI	95E E687	γ Be, $\bar{E}_\gamma = 220$ GeV

$\Gamma(K^+ \rho^0)/\Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{138}/\Gamma_{137}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.37 ± 0.14 ± 0.07	AITALA	97C E791	π^- nucleus, 500 GeV
---------------------------	--------	----------	--------------------------

$\Gamma(K^+ \rho^0)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{138}/Γ_{34}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0067	90	FRABETTI	95E E687	γ Be, $\bar{E}_\gamma = 220$ GeV
---------	----	----------	----------	---

$\Gamma(K^*(892)^0 \pi^+)/\Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{139}/\Gamma_{137}$

Unseen decay modes of the $K^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.53 ± 0.21 ± 0.02	AITALA	97C E791	π^- nucleus, 500 GeV
---------------------------	--------	----------	--------------------------

$\Gamma(K^*(892)^0 \pi^+)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{139}/Γ_{34}

Unseen decay modes of the $K^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0021	90	FRABETTI	95E E687	γ Be, $\bar{E}_\gamma = 220$ GeV
---------	----	----------	----------	---

$\Gamma(K^+ \pi^+ \pi^- \text{ nonresonant})/\Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{140}/\Gamma_{137}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.36 ± 0.14 ± 0.07	AITALA	97C E791	π^- nucleus, 500 GeV
---------------------------	--------	----------	--------------------------

$\Gamma(K^+ K^+ K^-)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{141}/Γ_{34}

A doubly Cabibbo-suppressed decay with no simple spectator process possible.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

<0.0016	90	63	FRABETTI	95F E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV
-------------------	----	----	----------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.057 ± 0.020 ± 0.007	13	ADAMOVICH	93 WA82	π^- 340 GeV
-----------------------	----	-----------	---------	-----------------

⁶³ Using the $\phi \pi^+$ mode to normalize, FRABETTI 95F gets $\Gamma(K^+ K^+ K^-)/\Gamma(\phi \pi^+) < 0.025$.

$\Gamma(\phi K^+)/\Gamma(\phi \pi^+)$ $\Gamma_{142}/\Gamma_{130}$

A doubly Cabibbo-suppressed decay with no simple spectator process possible.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

<0.021	90		FRABETTI	95F E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV
------------------	----	--	----------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.058 ^{+0.032} _{-0.026} ± 0.007	4	⁶⁴ ANJOS	92D E691	γ Be, $\bar{E}_\gamma = 145$ GeV
---	---	---------------------	----------	---

⁶⁴ The evidence of ANJOS 92D is a small excess of events ($4.5^{+2.4}_{-2.0}$).

$\Gamma(\pi^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{143}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

<6.6 × 10⁻⁵	90		AITALA	96 E791	π^- N 500 GeV
----------------------------------	----	--	--------	---------	-------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.1 × 10 ⁻⁴	90		FRABETTI	97B E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV
-------------------------	----	--	----------	----------	---

<2.5 × 10 ⁻³	90		WEIR	90B MRK2	$e^+ e^-$ 29 GeV
-------------------------	----	--	------	----------	------------------

<2.6 × 10 ⁻³	90	39	HAAS	88 CLEO	$e^+ e^-$ 10 GeV
-------------------------	----	----	------	---------	------------------

$\Gamma(\pi^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{144}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

<1.8 × 10⁻⁵	90		AITALA	96 E791	π^- N 500 GeV
----------------------------------	----	--	--------	---------	-------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<8.9 × 10 ⁻⁵	90		FRABETTI	97B E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV
-------------------------	----	--	----------	----------	---

<2.2 × 10 ⁻⁴	90	0	KODAMA	95 E653	π^- emulsion 600 GeV
-------------------------	----	---	--------	---------	--------------------------

<5.9 × 10 ⁻³	90		WEIR	90B MRK2	$e^+ e^-$ 29 GeV
-------------------------	----	--	------	----------	------------------

<2.9 × 10 ⁻³	90	36	HAAS	88 CLEO	$e^+ e^-$ 10 GeV
-------------------------	----	----	------	---------	------------------

$\Gamma(\rho^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{145}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<5.6 \times 10^{-4}$	90	0	KODAMA	95 E653	π^- emulsion 600 GeV

$\Gamma(K^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{146}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-4}$	90	FRABETTI	97B E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<4.8 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV
-----------------------	----	------	----------	------------------

$\Gamma(K^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{147}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<9.7 \times 10^{-5}$	90		FRABETTI	97B E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<3.2 \times 10^{-4}$	90	0	KODAMA	95 E653	π^- emulsion 600 GeV
$<9.2 \times 10^{-3}$	90		WEIR	90B MRK2	$e^+ e^-$ 29 GeV

$\Gamma(\pi^+ e^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{148}/Γ

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-4}$	90	FRABETTI	97B E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<3.3 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV
-----------------------	----	------	----------	------------------

$\Gamma(\pi^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{149}/Γ

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-4}$	90	FRABETTI	97B E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<3.3 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV
-----------------------	----	------	----------	------------------

$\Gamma(K^+ e^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{150}/Γ

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-4}$	90	FRABETTI	97B E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<3.4 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV
-----------------------	----	------	----------	------------------

$\Gamma(K^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{151}/Γ

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-4}$	90	FRABETTI	97B E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<3.4 \times 10^{-3}$	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV
-----------------------	----	------	----------	------------------

$\Gamma(\pi^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{152}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.1 × 10⁻⁴	90	FRABETTI	97B E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.8 × 10 ⁻³	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV
-------------------------	----	------	----------	------------------

$\Gamma(\pi^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{153}/Γ

A test of lepton-number conservation.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<8.7 × 10⁻⁵	90		FRABETTI	97B E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.2 × 10 ⁻⁴	90	0	KODAMA	95 E653	π^- emulsion 600 GeV
<6.8 × 10 ⁻³	90		WEIR	90B MRK2	$e^+ e^-$ 29 GeV

$\Gamma(\pi^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{154}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.1 × 10⁻⁴	90	FRABETTI	97B E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.7 × 10 ⁻³	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV
-------------------------	----	------	----------	------------------

$\Gamma(\rho^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{155}/Γ

A test of lepton-number conservation.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<5.6 × 10⁻⁴	90	0	KODAMA	95 E653	π^- emulsion 600 GeV

$\Gamma(K^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{156}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.2 × 10⁻⁴	90	FRABETTI	97B E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<9.1 × 10 ⁻³	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV
-------------------------	----	------	----------	------------------

$\Gamma(K^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{157}/Γ

A test of lepton-number conservation.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<1.2 × 10⁻⁴	90		FRABETTI	97B E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.2 × 10 ⁻⁴	90	0	KODAMA	95 E653	π^- emulsion 600 GeV
<4.3 × 10 ⁻³	90		WEIR	90B MRK2	$e^+ e^-$ 29 GeV

$\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{158}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.3 × 10⁻⁴	90	FRABETTI	97B E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.0 × 10 ⁻³	90	WEIR	90B MRK2	$e^+ e^-$ 29 GeV
-------------------------	----	------	----------	------------------

$\Gamma(K^*(892)^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{159}/Γ

A test of lepton-number conservation.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<8.5 \times 10^{-4}$	90	0	KODAMA	95 E653	π^- emulsion 600 GeV

D^\pm CP-VIOLATING DECAY-RATE ASYMMETRIES

$A_{CP}(K^+ K^- \pi^\pm)$ in $D^\pm \rightarrow K^+ K^- \pi^\pm$

This is the difference between D^+ and D^- partial widths for these modes divided by the sum of the widths.

VALUE	DOCUMENT ID	TECN	COMMENT
-0.017 ± 0.027 OUR AVERAGE			
-0.014 ± 0.029	⁶⁵ AITALA	97B E791	$-0.062 < A_{CP} < +0.034$ (90% CL)
-0.031 ± 0.068	⁶⁵ FRABETTI	94I E687	$-0.14 < A_{CP} < +0.081$ (90% CL)

⁶⁵ FRABETTI 94I and AITALA 97B measure $N(D^+ \rightarrow K^- K^+ \pi^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$, the ratio of numbers of events observed, and similarly for the D^- .

$A_{CP}(K^\pm K^{*0})$ in $D^+ \rightarrow K^+ \bar{K}^{*0}$ and $D^- \rightarrow K^- K^{*0}$

This is the difference between D^+ and D^- partial widths for these modes divided by the sum of the widths.

VALUE	DOCUMENT ID	TECN	COMMENT
-0.02 ± 0.05 OUR AVERAGE			
-0.010 ± 0.050	⁶⁶ AITALA	97B E791	$-0.092 < A_{CP} < +0.072$ (90% CL)
-0.12 ± 0.13	⁶⁶ FRABETTI	94I E687	$-0.33 < A_{CP} < +0.094$ (90% CL)

⁶⁶ FRABETTI 94I and AITALA 97B measure $N(D^+ \rightarrow K^+ \bar{K}^*(892)^0)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$, the ratio of numbers of events observed, and similarly for the D^- .

$A_{CP}(\phi \pi^\pm)$ in $D^\pm \rightarrow \phi \pi^\pm$

This is the difference between D^+ and D^- partial widths for these modes divided by the sum of the widths.

VALUE	DOCUMENT ID	TECN	COMMENT
-0.014 ± 0.033 OUR AVERAGE			
-0.028 ± 0.036	⁶⁷ AITALA	97B E791	$-0.087 < A_{CP} < +0.031$ (90% CL)
$+0.066 \pm 0.086$	⁶⁷ FRABETTI	94I E687	$-0.075 < A_{CP} < +0.21$ (90% CL)

⁶⁷ FRABETTI 94I and AITALA 97B measure $N(D^+ \rightarrow \phi \pi^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$, the ratio of numbers of events observed, and similarly for the D^- .

$A_{CP}(\pi^+ \pi^- \pi^\pm)$ in $D^\pm \rightarrow \pi^+ \pi^- \pi^\pm$

This is the difference between D^+ and D^- partial widths for these modes divided by the sum of the widths.

VALUE	DOCUMENT ID	TECN	COMMENT
-0.017 ± 0.042	⁶⁸ AITALA	97B E791	$-0.086 < A_{CP} < +0.052$ (90% CL)

⁶⁸ AITALA 97B measure $N(D^+ \rightarrow \pi^+ \pi^- \pi^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$, the ratio of numbers of events observed, and similarly for the D^- .

D^\pm PRODUCTION CROSS SECTION AT $\psi(3770)$

A compilation of the cross sections for the direct production of D^\pm mesons at or near the $\psi(3770)$ peak in e^+e^- production.

<u>VALUE (nanobarns)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4.2 ± 0.6 ± 0.3	⁶⁹ ADLER	88C MRK3	e^+e^- 3.768 GeV
5.5 ± 1.0	⁷⁰ PARTRIDGE	84 CBAL	e^+e^- 3.771 GeV
6.00 ± 0.72 ± 1.02	⁷¹ SCHINDLER	80 MRK2	e^+e^- 3.771 GeV
9.1 ± 2.0	⁷² PERUZZI	77 MRK1	e^+e^- 3.774 GeV

⁶⁹ This measurement compares events with one detected D to those with two detected D mesons, to determine the the absolute cross section. ADLER 88C measure the ratio of cross sections (neutral to charged) to be $1.36 \pm 0.23 \pm 0.14$. This measurement does not include the decays of the $\psi(3770)$ not associated with charmed particle production.

⁷⁰ This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. PARTRIDGE 84 measures 6.4 ± 1.15 nb for the cross section. We take the phase space division of neutral and charged D mesons in $\psi(3770)$ decay to be 1.33, and we assume that the $\psi(3770)$ is an isosinglet to evaluate the cross sections. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction.

⁷¹ This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. SCHINDLER 80 assume the phase space division of neutral and charged D mesons in $\psi(3770)$ decay to be 1.33, and that the $\psi(3770)$ is an isosinglet. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction.

⁷² This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. The phase space division of neutral and charged D mesons in $\psi(3770)$ decay is taken to be 1.33, and $\psi(3770)$ is assumed to be an isosinglet. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction. We exclude this measurement from the average because of uncertainties in the contamination from τ lepton pairs. Also see RAPIDIS 77.

$D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$ FORM FACTORS

$r_\nu \equiv V(0)/A_1(0)$ in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.82 ± 0.09 OUR AVERAGE				
1.45 ± 0.23 ± 0.07	763	ADAMOVICH	99 BEAT	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.90 ± 0.11 ± 0.09	3000	⁷³ AITALA	98B E791	$\bar{K}^*(892)^0 e^+ \nu_e$
1.84 ± 0.11 ± 0.09	3034	AITALA	98F E791	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.74 ± 0.27 ± 0.28	874	FRABETTI	93E E687	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
2.00 $^{+0.34}_{-0.32}$ ± 0.16	305	KODAMA	92 E653	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
2.0 ± 0.6 ± 0.3	183	ANJOS	90E E691	$\bar{K}^*(892)^0 e^+ \nu_e$

⁷³ This is slightly different from the AITALA 98B value: see ref. [5] in AITALA 98F.

$r_2 \equiv A_2(0)/A_1(0)$ in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.78±0.07 OUR AVERAGE				
1.00±0.15±0.03	763	ADAMOVICH	99 BEAT	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
0.71±0.08±0.09	3000	AITALA	98B E791	$\bar{K}^*(892)^0 e^+ \nu_e$
0.75±0.08±0.09	3034	AITALA	98F E791	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
0.78±0.18±0.10	874	FRABETTI	93E E687	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
0.82 ^{+0.22} _{-0.23} ±0.11	305	KODAMA	92 E653	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
0.0 ±0.5 ±0.2	183	ANJOS	90E E691	$\bar{K}^*(892)^0 e^+ \nu_e$

$r_3 \equiv A_3(0)/A_1(0)$ in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$


<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.04±0.33±0.29				
	3034	AITALA	98F E791	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$

Γ_L/Γ_T in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.14±0.08 OUR AVERAGE				
1.09±0.10±0.02	763	ADAMOVICH	99 BEAT	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.20±0.13±0.13	874	FRABETTI	93E E687	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.18±0.18±0.08	305	KODAMA	92 E653	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
1.8 ^{+0.6} _{-0.4} ±0.3	183	ANJOS	90E E691	$\bar{K}^*(892)^0 e^+ \nu_e$

Γ_+/Γ_- in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.21±0.04 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.				
0.28±0.05±0.02	763	ADAMOVICH	99 BEAT	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
0.16±0.05±0.02	305	KODAMA	92 E653	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$
0.15 ^{+0.07} _{-0.05} ±0.03	183	ANJOS	90E E691	$\bar{K}^*(892)^0 e^+ \nu_e$



$$\Gamma_+/\Gamma_- \text{ in } D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$$

D^\pm REFERENCES

ADAMOVICH	99	EPJ C6 35	M. Adamovich+	(CERN BEATRICE Collab.)
AITALA	98B	PRL 80 1393	+Amato, Anjos, Appel+	(FNAL E791 Collab.)
AITALA	98F	PL B440 435	E.M. Aitala+	(FNAL E791 Collab.)
BAI	98B	PL B429 188	J.Z. Bai+	(BEPC BES Collab.)
JESSOP	98	PR D58 052002	C.P. Jessop+	(CLEO Collab.)
PDG	98	EPJ C3 1	C. Caso+	
AITALA	97	PL B397 325	+Amato, Anjos, Appel+	(FNAL E791 Collab.)
AITALA	97B	PL B403 377	+Amato, Anjos, Appel+	(FNAL E791 Collab.)
AITALA	97C	PL B404 187	+Amato, Anjos, Appel+	(FNAL E791 Collab.)
BARTELT	97	PL B405 373	+Csorna, Jain, Marka+	(CLEO Collab.)
BISHAI	97	PRL 78 3261	+Fast, Gerndt, Hinson+	(CLEO Collab.)
FRABETTI	97	PL B391 235	+Cheung, Cumalat+	(FNAL E687 Collab.)
FRABETTI	97B	PL B398 239	+Cheung, Cumalat+	(FNAL E687 Collab.)
FRABETTI	97C	PL B401 131	+Cheung, Cumalat+	(FNAL E687 Collab.)
FRABETTI	97D	PL B407 79	+Cheung, Cumalat+	(FNAL E687 Collab.)
AITALA	96	PRL 76 364	+Amato, Anjos+	(FNAL E791 Collab.)
ALBRECHT	96C	PL B374 249	+Hamacher, Hofmann+	(ARGUS Collab.)
BIGI	95	PL B349 363	+Yamamoto	(NDAM, HARV)
FRABETTI	95	PL B346 199	+Cheung, Cumalat+	(FNAL E687 Collab.)
FRABETTI	95B	PL B351 591	+Cheung, Cumalat+	(FNAL E687 Collab.)
FRABETTI	95E	PL B359 403	+Cheung, Cumalat+	(FNAL E687 Collab.)
FRABETTI	95F	PL B363 259	+Cheung, Cumalat+	(FNAL E687 Collab.)
KODAMA	95	PL B345 85	+Ushida, Mokhtarani+	(FNAL E653 Collab.)
ALBRECHT	94I	ZPHY C64 375	+Hamacher, Hofmann+	(ARGUS Collab.)
ALEEV	94	PAN 57 1370	+Balandin+	(Serpukhov BIS-2 Collab.)

Translated from YF 57 1443.

BALEST	94	PRL 72 2328	+Cho, Daoudi, Ford+	(CLEO Collab.)
FRABETTI	94D	PL B323 459	+Cheung, Cumalat+	(FNAL E687 Collab.)
FRABETTI	94G	PL B331 217	+Cheung, Cumalat+	(FNAL E687 Collab.)
FRABETTI	94I	PR D50 R2953	+Cheung, Cumalat+	(FNAL E687 Collab.)
ABE	93E	PL B313 288	+Amako, Arai, Arima, Asano+	(VENUS Collab.)
ADAMOVICH	93	PL B305 177	+Alexandrov, Antinori+	(CERN WA82 Collab.)
AKERIB	93	PRL 71 3070	+Barish, Chadha, Chan+	(CLEO Collab.)
ALAM	93	PRL 71 1311	+Kim, Nemati, O'Neill+	(CLEO Collab.)
ANJOS	93	PR D48 56	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
BEAN	93C	PL B317 647	+Gronberg, Kutschke, Menary+	(CLEO Collab.)
FRABETTI	93E	PL B307 262	+Grim, Paolone, Yager+	(FNAL E687 Collab.)
KODAMA	93B	PL B313 260	+Ushida, Mokhtarani+	(FNAL E653 Collab.)
KODAMA	93C	PL B316 455	+Ushida, Mokhtarani+	(FNAL E653 Collab.)
SELEN	93	PRL 71 1973	+Sadoff, Ammar, Ball+	(CLEO Collab.)
ALBRECHT	92B	ZPHY C53 361	+Ehrlichmann, Hamacher, Krueger+	(ARGUS Collab.)
ALBRECHT	92F	PL B278 202	+Ehrlichmann, Hamacher+	(ARGUS Collab.)
ANJOS	92	PR D45 R2177	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ANJOS	92C	PR D46 1941	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ANJOS	92D	PRL 69 2892	+Appel, Bean, Bediaga+	(FNAL E691 Collab.)
BARLAG	92C	ZPHY C55 383	+Becker, Boezek, Boehringer+	(ACCMOR Collab.)
Also	90D	ZPHY C48 29	+Barlag, Becker, Boehringer, Bosman+	(ACCMOR Collab.)
COFFMAN	92B	PR D45 2196	+DeJongh, Dubois, Eigen+	(Mark III Collab.)
DAOUDI	92	PR D45 3965	+Ford, Johnson, Lingel+	(CLEO Collab.)
FRABETTI	92	PL B281 167	+Bogart, Cheung, Culy+	(FNAL E687 Collab.)
KODAMA	92	PL B274 246	+Ushida, Mokhtarani+	(FNAL E653 Collab.)
KODAMA	92C	PL B286 187	+Ushida, Mokhtarani+	(FNAL E653 Collab.)
ADAMOVICH	91	PL B268 142	+Alexandrov, Antinori, Barberis+	(WA82 Collab.)
ALBRECHT	91	PL B255 634	+Ehrlichmann, Hamacher, Krueger+	(ARGUS Collab.)
ALVAREZ	91	PL B255 639	+Barate, Bloch, Bonamy+	(CERN NA14/2 Collab.)
ALVAREZ	91B	ZPHY C50 11	+Barate, Bloch, Bonamy+	(CERN NA14/2 Collab.)
AMMAR	91	PR D44 3383	+Baringer, Coppage, Davis+	(CLEO Collab.)
ANJOS	91B	PR D43 R2063	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ANJOS	91C	PRL 67 1507	+Appel, Bean, Bracker+	(FNAL-TPS Collab.)
BAI	91	PRL 66 1011	+Bolton, Brown, Bunnell+	(Mark III Collab.)
COFFMAN	91	PL B263 135	+DeJongh, Dubois, Eigen, Hitlin+	(Mark III Collab.)
FRABETTI	91	PL B263 584	+Bogart, Cheung, Culy+	(FNAL E687 Collab.)
ALVAREZ	90	ZPHY C47 539	+Barate, Bloch, Bonamy+	(CERN NA14/2 Collab.)
ALVAREZ	90C	PL B246 261	+Barate, Bloch, Bonamy+	(CERN NA14/2 Collab.)
ANJOS	90C	PR D41 2705	+Appel, Bean+	(FNAL E691 Collab.)
ANJOS	90D	PR D42 2414	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ANJOS	90E	PRL 65 2630	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
BARLAG	90C	ZPHY C46 563	+Becker, Boehringer, Bosman+	(ACCMOR Collab.)
WEIR	90B	PR D41 1384	+Klein, Abrams, Adolphsen, Akerlof+	(Mark II Collab.)
ANJOS	89	PRL 62 125	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ANJOS	89B	PRL 62 722	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ANJOS	89E	PL B223 267	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ADLER	88B	PRL 60 1375	+Becker, Blaylock+	(Mark III Collab.)
ADLER	88C	PRL 60 89	+Becker, Blaylock+	(Mark III Collab.)
ALBRECHT	88I	PL B210 267	+Boeckmann, Glaeser+	(ARGUS Collab.)
ANJOS	88	PRL 60 897	+Appel+	(FNAL E691 Collab.)
AOKI	88	PL B209 113	+Arnold, Baroni+	(WA75 Collab.)
HAAS	88	PRL 60 1614	+Hempstead, Jensen+	(CLEO Collab.)
ONG	88	PRL 60 2587	+Weir, Abrams, Amidei+	(Mark II Collab.)
RAAB	88	PR D37 2391	+Anjos, Appel, Bracker+	(FNAL E691 Collab.)
ADAMOVICH	87	EPL 4 887	+Alexandrov, Bolta+	(Photon Emulsion Collab.)
ADLER	87	PL B196 107	+Becker, Blaylock, Bolton+	(Mark III Collab.)
AGUILAR-...	87D	PL B193 140	Aguilar-Benitez, Allison+	(LEBC-EHS Collab.)
Also	88B	ZPHY C40 321	Aguilar-Benitez, Allison, Bailly+	(LEBC-EHS Collab.)
AGUILAR-...	87E	ZPHY C36 551	Aguilar-Benitez, Allison+	(LEBC-EHS Collab.)
Also	88B	ZPHY C40 321	Aguilar-Benitez, Allison, Bailly+	(LEBC-EHS Collab.)
AGUILAR-...	87F	ZPHY C36 559	Aguilar-Benitez, Allison+	(LEBC-EHS Collab.)
Also	88	ZPHY C38 520 erratum		
BARLAG	87B	ZPHY C37 17	+Becker, Boehringer, Bosman+	(ACCMOR Collab.)
BARTEL	87	ZPHY C33 339	+Becker, Felst, Haidt+	(JADE Collab.)
CSORNA	87	PL B191 318	+Mestayer, Panvini, Word+	(CLEO Collab.)
PALKA	87B	ZPHY C35 151	+Bailey, Becker+	(ACCMOR Collab.)
ABE	86	PR D33 1	+	(SLAC Hybrid Facility Photon Collab.)
AGUILAR-...	86B	ZPHY C31 491	Aguilar-Benitez, Allison+	(LEBC-EHS Collab.)

BALTRUSAIT...	86E	PRL 56 2140	Baltrusaitis, Becker, Blaylock, Brown+	(Mark III Collab.)
PAL	86	PR D33 2708	+Atwood, Barish, Bonneaud+	(DELCO Collab.)
AIHARA	85	ZPHY C27 39	+Alston-Garnjost, Badtke, Bakken+	(TPC Collab.)
BALTRUSAIT...	85B	PRL 54 1976	Baltrusaitis, Becker, Blaylock, Brown+	(Mark III Collab.)
BALTRUSAIT...	85E	PRL 55 150	Baltrusaitis, Becker, Blaylock, Brown+	(Mark III Collab.)
BARTEL	85J	PL 163B 277	+Becker, Cords, Felst+	(JADE Collab.)
ADAMOVIH	84	PL 140B 119	+Alexandrov, Bolta, Bravo+	(CERN WA58 Collab.)
ALTHOFF	84G	ZPHY C22 219	+Braunschweig, Kirschfink+	(TASSO Collab.)
ALTHOFF	84J	PL 146B 443	+Branschweig, Kirschfink+	(TASSO Collab.)
DERRICK	84	PRL 53 1971	+Fernandez, Fries, Hyman+	(HRS Collab.)
KOOP	84	PRL 52 970	+Sakuda, Atwood, Baillon+	(DELCO Collab.)
PARTRIDGE	84	Thesis CALT-68-1150		(Crystal Ball Collab.)
AGUILAR-...	83B	PL 123B 98	Aguilar-Benitez, Allison+	(LEBC-EHS Collab.)
AUBERT	83	NP B213 31	+Bassompierre, Becks, Best+	(EMC Collab.)
PARTRIDGE	81	PRL 47 760	+Peck, Porter, Gu+	(Crystal Ball Collab.)
SCHINDLER	81	PR D24 78	+Alam, Boyarski, Breidenbach+	(Mark II Collab.)
TRILLING	81	PRPL 75 57		(LBL, UCB) J
BACINO	80	PRL 45 329	+Ferguson+	(DELCO Collab.)
SCHINDLER	80	PR D21 2716	+Siegrist, Alam, Boyarski+	(Mark II Collab.)
ZHOLENTZ	80	PL 96B 214	+Kurdadze, Lelchuk, Mishnev+	(NOVO)
Also	81	SJNP 34 814	Zholentz, Kurdadze, Lelchuk+	(NOVO)
		Translated from YAF 34 1471.		
BACINO	79	PRL 43 1073	+Ferguson, Nodulman+	(DELCO Collab.)
BRANDELIK	79	PL 80B 412	+Braunschweig, Martyn, Sander+	(DASP Collab.)
FELLER	78	PRL 40 274	+Litke, Madaras, Ronan+	(Mark I Collab.)
VUILLEMIN	78	PRL 41 1149	+Feldman, Feller+	(Mark I Collab.)
GOLDHABER	77	PL 69B 503	+Wiss, Abrams, Alam+	(Mark I Collab.)
PERUZZI	77	PRL 39 1301	+Piccolo, Feldman+	(Mark I Collab.)
PICCOLO	77	PL 70B 260	+Peruzzi, Luth, Nguyen, Wiss, Abrams+	(Mark I Collab.)
RAPIDIS	77	PRL 39 526	+Gobbi, Luke, Barbaro-Galtieri+	(Mark I Collab.)
PERUZZI	76	PRL 37 569	+Piccolo, Feldman, Nguyen, Wiss+	(Mark I Collab.)

OTHER RELATED PAPERS

RICHMAN	95	RMP 67 893	+Burchat	(UCSB, STAN)
ROSNER	95	CNPP 21 369		(CHIC)