

$\Delta(1940) D_{33}$

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^-) \text{ Status: } *$$

OMITTED FROM SUMMARY TABLE

 $\Delta(1940)$ BREIT-WIGNER MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
≈ 1940 OUR ESTIMATE			
2057 ± 110	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
2058.1 ± 34.5	CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$
1940 ± 100	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

 $\Delta(1940)$ BREIT-WIGNER WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
460 ± 320	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
198.4 ± 45.5	CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$
200 ± 100	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

 $\Delta(1940)$ POLE POSITION**REAL PART**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1900 ± 100	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
1915 or 1926	¹ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$

-2xIMAGINARY PART

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
200 ± 60	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
190 or 186	¹ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$

 $\Delta(1940)$ ELASTIC POLE RESIDUE**MODULUS $|r|$**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8 ± 3	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

PHASE θ

<u>VALUE ($^\circ$)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
135 ± 45	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

Δ(1940) DECAY MODES

Mode
Γ ₁ <i>N</i> π
Γ ₂ Σ <i>K</i>
Γ ₃ <i>N</i> ππ
Γ ₄ Δ(1232)π, <i>S</i> -wave
Γ ₅ Δ(1232)π, <i>D</i> -wave
Γ ₆ <i>N</i> ρ, <i>S</i> =3/2, <i>S</i> -wave
Γ ₇ <i>N</i> γ, helicity=1/2
Γ ₈ <i>N</i> γ, helicity=3/2

Δ(1940) BRANCHING RATIOS

Γ(<i>N</i> π)/Γ _{total}	Γ ₁ /Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.18±0.12	MANLEY 92 IPWA π <i>N</i> → π <i>N</i> & <i>N</i> ππ
0.18	CHEW 80 BPWA π ⁺ <i>p</i> → π ⁺ <i>p</i>
0.05±0.02	CUTKOSKY 80 IPWA π <i>N</i> → π <i>N</i>

(Γ _{<i>i</i>} Γ _{<i>f</i>}) ^{1/2} /Γ _{total} in <i>N</i> π → Δ(1940) → Σ <i>K</i>	(Γ ₁ Γ ₂) ^{1/2} /Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<0.015	CANDLIN 84 DPWA π ⁺ <i>p</i> → Σ ⁺ <i>K</i> ⁺

(Γ _{<i>i</i>} Γ _{<i>f</i>}) ^{1/2} /Γ _{total} in <i>N</i> π → Δ(1940) → Δ(1232)π, <i>S</i> -wave	(Γ ₁ Γ ₄) ^{1/2} /Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
+0.11±0.10	MANLEY 92 IPWA π <i>N</i> → π <i>N</i> & <i>N</i> ππ

(Γ _{<i>i</i>} Γ _{<i>f</i>}) ^{1/2} /Γ _{total} in <i>N</i> π → Δ(1940) → Δ(1232)π, <i>D</i> -wave	(Γ ₁ Γ ₅) ^{1/2} /Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
+0.27±0.16	MANLEY 92 IPWA π <i>N</i> → π <i>N</i> & <i>N</i> ππ

(Γ _{<i>i</i>} Γ _{<i>f</i>}) ^{1/2} /Γ _{total} in <i>N</i> π → Δ(1940) → <i>N</i> ρ, <i>S</i> =3/2, <i>S</i> -wave	(Γ ₁ Γ ₆) ^{1/2} /Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
+0.25±0.10	MANLEY 92 IPWA π <i>N</i> → π <i>N</i> & <i>N</i> ππ

Δ(1940) PHOTON DECAY AMPLITUDES

Δ(1940) → <i>N</i> γ, helicity-1/2 amplitude A _{1/2}			
<u>VALUE (GeV^{-1/2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.036±0.058	AWAJI 81	DPWA	γ <i>N</i> → π <i>N</i>

Δ(1940) → <i>N</i> γ, helicity-3/2 amplitude A _{3/2}			
<u>VALUE (GeV^{-1/2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.031±0.012	AWAJI 81	DPWA	γ <i>N</i> → π <i>N</i>

$\Delta(1940)$ FOOTNOTES

¹ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.

$\Delta(1940)$ REFERENCES

MANLEY	92	PR D45 4002	D.M. Manley, E.M. Saleski	(KENT) IJP
Also	84	PR D30 904	D.M. Manley <i>et al.</i>	(VPI)
CANDLIN	84	NP B238 477	D.J. Candlin <i>et al.</i>	(EDIN, RAL, LOWC)
AWAJI	81	Bonn Conf. 352	N. Awaji, R. Kajikawa	(NAGO)
Also	82	NP B197 365	K. Fujii <i>et al.</i>	(NAGO)
CHEW	80	Toronto Conf. 123	D.M. Chew	(LBL) IJP
CUTKOSKY	80	Toronto Conf. 19	R.E. Cutkosky <i>et al.</i>	(CMU, LBL) IJP
Also	79	PR D20 2839	R.E. Cutkosky <i>et al.</i>	(CMU, LBL)
LONGACRE	78	PR D17 1795	R.S. Longacre <i>et al.</i>	(LBL, SLAC)
