

# $f_1(1285)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

## $f_1(1285)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1281.9 ± 0.6 OUR AVERAGE</b>		Error includes scale factor of 1.7. See the ideogram below.		
1284 ± 6	1400	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
1281 ± 1		BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
1281 ± 1		BARBERIS	97C OMEG	450 $pp \rightarrow ppK_S^0 K^\pm \pi^\mp$
1280 ± 2		<sup>1</sup> ANTINORI	95 OMEG	300,450 $pp \rightarrow pp2(\pi^+ \pi^-)$
1282.2 ± 1.5		LEE	94 MPS2	18 $\pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$
1279 ± 5		FUKUI	91C SPEC	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
1278 ± 2	140	ARMSTRONG	89 OMEG	300 $pp \rightarrow K \bar{K} \pi pp$
1278 ± 2		ARMSTRONG	89G OMEG	85 $\pi^+ p \rightarrow 4\pi \pi p$ , $pp \rightarrow 4\pi pp$
1280.1 ± 2.1	60	RATH	89 MPS	21.4 $\pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
1285 ± 1	4750	<sup>2</sup> BIRMAN	88 MPS	8 $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
1280 ± 1	504	BITYUKOV	88 SPEC	32.5 $\pi^- p \rightarrow K^+ K^- \pi^0 n$
1280 ± 4		ANDO	86 SPEC	8 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
1277 ± 2	420	REEVES	86 SPEC	6.6 $p\bar{p} \rightarrow K K \pi X$
1285 ± 2		CHUNG	85 SPEC	8 $\pi^- p \rightarrow N K \bar{K} \pi$
1279 ± 2	604	ARMSTRONG	84 OMEG	85 $\pi^+ p \rightarrow K \bar{K} \pi \pi p$ , $pp \rightarrow K \bar{K} \pi pp$
1286 ± 1		CHAUVAT	84 SPEC	ISR 31.5 $pp$
1278 ± 4		EVANGELISTA	81 OMEG	12 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$
1283 ± 3	103	DIONISI	80 HBC	4 $\pi^- p \rightarrow K \bar{K} \pi n$
1282 ± 2	320	NACASCH	78 HBC	0.7,0.76 $\bar{p}p \rightarrow K \bar{K} 3\pi$
1279 ± 5	210	GRASSLER	77 HBC	16 $\pi^\mp p$
1286 ± 3	180	DUBOC	72 HBC	1.2 $\bar{p}p \rightarrow 2K 4\pi$
1283 ± 5		DAHL	67 HBC	1.6–4.2 $\pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1281.9 ± 0.5		<sup>3</sup> SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-) p_{\text{fast}}$

1282.8 ± 0.6		<sup>3</sup> SOSA	99	SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+) p_{\text{fast}}$
1270 ± 10		AMELIN	95	VES	$37 \pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$
1280 ± 2		ABATZIS	94	OMEG	$450 pp \rightarrow pp2(\pi^+ \pi^-)$
1282 ± 4		ARMSTRONG	93C	E760	$\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
1270 ± 6 ± 10		ARMSTRONG	92C	OMEG	$300 pp \rightarrow pp\pi^+ \pi^- \gamma$
1264 ± 8		AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
1281 ± 1		ARMSTRONG	89E	OMEG	$300 pp \rightarrow pp2(\pi^+ \pi^-)$
1279 ± 6 ± 10	16	BECKER	87	MRK3	$e^+ e^- \rightarrow \phi K \bar{K} \pi$
1286 ± 9		GIDAL	87	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
1287 ± 5	353	BITYUKOV	84B	SPEC	$32 \pi^- p \rightarrow K^+ K^- \pi^0 n$
~ 1279		<sup>4</sup> TORNQVIST	82B	RVUE	
1275 ± 6	31	BROMBERG	80	SPEC	$100 \pi^- p \rightarrow K \bar{K} \pi X$
1288 ± 9	200	GURTU	79	HBC	$4.2 K^- p \rightarrow n \eta 2\pi$
~ 1275.0	46	<sup>5</sup> STANTON	79	CNTR	$8.5 \pi^- p \rightarrow n 2\gamma 2\pi$
1271 ± 10	34	CORDEN	78	OMEG	$12-15 \pi^- p \rightarrow K^+ K^- \pi n$
1295 ± 12	85	CORDEN	78	OMEG	$12-15 \pi^- p \rightarrow n 5\pi$
1292 ± 10	150	DEFOIX	72	HBC	$0.7 \bar{p}p \rightarrow 7\pi$
1280 ± 3	500	<sup>6</sup> THUN	72	MMS	$13.4 \pi^- p$
1303 ± 8		BARDADIN-...	71	HBC	$8 \pi^+ p \rightarrow p 6\pi$
1283 ± 6		BOESEBECK	71	HBC	$16.0 \pi p \rightarrow p 5\pi$
1270 ± 10		CAMPBELL	69	DBC	$2.7 \pi^+ d$
1285 ± 7		LORSTAD	69	HBC	$0.7 \bar{p}p, 4,5\text{-body}$
1290 ± 7		D'ANDLAU	68	HBC	$1.2 \bar{p}p, 5-6 \text{ body}$

<sup>1</sup> Supersedes ABATZIS 94, ARMSTRONG 89E.


<sup>2</sup> From partial wave analysis of  $K^+ \bar{K}^0 \pi^-$  system.

<sup>3</sup> No systematic error given.

<sup>4</sup> From a unitarized quark-model calculation.

<sup>5</sup> From phase shift analysis of  $\eta \pi^+ \pi^-$  system.

<sup>6</sup> Seen in the missing mass spectrum.



$f_1(1285)$  mass (MeV)

### $f_1(1285)$ WIDTH

Only experiments giving width error less than 20 MeV are kept for averaging.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>24.0± 1.2 OUR AVERAGE</b>		Error includes scale factor of 1.4. See the ideogram below.		
55 ±18	1400	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
24 ± 3		BARBERIS	97B OMEG	450 $pp \rightarrow$ $pp2(\pi^+ \pi^-)$
20 ± 2		BARBERIS	97C OMEG	450 $pp \rightarrow$ $ppK_S^0 K^\pm \pi^\mp$
36 ± 5		<sup>7</sup> ANTINORI	95 OMEG	300,450 $pp \rightarrow$ $pp2(\pi^+ \pi^-)$
29.0± 4.1		LEE	94 MPS2	18 $\pi^- p \rightarrow$ $K^+ \bar{K}^0 2\pi^- p$
25 ± 4	140	ARMSTRONG	89 OMEG	300 $pp \rightarrow K \bar{K} \pi pp$
22 ± 2	4750	<sup>8</sup> BIRMAN	88 MPS	8 $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
25 ± 4	504	BITYUKOV	88 SPEC	32.5 $\pi^- p \rightarrow$ $K^+ K^- \pi^0 n$

19 ± 5		ANDO	86	SPEC	8 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
32 ± 8	420	REEVES	86	SPEC	6.6 $p \bar{p} \rightarrow K K \pi X$
22 ± 2		CHUNG	85	SPEC	8 $\pi^- p \rightarrow N K \bar{K} \pi$
32 ± 3	604	ARMSTRONG	84	OMEG	85 $\pi^+ p \rightarrow K \bar{K} \pi \pi p$ , $p p \rightarrow K \bar{K} \pi p p$
24 ± 3		CHAUVAT	84	SPEC	ISR 31.5 $p p$
29 ± 10	103	DIONISI	80	HBC	4 $\pi^- p \rightarrow K \bar{K} \pi n$
28.3 ± 6.7	320	NACASCH	78	HBC	0.7, 0.76 $\bar{p} p \rightarrow K \bar{K} 3\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
18.2 ± 1.2		<sup>9</sup> SOSA	99	SPEC	$p p \rightarrow p_{\text{slow}}$ $(K_S^0 K^+ \pi^-) p_{\text{fast}}$
19.4 ± 1.5		<sup>9</sup> SOSA	99	SPEC	$p p \rightarrow p_{\text{slow}}$ $(K_S^0 K^- \pi^+) p_{\text{fast}}$
40 ± 5		ABATZIS	94	OMEG	450 $p p \rightarrow$ $p p 2(\pi^+ \pi^-)$
44 ± 20		AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
31 ± 5		ARMSTRONG	89E	OMEG	300 $p p \rightarrow$ $p p 2(\pi^+ \pi^-)$
41 ± 12		ARMSTRONG	89G	OMEG	85 $\pi^+ p \rightarrow 4\pi \pi p$ , $p p \rightarrow 4\pi p p$
17.9 ± 10.9	60	RATH	89	MPS	21.4 $\pi^- p \rightarrow$ $K_S^0 K_S^0 \pi^0 n$
14 $\begin{smallmatrix} +20 \\ -14 \end{smallmatrix}$ ± 10	16	BECKER	87	MRK3	$e^+ e^- \rightarrow \phi K \bar{K} \pi$
26 ± 12		EVANGELISTA	81	OMEG	12 $\pi^- p \rightarrow$ $\eta \pi^+ \pi^- \pi^- p$
25 ± 15	200	GURTU	79	HBC	4.2 $K^- p \rightarrow n \eta 2\pi$
~ 10		<sup>10</sup> STANTON	79	CNTR	8.5 $\pi^- p \rightarrow n 2\gamma 2\pi$
24 ± 18	210	GRASSLER	77	HBC	16 $\pi^\mp p$
28 ± 5	150	<sup>11</sup> DEFOIX	72	HBC	0.7 $\bar{p} p \rightarrow 7\pi$
46 ± 9	180	<sup>11</sup> DUBOC	72	HBC	1.2 $\bar{p} p \rightarrow 2K 4\pi$
37 ± 5	500	<sup>12</sup> THUN	72	MMS	13.4 $\pi^- p$
10 ± 10		BOESEBECK	71	HBC	16.0 $\pi p \rightarrow p 5\pi$
30 ± 15		CAMPBELL	69	DBC	2.7 $\pi^+ d$
60 ± 15		<sup>11</sup> LORSTAD	69	HBC	0.7 $\bar{p} p$ , 4,5-body
35 ± 10		<sup>11</sup> DAHL	67	HBC	1.6–4.2 $\pi^- p$

<sup>7</sup> Supersedes ABATZIS 94, ARMSTRONG 89E.


<sup>8</sup> From partial wave analysis of  $K^+ \bar{K}^0 \pi^-$  system.

<sup>9</sup> No systematic error given.

<sup>10</sup> From phase shift analysis of  $\eta \pi^+ \pi^-$  system.

<sup>11</sup> Resolution is not unfolded.

<sup>12</sup> Seen in the missing mass spectrum.



$f_1(1285)$  width (MeV)

### $f_1(1285)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $4\pi$	$(33.1^{+2.1}_{-1.8})\%$	S=1.3
$\Gamma_2$ $\pi^0\pi^0\pi^+\pi^-$	$(22.0^{+1.4}_{-1.2})\%$	S=1.3
$\Gamma_3$ $2\pi^+2\pi^-$	$(11.0^{+0.7}_{-0.6})\%$	S=1.3
$\Gamma_4$ $\rho^0\pi^+\pi^-$	$(11.0^{+0.7}_{-0.6})\%$	S=1.3
$\Gamma_5$ $4\pi^0$	$< 7 \times 10^{-4}$	CL=90%
$\Gamma_6$ $\eta\pi\pi$	$(52 \pm 16)\%$	
$\Gamma_7$ $a_0(980)\pi$ [ignoring $a_0(980) \rightarrow K\bar{K}$ ]	$(36 \pm 7)\%$	
$\Gamma_8$ $\eta\pi\pi$ [excluding $a_0(980)\pi$ ]	$(16 \pm 7)\%$	
$\Gamma_9$ $K\bar{K}\pi$	$(9.0 \pm 0.4)\%$	S=1.1
$\Gamma_{10}$ $K\bar{K}^*(892)$	not seen	
$\Gamma_{11}$ $\gamma\rho^0$	$(5.5 \pm 1.3)\%$	S=2.8
$\Gamma_{12}$ $\phi\gamma$	$(7.4 \pm 2.6) \times 10^{-4}$	
$\Gamma_{13}$ $\gamma\gamma^*$		
$\Gamma_{14}$ $\gamma\gamma$		

## CONSTRAINED FIT INFORMATION

An overall fit to 7 branching ratios uses 16 measurements and one constraint to determine 5 parameters. The overall fit has a  $\chi^2 = 24.7$  for 12 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_7$	-17			
$x_8$	-8	-95		
$x_9$	46	-9	-4	
$x_{11}$	-36	-4	-2	-34
	$x_1$	$x_7$	$x_8$	$x_9$

### $f_1(1285) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_6\Gamma_{14}/\Gamma = (\Gamma_7+\Gamma_8)\Gamma_{14}/\Gamma$
VALUE (keV)	CL% DOCUMENT ID TECN COMMENT
<b>&lt;0.62</b>	95 GIDAL 87 MRK2 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma^*)/\Gamma_{\text{total}}$	$\Gamma_6\Gamma_{13}/\Gamma = (\Gamma_7+\Gamma_8)\Gamma_{13}/\Gamma$
VALUE (keV)	EVTS DOCUMENT ID TECN COMMENT
<b>1.4 ± 0.4 OUR AVERAGE</b>	Error includes scale factor of 1.4.
1.18 ± 0.25 ± 0.20	26 <sup>13,14</sup> AIHARA 88B TPC $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
2.30 ± 0.61 ± 0.42	13,15 GIDAL 87 MRK2 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$

<sup>13</sup> Assuming a  $\rho$ -pole form factor.

<sup>14</sup> Published value multiplied by  $\eta\pi\pi$  branching ratio 0.49.

<sup>15</sup> Published value divided by 2 and multiplied by the  $\eta\pi\pi$  branching ratio 0.49.

### $f_1(1285) \text{ BRANCHING RATIOS}$

$\Gamma(K\bar{K}\pi)/\Gamma(4\pi)$	$\Gamma_9/\Gamma_1$
VALUE	DOCUMENT ID TECN COMMENT
<b>0.271 ± 0.016 OUR FIT</b>	Error includes scale factor of 1.3.
<b>0.271 ± 0.016 OUR AVERAGE</b>	Error includes scale factor of 1.2.
0.265 ± 0.014	<sup>16</sup> BARBERIS 97C OMEG 450 $pp \rightarrow ppK_S^0 K^\pm \pi^\mp$
0.28 ± 0.05	<sup>17</sup> ARMSTRONG 89E OMEG 300 $pp \rightarrow pp f_1(1285)$
0.37 ± 0.03 ± 0.05	<sup>18</sup> ARMSTRONG 89G OMEG 85 $\pi p \rightarrow 4\pi X$

<sup>16</sup> Using  $2(\pi^+\pi^-)$  data from BARBERIS 97B.

<sup>17</sup> Assuming  $\rho\pi\pi$  and  $a_0(980)\pi$  intermediate states.

<sup>18</sup>  $4\pi$  consistent with being entirely  $\rho\pi\pi$ .

$$\Gamma(\pi^0 \pi^0 \pi^+ \pi^-) / \Gamma_{\text{total}} \qquad \Gamma_2 / \Gamma = \frac{2}{3} \Gamma_1 / \Gamma$$

VALUE DOCUMENT ID  
**0.220<sup>+0.014</sup><sub>-0.012</sub> OUR FIT** Error includes scale factor of 1.3.

$$\Gamma(2\pi^+ 2\pi^-) / \Gamma_{\text{total}} \qquad \Gamma_3 / \Gamma = \frac{1}{3} \Gamma_1 / \Gamma$$

VALUE DOCUMENT ID  
**0.110<sup>+0.007</sup><sub>-0.006</sub> OUR FIT** Error includes scale factor of 1.3.

$$\Gamma(\rho^0 \pi^+ \pi^-) / \Gamma_{\text{total}} \qquad \Gamma_4 / \Gamma = \frac{1}{3} \Gamma_1 / \Gamma$$

VALUE DOCUMENT ID  
**0.110<sup>+0.007</sup><sub>-0.006</sub> OUR FIT** Error includes scale factor of 1.3.

$$\Gamma(K \bar{K} \pi) / \Gamma(\eta \pi \pi) \qquad \Gamma_9 / \Gamma_6 = \Gamma_9 / (\Gamma_7 + \Gamma_8)$$

VALUE DOCUMENT ID TECN COMMENT  
**0.171 $\pm$ 0.013 OUR FIT** Error includes scale factor of 1.1.  
**0.170 $\pm$ 0.012 OUR AVERAGE**

0.166 $\pm$ 0.01 $\pm$ 0.008	BARBERIS	98C	OMEG	450 $p p \rightarrow p_f f_1(1285) p_s$
0.42 $\pm$ 0.15	GURTU	79	HBC	4.2 $K^- p$
0.5 $\pm$ 0.2	CORDEN	78	OMEG	12-15 $\pi^- p$
0.20 $\pm$ 0.08	<sup>19</sup> DEFOIX	72	HBC	0.7 $\bar{p} p \rightarrow 7\pi$
0.16 $\pm$ 0.08	CAMPBELL	69	DBC	2.7 $\pi^+ d$

<sup>19</sup>  $K \bar{K}$  system characterized by the  $l = 1$  threshold enhancement. (See under  $a_0(980)$ ).

$$\Gamma(a_0(980) \pi \text{ [ignoring } a_0(980) \rightarrow K \bar{K}]) / \Gamma(\eta \pi \pi) \qquad \Gamma_7 / \Gamma_6 = \Gamma_7 / (\Gamma_7 + \Gamma_8)$$

VALUE EVTS DOCUMENT ID TECN COMMENT  
**0.69 $\pm$ 0.13 OUR FIT**

**0.69<sup>+0.13</sup><sub>-0.12</sub> OUR AVERAGE**

0.72 $\pm$ 0.15	GURTU	79	HBC	4.2 $K^- p$
0.6 <sup>+0.3</sup> <sub>-0.2</sub>	CORDEN	78	OMEG	12-15 $\pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.28 $\pm$ 0.07	1400	ALDE	97B	GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
1.0 $\pm$ 0.3		GRASSLER	77	HBC	16 $\pi^\mp p$

$$\Gamma(4\pi) / \Gamma(\eta \pi \pi) \qquad \Gamma_1 / \Gamma_6 = \Gamma_1 / (\Gamma_7 + \Gamma_8)$$

VALUE DOCUMENT ID TECN COMMENT  
**0.63 $\pm$ 0.06 OUR FIT** Error includes scale factor of 1.2.

**0.41 $\pm$ 0.14 OUR AVERAGE**

0.37 $\pm$ 0.11 $\pm$ 0.11	BOLTON	92	MRK3	$J/\psi \rightarrow \gamma f_1(1285)$
0.64 $\pm$ 0.40	GURTU	79	HBC	4.2 $K^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.93 $\pm$ 0.30	<sup>20</sup> GRASSLER	77	HBC	16 $\pi^\mp p$
-----------------	------------------------	----	-----	----------------

<sup>20</sup> Assuming  $\rho \pi \pi$  and  $a_0(980) \pi$  intermediate states.

$\Gamma(K\bar{K}^*(892))/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>not seen</b>	NACASCH	78 HBC	0.7,0.76 $\bar{p}p \rightarrow K\bar{K}3\pi$

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma(2\pi^+2\pi^-)$   $\Gamma_4/\Gamma_3$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.0 \pm 0.4$	GRASSLER	77 HBC	16 GeV $\pi^\pm p$

$\Gamma(4\pi^0)/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;7</b>	90	ALDE	87 GAM4	100 $\pi^- p \rightarrow 4\pi^0 n$

$\Gamma(\phi\gamma)/\Gamma(K\bar{K}\pi)$   $\Gamma_{12}/\Gamma_9$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.82 \pm 0.21 \pm 0.20</math></b>		19	BITYUKOV	88 SPEC	32.5 $\pi^- p \rightarrow K^+ K^- \pi^0 n$
$<0.50$	95		BARBERIS	98C OMEG	450 $pp \rightarrow \rho_f f_1(1285) p_S$
$<0.93$	95		AMELIN	95 VES	37 $\pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$

$\Gamma(\gamma\rho^0)/\Gamma(K\bar{K}\pi)$   $\Gamma_{11}/\Gamma_9$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$>0.035$	90	<sup>21</sup> COFFMAN	90 MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
<sup>21</sup> Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma K\bar{K}\pi) < 0.72 \times 10^{-3}$ .				

$\Gamma(\gamma\rho^0)/\Gamma(2\pi^+2\pi^-)$   $\Gamma_{11}/\Gamma_3 = \Gamma_{11}/\frac{1}{3}\Gamma_1$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.50 \pm 0.13</math> OUR FIT</b>	Error includes scale factor of 2.5.		
<b><math>0.45 \pm 0.18</math></b>	<sup>22</sup> COFFMAN	90 MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
<sup>22</sup> Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma 2\pi^+ 2\pi^-) = 0.55 \times 10^{-4}$ given by MIR 88.			

$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.055 \pm 0.013</math> OUR FIT</b>	Error includes scale factor of 2.8.			
<b><math>0.028 \pm 0.007 \pm 0.006</math></b>		AMELIN	95 VES	37 $\pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$
$<0.05$	95	BITYUKOV	91B SPEC	32 $\pi^- p \rightarrow \pi^+ \pi^- \gamma n$





————— **OTHER RELATED PAPERS** —————

AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2 $\gamma$ Collab.) JPC
ASTON	85	PR D32 2255	D. Aston <i>et al.</i>	(SLAC, CARL, CNRC)
ATKINSON	84E	PL 138B 459	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
GAVILLET	82	ZPHY C16 119	P. Gavillet <i>et al.</i>	(CERN, CDEF, PADO+)
D'ANDLAU	65	PL 17 347	C. d'Andlau <i>et al.</i>	(CDEF, CERN, IRAD+)
MILLER	65	PRL 14 1074	D.H. Miller <i>et al.</i>	(LRL, UCB)

---