



$$I(J^P) = 0(\frac{1}{2}^+)$$

$$\text{Charge} = \frac{2}{3} e \quad \text{Top} = +1$$

THE TOP QUARK

Revised April 2000 by M. Mangano (CERN) and T. Trippe (LBNL).

A. Introduction: The top quark is the $Q = 2/3$, $T_3 = +1/2$ member of the weak-isospin doublet containing the bottom quark (see our review on the “Standard Model of Electroweak Interactions” for more information). This note summarizes its currently measured properties, and provides a discussion of the experimental and theoretical issues involved in the determination of its parameters (mass, production cross section, decay branching ratios, *etc.*); it also comments on prospects for future improvements.

B. Top quark production at the Tevatron: All direct measurements of top quark production and decay have been made by the CDF and DØ experiments at the Fermilab Tevatron collider in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. Here top quarks are produced dominantly in pairs from the QCD processes $q\bar{q} \rightarrow t\bar{t}$ and $gg \rightarrow t\bar{t}$. At this energy, the production cross section in these channels is expected to be approximately 5 pb for $m_t = 175$ GeV/ c^2 , with a 90% contribution from $q\bar{q}$ annihilation. Smaller contributions are expected from electroweak single-top production mechanisms, namely $q\bar{q}' \rightarrow W^* \rightarrow t\bar{b}$ and $qg \rightarrow q't\bar{b}$, the latter mediated by virtual- W exchange (“ W -gluon fusion”). The combined rate from these processes is approximately 2.5 pb at $m_t = 175$ GeV/ c^2 (see Ref. 1 and references therein). The expected contribution of these channels is further reduced relative to the dominant pair-production mechanisms because of larger backgrounds and poor detection efficiency.

With a mass above the Wb threshold, the decay width of the top quark is expected to be dominated by the two-body channel $t \rightarrow Wb$. Neglecting terms of order m_b^2/m_t^2 , α_s^2 and those of order $(\alpha_s/\pi)m_W^2/m_t^2$, this is predicted in the Standard Model to be [2]:

$$\Gamma_t = \frac{G_F m_t^3}{8\pi\sqrt{2}} \left(1 - \frac{M_W^2}{m_t^2}\right)^2 \left(1 + 2\frac{M_W^2}{m_t^2}\right) \left[1 - \frac{2\alpha_s}{3\pi} \left(\frac{2\pi^2}{3} - \frac{5}{2}\right)\right]. \quad (1)$$

The use of G_F in this equation accounts for the largest part of the one-loop electroweak radiative corrections, providing an expression accurate to better than 2%. The width increases with mass, going for example from 1.02 GeV/c² at $m_t = 160$ GeV/c² to 1.56 GeV/c² at $m_t = 180$ GeV/c² (we used $\alpha_s(M_Z) = 0.118$). With such a correspondingly short lifetime, the top quark is expected to decay before top-flavored hadrons or $t\bar{t}$ -quarkonium bound states can form [3]. Recently, the order α_s^2 QCD corrections to Γ_t have also been calculated [4], thereby improving the overall theoretical accuracy to better than 1%.

In top decay, the Ws and Wd final states are expected to be suppressed relative to Wb by the square of the CKM matrix elements V_{ts} and V_{td} , whose values can be estimated under the assumption of unitarity of the three-generation CKM matrix to be less than 0.043 and 0.014, respectively (see our review “The Cabibbo-Kobayashi-Maskawa Mixing Matrix” in the current edition for more information). Typical final states for the leading pair-production process therefore belong to three classes:

- A. $t\bar{t} \rightarrow W b W \bar{b} \rightarrow q \bar{q}' b q'' \bar{q}''' \bar{b}$,
- B. $t\bar{t} \rightarrow W b W \bar{b} \rightarrow q \bar{q}' b \ell \bar{\nu}_\ell \bar{b} + \bar{\ell} \nu_\ell b q \bar{q}' \bar{b}$,
- C. $t\bar{t} \rightarrow W b W \bar{b} \rightarrow \bar{\ell} \nu_\ell b \ell' \bar{\nu}_{\ell'} \bar{b}$,

where A, B, and C are referred to as the all-jets, lepton + jets, and dilepton channels, respectively.

The final state quarks can emit radiation and eventually evolve into jets of hadrons. The precise number of jets reconstructed by the detectors varies event by event, as it depends on the decay kinematics, as well as on the precise definition of jet used in the analysis. (Additional gluon radiation can also be emitted from the initial states.) The transverse momenta of the neutrinos are reconstructed via the large imbalance in detected transverse momentum of the event (missing E_T).

The observation of $t\bar{t}$ pairs has been reported in all of the above decay modes. As discussed below, the production and decay properties of the top quark extracted from the above three decay channels are all consistent with each other within experimental uncertainty. In particular, the $t \rightarrow Wb$ decay mode is supported through the reconstruction of the $W \rightarrow jj$ invariant mass in the $\ell\bar{\nu}_\ell b\bar{b}jj$ final state [5].

The extraction of top-quark properties from Tevatron data requires a good understanding of the production and decay mechanisms of the top, as well as of the large background processes. Because only leading order QCD calculations are available for most of the relevant processes ($W+3$ and 4 jets, or $WW+2$ jets), theoretical estimates of the backgrounds have large uncertainties. While this limitation affects estimates of the overall $t\bar{t}$ production rates, it is believed that the LO determination of the event kinematics and of the fraction of W + multi-jet events containing b quarks is relatively accurate. In particular, for the background one expects the E_T spectrum of jets to fall rather steeply, the jet direction to peak at small angles to the beams, and the fraction of events with b quarks to be of the order of a few percent. On the contrary, for the top signal, the b fraction is $\sim 100\%$ and the jets are rather

energetic, since they come from the decay of a massive object. It is therefore possible to improve the S/B ratio either by requiring the presence of a b quark, or by selecting very energetic and central kinematic configurations.

A detailed study of control samples with features similar to those of the relevant backgrounds, but free from possible top contamination, is required to provide a reliable check on background estimates.

C. Measured top properties: Current measurements of top properties are based on the full Run I integrated luminosity of 109 pb^{-1} for CDF and 125 pb^{-1} for DØ. DØ and CDF determine the $t\bar{t}$ cross section $\sigma_{t\bar{t}}$ from their number of observed top candidates, estimated background, $t\bar{t}$ acceptance, and integrated luminosity, assuming the Standard-Model decay $t \rightarrow Wb$ with unity branching ratio. Table 1 shows the measured cross sections from DØ and CDF along with the range of theoretical expectations, evaluated at the m_t values used by the experiments in calculating their acceptances. The DØ results have been updated in conference proceedings [7] to adjust to the current DØ value of the top mass. The CDF results have been updated in conference proceedings [16] to include improvements in their Monte Carlo determination of secondary-vertex tagging efficiency, calibration of the background estimate of the heavy-flavor fraction in inclusive W +jets events, and an updated total luminosity. This has brought the CDF cross section into better agreement with theoretical expectations. The agreement of both DØ and CDF $t\bar{t}$ cross sections with theory supports the hypothesis that the excess of events over background in all of these channels can be attributed to $t\bar{t}$ production.

Table 1: Cross section for $t\bar{t}$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV from DØ ($m_t = 172.1$ GeV/ c^2), CDF ($m_t = 175$ GeV/ c^2), and theory.

$\sigma_{t\bar{t}}(pb)$	Source	Ref.	Method
4.1 ± 2.1	DØ	[6,7]	ℓ + jets/topological
8.3 ± 3.5	DØ	[6,7]	ℓ + jets/soft μ b-tag
6.4 ± 3.3	DØ	[6,7]	$\ell\ell$ + $e\nu$
7.1 ± 3.2	DØ	[8]	all jets
5.9 ± 1.7	DØ	[8]	all combined
$5.2 - 6.0$	Theory	[9–12]	$m_t = 172.1$ GeV/ c^2
5.1 ± 1.5	CDF	[13,16]	ℓ + jets/vtx b-tag
9.2 ± 4.3	CDF	[13,16]	ℓ + jets/soft ℓ b-tag
$8.4^{+4.5}_{-3.5}$	CDF	[14,16]	$\ell\ell$
$7.6^{+3.5}_{-2.7}$	CDF	[15,16]	all jets
$6.5^{+1.7}_{-1.4}$	CDF	[16]	all combined
$4.75 - 5.5$	Theory	[9–12]	$m_t = 175$ GeV/ c^2

More precise measurements of the top production cross section will test current understanding of the production mechanisms [9–12]. This is important for the extrapolation to higher energies of colliders such as the LHC, where the larger expected cross section will permit more extensive studies [17]. Discrepancies in rate between theory and data, even at the Tevatron, would be quite exciting, and might indicate the presence of exotic production or decay channels, as predicted in certain models. Such new sources of top would lead to a modification of kinematic distributions such as the invariant mass of the top pair or the transverse momentum of the top quark. Studies by

CDF of the former [18] and of the latter [19] distributions, show no deviation from expected QCD behavior. DØ [20] also finds these kinematic distributions consistent with Standard Model expectations.

The top mass has been measured in the lepton + jets and dilepton channels by both DØ and CDF, and in the all-jets channel by CDF. At present, the most precise measurements come from the lepton + jets channel, with four or more jets and large missing E_T . In this channel, each event is subjected to a two-constraint kinematic fit to the hypothesis $t\bar{t} \rightarrow W^+ b W^- \bar{b} \rightarrow \ell \nu_\ell q \bar{q}' b \bar{b}$, assuming that the four highest E_T jets are the quarks from $t\bar{t}$ decay. The shape of the distribution of fitted top masses from these events is compared to templates expected from a mixture of background and signal distributions for a series of assumed top masses. This comparison yields values of the likelihood as a function of top mass, from which a best value of the top mass and its uncertainty can be obtained. The results are shown in Table 2. The systematic uncertainty (second uncertainty shown) is comparable to the statistical uncertainty, and is primarily due to uncertainties in the jet energy scale and in the Monte Carlo modeling.

Less precise determinations of the top mass come from the dilepton channel with two or more jets and large missing E_T , and from the all-jets channel. In the dilepton channel, a kinematically constrained fit is not possible because there are two missing neutrinos, so experiments must use other mass estimators than the reconstructed top mass. In principle, any quantity which is correlated with the top mass can be used as such an estimator. The DØ method uses the fact that if a value for m_t is assumed, the $t\bar{t}$ system can be reconstructed (up to a four-fold ambiguity). They compare the resulting kinematic

configurations to expectations from $t\bar{t}$ production, and obtain an m_t -dependent weight curve for each event, which they histogram in five bins to obtain four shape-sensitive quantities as their multidimensional mass estimator. This method yields a significant increase in precision over one-dimensional estimators. CDF has employed a similar method, thereby reducing their previous systematic uncertainty in the $\ell\ell + \text{jets}$ channel by a factor of two. DØ and CDF obtain the top mass and uncertainty from these mass estimators using the same type of template likelihood method as for the lepton + jets channel. CDF also measures the mass in the all-jets channel using events with six or more jets, at least one of which is tagged as a b jet through the detection of a secondary vertex.

Table 2: Top mass measurements from DØ and CDF.

m_t (GeV/c ²)	Source	Ref.	Method
$173.3 \pm 5.6 \pm 5.5$	DØ	[20]	$\ell + \text{jets}$
$168.4 \pm 12.3 \pm 3.6$	DØ	[21]	$\ell\ell$
$172.1 \pm 5.2 \pm 4.9$	DØ	[20]	DØ comb.
$175.9 \pm 4.8 \pm 5.3$	CDF	[22,23]	$\ell + \text{jet}$
$167.4 \pm 10.3 \pm 4.8$	CDF	[22]	$\ell\ell$
$186.0 \pm 10.0 \pm 5.7$	CDF	[22,15]	all jets
$176.0 \pm 4.0 \pm 5.1$	CDF	[22]	CDF comb.
$174.3 \pm 3.2 \pm 4.0$ *	DØ & CDF	[24]	PDG best

* PDG uses this Top Averaging Group result as its best value

As seen in Table 2, all results are in good agreement with a unique mass for the top quark, giving further support to the hypothesis that these events are due to $t\bar{t}$ production. The Top Averaging Group, a joint CDF/DØ working group, produced the combined CDF/DØ average top mass in Table 2, taking into account correlations between systematic uncertainties in different measurements. They assume that the uncertainty in jet energy scale is completely correlated within CDF and within DØ but uncorrelated between the two experiments, and that the signal model and Monte Carlo generator uncertainties are completely correlated between all measurements. The uncertainties from uranium noise and multiple interactions relate only to DØ and are assumed completely correlated between their two measurements. The uncertainty on the background model is taken to be completely correlated between the CDF and the DØ ℓ +jets measurements, and similarly for the $\ell\ell$ measurements. The Particle Data Group uses this combined top mass, $m_t = 174.3 \pm 5.1$ GeV/c² (statistical and systematic uncertainties combined in quadrature), as our PDG best value.

Given the experimental technique used to extract the top mass, these mass values should be taken as representing the top *pole mass* (see our review “Note on Quark Masses” in the current edition for more information).

With a smaller uncertainty on the top mass, and with improved measurements of other electroweak parameters, it will be possible to get important constraints on the value of the Higgs mass. Current global fits performed within the Standard Model and its minimal supersymmetric extension provide indications for a relatively light Higgs (see the review “ H^0 Indirect Mass Limits from Electroweak Analysis” in the Particle Listings of the current edition for more information).

Other properties of top decays are being studied. CDF reports a direct measurement of the $t \rightarrow Wb$ branching ratio [25]. Their preliminary result, obtained by comparing the number of events with 0, 1 and 2 tagged b jets and using the known b -tagging efficiency, is: $R = B(t \rightarrow Wb) / \sum_{q=d,s,b} B(t \rightarrow Wq) = 0.99 \pm 0.29$ where statistical and systematic uncertainties are included, or as a lower limit, $R > 0.58$ at 95% CL. Assuming that non- W decays of top can be neglected, that only three generations of fermions exist, and that the CKM matrix is unitary, they extract a CKM matrix-element $|V_{tb}| = 0.99 \pm 0.15$ or $|V_{tb}| > 0.76$ at 95% CL. A more direct measurement of the Wtb coupling constant will be possible when enough data are accumulated to detect the less frequent single-top production processes, such as $q\bar{q}' \rightarrow W^* \rightarrow t\bar{b}$ (a.k.a. s -channel W exchange) and $qb \rightarrow q't$ via W exchange (a.k.a. Wg fusion). The cross sections for these processes are proportional to $|V_{tb}|^2$, and there is no assumption needed on the number of families or the unitarity of the CKM matrix in the extraction of $|V_{tb}|$. Preliminary CDF results [19] give 95% CL limits of 15.8 and 15.4 pb for the single-top production rates in the s -channel and Wg -fusion channels, respectively. Comparison with the expected Standard Model rates of 0.73 ± 0.10 pb and 1.70 ± 0.30 pb, respectively, shows that far better statistics will be required before significant measurements can be achieved. For the prospects of these measurements at the LHC, see [17].

Both CDF and DØ have searched for non-Standard Model top decays [26,27], particularly those expected in supersymmetric models. These studies search for $t \rightarrow H^+b$, followed by $H^+ \rightarrow \tau\nu$ or $c\bar{s}$. The $t \rightarrow H^+b$ branching ratio is a minimum at $\tan\beta = \sqrt{m_t/m_b} \simeq 6$ and is large in the region of either $\tan\beta \ll 6$ or $\tan\beta \gg 6$. In the former range $H^+ \rightarrow c\bar{s}$ is the

dominant decay, while $H^+ \rightarrow \tau\nu$ dominates in the latter range. These studies are based either on direct searches for these final states, or on top disappearance. In the standard lepton + jets or dilepton cross section analyses, the charged Higgs decays are not detected as efficiently as $t \rightarrow W^\pm b$, primarily because the selection criteria are optimized for the standard decays, and because of the absence of energetic isolated leptons in the Higgs decays. With a significant $t \rightarrow H^+ b$ contribution, this would give rise to measured cross sections lower than the prediction from the Standard Model (assuming that non-Standard contributions to $t\bar{t}$ production are negligible). More details, and the results of these studies, can be found in the review “Search for Higgs bosons” and in the “ H^+ Mass Limits” section of the Higgs Particle Listings of the current edition.

CDF reports a search for flavor changing neutral current (FCNC) decays of the top quark $t \rightarrow q\gamma$ and $t \rightarrow qZ$ [28], for which the Standard Model predicts such small rates that their observation here would indicate new physics. They assume that one top decays via FCNC while the other decays via Wb . For the $t \rightarrow q\gamma$ search, they examine two signatures, depending on whether the W decays leptonically or hadronically. For leptonic W decay, the signature is $\gamma\ell$ and missing E_T and two or more jets, while for hadronic W decay, it is γ plus four or more jets, one with a secondary vertex b tag. They observe one event ($\mu\gamma$) with an expected background of less than half an event, giving an upper limit on the top branching ratio of $B(t \rightarrow q\gamma) < 3.2\%$ at 95% CL.

For the $t \rightarrow qZ$ FCNC search, they look for $Z \rightarrow \mu\mu$ or ee and $W \rightarrow$ hadrons, giving a $Z +$ four jets signature. They observe one $\mu\mu$ event with an expected background of 1.2 events, giving an upper limit on the top branching ratio of

$B(t \rightarrow qZ) < 33\%$ at 95% CL. Both the γ and Z limits are non-background subtracted (i.e. conservative) estimates.

Indirect constraints on FCNC couplings of the top quark can be obtained from single-top production in e^+e^- collisions, via the process $e^+e^- \rightarrow \gamma, Z^* \rightarrow t\bar{q}$ and its charge-conjugate ($q = u, c$). Limits on the cross-section for this reaction have been obtained by DELPHI [29] using LEP2 data at energies between 183 and 189 GeV. When interpreted in terms of top decay branching ratios [30,17], these limits lead to a bound of $B(t \rightarrow qZ) < 22\%$ at 95% CL, which is stronger than the direct CDF limit.

Studies of the decay angular distributions allow a direct analysis of the $V-A$ nature of the Wtb coupling, and provide information on the relative coupling of longitudinal and transverse W bosons to the top quark. In the Standard Model, the fraction of decays to longitudinally polarized W bosons is expected to be $\mathcal{F}_0^{\text{SM}} = x/(1+x)$, $x = m_t^2/2M_W^2$ ($\mathcal{F}_0^{\text{SM}} \sim 70\%$ for $m_t = 175 \text{ GeV}/c^2$). Deviations from this value would bring into question the validity of the Higgs mechanism of spontaneous symmetry breaking. CDF has recently measured $\mathcal{F}_0^{\text{SM}} = 0.91 \pm 0.37_{\text{stat}} \pm 0.13_{\text{syst}}$ [31], in agreement with the expectations.

$D\bar{O}$ has studied $t\bar{t}$ spin correlation [32]. Top quark pairs produced at the Tevatron are expected to be unpolarized but to have correlated spins. Since top quarks decay before hadronizing, their spins are transmitted to their decay daughters. Spin correlation is studied by analyzing the joint decay angular distribution of one t daughter and one \bar{t} daughter. The sensitivity to top spin is greatest when the daughters are charged leptons or d -type quarks, in which case, the joint distribution is

$$\frac{1}{\sigma} \frac{d^2\sigma}{d(\cos\theta_+)d(\cos\theta_-)} = \frac{1 + \kappa \cos\theta_+ \cos\theta_-}{4}, \quad (2)$$

where θ_+ and θ_- are the angles of the daughters in the top rest frames with respect to a particular quantization axis, the optimal off-diagonal basis [33]. In this basis, the Standard Model predicts maximum correlation with $\kappa = 0.88$ at the Tevatron. DØ analyzes their six dilepton events and obtains a likelihood as a function of κ which weakly favors the Standard Model ($\kappa = 0.88$) over no correlation ($\kappa = 0$) or anticorrelation ($\kappa = -1$, as would be expected for $t\bar{t}$ produced via an intermediate scalar). They quote a limit $\kappa > -0.25$ at 68% CL. With improved statistics, an observation of $t\bar{t}$ spin correlation could yield a lower limit on $|V_{tb}|$, independent of the assumption of three quark families [34].

References

1. T. Stelzer, Z. Sullivan, and S. Willenbrock, Phys. Rev. **D56**, 5919 (1997).
2. M. Jeżabek and J.H. Kühn, Nucl. Phys. **B314**, 1 (1989).
3. I.I.Y. Bigi, Yu.L. Dokshitzer, V. Khoze, J.H. Kühn, P. Zerwas, Phys. Lett. **B181**, 157 (1986).
4. A. Czarnecki, K. Melnikov, Nucl. Phys. **B544**, 520 (1999); K.G. Chetyrkin, R. Harlander, T. Seidensticker, M. Steinhauser, Phys. Rev. **D60**, 114015 (1999).
5. F. Abe *et al.*, CDF Collab., Phys. Rev. Lett. **80**, 5720 (1998).
6. S. Abachi *et al.*, DØ Collab., Phys. Rev. Lett. **79**, 1203 (1997).
7. R. Partridge, 29th Intl. Conf. on High Energy Physics (ICHEP 98), Vancouver, Canada, 23-29 July, 1998, hep-ex/9811035.
8. B. Abbott *et al.*, DØ Collab., Phys. Rev. Lett. **83**, 1908 (1999);
B. Abbott *et al.*, DØ Collab., Phys. Rev. **D60**, 012001 (1999).

9. P. Nason, S. Dawson, and R.K. Ellis, Nucl. Phys. **B303**, 607 (1988);
W. Beenakker, H. Kuijf, W.L. van Neerven, and J. Smith, Phys. Rev. **D40**, 54 (1989).
10. E. Berger and H. Contopanagos, Phys. Lett. **B361**, 115 (1995).
11. E. Laenen, J. Smith, and W. van Neerven, Phys. Lett. **B321**, 254 (1994).
12. S. Catani, M. Mangano, P. Nason, and L. Trentadue, Phys. Lett. **B378**, 329 (1996);
R. Bonciani, S. Catani, M.L. Mangano, P. Nason, Nucl. Phys. **B529**, 424 (1998);
M.L. Mangano, [hep-ph/9911256](#), to appear in Procs. of Intl. Europhysics Conf. on High Energy Physics (EPS-HEP 99), Tampere, Finland, 15-21 Jul 1999.
13. F. Abe *et al.*, CDF Collab., Phys. Rev. Lett. **80**, 2773 (1998).
14. F. Abe *et al.*, CDF Collab., Phys. Rev. Lett. **80**, 2779 (1998).
15. F. Abe *et al.*, CDF Collab., Phys. Rev. Lett. **79**, 1992 (1997).
16. F. Ptohos, Representing the CDF Collab., Procs. of Intl. Europhysics Conf. on High Energy Physics (EPS-HEP 99), Tampere, Finland, 15-21 July 1999, to be publ.
17. M. Beneke, I. Efthymiopoulos, M.L. Mangano, J. Womersley *et al.*, [hep-ph/0003033](#), to appear in Proceedings of 1999 CERN Workshop on Standard Model Physics (and more) at the LHC, G. Altarelli and M.L. Mangano eds.
18. T. Affolder *et al.*, CDF Collab., FERMILAB-PUB-00-051, [hep-ex/0003005](#).
19. P. Koehn, for the CDF Collab., FERMILAB-CONF-99-306-E, to be publ. in Procs. of Intl. Europhysics Conf. on High-Energy Physics (EPS-HEP 99), Tampere, Finland, 15-21 Jul 1999.
20. B. Abbott *et al.*, DØ Collab., Phys. Rev. **D58**, 052001 (1998);

- S. Abachi *et al.*, DØ Collab., Phys. Rev. Lett. **79**, 1197 (1997).
21. B. Abbott *et al.*, DØ Collab., Phys. Rev. **D60**, 052001 (1999);
B. Abbott *et al.*, DØ Collab., Phys. Rev. Lett. **80**, 2063 (1998).
22. F. Abe *et al.*, CDF Collab., Phys. Rev. Lett. **82**, 271 (1999).
23. F. Abe *et al.*, CDF Collab., Phys. Rev. Lett. **80**, 2767 (1998).
24. L. Demortier *et al.*, The Top Averaging Group, For the CDF and DØ Collaborations, FERMILAB-TM-2084, September, 1999.
25. G. Chiarelli, Int. J. Mod. Phys. **A13**, 2883 (1998).
26. F. Abe *et al.*, CDF Collab., Phys. Rev. Lett. **79**, 357 (1997);
B. Bevensee, for the CDF Collab., FERMILAB-CONF-98/155-E;
T. Affolder *et al.*, CDF Collab., hep-ex/9912013.
27. B. Abbott *et al.*, Phys. Rev. Lett. **82**, 4975 (1999).
28. F. Abe *et al.*, CDF Collab., Phys. Rev. Lett. **80**, 2525 (1998).
29. P. Abreu *et al.*, DELPHI Collab., Phys. Lett. **B446**, 62 (1998);
P. Abreu *et al.*, DELPHI Collab., DELPHI Note 99-85, submitted to the Intl. Europhysics Conference on High Energy Physics, Tampere, Finland, 15-21 July 1999.
30. V.F. Obraztsov, S.R. Slabospitsky, O.P. Yushchenko, Phys. Lett. **B426**, 393 (1998).
31. T. Affolder *et al.*, CDF Collab., Phys. Rev. Lett. **84**, 216 (2000).
32. B. Abbott *et al.*, DØ Collaboration, FERMILAB-Pub-00/046-E, submitted to Phys. Rev. Lett., hep-ex/0002058.
33. G. Mahlon and S. Parke, Phys. Rev. **D53**, 4886 (1996); G. Mahlon and S. Parke, Phys. Lett. **B411**, 173 (1997).

34. T. Stelzer and S. Willenbrock, Phys. Lett. **B374**, 169 (1996).***t*-Quark Mass in $p\bar{p}$ Collisions**

The t quark has been observed. Its mass is sufficiently high that decay is expected to occur before hadronization. OUR EVALUATION is an AVERAGE which incorporates correlations between systematic errors of the five different measurements. The average was done by a joint CDF/DØ working group and is reported in DEMORTIER 99, an FNAL Technical Memo. They report $174.3 \pm 3.2 \pm 4.0$ GeV, which yields "OUR EVALUATION" when statistical and systematic errors are combined.

For earlier search limits see the *Review of Particle Physics*, Phys. Rev. **D54**,1 (1996).

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
174.3 ± 5.1 OUR EVALUATION			
167.4 ± 10.3 ± 4.8	1 ABE	99B CDF	dilepton
168.4 ± 12.3 ± 3.6	2 ABBOTT	98D D0	dilepton
173.3 ± 5.6 ± 5.5	2 ABBOTT	98F D0	lepton + jets
175.9 ± 4.8 ± 5.3	1,3 ABE	98E CDF	lepton + jets
186 ± 10 ± 5.7	1,4 ABE	97R CDF	6 or more jets
• • • We do not use the following data for averages, fits, limits, etc. • • •			
172.1 ± 5.2 ± 4.9	5 ABBOTT	99G D0	di-lepton, lepton+jets
176.0 ± 6.5	6 ABE	99B CDF	dilepton, lepton+jets, and all jets
161 ± 17 ± 10	1 ABE	98F CDF	dilepton
172.1 ± 5.2 ± 4.9	7 BHAT	98B RVUE	dilepton and lepton+jets
173.8 ± 5.0	8 BHAT	98B RVUE	dilepton, lepton+jets, and all jets
173.3 ± 5.6 ± 6.2	2 ABACHI	97E D0	lepton + jets
199 $\begin{smallmatrix} +19 \\ -21 \end{smallmatrix}$ ± 22	ABACHI	95 D0	lepton + jets
176 ± 8 ± 10	ABE	95F CDF	lepton + b -jet
174 ± 10 $\begin{smallmatrix} +13 \\ -12 \end{smallmatrix}$	ABE	94E CDF	lepton + b -jet

¹ Result is based on $109 \pm 7 \text{ pb}^{-1}$ of data at $\sqrt{s} = 1.8$ TeV.

² Result is based on $125 \pm 7 \text{ pb}^{-1}$ of data at $\sqrt{s} = 1.8$ TeV.

³ The updated systematic error is listed. See ABE 99B.

⁴ ABE 97R result is based on the first observation of all hadronic decays of $t\bar{t}$ pairs. Single b -quark tagging with jet-shape variable constraints was used to select signal enriched multi-jet events. The updated systematic error is listed. See ABE 99B.

⁵ ABBOTT 99G result is obtained by combining the D0 result m_t (GeV) = $168.4 \pm 12.3 \pm 3.6$ from 6 di-lepton events (see also ABBOTT 98D) and m_t (GeV) = $173.3 \pm 5.6 \pm 5.5$ from lepton+jet events (ABBOTT 98F).

⁶ ABE 99B result is obtained by combining the CDF results of m_t (GeV)= $167.4 \pm 10.3 \pm 4.8$ from 8 dilepton events, m_t (GeV)= $175.9 \pm 4.8 \pm 5.3$ from lepton+jet events (ABE 98E), and m_t (GeV)= $186.0 \pm 10.0 \pm 5.7$ from all-jet events (ABE 97R). The systematic errors in the latter two measurements are changed in this paper.

⁷ BHAT 98B result is obtained by combining the DØ results of m_t (GeV)= $168.4 \pm 12.3 \pm 3.6$ from 6 dilepton events and m_t (GeV)= $173.3 \pm 5.6 \pm 5.5$ from 77 lepton+jet events.

⁸ BHAT 98B result is obtained by combining the DØ results from dilepton and lepton+jet events, and the CDF results (ABE 99B) from dilepton, lepton+jet events, and all-jet events.

Indirect t -Quark Mass from Standard Model Electroweak Fit

“OUR EVALUATION” below is from the fit to electroweak data described in the “Electroweak Model and Constraints on New Physics” section of this Review. This fit result does not include direct measurements of m_t .

The RVUE values are based on the data described in the footnotes. RVUE's published before 1994 and superseded analyses are now omitted. For more complete listings of earlier results, see the 1994 edition (Physical Review **D50** 1173 (1994)).

<u>VALUE (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
168.2^{+9.6}_{-7.4} OUR EVALUATION			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
171.2 ^{+3.7} _{-3.8}	⁹ FIELD	99 RVUE	Z parameters without b jet + Direct
172.0 ^{+5.8} _{-5.7}	¹⁰ DEBOER	97B RVUE	Electroweak + Direct
157 ⁺¹⁶ ₋₁₂	¹¹ ELLIS	96C RVUE	Z parameters, m_W , low energy
175 ± 11 ⁺¹⁷ ₋₁₉	¹² ERLER	95 RVUE	Z parameters, m_W , low energy
180 ± 9 ⁺¹⁹ ₋₂₁ ± 2.6 ± 4.8	¹³ MATSUMOTO	95 RVUE	
157 ⁺³⁶ ₋₄₈ ⁺¹⁹ ₋₂₀	¹⁴ ABREU	94 DLPH	Z parameters
158 ⁺³² ₋₄₀ ± 19	¹⁵ ACCIARRI	94 L3	Z parameters
132 ⁺⁴¹ ₋₄₈ ⁺²⁴ ₋₁₈	¹⁶ AKERS	94 OPAL	Z parameters
190 ⁺³⁹ ₋₄₈ ⁺¹² ₋₁₄	¹⁷ ARROYO	94 CCFR	ν_μ iron scattering
184 ⁺²⁵ ₋₂₉ ⁺¹⁷ ₋₁₈	¹⁸ BUSKULIC	94 ALEP	Z parameters
153 ± 15	¹⁹ ELLIS	94B RVUE	Electroweak
177 ± 9 ⁺¹⁶ ₋₂₀	²⁰ GURTU	94 RVUE	Electroweak
174 ⁺¹¹ ₋₁₃ ⁺¹⁷ ₋₁₈	²¹ MONTAGNA	94 RVUE	Electroweak
171 ± 12 ⁺¹⁵ ₋₂₁	²² NOVIKOV	94B RVUE	Electroweak
160 ⁺⁵⁰ ₋₆₀	²³ ALITTI	92B UA2	m_W , m_Z

⁹ FIELD 99 result is from the two-parameter fit with free m_t and m_H , yielding also $m_H = 47.2^{+29.8}_{-24.5}$ GeV. Only the lepton and charm-jet asymmetry data are used together with the direct measurement constraint $m_t = 173.8 \pm 5.0$ GeV, and $1/\alpha(m_Z) = 128.896$.

¹⁰ DEBOER 97B result is from the five-parameter fit which varies m_Z , m_t , m_H , α_s , and $\alpha(m_Z)$ under the constraints: $m_t = 175 \pm 6$ GeV, $1/\alpha(m_Z) = 128.896 \pm 0.09$. They found $m_H = 141^{+140}_{-77}$ GeV and $\alpha_s(m_Z) = 0.1197 \pm 0.0031$.

¹¹ ELLIS 96C result is a the two-parameter fit with free m_t and m_H , yielding also $m_H = 65^{+117}_{-37}$ GeV.

¹² ERLER 95 result is from fit with free m_t and $\alpha_s(m_Z)$, yielding $\alpha_s(m_Z) = 0.127(5)(2)$.

¹³ MATSUMOTO 95 result is from fit with free m_t to Z parameters, M_W , and low-energy neutral-current data. The second error is for $m_H = 300^{+700}_{-240}$ GeV, the third error is for $\alpha_s(m_Z) = 0.116 \pm 0.005$, the fourth error is for $\delta\alpha_{\text{had}} = 0.0283 \pm 0.0007$.

- ¹⁴ ABREU 94 value is for $\alpha_s(m_Z)$ constrained to 0.123 ± 0.005 . The second error corresponds to $m_H = 300^{+700}_{-240}$ GeV.
- ¹⁵ ACCIARRI 94 value is for $\alpha_s(m_Z)$ constrained to 0.124 ± 0.006 . The second error corresponds to $m_H = 300^{+700}_{-240}$ GeV.
- ¹⁶ AKERS 94 result is from fit with free α_s . The second error corresponds to $m_H = 300^{+700}_{-240}$ GeV. The 95%CL limit is $m_t < 210$ GeV.
- ¹⁷ ARROYO 94 measures the ratio of the neutral-current and charged-current deep inelastic scattering of ν_μ on an iron target. By assuming the SM electroweak correction, they obtain $1 - m_W^2/m_Z^2 = 0.2218 \pm 0.0059$, yielding the quoted m_t value. The second error corresponds to $m_H = 300^{+700}_{-240}$ GeV.
- ¹⁸ BUSKULIC 94 result is from fit with free α_s . The second error is from $m_H = 300^{+700}_{-240}$ GeV.
- ¹⁹ ELLIS 94B result is fit to electroweak data available in spring 1994, including the 1994 A_{LR} data from SLD. m_t and m_H are two free parameters of the fit for $\alpha_s(m_Z) = 0.118 \pm 0.007$ yielding m_t above, and $m_H = 35^{+70}_{-22}$ GeV. ELLIS 94B also give results for fits including constraints from CDF's direct measurement of m_t and CDF's and DØ 's production cross-section measurements. Fits excluding the A_{LR} data from SLD are also given.
- ²⁰ GURTU 94 result is from fit with free m_t and $\alpha_s(m_Z)$, yielding m_t above and $\alpha_s(m_Z) = 0.125 \pm 0.005^{+0.003}_{-0.001}$. The second errors correspond to $m_H = 300^{+700}_{-240}$ GeV. Uses LEP, M_W , νN , and SLD electroweak data available in spring 1994.
- ²¹ MONTAGNA 94 result is from fit with free m_t and $\alpha_s(m_Z)$, yielding m_t above and $\alpha_s(m_Z) = 0.124$. The second errors correspond to $m_H = 300^{+700}_{-240}$ GeV. Errors in $\alpha(m_Z)$ and m_b are taken into account in the fit. Uses LEP, SLC, and M_W/M_Z data available in spring 1994.
- ²² NOVIKOV 94B result is from fit with free m_t and $\alpha_s(m_Z)$, yielding m_t above and $\alpha_s(m_Z) = 0.125 \pm 0.005 \pm 0.002$. The second errors correspond to $m_H = 300^{+700}_{-240}$ GeV. Uses LEP and CDF electroweak data available in spring 1994.
- ²³ ALITTI 92B assume $m_H = 100$ GeV. The 95%CL limit is $m_t < 250$ GeV for $m_H < 1$ TeV.

t DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 Wb		
Γ_2 $\ell\nu_\ell$ anything	[a,b] (9.4±2.4) %	
Γ_3 $\tau\nu_\tau b$		
Γ_4 $\gamma q(q=u,c)$	[c] < 3.2 %	95%
$\Delta T = 1$ weak neutral current (T1) modes		
Γ_5 $Z q(q=u,c)$	T1 [d] < 33 %	95%

[a] ℓ means e or μ decay mode, not the sum over them.

[b] Assumes lepton universality and W -decay acceptance.

[c] This limit is for $\Gamma(t \rightarrow \gamma q)/\Gamma(t \rightarrow Wb)$.

[d] This limit is for $\Gamma(t \rightarrow Z q)/\Gamma(t \rightarrow Wb)$.

t BRANCHING RATIOS

$\Gamma(\ell\nu_\ell \text{ anything})/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN
0.094 ± 0.024	²⁴ ABE	98X CDF

²⁴ ℓ means e or μ decay mode, not the sum. Assumes lepton universality and W -decay acceptance.

$\Gamma(\tau\nu_\tau b)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

²⁵ ABE 97V CDF $\ell\tau + \text{jets}$

²⁵ ABE 97V searched for $t\bar{t} \rightarrow (\ell\nu_\ell)(\tau\nu_\tau)b\bar{b}$ events in 109 pb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.8 \text{ TeV}$. They observed 4 candidate events where one expects ~ 1 signal and ~ 2 background events. Three of the four observed events have jets identified as b candidates.

$\Gamma(\gamma q(q=u,c))/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	CL%	DOCUMENT ID	TECN
<0.032	95	²⁶ ABE	98G CDF

²⁶ ABE 98G looked for $t\bar{t}$ events where one t decays into $q\gamma$ while the other decays into bW . The quoted bound is for $\Gamma(\gamma q)/\Gamma(Wb)$.

$\Gamma(Z q(q=u,c))/\Gamma_{\text{total}}$ Γ_5/Γ

Test for $\Delta T=1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE	CL%	DOCUMENT ID	TECN
<0.33	95	²⁷ ABE	98G CDF

²⁷ ABE 98G looked for $t\bar{t}$ events where one t decays into three jets and the other decays into qZ with $Z \rightarrow \ell\ell$. The quoted bound is for $\Gamma(Zq)/\Gamma(Wb)$.

t-Quark REFERENCES

ABBOTT	99G	PR D60 052001	B. Abbott <i>et al.</i>	(D0 Collab.)
ABE	99B	PRL 82 271	F. Abe <i>et al.</i>	(CDF Collab.)
Also	99G	PRL 82 2808 (erratum)	F. Abe <i>et al.</i>	(CDF Collab.)
DEMORTIER	99	FNAL-TM-2084	L. Demortier <i>et al.</i>	(CDF/D0 Working Group)
FIELD	99	MPL A14 1815	J.H. Field	
ABBOTT	98D	PRL 80 2063	B. Abbott <i>et al.</i>	(D0 Collab.)
ABBOTT	98F	PR D58 052001	B. Abbott <i>et al.</i>	(D0 Collab.)
ABE	98E	PRL 80 2767	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	98F	PRL 80 2779	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	98G	PRL 80 2525	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	98X	PRL 80 2773	F. Abe <i>et al.</i>	(CDF Collab.)
BHAT	98B	IJMP A13 5113	P.C. Bhat, H.B. Prosper, S.S. Snyder	
ABACHI	97E	PRL 79 1197	S. Abachi <i>et al.</i>	(D0 Collab.)
ABE	97R	PRL 79 1992	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	97V	PRL 79 3585	F. Abe <i>et al.</i>	(CDF Collab.)
DEBOER	97B	ZPHY C75 627	W. de Boer <i>et al.</i>	
ELLIS	96C	PL B389 321	J. Ellis, G.L. Fogli, E. Lisi	(CERN, BARI)

ABACHI	95	PRL 74 2632	S. Abachi <i>et al.</i>	(D0 Collab.)
ABE	95F	PRL 74 2626	F. Abe <i>et al.</i>	(CDF Collab.)
ERLER	95	PR D52 441	J. Erler, P. Langacker	(PENN)
MATSUMOTO	95	MPL A10 2553	S. Matsumoto	(KEK)
ABE	94E	PR D50 2966	F. Abe <i>et al.</i>	(CDF Collab.)
Also	94F	PRL 73 225	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	94	NP B418 403	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	94	ZPHY C62 551	M. Acciarri <i>et al.</i>	(L3 Collab.)
AKERS	94	ZPHY C61 19	R. Akers <i>et al.</i>	(OPAL Collab.)
ARROYO	94	PRL 72 3452	C.G. Arroyo <i>et al.</i>	(COLU, CHIC, FNAL+)
BUSKULIC	94	ZPHY C62 539	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ELLIS	94B	PL B333 118	J. Ellis, G.L. Fogli, E. Lisi	(CERN, BARI)
GURTU	94	MPL A9 3301	A. Gurtu	(TATA)
MONTAGNA	94	PL B335 484	G. Montagna <i>et al.</i>	(INFN, PAVI, CERN+)
NOVIKOV	94B	MPL A9 2641	V.A. Novikov <i>et al.</i>	(GUEL, CERN, ITEP)
PDG	94	PR D50 1173	L. Montanet <i>et al.</i>	(CERN, LBL, BOST+)
ALITTI	92B	PL B276 354	J. Alitti <i>et al.</i>	(UA2 Collab.)
