

$\eta_c(2S)$

$$J^{PC} = 0^+(0^-+)$$

OMITTED FROM SUMMARY TABLE

Needs confirmation. Quantum numbers are quark model predictions.

$\eta_c(2S)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3654 ± 6 ± 8	39 ± 11	CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3622 ± 12	42	¹ ABE,K	02 BELL	$10.6 e^+ e^- \rightarrow J/\psi +$
3594 ± 5		² EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$
¹ From a fit of the J/ψ recoil mass spectrum. Systematic errors not estimated.				
² Assuming mass of $\psi(2S) = 3686$ MeV.				

$\eta_c(2S)$ WIDTH

<u>VALUE (MeV)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 55	90	39 ± 11	³ CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
< 8.0	95		⁴ EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$
³ For a mass value of 3654 ± 6 MeV					
⁴ For a mass value of 3594 ± 5 MeV					

$\eta_c(2S)$ DECAY MODES

Mode
Γ_1 hadrons
Γ_2 $K \bar{K} \pi$
Γ_3 $p \bar{p}$
Γ_4 $\gamma \gamma$

$\eta_c(2S)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

<u>VALUE (units 10^{-8})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_3 \Gamma_4 / \Gamma^2$
< 5.6	90	^{5,6,7} AMBROGIANI	01 E835	$\bar{p} p \rightarrow \gamma \gamma$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 8.0	90	^{5,6,8} AMBROGIANI	01 E835	$\bar{p} p \rightarrow \gamma \gamma$	
< 12.0	90	^{6,8} AMBROGIANI	01 E835	$\bar{p} p \rightarrow \gamma \gamma$	
⁵ Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.					
⁶ For a total width $\Gamma=5$ MeV.					
⁷ For the resonance mass region 3589–3599 MeV/ c^2 .					
⁸ For the resonance mass region 3575–3660 MeV/ c^2 .					

$\eta_c(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ Γ_1/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
not seen	ABREU	980 DLPH	$e^+e^- \rightarrow e^+e^- + \text{hadrons}$
seen	⁹ EDWARDS	82C CBAL	$e^+e^- \rightarrow \gamma X$

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_2/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	39 ± 11	¹⁰ CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.01	90	LEE	85 CBAL	$\psi' \rightarrow \text{photons}$
⁹ For a mass value of 3594 ± 5 MeV				
¹⁰ For a mass value of 3654 ± 6 MeV				

$\eta_c(2S)$ REFERENCES

ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU	980	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)

OTHER RELATED PAPERS

BADALIAN	03	PR D67 071901	A.M. Badalian, B.L.G. Bakker	
EICHTEN	02	PRL 89 162002	E.J. Eichten, K. Lane, C. Quigg	
ACCIARRI	99T	PL B461 155	M. Acciarri <i>et al.</i>	(L3 Collab.)
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
PORTER	81	SLAC Summer Inst. 355	F.C. Porter <i>et al.</i>	(CIT, HARV, PRIN+)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)