# *ρ*(2150)

$$I^{G}(J^{PC}) = 1^{+}(1^{--})$$

### OMITTED FROM SUMMARY TABLE This entry was previously called $T_1(2190)$ .

# ρ(2150) MASS

$e^+e^- \rightarrow \pi^+\pi^-$ , $K^+K^-$ , $6\pi$							
VALUE (MeV)	DOCUMENT IL	)	TECN	CHG	COMMENT		
2149 $\pm$ 17 OUR AVERAGE	Includes data from th	e data	block that	at follo	ows this one.		
2153±37	BIAGINI	91	RVUE		$e^+e^- \rightarrow$		
2110±50	<sup>2</sup> CLEGG	90	RVUE	0	$ \begin{array}{c} \pi^{+}\pi^{-}, \\ K^{+}K^{-} \\ e^{+}e^{-} \rightarrow \\ 3(\pi^{+}\pi^{-}), \\ 2(\pi^{+}\pi^{-}\pi^{0}) \end{array} $		

VALUE (MeV)	DOCUMENT ID		TECN	COMMENT
ullet $ullet$ $ullet$ We do not use the following	data for averages	, fits	, limits,	etc. • • •
$\sim$ 2191	HASAN	94	RVUE	$\overline{p}p \rightarrow \pi\pi$
$\sim$ 1988	HASAN	94	RVUE	$\overline{p}p \rightarrow \pi\pi$
$\sim$ 2070	<sup>1</sup> OAKDEN	94	RVUE	0.36–1.55 $\overline{p}p \rightarrow \pi\pi$
$\sim$ 2170	<sup>3</sup> MARTIN	<b>80</b> B	RVUE	
$\sim$ 2100	<sup>3</sup> MARTIN	80C	RVUE	

<sup>1</sup>See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to J = 3 to be important but not significantly resonant.

# S-CHANNEL $\overline{N}N$

 $\overline{p}p \rightarrow \pi\pi$ 

VALUE (MeV)	DOCUMENT ID		TECN	CHG	COMMENT
$\bullet \bullet \bullet$ We do not use the follow	ving data for averages	s, fits	, limits,	etc. •	• •
$2110\pm35$	<sup>4</sup> ANISOVICH	02	SPEC		$\begin{array}{ccc} 0.61.9 \ p \overline{p} \rightarrow \\ & \omega  \pi^0, \ \omega  \eta  \pi^0, \end{array}$
$\sim$ 2190	<sup>5</sup> CUTTS	<b>78</b> B	CNTR		$ \begin{array}{c} \pi^{+}\pi^{-} \\ 0.97-3 \overline{p}p \rightarrow \\ \overline{N}N \end{array} $
$2155 \pm 15$	<sup>5,6</sup> COUPLAND	77	CNTR	0	$0.7-2.4 \ \overline{p}p \rightarrow \overline{p}p$
$2193\pm~2$	<sup>5,7</sup> ALSPECTOR	73	CNTR		p S channel
$2190 \pm 10$	<sup>8</sup> ABRAMS	70	CNTR		S channel <del>p</del> N
0					

## $\pi^- p \rightarrow \omega \pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
The data in this block is included in	the average printed f	or a prev	vious datablock.

#### 2155 $\pm$ 21 OUR AVERAGE

$2140 \pm 30$	ALDE	95 GAM2	$38 \pi^- p \rightarrow \omega \pi^0 n$
$2170\!\pm\!30$	ALDE	92c GAM4	$100 \ \pi^- p \rightarrow \omega \pi^0 n$

HTTP://PDG.LBL.GOV

<sup>2</sup> Includes ATKINSON 85. <sup>3</sup>  $I(J^P) = 1(1^-)$  from simultaneous analysis of  $p\overline{p} \rightarrow \pi^- \pi^+$  and  $\pi^0 \pi^0$ . <sup>4</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.  $^{5}$  Isospins 0 and 1 not separated. <sup>6</sup> From a fit to the total elastic cross section. <sup>7</sup> Referred to as T or T region by ALSPECTOR 73.

<sup>8</sup>Seen as bump in I = 1 state. See also COOPER 68. PEASLEE 75 confirm  $\overline{p}p$  results of ABRAMS 70, no narrow structure.

# ρ(2150) WIDTH

$e^+e^- \rightarrow \pi^+\pi^-, K^+K^-, 6\pi$						
VALUE (MeV)	DOCUMENT ID		TECN	CHG	COMMENT	
$363 \pm 50$ OUR AVERAGE	Includes data from the	datal	block that	at follo	ows this one.	
389± 79 410±100	BIAGINI <sup>10</sup> CLEGG	91 90	RVUE RVUE	0	$e^{+}e^{-} \rightarrow \\\pi^{+}\pi^{-}, \\e^{+}e^{-} \rightarrow \\3(\pi^{+}\pi^{-}),$	
					$2(\pi^{+}\pi^{-}\pi^{0})$	

 $\overline{p}p \rightarrow \pi\pi$ 

VALUE (MeV)	DOCUMENT ID	TECN COMMENT
$\bullet$ $\bullet$ We do not use the follow	ing data for average	es, fits, limits, etc. • • •
$\sim 296$	HASAN	94 RVUE $\overline{p}p \rightarrow \pi\pi$
$\sim 244$	HASAN	94 RVUE $\overline{p}p \rightarrow \pi\pi$
$\sim$ 40	<sup>9</sup> OAKDEN	94 RVUE 0.36–1.55 $\overline{p}p \rightarrow \pi\pi$
$\sim 250$	<sup>11</sup> MARTIN	80B RVUE
$\sim 200$	$^{11}$ MARTIN	80C RVUE

<sup>9</sup>See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to J=3 to be important but not significantly resonant.

# S-CHANNEL NN

VALUE (MeV)	DOCUMENT ID		TECN	CHG	COMMENT
$\bullet \bullet \bullet$ We do not use the fo	ollowing data for averages	, fits	, limits, e	tc. •	••
230±50	<sup>12</sup> ANISOVICH	02	SPEC		$0.6-1.9 \ p\overline{p} \rightarrow \\ \omega \pi^{0}, \ \omega \eta \pi^{0}, \\ \pi^{+} \pi^{-}$
$135 \pm 75 \\ 98 \pm 8 \\ \sim 85$	<sup>13,14</sup> COUPLAND <sup>14</sup> ALSPECTOR <sup>15</sup> ABRAMS	77 73 70	CNTR ( CNTR CNTR	C	$\begin{array}{l} 0.7 - 2.4 \ \overline{p}p \rightarrow \overline{p}p \\ \overline{p}p \ S \ \text{channel} \\ S \ \text{channel} \ \overline{p}N \end{array}$

# $\pi^- p \rightarrow \omega \pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
The data in this block is included in	the average printed	for a pre	vious datablock.

95 GAM2 38  $\pi^- p \rightarrow \omega \pi^0 n$ ALDE  $320\pm70$ • • • We do not use the following data for averages, fits, limits, etc. • • • 92C GAM4 100  $\pi^- p \rightarrow \omega \pi^0 n$ ALDE  $\sim 300$ 

HTTP://PDG.LBL.GOV Page 2

Citation: S. Eidelman et al. (Particle Data Group), Phys. Lett. B 592, 1 (2004) and 2005 partial update for edition 2006 (URL: http://pdg.lbl.gov)

<sup>10</sup> Includes ATKINSON 85. <sup>11</sup>  $I(J^P) = 1(1^-)$  from simultaneous analysis of  $p\overline{p} \rightarrow \pi^- \pi^+$  and  $\pi^0 \pi^0$ . <sup>12</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02. <sup>13</sup> From a fit to the total elastic cross section. <sup>14</sup> Isospins 0 and 1 not separated.

NP B96 109

PL 29B 451

PRL 18 1209

- <sup>15</sup>Seen as bump in I = 1 state. See also COOPER 68. PEASLEE 75 confirm  $\overline{p}p$  results of ABRAMS 70, no narrow structure.

# $\rho(2150)$ REFERENCES

ANISOVICH	02	PL B542 8	A.V. Anisovich et al.	
ANISOVICH	01D	PL B508 6	A.V. Anisovich et al.	
ANISOVICH	01E	PL B513 281	A.V. Anisovich et al.	
ANISOVICH	00 J	PL B491 47	A.V. Anisovich et al.	
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
ALDE	95	ZPHY C66 379	D.M. Alde <i>et al.</i>	(GAMS Collab.) JP
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Penningt	con (DURH)
ALDE	92C	ZPHY C54 553	D.M. Alde <i>et al.</i> (	BELG, SERP, KEK, LANL+)
BIAGINI	91	NC 104A 363	M.E. Biagini <i>et al.</i>	(FRAS, PRAG)
CLEGG	90	ZPHY C45 677	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ATKINSON	85	ZPHY C29 333	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Penningto	n (DURH) JP
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
PEASLEE	75	PL 57B 189	D.C. Peaslee <i>et al.</i>	(CANB, BARI, BROW+)
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)
COOPER	68	PRL 20 1059	W.A. Cooper et al.	(ANL)
		OTHER	RELATED PAPERS	
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)

E. Eisenhandler et al.

C. Bricman et al.

R.J. Abrams et al.

EISENHAND... 75

69

67C

BRICMAN

ABRAMS

(LOQM, LIVP, DARE+) (CERN, CAEN, SACL)

(BNL)