

**$K_3^*(1780)$** 

$$I(J^P) = \frac{1}{2}(3^-)$$

 **$K_3^*(1780)$  MASS**

| <u>VALUE (MeV)</u>  | <u>EVTS</u> | <u>DOCUMENT ID</u>                  | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u>                                      |
|---|-------------|-------------------------------------|-------------|------------|---|
| <b>1776 ± 7 OUR AVERAGE</b>   |             | Error includes scale factor of 1.1. |             |            |   |
| 1781 ± 8 ± 4  |             | <sup>1</sup> ASTON                  | 88 LASS     | 0          | 11 $K^- p \rightarrow K^- \pi^+ n$                  |
| 1740 ± 14 ± 15  |             | <sup>1</sup> ASTON                  | 87 LASS     | 0          | 11 $K^- p \rightarrow$<br>$\bar{K}^0 \pi^+ \pi^- n$ |
| 1779 ± 11   |             | <sup>2</sup> BALDI                  | 76 SPEC     | +          | 10 $K^+ p \rightarrow K^0 \pi^+ p$                  |
| 1776 ± 26   |             | <sup>3</sup> BRANDENB...            | 76D ASPK    | 0          | 13 $K^\pm p \rightarrow K^\pm \pi^\mp N$            |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |                                     |             |            |   |
| 1720 ± 10 ± 15  | 6111        | <sup>4</sup> BIRD                   | 89 LASS     | -          | 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$            |
| 1749 ± 10   |             | ASTON                               | 88B LASS    | -          | 11 $K^- p \rightarrow K^- \eta p$                   |
| 1780 ± 9  | 300         | BAUBILLIER                          | 84B HBC     | -          | 8.25 $K^- p \rightarrow$<br>$\bar{K}^0 \pi^- p$     |
| 1790 ± 15   |             | BAUBILLIER                          | 82B HBC     | 0          | 8.25 $K^- p \rightarrow$<br>$K_S^0 2\pi N$          |
| 1784 ± 9  | 2060        | CLELAND                             | 82 SPEC     | ±          | 50 $K^+ p \rightarrow K_S^0 \pi^\pm p$              |
| 1786 ± 15   |             | <sup>5</sup> ASTON                  | 81D LASS    | 0          | 11 $K^- p \rightarrow K^- \pi^+ n$                  |
| 1762 ± 9  | 190         | TOAFF                               | 81 HBC      | -          | 6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$           |
| 1850 ± 50   |             | ETKIN                               | 80 MPS      | 0          | 6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^-$         |
| 1812 ± 28   |             | BEUSCH                              | 78 OMEG     |            | 10 $K^- p \rightarrow$<br>$\bar{K}^0 \pi^+ \pi^- n$ |
| 1786 ± 8  |             | CHUNG                               | 78 MPS      | 0          | 6 $K^- p \rightarrow K^- \pi^+ n$                   |

<sup>1</sup> From energy-independent partial-wave analysis.<sup>2</sup> From a fit to  $Y_6^2$  moment.  $J^P = 3^-$  found.<sup>3</sup> Confirmed by phase shift analysis of ESTABROOKS 78, yields  $J^P = 3^-$ .<sup>4</sup> From a partial wave amplitude analysis.<sup>5</sup> From a fit to the  $Y_6^0$  moment. **$K_3^*(1780)$  WIDTH**

| <u>VALUE (MeV)</u>  | <u>EVTS</u> | <u>DOCUMENT ID</u>  | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u>                                      |
|---|-------------|---|-------------|------------|---|
| <b>159 ± 21 OUR AVERAGE</b>   |             | Error includes scale factor of 1.3. See the ideogram below. |             |            |   |
| 203 ± 30 ± 8  |             | <sup>6</sup> ASTON  | 88 LASS     | 0          | 11 $K^- p \rightarrow K^- \pi^+ n$                  |
| 171 ± 42 ± 20   |             | <sup>6</sup> ASTON  | 87 LASS     | 0          | 11 $K^- p \rightarrow$<br>$\bar{K}^0 \pi^+ \pi^- n$ |
| 135 ± 22  |             | <sup>7</sup> BALDI  | 76 SPEC     | +          | 10 $K^+ p \rightarrow K^0 \pi^+ p$                  |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |   |             |            |   |
| 187 ± 31 ± 20   | 6111        | <sup>8</sup> BIRD   | 89 LASS     | -          | 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$            |
| 193 <sup>+51</sup> <sub>-37</sub>   |             | ASTON   | 88B LASS    | -          | 11 $K^- p \rightarrow K^- \eta p$                   |
| 99 ± 30   | 300         | BAUBILLIER  | 84B HBC     | -          | 8.25 $K^- p \rightarrow$<br>$\bar{K}^0 \pi^- p$     |
| ~ 130   |             | BAUBILLIER  | 82B HBC     | 0          | 8.25 $K^- p \rightarrow$<br>$K_S^0 2\pi N$          |

|        |      |                           |     |      |   |     |   |
|--------|------|---------------------------|-----|------|---|-----|---|
| 191±24 | 2060 | CLELAND                   | 82  | SPEC | ± | 50  | $K^+ p \rightarrow K_S^0 \pi^\pm p$         |
| 225±60 |      | <sup>9</sup> ASTON        | 81D | LASS | 0 | 11  | $K^- p \rightarrow K^- \pi^+ n$             |
| ~ 80   | 190  | TOAFF                     | 81  | HBC  | - | 6.5 | $K^- p \rightarrow \bar{K}^0 \pi^- p$       |
| 240±50 |      | ETKIN                     | 80  | MPS  | 0 | 6   | $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^-$   |
| 181±44 |      | <sup>10</sup> BEUSCH      | 78  | OMEG |   | 10  | $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$ |
| 96±31  |      | CHUNG                     | 78  | MPS  | 0 | 6   | $K^- p \rightarrow K^- \pi^+ n$             |
| 270±70 |      | <sup>11</sup> BRANDENB... | 76D | ASPK | 0 | 13  | $K^\pm p \rightarrow K^\pm \pi^\mp N$       |

<sup>6</sup> From energy-independent partial-wave analysis.


<sup>7</sup> From a fit to  $Y_6^2$  moment.  $J^P = 3^-$  found.

<sup>8</sup> From a partial wave amplitude analysis.

<sup>9</sup> From a fit to  $Y_6^0$  moment.

<sup>10</sup> Errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

<sup>11</sup> ESTABROOKS 78 find that BRANDENBURG 76D data are consistent with 175 MeV width. Not averaged.



### $K_3^*(1780)$ DECAY MODES

| Mode                        | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level |
|-----------------------------|--------------------------------|------------------|
| $\Gamma_1$ $K\rho$          | (31 ± 9) %                     |                  |
| $\Gamma_2$ $K^*(892)\pi$    | (20 ± 5) %                     |                  |
| $\Gamma_3$ $K\pi$           | (18.8 ± 1.0) %                 |                  |
| $\Gamma_4$ $K\eta$          | (30 ± 13) %                    |                  |
| $\Gamma_5$ $K_2^*(1430)\pi$ | < 16 %                         | 95%              |

**CONSTRAINED FIT INFORMATION**

An overall fit to 3 branching ratios uses 4 measurements and one constraint to determine 4 parameters. The overall fit has a  $\chi^2 = 0.0$  for 1 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

|       |       |       |       |
|-------|-------|-------|-------|
| $x_2$ | 85    |       |       |
| $x_3$ | 18    | 21    |       |
| $x_4$ | -98   | -94   | -27   |
|       | $x_1$ | $x_2$ | $x_3$ |

 **$K_3^*(1780)$  BRANCHING RATIOS** **$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$   $\Gamma_1/\Gamma_2$** 

| VALUE                    | DOCUMENT ID | TECN | CHG  | COMMENT |  |
|--------------------------|-------------|------|------|---------|--|
| <b>1.52±0.23 OUR FIT</b> |             |      |      |         |  |
| <b>1.52±0.21±0.10</b>    | ASTON       | 87   | LASS | 0       | 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$ |

 **$\Gamma(K^*(892)\pi)/\Gamma(K\pi)$   $\Gamma_2/\Gamma_3$** 

| VALUE                    | DOCUMENT ID | TECN | CHG  | COMMENT |   |
|--------------------------|-------------|------|------|---------|---|
| <b>1.09±0.26 OUR FIT</b> |             |      |      |         |   |
| <b>1.09±0.26</b>         | ASTON       | 84B  | LASS | 0       | 11 $K^- p \rightarrow \bar{K}^0 2\pi n$ |

 **$\Gamma(K\pi)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$** 

| VALUE                          | DOCUMENT ID | TECN | CHG  | COMMENT |                                    |
|--------------------------------|-------------|------|------|---------|------------------------------------|
| <b>0.188±0.010 OUR FIT</b>     |             |      |      |         |                                    |
| <b>0.188±0.010 OUR AVERAGE</b> |             |      |      |         |                                    |
| 0.187±0.008±0.008              | ASTON       | 88   | LASS | 0       | 11 $K^- p \rightarrow K^- \pi^+ n$ |
| 0.19 ±0.02                     | ESTABROOKS  | 78   | ASPK | 0       | 13 $K^\pm p \rightarrow K\pi N$    |

 **$\Gamma(K\eta)/\Gamma(K\pi)$   $\Gamma_4/\Gamma_3$** 

| VALUE                   | DOCUMENT ID | TECN | CHG | COMMENT |
|-------------------------|-------------|------|-----|---------|
| <b>1.6 ±0.7 OUR FIT</b> |             |      |     |         |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|            |                    |     |      |   |  |
|------------|--------------------|-----|------|---|--|
| 0.41±0.050 | <sup>12</sup> BIRD | 89  | LASS | - | 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$ |
| 0.50±0.18  | ASTON              | 88B | LASS | - | 11 $K^- p \rightarrow K^- \eta p$        |

<sup>12</sup> This result supersedes ASTON 88B.

 **$\Gamma(K_2^*(1430)\pi)/\Gamma(K^*(892)\pi)$   $\Gamma_5/\Gamma_2$** 

| VALUE           | CL% | DOCUMENT ID | TECN | CHG  | COMMENT |  |
|-----------------|-----|-------------|------|------|---------|--|
| <b>&lt;0.78</b> | 95  | ASTON       | 87   | LASS | 0       | 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$ |

### $K_3^*(1780)$ REFERENCES

|             |     |             |                                |                             |
|-------------|-----|-------------|--------------------------------|-----------------------------|
| BIRD        | 89  | SLAC-332    | P.F. Bird                      | (SLAC)                      |
| ASTON       | 88  | NP B296 493 | D. Aston <i>et al.</i>         | (SLAC, NAGO, CINC, INUS)    |
| ASTON       | 88B | PL B201 169 | D. Aston <i>et al.</i>         | (SLAC, NAGO, CINC, INUS) JP |
| ASTON       | 87  | NP B292 693 | D. Aston <i>et al.</i>         | (SLAC, NAGO, CINC, INUS)    |
| ASTON       | 84B | NP B247 261 | D. Aston <i>et al.</i>         | (SLAC, CARL, OTTA)          |
| BAUBILLIER  | 84B | ZPHY C26 37 | M. Baubillier <i>et al.</i>    | (BIRM, CERN, GLAS+)         |
| BAUBILLIER  | 82B | NP B202 21  | M. Baubillier <i>et al.</i>    | (BIRM, CERN, GLAS+)         |
| CLELAND     | 82  | NP B208 189 | W.E. Cleland <i>et al.</i>     | (DURH, GEVA, LAUS+)         |
| ASTON       | 81D | PL 99B 502  | D. Aston <i>et al.</i>         | (SLAC, CARL, OTTA) JP       |
| TOAFF       | 81  | PR D23 1500 | S. Toaff <i>et al.</i>         | (ANL, KANS)                 |
| ETKIN       | 80  | PR D22 42   | A. Etkin <i>et al.</i>         | (BNL, CUNY) JP              |
| BEUSCH      | 78  | PL 74B 282  | W. Beusch <i>et al.</i>        | (CERN, AACH3, ETH) JP       |
| CHUNG       | 78  | PRL 40 355  | S.U. Chung <i>et al.</i>       | (BNL, BRAN, CUNY+) JP       |
| ESTABROOKS  | 78  | NP B133 490 | P.G. Estabrooks <i>et al.</i>  | (MCGI, CARL, DURH+) JP      |
| Also        |     | PR D17 658  | P.G. Estabrooks <i>et al.</i>  | (MCGI, CARL, DURH+)         |
| BALDI       | 76  | PL 63B 344  | R. Baldi <i>et al.</i>         | (GEVA) JP                   |
| BRANDENB... | 76D | PL 60B 478  | G.W. Brandenburg <i>et al.</i> | (SLAC) JP                   |

### OTHER RELATED PAPERS

|             |    |             |                                       |                   |
|-------------|----|-------------|---------------------------------------|-------------------|
| AGUILAR-... | 73 | PRL 30 672  | M. Aguilar-Benitez <i>et al.</i>      | (BNL)             |
| WALUCH      | 73 | PR D8 2837  | V. Waluch, S.M. Flatte, J.H. Friedman | (LBL)             |
| CARMONY     | 71 | PRL 27 1160 | D.D. Carmony <i>et al.</i>            | (PURD, UCD, IUPU) |
| FIRESTONE   | 71 | PL 36B 513  | A. Firestone <i>et al.</i>            | (LBL)             |