

# $\chi_{b2}(2P)$

$$J^G(J^{PC}) = 0^+(2^{++})$$

$J$  needs confirmation.

Observed in radiative decay of the  $\Upsilon(3S)$ , therefore  $C = +$ . Branching ratio requires E1 transition, M1 is strongly disfavored, therefore  $P = +$ .

## $\chi_{b2}(2P)$ MASS

| VALUE (GeV)  | DOCUMENT ID   |
|--|---|
| <b>10.26865 ± 0.00022 ± 0.00050 OUR EVALUATION</b> | From $\gamma$ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeV |

## $m_{\chi_{b2}(2P)} - m_{\chi_{b1}(2P)}$

| VALUE (MeV)   | DOCUMENT ID         | TECN    | COMMENT   |
|---|---------------------|---------|---|
| <b>13.5 ± 0.4 ± 0.5</b>   | <sup>1</sup> HEINTZ | 92 CSB2 | $e^+e^- \rightarrow \gamma X, \ell^+\ell^-\gamma\gamma$ |
| <sup>1</sup> From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91. |                     |         |   |

## $\gamma$ ENERGY IN $\Upsilon(3S)$ DECAY

| VALUE (MeV)   | EVTS  | DOCUMENT ID         | TECN     | COMMENT                                       |
|---|-------|---------------------|----------|---|
| <b>86.19 ± 0.22 OUR EVALUATION</b>  |       |                     |          | Treating systematic errors as correlated      |
| <b>86.40 ± 0.18 OUR AVERAGE</b>   |       |                     |          |   |
| 86.04 ± 0.06 ± 0.27   |       | ARTUSO              | 05 CLEO  | $\Upsilon(3S) \rightarrow \gamma X$           |
| 86 ± 1  | 101   | CRAWFORD            | 92B CLE2 | $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$ |
| 86.7 ± 0.4  | 10319 | <sup>2</sup> HEINTZ | 92 CSB2  | $e^+e^- \rightarrow \gamma X$                 |
| 86.9 ± 0.4  | 157   | <sup>3</sup> HEINTZ | 92 CSB2  | $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$ |
| 86.4 ± 0.1 ± 0.4  | 30741 | MORRISON            | 91 CLE2  | $e^+e^- \rightarrow \gamma X$                 |
| <sup>2</sup> A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91. |       |                     |          |   |
| <sup>3</sup> A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91. |       |                     |          |   |

## $\chi_{b2}(2P)$ DECAY MODES

| Mode                                 | Fraction ( $\Gamma_i/\Gamma$ )               |
|--------------------------------------|--|
| $\Gamma_1 \quad \omega \Upsilon(1S)$ | ( 1.10 <sup>+0.34</sup> <sub>-0.30</sub> ) % |
| $\Gamma_2 \quad \gamma \Upsilon(2S)$ | ( 16.2 ± 2.4 ) %                             |
| $\Gamma_3 \quad \gamma \Upsilon(1S)$ | ( 7.1 ± 1.0 ) %                              |
| $\Gamma_4 \quad \pi\pi\chi_{b2}(1P)$ | ( 6.0 ± 2.1 ) × 10 <sup>-3</sup>             |

**$\chi_{b2}(2P)$  BRANCHING RATIOS** **$\Gamma(\omega \Upsilon(1S))/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$** 

| <u>VALUE (units <math>10^{-2}</math>)</u>          | <u>EVTS</u>          | <u>DOCUMENT ID</u>          | <u>TECN</u> | <u>COMMENT</u>  |
|--|----------------------|-----------------------------|-------------|---|
| <b><math>1.10^{+0.32+0.11}_{-0.28-0.10}</math></b> | $20.1^{+5.8}_{-5.1}$ | <sup>4</sup> CRONIN-HEN..04 | CLE3        | $\Upsilon(3S) \rightarrow \gamma \omega \Upsilon(1S)$ |

<sup>4</sup> Using  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.4 \pm 0.8)\%$  and  $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 2 (2.48 \pm 0.06)\%$ .

 **$\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$** 

| <u>VALUE</u>                                    | <u>DOCUMENT ID</u>        | <u>TECN</u> | <u>COMMENT</u>                                    |
|---|---------------------------|-------------|---|
| <b><math>0.162 \pm 0.024</math> OUR AVERAGE</b> |                           |             |   |
| $0.135 \pm 0.025 \pm 0.035$                     | <sup>5</sup> CRAWFORD 92B | CLE2        | $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$ |
| $0.173 \pm 0.021 \pm 0.019$                     | <sup>6</sup> HEINTZ 92    | CSB2        | $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$ |

<sup>5</sup> Using  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.37 \pm 0.26)\%$ ,  $B(\Upsilon(3S) \rightarrow \gamma \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (4.98 \pm 0.94 \pm 0.62) \times 10^{-4}$ , and  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 0.135 \pm 0.003 \pm 0.017$ .

<sup>6</sup> Using  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$ ,  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.1 \pm 0.5 \pm 0.4)\%$  and assuming  $e\mu$  universality. Supersedes HEINTZ 91.

 **$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$** 

| <u>VALUE</u>                                    | <u>DOCUMENT ID</u>        | <u>TECN</u> | <u>COMMENT</u>                                    |
|---|---------------------------|-------------|---|
| <b><math>0.071 \pm 0.010</math> OUR AVERAGE</b> |                           |             |   |
| $0.072 \pm 0.014 \pm 0.013$                     | <sup>7</sup> CRAWFORD 92B | CLE2        | $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$ |
| $0.070 \pm 0.010 \pm 0.006$                     | <sup>8</sup> HEINTZ 92    | CSB2        | $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$ |

<sup>7</sup> Using  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$ ,  $B(\Upsilon(3S) \rightarrow \gamma \gamma \Upsilon(2S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (5.03 \pm 0.94 \pm 0.63) \times 10^{-4}$ , and  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 0.135 \pm 0.003 \pm 0.017$ .

<sup>8</sup> Using  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$ ,  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.1 \pm 0.5 \pm 0.4)\%$  and assuming  $e\mu$  universality. Supersedes HEINTZ 91.

 **$\Gamma(\pi \pi \chi_{b2}(1P))/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$** 

| <u>VALUE (units <math>10^{-3}</math>)</u> | <u>DOCUMENT ID</u>        | <u>TECN</u> | <u>COMMENT</u>                                |
|---|---------------------------|-------------|---|
| <b><math>6.0 \pm 1.6 \pm 1.4</math></b>   | <sup>9</sup> CAWLFIELD 06 | CLE3        | $\Upsilon(3S) \rightarrow 2(\gamma \pi \ell)$ |

<sup>9</sup> CAWLFIELD 06 quote  $\Gamma(\chi_b(2P) \rightarrow \pi \pi \chi_b(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$  keV assuming l-spin conservation, no D-wave contribution,  $\Gamma(\chi_{b1}(2P)) = 96 \pm 16$  keV, and  $\Gamma(\chi_{b2}(2P)) = 138 \pm 19$  keV.

 **$\chi_{b2}(2P)$  REFERENCES**

|                 |               |                                  |                   |
|-----------------|---------------|----------------------------------|-------------------|
| CAWLFIELD 06    | PR D73 012003 | C. Cawlfeld <i>et al.</i>        | (CLEO Collab.)    |
| ARTUSO 05       | PRL 94 032001 | M. Artuso <i>et al.</i>          | (CLEO Collab.)    |
| CRONIN-HEN...04 | PRL 92 222002 | D. Cronin-Hennessy <i>et al.</i> | (CLEO3 Collab.)   |
| CRAWFORD 92B    | PL B294 139   | G. Crawford, R. Fulton           | (CLEO Collab.)    |
| HEINTZ 92       | PR D46 1928   | U. Heintz <i>et al.</i>          | (CUSB II Collab.) |
| HEINTZ 91       | PRL 66 1563   | U. Heintz <i>et al.</i>          | (CUSB Collab.)    |
| MORRISON 91     | PRL 67 1696   | R.J. Morrison <i>et al.</i>      | (CLEO Collab.)    |
| NARAIN 91       | PRL 66 3113   | M. Narain <i>et al.</i>          | (CUSB Collab.)    |

**OTHER RELATED PAPERS**

|          |             |                        |                |
|----------|-------------|------------------------|----------------|
| EIGEN 82 | PRL 49 1616 | G. Eigen <i>et al.</i> | (CUSB Collab.) |
| HAN 82   | PRL 49 1612 | K. Han <i>et al.</i>   | (CUSB Collab.) |