

# Supersymmetric Particle Searches

## SUPERSYMMETRY, PART I (THEORY)

Revised April 2006 by Howard E. Haber (Univ. of California, Santa Cruz)

**I.1. Introduction:** Supersymmetry (SUSY) is a generalization of the space-time symmetries of quantum field theory that transforms fermions into bosons and vice versa. The existence of such a non-trivial extension of the Poincaré symmetry of ordinary quantum field theory was initially surprising, and its form is highly constrained by theoretical principles [1]. Supersymmetry also provides a framework for the unification of particle physics and gravity [2–5], which is governed by the Planck energy scale,  $M_P \approx 10^{19}$  GeV (where the gravitational interactions become comparable in magnitude to the gauge interactions). In particular, it is possible that supersymmetry will ultimately explain the origin of the large hierarchy of energy scales from the  $W$  and  $Z$  masses to the Planck scale [6–9]. This is the so-called *gauge hierarchy*. The stability of the gauge hierarchy in the presence of radiative quantum corrections is not possible to maintain in the Standard Model, but can be maintained in supersymmetric theories.

If supersymmetry were an exact symmetry of nature, then particles and their superpartners (which differ in spin by half a unit) would be degenerate in mass. Since superpartners have not (yet) been observed, supersymmetry must be a broken symmetry. Nevertheless, the stability of the gauge hierarchy can still be maintained if the supersymmetry breaking is *soft* [10] and the corresponding supersymmetry-breaking mass parameters are no larger than a few TeV. (In this context, soft supersymmetry-breaking terms are non-supersymmetric terms

in the Lagrangian that are either linear, quadratic or cubic in the fields, with some restrictions elucidated in Ref. [10]. The impact of such terms becomes negligible at energy scales much larger than the size of the supersymmetry-breaking masses.) The most interesting theories of this type are theories of “low-energy” (or “weak-scale”) supersymmetry, where the effective scale of supersymmetry breaking is tied to the scale of electroweak symmetry breaking [6–9]. The latter is characterized by the Standard Model Higgs vacuum expectation value,  $v = 246$  GeV.

Although there are no unambiguous experimental results (at present) that require the existence of new physics at the TeV-scale, expectations of the latter are primarily based on three theoretical arguments. First, a *natural* explanation (*i.e.*, one that is stable with respect to quantum corrections) of the gauge hierarchy demands new physics at the TeV-scale [9]. Second, the unification of the three gauge couplings at a very high energy close to the Planck scale does not occur in the Standard Model. However, unification can be achieved with the addition of new physics that can modify the way gauge couplings run above the electroweak scale. A simple example of successful unification arises in the minimal supersymmetric extension of the Standard Model, where supersymmetric masses lie below a few TeV [11]. Third, the existence of dark matter which makes up approximately one quarter of the energy density of the universe, cannot be explained within the Standard Model of particle physics [12]. It is tempting to attribute the dark matter to the existence of a neutral stable thermal relic (*i.e.*, a particle that was in thermal equilibrium with all other fundamental particles in the early universe at temperatures above the particle mass). Remarkably, the existence of such

a particle could yield the observed density of dark matter if its mass and interaction rate were governed by new physics associated with the TeV-scale. The lightest supersymmetric particle is a promising (although not the unique) candidate for the dark matter [13].

Low-energy supersymmetry has traditionally been motivated by the three theoretical arguments just presented. More recently, some theorists [14,15] have argued that the explanation for the gauge hierarchy could lie elsewhere, in which case the effective TeV-scale theory would appear to be highly *unnatural*. Nevertheless, even without the naturalness argument, supersymmetry is expected to be a necessary ingredient of the ultimate theory at the Planck scale that unifies gravity with the other fundamental forces. Moreover, one can imagine that some remnant of supersymmetry does survive down to the TeV-scale. For example, in models of *split-supersymmetry* [15,16], some fraction of the supersymmetric spectrum remains light enough (with masses near the TeV scale) to provide successful gauge coupling unification and a viable dark matter candidate. If experimentation at future colliders uncovers evidence for (any remnant of) supersymmetry at low-energies, this would have a profound effect on the study of TeV-scale physics, and the development of a more fundamental theory of mass and symmetry-breaking phenomena in particle physics.

***1.2. Structure of the MSSM:*** The minimal supersymmetric extension of the Standard Model (MSSM) consists of taking the fields of the two-Higgs-doublet extension of the Standard Model and adding the corresponding supersymmetric partners [4,17]. The corresponding field content of the MSSM and their gauge quantum numbers are shown in Table 1. The electric charge

$Q = T_3 + \frac{1}{2}Y$  is determined in terms of the third component of the weak isospin ( $T_3$ ) and the U(1) hypercharge ( $Y$ ).

**Table 1:** The fields of the MSSM and their  $SU(3) \times SU(2) \times U(1)$  quantum numbers are listed. Only one generation of quarks and leptons is exhibited. For each lepton, quark and Higgs super-multiplet, there is a corresponding anti-particle multiplet of charge-conjugated fermions and their associated scalar partners.

Field Content of the MSSM					
Super-Multiplets	Boson Fields	Fermionic Partners	SU(3)	SU(2)	U(1)
gluon/gluino gauge/ gaugino	$g$	$\tilde{g}$	8	0	0
	$W^\pm, W^0$	$\tilde{W}^\pm, \tilde{W}^0$	1	3	0
	$B$	$\tilde{B}$	1	1	0
slepton/ lepton	$(\tilde{\nu}, \tilde{e}^-)_L$	$(\nu, e^-)_L$	1	2	-1
	$\tilde{e}_R^-$	$e_R^-$	1	1	-2
squark/ quark	$(\tilde{u}_L, \tilde{d}_L)$	$(u, d)_L$	3	2	1/3
	$\tilde{u}_R$	$u_R$	3	1	4/3
	$\tilde{d}_R$	$d_R$	3	1	-2/3
Higgs/ higgsino	$(H_d^0, H_d^-)$	$(\tilde{H}_d^0, \tilde{H}_d^-)$	1	2	-1
	$(H_u^+, H_u^0)$	$(\tilde{H}_u^+, \tilde{H}_u^0)$	1	2	1

The gauge super-multiplets consist of the gluons and their *gluino* fermionic superpartners and the  $SU(2) \times U(1)$  gauge bosons and their *gaugino* fermionic superpartners. The Higgs multiplets consist of two complex doublets of Higgs fields, their *higgsino* fermionic superpartners and the corresponding antiparticle fields. The matter super-multiplets consist of three generations of left-handed and right-handed quarks and lepton

fields, their scalar superpartners (squark and slepton fields) and the corresponding antiparticle fields.

The enlarged Higgs sector of the MSSM constitutes the minimal structure needed to guarantee the cancellation of anomalies from the introduction of the higgsino superpartners. Moreover, without a second Higgs doublet, one cannot generate mass for both “up”-type and “down”-type quarks (and charged leptons) in a way consistent with the supersymmetry [18–20]. The (renormalizable) MSSM Lagrangian is then constructed by including all possible interaction terms (of dimension four or less) that satisfy the spacetime supersymmetry algebra,  $SU(3)\times SU(2)\times U(1)$  gauge invariance and  $B-L$  conservation ( $B$  =baryon number and  $L$  =lepton number). Finally, the most general soft-supersymmetry-breaking terms are added [10]. To generate nonzero neutrino masses, extra structure is needed as discussed in section I.8.

### ***I.2.1. Constraints on supersymmetric parameters:***

If supersymmetry is associated with the origin of the electroweak scale, then the mass parameters introduced by the soft-supersymmetry-breaking must be generally of order 1 TeV or below [21] (although models have been proposed in which some supersymmetric particle masses can be larger, in the range of 1–10 TeV [22]) . Some lower bounds on these parameters exist due to the absence of supersymmetric-particle production at current accelerators [23]. Additional constraints arise from limits on the contributions of virtual supersymmetric particle exchange to a variety of Standard Model processes [24,25].

For example, the Standard Model global fit to precision electroweak data is quite good [26]. If all supersymmetric particle masses are significantly heavier than  $m_Z$  (in practice, masses greater than 300 GeV are sufficient [27]) , then the effects of the

supersymmetric particles decouple in loop-corrections to electroweak observables [28]. In this case, the Standard Model global fit to precision data and the corresponding MSSM fit yield similar results. On the other hand, regions of parameter space with light supersymmetric particle masses (just above the present day experimental limits) can in some cases generate significant one-loop corrections, resulting in a slight improvement or worsening of the overall global fit to the electroweak data depending on the choice of the MSSM parameters [29]. Thus, the precision electroweak data provide some constraints on the magnitude of the soft-supersymmetry-breaking terms.

There are a number of other low-energy measurements that are especially sensitive to the effects of new physics through virtual loops. For example, the virtual exchange of supersymmetric particles can contribute to the muon anomalous magnetic moment,  $a_\mu \equiv \frac{1}{2}(g-2)_\mu$ , and to the inclusive decay rate for  $b \rightarrow s\gamma$ . The most recent theoretical analysis of  $(g-2)_\mu$  finds a small deviation (less than three standard deviations) of the theoretical prediction from the experimentally observed value [30]. The theoretical prediction for  $\Gamma(b \rightarrow s\gamma)$  agrees quite well (within the error bars) to the experimental observation [31]. In both cases, supersymmetric corrections could have generated an observable shift from the Standard Model prediction in some regions of the MSSM parameter space [31–33]. The absence of a significant deviation places interesting constraints on the low-energy supersymmetry parameters.

***1.2.2. R-Parity and the lightest supersymmetric particle:*** As a consequence of  $B-L$  invariance, the MSSM possesses a multiplicative R-parity invariance, where  $R = (-1)^{3(B-L)+2S}$  for a particle of spin  $S$  [34]. Note that this implies that all the ordinary Standard Model particles

have even R parity, whereas the corresponding supersymmetric partners have odd R parity. The conservation of R parity in scattering and decay processes has a crucial impact on supersymmetric phenomenology. For example, starting from an initial state involving ordinary (R-even) particles, it follows that supersymmetric particles must be produced in pairs. In general, these particles are highly unstable and decay into lighter states. However, R-parity invariance also implies that the lightest supersymmetric particle (LSP) is absolutely stable, and must eventually be produced at the end of a decay chain initiated by the decay of a heavy unstable supersymmetric particle.

In order to be consistent with cosmological constraints, a stable LSP is almost certainly electrically and color neutral [35]. (There are some model circumstances in which a colored gluino LSP is allowed [36], but we do not consider this possibility further here.) Consequently, the LSP in an R-parity-conserving theory is weakly interacting with ordinary matter, *i.e.*, it behaves like a stable heavy neutrino and will escape collider detectors without being directly observed. Thus, the canonical signature for conventional R-parity-conserving supersymmetric theories is missing (transverse) energy, due to the escape of the LSP. Moreover, the LSP is a prime candidate for “cold dark matter” [13], an important component of the non-baryonic dark matter that is required in many models of cosmology and galaxy formation [37]. Further aspects of dark matter can be found in Ref. [38].

***1.2.3. The goldstino and gravitino:*** In the MSSM, supersymmetry breaking is accomplished by including the most general renormalizable soft-supersymmetry-breaking terms consistent with the  $SU(3)\times SU(2)\times U(1)$  gauge symmetry and R-parity invariance. These terms parameterize our ignorance of

the fundamental mechanism of supersymmetry breaking. If supersymmetry breaking occurs spontaneously, then a massless Goldstone fermion called the *goldstino* ( $\tilde{G}$ ) must exist. The goldstino would then be the LSP and could play an important role in supersymmetric phenomenology [39]. However, the goldstino is a physical degree of freedom only in models of spontaneously-broken global supersymmetry. If supersymmetry is a local symmetry, then the theory must incorporate gravity; the resulting theory is called supergravity [40]. In models of spontaneously-broken supergravity, the goldstino is “absorbed” by the *gravitino* ( $\tilde{g}_{3/2}$ ), the spin-3/2 partner of the graviton [41]. By this super-Higgs mechanism, the goldstino is removed from the physical spectrum and the gravitino acquires a mass ( $m_{3/2}$ ).

***1.2.4. Hidden sectors and the structure of supersymmetry breaking:*** It is very difficult (perhaps impossible) to construct a realistic model of spontaneously-broken low-energy supersymmetry where the supersymmetry breaking arises solely as a consequence of the interactions of the particles of the MSSM. A more viable scheme posits a theory consisting of at least two distinct sectors: a “hidden” sector consisting of particles that are completely neutral with respect to the Standard Model gauge group, and a “visible” sector consisting of the particles of the MSSM. There are no renormalizable tree-level interactions between particles of the visible and hidden sectors. Supersymmetry breaking is assumed to occur in the hidden sector, and to then be transmitted to the MSSM by some mechanism. Two theoretical scenarios have been examined in detail: gravity-mediated and gauge-mediated supersymmetry breaking.

Supergravity models provide a natural mechanism for transmitting the supersymmetry breaking of the hidden sector to the



particle spectrum of the MSSM. In models of *gravity-mediated* supersymmetry breaking, gravity is the messenger of supersymmetry breaking [42–44]. More precisely, supersymmetry breaking is mediated by effects of gravitational strength (suppressed by an inverse power of the Planck mass). In this scenario, the gravitino mass is of order the electroweak-symmetry-breaking scale, while its couplings are roughly gravitational in strength [2,45]. Such a gravitino would play no role in supersymmetric phenomenology at colliders.

In *gauge-mediated* supersymmetry breaking, supersymmetry breaking is transmitted to the MSSM via gauge forces. A typical structure of such models involves a hidden sector where supersymmetry is broken, a “messenger sector” consisting of particles (messengers) with  $SU(3) \times SU(2) \times U(1)$  quantum numbers, and the visible sector consisting of the fields of the MSSM [46,47]. The direct coupling of the messengers to the hidden sector generates a supersymmetry-breaking spectrum in the messenger sector. Finally, supersymmetry breaking is transmitted to the MSSM via the virtual exchange of the messengers. If this approach is extended to incorporate gravitational phenomena, then supergravity effects will also contribute to supersymmetry breaking. However, in models of gauge-mediated supersymmetry breaking, one usually chooses the model parameters in such a way that the virtual exchange of the messengers dominates the effects of the direct gravitational interactions between the hidden and visible sectors. In this scenario, the gravitino mass is typically in the eV to keV range, and is therefore the LSP. The helicity  $\pm \frac{1}{2}$  components of  $\tilde{g}_{3/2}$  behave approximately like the goldstino; its coupling to the particles of the MSSM is significantly stronger than a coupling of gravitational strength.

***1.2.5. Supersymmetry and extra dimensions:*** During the last few years, new approaches to supersymmetry breaking have been proposed, based on theories in which the number of space dimensions is greater than three. This is not a new idea—consistent superstring theories are formulated in ten spacetime dimensions, and the associated  $M$ -theory is based in eleven spacetime dimensions [48]. Nevertheless, in all approaches considered above, the string scale and the inverse size of the extra dimensions are assumed to be at or near the Planck scale, below which an effective four spacetime dimensional broken supersymmetric field theory emerges. More recently, a number of supersymmetry-breaking mechanisms have been proposed that are inherently extra-dimensional [49]. The size of the extra dimensions can be significantly larger than  $M_{\text{P}}^{-1}$ : in some cases of order  $(\text{TeV})^{-1}$  or even larger [50,51]. For example, in one approach, the fields of the MSSM live on some brane (a lower-dimensional manifold embedded in a higher dimensional spacetime), while the sector of the theory that breaks supersymmetry lives on a second separated brane. Two examples of this approach are anomaly-mediated supersymmetry breaking of Ref. [52] and gaugino-mediated supersymmetry breaking of Ref. [53]; in both cases supersymmetry-breaking is transmitted through fields that live in the bulk (the higher dimensional space between the two branes). This setup has some features in common with both gravity-mediated and gauge-mediated supersymmetry breaking (*e.g.*, a hidden and visible sector and messengers).

Alternatively, one can consider a higher dimensional theory that is compactified to four spacetime dimensions. In this approach, supersymmetry is broken by boundary conditions on the compactified space that distinguish between fermions and

bosons. This is the so-called Scherk-Schwarz mechanism [54]. The phenomenology of such models can be strikingly different from that of the usual MSSM [55]. All these extra-dimensional ideas clearly deserve further investigation, although they will not be discussed further here.

***1.2.6. Split-supersymmetry:*** If supersymmetry is not connected with the origin of the electroweak scale, string theory suggests that supersymmetry still plays a significant role in Planck-scale physics. However, it may still be possible that some remnant of the superparticle spectrum survives down to the TeV-scale or below. This is the idea of split-supersymmetry [15], in which supersymmetric scalar partners of the quarks and leptons are significantly heavier (perhaps by many orders of magnitude) than 1 TeV, whereas the fermionic partners of the gauge and Higgs bosons have masses of order 1 TeV or below (presumably protected by some chiral symmetry). With the exception of a single light neutral scalar whose properties are indistinguishable from those of the Standard Model Higgs boson, all other Higgs bosons are also taken to be very heavy.

The supersymmetry-breaking required to produce such a scenario would destabilize the gauge hierarchy. In particular, split-supersymmetry cannot provide a natural explanation for the existence of the light Standard Model-like Higgs boson whose mass lies orders below the the mass scale of the heavy scalars. Nevertheless, models of split-supersymmetry can account for the dark matter (which is assumed to be the LSP) and gauge coupling unification. Thus, there is some motivation for pursuing the phenomenology of such approaches [16]. One notable difference from the usual MSSM phenomenology is the existence of a long-lived gluino [56].

**I.3. Parameters of the MSSM:** The parameters of the MSSM are conveniently described by considering separately the supersymmetry-conserving sector and the supersymmetry-breaking sector. A careful discussion of the conventions used in defining the tree-level MSSM parameters can be found in Ref. [57]. (Additional fields and parameters must be introduced if one wishes to account for non-zero neutrino masses. We shall not pursue this here; see section I.8 for a discussion of supersymmetric approaches that incorporate neutrino masses.) For simplicity, consider first the case of one generation of quarks, leptons, and their scalar superpartners.

**I.3.1. The supersymmetric-conserving parameters:**

The parameters of the supersymmetry-conserving sector consist of: (i) gauge couplings:  $g_s$ ,  $g$ , and  $g'$ , corresponding to the Standard Model gauge group  $SU(3) \times SU(2) \times U(1)$  respectively; (ii) a supersymmetry-conserving higgsino mass parameter  $\mu$ ; and (iii) Higgs-fermion Yukawa coupling constants:  $\lambda_u$ ,  $\lambda_d$ , and  $\lambda_e$  (corresponding to the coupling of one generation of left and right-handed quarks and leptons and their superpartners to the Higgs bosons and higgsinos). Because there is no right-handed neutrino (and its superpartner) in the MSSM as defined here, one cannot introduce a Yukawa coupling  $\lambda_\nu$ .

**I.3.2. The supersymmetric-breaking parameters:**

The supersymmetry-breaking sector contains the following set of parameters: (i) gaugino Majorana masses  $M_3$ ,  $M_2$ , and  $M_1$  associated with the  $SU(3)$ ,  $SU(2)$ , and  $U(1)$  subgroups of the Standard Model; (ii) five scalar squared-mass parameters for the squarks and sleptons,  $M_Q^2$ ,  $M_U^2$ ,  $M_D^2$ ,  $M_L^2$ , and  $M_E^2$  [corresponding to the five electroweak gauge multiplets, *i.e.*, superpartners of  $(u, d)_L$ ,  $u_L^c$ ,  $d_L^c$ ,  $(\nu, e^-)_L$ , and  $e_L^c$ , where the superscript

$c$  indicates a charge-conjugated fermion]; and (iii) Higgs-squark-squark and Higgs-slepton-slepton trilinear interaction terms, with coefficients  $\lambda_u A_U$ ,  $\lambda_d A_D$ , and  $\lambda_e A_E$  (which define the so-called “ $A$ -parameters”). It is traditional to factor out the Yukawa couplings in the definition of the  $A$ -parameters (originally motivated by a simple class of gravity-mediated supersymmetry-breaking models [2,4]). If the  $A$ -parameters defined in this way are parametrically of the same order (or smaller) as compared to other supersymmetry-breaking mass parameters, then only the  $A$ -parameters of the third generation will be phenomenologically relevant.

Finally, we add: (iv) three scalar squared-mass parameters—two of which ( $m_1^2$  and  $m_2^2$ ) contribute to the diagonal Higgs squared-masses, given by  $m_1^2 + |\mu|^2$  and  $m_2^2 + |\mu|^2$ , and a third which contributes to the off-diagonal Higgs squared-mass term,  $m_{12}^2 \equiv B\mu$  (which defines the “ $B$ -parameter”). The breaking of the electroweak symmetry  $SU(2) \times U(1)$  to  $U(1)_{EM}$  is only possible after introducing the supersymmetry-breaking Higgs squared-mass parameters. Minimizing the resulting Higgs scalar potential, these three squared-mass parameters can be re-expressed in terms of the two Higgs vacuum expectation values,  $v_d$  and  $v_u$  (also called  $v_1$  and  $v_2$ , respectively, in the literature), and one physical Higgs mass. Here,  $v_d$  [ $v_u$ ] is the vacuum expectation value of the neutral component of the Higgs field  $H_d$  [ $H_u$ ] that couples exclusively to down-type (up-type) quarks and leptons. Note that  $v_d^2 + v_u^2 = 4m_W^2/g^2 = (246 \text{ GeV})^2$  is fixed by the  $W$  mass and the gauge coupling, whereas the ratio

$$\tan \beta = v_u/v_d \tag{1}$$

is a free parameter of the model. By convention, the Higgs field phases are chosen such that  $0 \leq \beta \leq \pi/2$ .

**I.3.3. MSSM-124:** The total number of degrees of freedom of the MSSM is quite large, primarily due to the parameters of the soft-supersymmetry-breaking sector. In particular, in the case of three generations of quarks, leptons, and their superpartners,  $M_{\tilde{Q}}^2$ ,  $M_{\tilde{U}}^2$ ,  $M_{\tilde{D}}^2$ ,  $M_{\tilde{L}}^2$ , and  $M_{\tilde{E}}^2$  are hermitian  $3 \times 3$  matrices, and  $A_U$ ,  $A_D$  and  $A_E$  are complex  $3 \times 3$  matrices. In addition,  $M_1$ ,  $M_2$ ,  $M_3$ ,  $B$ , and  $\mu$  are in general complex. Finally, as in the Standard Model, the Higgs-fermion Yukawa couplings,  $\lambda_f$  ( $f = u, d$ , and  $e$ ), are complex  $3 \times 3$  matrices that are related to the quark and lepton mass matrices via:  $M_f = \lambda_f v_f / \sqrt{2}$ , where  $v_e \equiv v_d$  (with  $v_u$  and  $v_d$  as defined above). However, not all these parameters are physical. Some of the MSSM parameters can be eliminated by expressing interaction eigenstates in terms of the mass eigenstates, with an appropriate redefinition of the MSSM fields to remove unphysical degrees of freedom. The analysis of Ref. [58] shows that the MSSM possesses 124 independent parameters. Of these, 18 parameters correspond to Standard Model parameters (including the QCD vacuum angle  $\theta_{\text{QCD}}$ ), one corresponds to a Higgs sector parameter (the analogue of the Standard Model Higgs mass), and 105 are genuinely new parameters of the model. The latter include: five real parameters and three  $CP$ -violating phases in the gaugino/higgsino sector, 21 squark and slepton masses, 36 real mixing angles to define the squark and slepton mass eigenstates, and 40  $CP$ -violating phases that can appear in squark and slepton interactions. The most general R-parity-conserving minimal supersymmetric extension of the Standard Model (without additional theoretical assumptions) will be denoted henceforth as MSSM-124 [59].

**I.4. The supersymmetric-particle sector:** Consider the sector of supersymmetric particles (*sparticles*) in the MSSM.

The supersymmetric partners of the gauge and Higgs bosons are fermions, whose names are obtained by appending “ino” at the end of the corresponding Standard Model particle name. The gluino is the color octet Majorana fermion partner of the gluon with mass  $M_{\tilde{g}} = |M_3|$ . The supersymmetric partners of the electroweak gauge and Higgs bosons (the gauginos and higgsinos) can mix. As a result, the physical states of definite mass are model-dependent linear combinations of the charged and neutral gauginos and higgsinos, called *charginos* and *neutralinos*, respectively. Like the gluino, the neutralinos are also Majorana fermions, which provide for some distinctive phenomenological signatures [60,61].

**I.4.1. The charginos and neutralinos:** The mixing of the charged gauginos ( $\widetilde{W}^\pm$ ) and charged higgsinos ( $H_u^+$  and  $H_d^-$ ) is described (at tree-level) by a  $2 \times 2$  complex mass matrix [62–64]:

$$M_C \equiv \begin{pmatrix} M_2 & \frac{1}{\sqrt{2}} g v_u \\ \frac{1}{\sqrt{2}} g v_d & \mu \end{pmatrix}. \quad (2)$$

To determine the physical chargino states and their masses, one must perform a singular value decomposition [65] of the complex matrix  $M_C$ :

$$U^* M_C V^{-1} = \text{diag}(M_{\tilde{\chi}_1^+}, M_{\tilde{\chi}_2^+}), \quad (3)$$

where  $U$  and  $V$  are unitary matrices and the right hand side of Eq. (3) is the diagonal matrix of (non-negative) chargino masses. The physical chargino states are denoted by  $\tilde{\chi}_1^\pm$  and  $\tilde{\chi}_2^\pm$ . These are linear combinations of the charged gaugino and higgsino states determined by the matrix elements of  $U$  and  $V$  [62–64].

The chargino masses correspond to the *singular values* [65] of  $M_C$ , *i.e.*, the positive square roots of the eigenvalues of  $M_C^\dagger M_C$ :

$$M_{\tilde{\chi}_1^\pm, \tilde{\chi}_2^\pm}^2 = \frac{1}{2} \left\{ |\mu|^2 + |M_2|^2 + 2m_W^2 \mp \left[ (|\mu|^2 + |M_2|^2 + 2m_W^2)^2 - 4|\mu|^2|M_2|^2 - 4m_W^4 \sin^2 2\beta + 8m_W^2 \sin 2\beta \operatorname{Re}(\mu M_2) \right]^{1/2} \right\}, \quad (4)$$

where the states are ordered such that  $M_{\tilde{\chi}_1^\pm} \leq M_{\tilde{\chi}_2^\pm}$ . It is often convenient to choose a convention where  $\tan\beta$  and  $M_2$  are real and positive. Note that the relative phase of  $M_2$  and  $\mu$  is meaningful. (If  $CP$ -violating effects are neglected, then  $\mu$  can be chosen real but may be either positive or negative.) The sign of  $\mu$  is convention-dependent; the reader is warned that both sign conventions appear in the literature. The sign convention for  $\mu$  in Eq. (2) is used by the LEP collaborations [23] in their plots of exclusion contours in the  $M_2$  vs.  $\mu$  plane derived from the non-observation of  $e^+e^- \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_1^-$ .

The mixing of the neutral gauginos ( $\tilde{B}$  and  $\tilde{W}^0$ ) and neutral higgsinos ( $\tilde{H}_d^0$  and  $\tilde{H}_u^0$ ) is described (at tree-level) by a  $4 \times 4$  complex symmetric mass matrix [62,63,66,67]:

$$M_N \equiv \begin{pmatrix} M_1 & 0 & -\frac{1}{2}g'v_d & \frac{1}{2}g'v_u \\ 0 & M_2 & \frac{1}{2}gv_d & -\frac{1}{2}gv_u \\ -\frac{1}{2}g'v_d & \frac{1}{2}gv_d & 0 & -\mu \\ \frac{1}{2}g'v_u & -\frac{1}{2}gv_u & -\mu & 0 \end{pmatrix}. \quad (5)$$

To determine the physical neutralino states and their masses, one must perform a Takagi factorization [65,68] of the complex symmetric matrix  $M_N$ :

$$W^T M_N W = \operatorname{diag}(M_{\tilde{\chi}_1^0}, M_{\tilde{\chi}_2^0}, M_{\tilde{\chi}_3^0}, M_{\tilde{\chi}_4^0}), \quad (6)$$



where  $W$  is a unitary matrix and the right hand side of Eq. (6) is the diagonal matrix of (non-negative) neutralino masses. The physical neutralino states are denoted by  $\tilde{\chi}_i^0$  ( $i = 1, \dots, 4$ ), where the states are ordered such that  $M_{\tilde{\chi}_1^0} \leq M_{\tilde{\chi}_2^0} \leq M_{\tilde{\chi}_3^0} \leq M_{\tilde{\chi}_4^0}$ . The  $\tilde{\chi}_i^0$  are the linear combinations of the neutral gaugino and higgsino states determined by the matrix elements of  $W$  (in Ref. [62],  $W = N^{-1}$ ). The neutralino masses correspond to the singular values of  $M_N$  (*i.e.*, the positive square roots of the eigenvalues of  $M_N^\dagger M_N$ ). Exact formulae for these masses can be found in Ref. [66,69].

If a chargino or neutralino state approximates a particular gaugino or higgsino state, it is convenient to employ the corresponding nomenclature. Specifically, if  $M_1$  and  $M_2$  are small compared to  $m_Z$  and  $|\mu|$ , then the lightest neutralino  $\tilde{\chi}_1^0$  would be nearly a pure *photino*,  $\tilde{\gamma}$ , the supersymmetric partner of the photon. If  $M_1$  and  $m_Z$  are small compared to  $M_2$  and  $|\mu|$ , then the lightest neutralino would be nearly a pure *bin*o,  $\tilde{B}$ , the supersymmetric partner of the weak hypercharge gauge boson. If  $M_2$  and  $m_Z$  are small compared to  $M_1$  and  $|\mu|$ , then the lightest chargino pair and neutralino would constitute a triplet of roughly mass-degenerate pure *wino*s,  $\tilde{W}^\pm$ , and  $\tilde{W}_3^0$ , the supersymmetric partners of the weak SU(2) gauge bosons. Finally, if  $|\mu|$  and  $m_Z$  are small compared to  $M_1$  and  $M_2$ , then the lightest neutralino would be nearly a pure *higgsino*. Each of the above cases leads to a strikingly different phenomenology.

***1.4.2. The squarks, sleptons and sneutrinos:*** The supersymmetric partners of the quarks and leptons are spin-zero bosons: the *squarks*, charged *sleptons*, and *sneutrinos*. For a given fermion  $f$ , there are two supersymmetric partners,  $\tilde{f}_L$  and  $\tilde{f}_R$ , which are scalar partners of the corresponding left- and right-handed fermion. (There is no  $\tilde{\nu}_R$  in the MSSM.)

However, in general,  $\tilde{f}_L$  and  $\tilde{f}_R$  are not mass-eigenstates, since there is  $\tilde{f}_L$ - $\tilde{f}_R$  mixing. For three generations of squarks, one must in general diagonalize  $6 \times 6$  matrices corresponding to the basis  $(\tilde{q}_{iL}, \tilde{q}_{iR})$ , where  $i = 1, 2, 3$  are the generation labels. For simplicity, only the one-generation case is illustrated in detail below (using the notation of the third family). In this case, the tree-level squark squared-mass matrix is given by [70]

$$M_F^2 = \begin{pmatrix} M_Q^2 + m_q^2 + L_q & m_q X_q^* \\ m_q X_q & M_R^2 + m_q^2 + R_q \end{pmatrix}, \quad (7)$$

where

$$X_q \equiv A_q - \mu^* (\cot \beta)^{2T_{3q}}, \quad (8)$$

and  $T_{3q} = \frac{1}{2} [-\frac{1}{2}]$  for  $q = t$  [ $b$ ]. The diagonal squared-masses are governed by soft-supersymmetry breaking squared-masses  $M_Q^2$  and  $M_R^2 \equiv M_U^2$  [ $M_D^2$ ] for  $q = t$  [ $b$ ], the corresponding quark masses  $m_t$  [ $m_b$ ], and electroweak correction terms:

$$\begin{aligned} L_q &\equiv (T_{3q} - e_q \sin^2 \theta_W) m_Z^2 \cos 2\beta, \\ R_q &\equiv e_q \sin^2 \theta_W m_Z^2 \cos 2\beta, \end{aligned} \quad (9)$$

where  $e_q = \frac{2}{3} [-\frac{1}{3}]$  for  $q = t$  [ $b$ ]. The off-diagonal squared squark masses are proportional to the corresponding quark masses and depend on  $\tan \beta$  [Eq. (1)], the soft-supersymmetry-breaking  $A$ -parameters and the higgsino mass parameter  $\mu$ . The signs of the  $A$  and  $\mu$  parameters are convention-dependent; other choices appear frequently in the literature. Due to the appearance of the *quark* mass in the off-diagonal element of the squark squared-mass matrix, one expects the  $\tilde{q}_L$ - $\tilde{q}_R$  mixing to be small, with the possible exception of the third-generation, where mixing can be enhanced by factors of  $m_t$  and  $m_b \tan \beta$ .

In the case of third generation  $\tilde{q}_L$ - $\tilde{q}_R$  mixing, the mass eigenstates (usually denoted by  $\tilde{q}_1$  and  $\tilde{q}_2$ , with  $m_{\tilde{q}_1} < m_{\tilde{q}_2}$ ) are determined by diagonalizing the  $2 \times 2$  matrix  $M_F^2$  given by Eq. (7). The corresponding squared-masses and mixing angle are given by [70]:

$$m_{\tilde{q}_{1,2}}^2 = \frac{1}{2} \left[ \text{Tr } M_F^2 \pm \sqrt{(\text{Tr } M_F^2)^2 - 4 \det M_F^2} \right],$$

$$\sin 2\theta_{\tilde{q}} = \frac{2m_q |X_q|}{m_{\tilde{q}_2}^2 - m_{\tilde{q}_1}^2}. \quad (10)$$

The one-generation results above also apply to the charged sleptons, with the obvious substitutions:  $q \rightarrow \tau$  with  $T_{3\tau} = -\frac{1}{2}$  and  $e_\tau = -1$ , and the replacement of the supersymmetry-breaking parameters:  $M_Q^2 \rightarrow M_L^2$ ,  $M_D^2 \rightarrow M_E^2$  and  $A_q \rightarrow A_\tau$ . For the neutral sleptons,  $\tilde{\nu}_R$  does not exist in the MSSM, so  $\tilde{\nu}_L$  is a mass-eigenstate.

In the case of three generations, the supersymmetry-breaking scalar squared-masses [ $M_Q^2$ ,  $M_U^2$ ,  $M_D^2$ ,  $M_L^2$  and  $M_E^2$ ] and the  $A$ -parameters that parameterize the Higgs couplings to up and down-type squarks and charged sleptons (henceforth denoted by  $A_U$ ,  $A_D$  and  $A_E$ , respectively) are now  $3 \times 3$  matrices as noted in Section I.3. The diagonalization of the  $6 \times 6$  squark mass matrices yields  $\tilde{f}_{iL}$ - $\tilde{f}_{jR}$  mixing (for  $i \neq j$ ). In practice, since the  $\tilde{f}_L$ - $\tilde{f}_R$  mixing is appreciable only for the third generation, this additional complication can usually be neglected.

Radiative loop corrections will modify all tree-level results for masses quoted in this section. These corrections must be included in any precision study of supersymmetric phenomenology [71]. Beyond tree-level, the definition of the supersymmetric parameters becomes convention-dependent. For example, one can define physical couplings or running couplings,

which differ beyond tree-level. This provides a challenge to any effort that attempts to extract supersymmetric parameters from data. The supersymmetric parameter analysis (SPA) project proposes a set of conventions [72] based on a consistent set of conventions and input parameters. dimensional reduction scheme for the regularization of higher-order loop corrections in supersymmetric theories recently advocated in Ultimately, these efforts will facilitate the reconstruction of the fundamental supersymmetric theory (and its breaking mechanism) from high precision studies of supersymmetric phenomena at future colliders.

***1.5. The Higgs sector of the MSSM:*** Next, consider the MSSM Higgs sector [19,20,73]. Despite the large number of potential  $CP$ -violating phases among the MSSM-124 parameters, the tree-level MSSM Higgs sector is automatically  $CP$ -conserving. That is, unphysical phases can be absorbed into the definition of the Higgs fields such that  $\tan\beta$  is a real parameter (conventionally chosen to be positive). Moreover, the physical neutral Higgs scalars are  $CP$  eigenstates. The model contains five physical Higgs particles: a charged Higgs boson pair ( $H^\pm$ ), two  $CP$ -even neutral Higgs bosons (denoted by  $h^0$  and  $H^0$  where  $m_h \leq m_H$ ), and one  $CP$ -odd neutral Higgs boson ( $A^0$ ).

***1.5.1 The Tree-level MSSM Higgs sector:*** The properties of the Higgs sector are determined by the Higgs potential, which is made up of quadratic terms [whose squared-mass coefficients were mentioned above Eq. (1)] and quartic interaction terms whose coefficients are dimensionless couplings. The quartic interaction terms are manifestly supersymmetric at tree-level (and are modified by supersymmetry-breaking effects only at the loop level). In general, the quartic couplings arise

from two sources: (i) the supersymmetric generalization of the scalar potential (the so-called “ $F$ -terms”), and (ii) interaction terms related by supersymmetry to the coupling of the scalar fields and the gauge fields, whose coefficients are proportional to the corresponding gauge couplings (the so-called “ $D$ -terms”). In the MSSM,  $F$ -term contributions to the quartic couplings are absent (although such terms may be present in extensions of the MSSM, *e.g.*, models with Higgs singlets). As a result, the strengths of the MSSM quartic Higgs interactions are fixed in terms of the gauge couplings. Due to the resulting constraint on the form of the two-Higgs-doublet scalar potential, all the tree-level MSSM Higgs-sector parameters depend only on two quantities:  $\tan\beta$  [defined in Eq. (1)] and one Higgs mass (usually taken to be  $m_A$ ). From these two quantities, one can predict the values of the remaining Higgs boson masses, an angle  $\alpha$  (which measures the component of the original  $Y = \pm 1$  Higgs doublet states in the physical  $CP$ -even neutral scalars), and the Higgs boson self-couplings.

***1.5.2 The radiatively-corrected MSSM Higgs sector:*** When radiative corrections are incorporated, additional parameters of the supersymmetric model enter via virtual loops. The impact of these corrections can be significant [74]. For example, the tree-level MSSM-124 prediction for the upper bound of the lightest  $CP$ -even Higgs mass,  $m_h \leq m_Z |\cos 2\beta| \leq m_Z$  [19,20], can be substantially modified when radiative corrections are included. The qualitative behavior of these radiative corrections can be most easily seen in the large top-squark mass limit, where in addition, both the splitting of the two diagonal entries and the two off-diagonal entries of the top-squark squared-mass matrix [Eq. (7)] are small in comparison to the average of the two top-squark

squared-masses,  $M_S^2 \equiv \frac{1}{2}(M_{\tilde{t}_1}^2 + M_{\tilde{t}_2}^2)$ . In this case (assuming  $m_A > m_Z$ ), the predicted upper bound for  $m_h$  (which reaches its maximum at large  $\tan\beta$ ) is approximately given by

$$m_h^2 \lesssim m_Z^2 + \frac{3g^2 m_t^4}{8\pi^2 m_W^2} \left\{ \ln(M_S^2/m_t^2) + \frac{X_t^2}{M_S^2} \left( 1 - \frac{X_t^2}{12M_S^2} \right) \right\}, \quad (11)$$

where  $X_t \equiv A_t - \mu \cot\beta$  is the top-squark mixing factor [see Eq. (7)]. A more complete treatment of the radiative corrections [75] shows that Eq. (11) somewhat overestimates the true upper bound of  $m_h$ . These more refined computations, which incorporate renormalization group improvement and the leading two-loop contributions, yield  $m_h \lesssim 135$  GeV (with an accuracy of a few GeV) for  $m_t = 175$  GeV and  $M_S \lesssim 2$  TeV [75]. This Higgs mass upper bound can be relaxed somewhat in non-minimal extensions of the MSSM, as noted in Section I.9.

In addition, one-loop radiative corrections can introduce  $CP$ -violating effects in the Higgs sector, which depend on some of the  $CP$ -violating phases among the MSSM-124 parameters [76]. Although these effects are more model-dependent, they can have a non-trivial impact on the Higgs searches at future colliders. A summary of the current MSSM Higgs mass limits can be found in Ref. [77].

***I.6. Restricting the MSSM parameter freedom:*** In Sections I.4 and I.5 we surveyed the parameters that comprise the MSSM-124. However in its most general form, the MSSM-124 is not a phenomenologically-viable theory over most of its parameter space. This conclusion follows from the observation that a generic point in the MSSM-124 parameter space exhibits: (i) no conservation of the separate lepton numbers  $L_e$ ,  $L_\mu$ , and  $L_\tau$ ; (ii) unsuppressed FCNC's; and (iii) new sources

of  $CP$  violation that are inconsistent with the experimental bounds.

For example, the MSSM contains many new sources of  $CP$  violation [78]. In particular, some combinations of the complex phases of the gaugino-mass parameters, the  $A$  parameters, and  $\mu$  must be less than of order  $10^{-2}$ – $10^{-3}$  (for a supersymmetry-breaking scale of 100 GeV) to avoid generating electric dipole moments for the neutron, electron, and atoms in conflict with observed data [79–81]. The non-observation of FCNC's [24,25] places additional strong constraints on the off-diagonal matrix elements of the squark and slepton soft-supersymmetry-breaking squared masses and  $A$ -parameters (see Section I.3.3). As a result of the phenomenological deficiencies listed above, almost the entire MSSM-124 parameter space is ruled out! This theory is viable only at very special “exceptional” regions of the full parameter space.

The MSSM-124 is also theoretically incomplete since it provides no explanation for the origin of the supersymmetry-breaking parameters (and in particular, why these parameters should conform to the exceptional points of the parameter space mentioned above). Moreover, there is no understanding of the choice of parameters that leads to the breaking of the electroweak symmetry. What is needed ultimately is a fundamental theory of supersymmetry breaking, which would provide a rationale for some set of soft-supersymmetry breaking terms that would be consistent with the phenomenological constraints referred to above. Presumably, the number of independent parameters characterizing such a theory would be considerably less than 124.

***1.6.1. Bottom-up approach for constraining the parameters of the MSSM:*** In the absence of a fundamental

theory of supersymmetry breaking, there are two general approaches for reducing the parameter freedom of MSSM-124. In the low-energy approach, an attempt is made to elucidate the nature of the exceptional points in the MSSM-124 parameter space that are phenomenologically viable. Consider the following two possible choices. First, one can assume that  $M_{\tilde{Q}}^2$ ,  $M_{\tilde{U}}^2$ ,  $M_{\tilde{D}}^2$ ,  $M_{\tilde{L}}^2$ ,  $M_{\tilde{E}}^2$ , and  $A_U$ ,  $A_D$ ,  $A_E$  are generation-independent (horizontal universality [7,58,82]). Alternatively, one can simply require that all the aforementioned matrices are flavor diagonal in a basis where the quark and lepton mass matrices are diagonal (flavor alignment [83]). In either case,  $L_e$ ,  $L_\mu$ , and  $L_\tau$  are separately conserved, while tree-level FCNC's are automatically absent. In both cases, the number of free parameters characterizing the MSSM is substantially less than 124. Both scenarios are phenomenologically viable, although there is no strong theoretical basis for either scenario.

***1.6.2. Top-down approach for constraining the parameters of the MSSM:*** In the high-energy approach, one imposes a particular structure on the soft-supersymmetry-breaking terms at a common high-energy scale (such as the Planck scale,  $M_P$ ). Using the renormalization group equations, one can then derive the low-energy MSSM parameters relevant for collider physics. The initial conditions (at the appropriate high-energy scale) for the renormalization group equations depend on the mechanism by which supersymmetry breaking is communicated to the effective low energy theory. Examples of this scenario are provided by models of gravity-mediated and gauge-mediated supersymmetry breaking (see Section I.2). One bonus of such an approach is that one of the diagonal Higgs squared-mass parameters is typically driven negative by renormalization group evolution [84]. Thus, electroweak



symmetry breaking is generated radiatively, and the resulting electroweak symmetry-breaking scale is intimately tied to the scale of low-energy supersymmetry breaking.

One prediction of the high-energy approach that arises in most grand unified supergravity models and gauge-mediated supersymmetry-breaking models is the unification of the (tree-level) gaugino mass parameters at some high-energy scale  $M_X$ :

$$M_1(M_X) = M_2(M_X) = M_3(M_X) = m_{1/2}. \quad (12)$$

Consequently, the effective low-energy gaugino mass parameters (at the electroweak scale) are related:

$$M_3 = (g_s^2/g^2)M_2, \quad M_1 = (5g'^2/3g^2)M_2 \simeq 0.5M_2. \quad (13)$$

In this case, the chargino and neutralino masses and mixing angles depend only on three unknown parameters: the gluino mass,  $\mu$ , and  $\tan\beta$ . If in addition  $|\mu| \gg M_1 \gtrsim m_Z$ , then the lightest neutralino is nearly a pure bino, an assumption often made in supersymmetric particle searches at colliders.

### ***I.6.3. Anomaly-mediated supersymmetry-breaking:***

In some supergravity models, tree-level masses for the gauginos are absent. The gaugino mass parameters arise at one-loop and do not satisfy Eq. (13). In this case, one finds a model-independent contribution to the gaugino mass whose origin can be traced to the super-conformal (super-Weyl) anomaly, which is common to all supergravity models [52]. This approach is called *anomaly-mediated* supersymmetry breaking (AMSB). Eq. (13) is then replaced (in the one-loop approximation) by:

$$M_i \simeq \frac{b_i g_i^2}{16\pi^2} m_{3/2}, \quad (14)$$

where  $m_{3/2}$  is the gravitino mass (assumed to be of order 1 TeV), and  $b_i$  are the coefficients of the MSSM gauge beta-functions corresponding to the corresponding U(1), SU(2) and SU(3) gauge groups:  $(b_1, b_2, b_3) = (\frac{33}{5}, 1, -3)$ . Eq. (14) yields  $M_1 \simeq 2.8M_2$  and  $M_3 \simeq -8.3M_2$ , which implies that the lightest chargino pair and neutralino comprise a nearly mass-degenerate triplet of winos,  $\widetilde{W}^\pm, \widetilde{W}^0$  (*c.f.* Table 1), over most of the MSSM parameter space. (For example, if  $|\mu| \gg m_Z$ , then Eq. (14) implies that  $M_{\widetilde{\chi}_1^\pm} \simeq M_{\widetilde{\chi}_1^0} \simeq M_2$  [85].) The corresponding supersymmetric phenomenology differs significantly from the standard phenomenology based on Eq. (13), and is explored in detail in Ref. [86]. Anomaly-mediated supersymmetry breaking also generates (approximate) flavor-diagonal squark and slepton mass matrices. However, this yields negative squared-mass contributions for the sleptons in the MSSM. This fatal flaw may be possible to cure in approaches beyond the minimal supersymmetric model [87]. Alternatively, one may conclude that anomaly-mediation is not the sole source of supersymmetry-breaking in the slepton sector.

***I.7. The constrained MSSMs: mSUGRA, GMSB, and SGUTs:*** One way to guarantee the absence of significant FCNC's mediated by virtual supersymmetric-particle exchange is to posit that the diagonal soft-supersymmetry-breaking scalar squared-masses are universal at some energy scale.

***I.7.1. The minimal supergravity (mSUGRA) model:*** In the *minimal* supergravity (mSUGRA) framework [2–4], the soft-supersymmetry-breaking parameters at the Planck scale take a particularly simple form in which the scalar squared-masses and the  $A$ -parameters are flavor-diagonal

and universal [43]:

$$\begin{aligned}
 M_{\tilde{Q}}^2(M_{\text{P}}) &= M_{\tilde{U}}^2(M_{\text{P}}) = M_{\tilde{D}}^2(M_{\text{P}}) = m_0^2 \mathbf{1}, \\
 M_{\tilde{L}}^2(M_{\text{P}}) &= M_{\tilde{E}}^2(M_{\text{P}}) = m_0^2 \mathbf{1}, \\
 m_1^2(M_{\text{P}}) &= m_2^2(M_{\text{P}}) = m_0^2, \\
 A_U(M_{\text{P}}) &= A_D(M_{\text{P}}) = A_E(M_{\text{P}}) = A_0 \mathbf{1},
 \end{aligned}
 \tag{15}$$

where  $\mathbf{1}$  is a  $3 \times 3$  identity matrix in generation space. Renormalization group evolution is then used to derive the values of the supersymmetric parameters at the low-energy (electroweak) scale. For example, to compute squark masses, one must use the *low-energy* values for  $M_{\tilde{Q}}^2$ ,  $M_{\tilde{U}}^2$  and  $M_{\tilde{D}}^2$  in Eq. (7). Through the renormalization group running with boundary conditions specified in Eq. (13) and Eq. (15), one can show that the low-energy values of  $M_{\tilde{Q}}^2$ ,  $M_{\tilde{U}}^2$  and  $M_{\tilde{D}}^2$  depend primarily on  $m_0^2$  and  $m_{1/2}^2$ . A number of useful approximate analytic expressions for superpartner masses in terms of the mSUGRA parameters can be found in Ref. [88].

Clearly, in the mSUGRA approach, the MSSM-124 parameter freedom has been significantly reduced. Typical mSUGRA models give low-energy values for the scalar mass parameters that satisfy  $M_{\tilde{L}} \approx M_{\tilde{E}} < M_{\tilde{Q}} \approx M_{\tilde{U}} \approx M_{\tilde{D}}$ , with the squark mass parameters somewhere between a factor of 1–3 larger than the slepton mass parameters (*e.g.*, see Ref. [88]). More precisely, the low-energy values of the squark mass parameters of the first two generations are roughly degenerate, while  $M_{\tilde{Q}_3}$  and  $M_{\tilde{U}_3}$  are typically reduced by a factor of 1–3 from the values of the first and second generation squark mass parameters, because of renormalization effects due to the heavy top-quark mass.

As a result, one typically finds that four flavors of squarks (with two squark eigenstates per flavor) and  $\tilde{b}_R$  are nearly mass-degenerate. The  $\tilde{b}_L$  mass and the diagonal  $\tilde{t}_L$  and  $\tilde{t}_R$  masses are reduced compared to the common squark mass of the first two generations. In addition, there are six flavors of nearly mass-degenerate sleptons (with two slepton eigenstates per flavor for the charged sleptons and one per flavor for the sneutrinos); the sleptons are expected to be somewhat lighter than the mass-degenerate squarks. Finally, third generation squark masses and tau-slepton masses are sensitive to the strength of the respective  $\tilde{f}_L$ - $\tilde{f}_R$  mixing, as discussed below Eq. (7). If  $\tan\beta \gg 1$ , then the pattern of third generation squark masses is somewhat altered, as discussed in Ref. [89].

In mSUGRA models, the LSP is typically the lightest neutralino,  $\tilde{\chi}_1^0$ , which is dominated by its bino component. In particular, one can reject those mSUGRA parameter regimes in which the LSP is a chargino or the  $\tilde{\tau}_1$  (the lightest scalar superpartner of the  $\tau$ -lepton). In general, if one imposes the constraints of supersymmetric particle searches and those of cosmology (say, by requiring the LSP to be a suitable dark matter candidate), one obtains significant restrictions to the mSUGRA parameter space [90].


One can count the number of independent parameters in the mSUGRA framework. In addition to 18 Standard Model parameters (excluding the Higgs mass), one must specify  $m_0$ ,  $m_{1/2}$ ,  $A_0$ , and Planck-scale values for  $\mu$  and  $B$ -parameters (denoted by  $\mu_0$  and  $B_0$ ). In principle,  $A_0$ ,  $B_0$ , and  $\mu_0$  can be complex, although in the mSUGRA approach, these parameters are taken (arbitrarily) to be real. As previously noted, renormalization group evolution is used to compute the low-energy values of the mSUGRA parameters, which then fixes all the parameters of the low-energy MSSM. In particular, the two Higgs

vacuum expectation values (or equivalently,  $m_Z$  and  $\tan\beta$ ) can be expressed as a function of the Planck-scale supergravity parameters. The simplest procedure is to remove  $\mu_0$  and  $B_0$  in favor of  $m_Z$  and  $\tan\beta$  [the sign of  $\mu_0$ , denoted  $\text{sgn}(\mu_0)$  below, is not fixed in this process]. In this case, the MSSM spectrum and its interaction strengths are determined by five parameters:

$$m_0, A_0, m_{1/2}, \tan\beta, \text{ and } \text{sgn}(\mu_0), \quad (16)$$

in addition to the 18 parameters of the Standard Model. However, the mSUGRA approach is probably too simplistic. Theoretical considerations suggest that the universality of Planck-scale soft-supersymmetry-breaking parameters is not generic [91]. In particular, it is easy to write down effective operators at the Planck scale that do not respect flavor universality, and it is difficult to find a theoretical principle that would forbid them.

In order to facilitate studies of supersymmetric phenomenology at colliders, it has been a valuable exercise to compile a set of benchmark supersymmetric parameters, from which supersymmetric spectra and couplings can be derived [92]. A compilation of benchmark mSUGRA points consistent with present data from particle physics and cosmology can be found in Ref. [93]. One particular well-studied benchmark points, the so-called SPS 1a' reference point [72] (this is a slight modification of the SPS 1a point of Ref. [92], which incorporates the latest constraints from collider data and cosmology) has been especially useful in experimental studies of supersymmetric phenomena at future colliders. The supersymmetric particle spectrum for the SPS 1a' reference point is exhibited in Figure 1. However, it is important to keep in mind that even within the mSUGRA framework, the resulting supersymmetric



**Figure 1:** Mass spectrum of supersymmetric particles and Higgs bosons for the mSUGRA reference point SPS 1a'. The masses of the first and second generation squarks, sleptons and sneutrinos are denoted collectively by  $\tilde{q}$ ,  $\tilde{\ell}$  and  $\tilde{\nu}_\ell$ , respectively. Taken from Ref. [72]. See full-color version on color pages at end of book.

theory and its attendant phenomenology can be quite different from the SPS 1a' reference point.

### ***I.7.2. Gauge-mediated supersymmetry breaking:***

In contrast to models of gravity-mediated supersymmetry breaking, the universality of the fundamental soft-supersymmetry-breaking squark and slepton squared-mass parameters is guaranteed in gauge-mediated supersymmetry-breaking because the supersymmetry-breaking is communicated to the sector of MSSM fields via gauge interactions. In the minimal gauge-mediated supersymmetry-breaking (GMSB) approach, there is one effective mass scale,  $\Lambda$ , that determines all low-energy scalar

and gaugino mass parameters through loop-effects (while the resulting  $A$  parameters are suppressed). In order that the resulting superpartner masses be of order 1 TeV or less, one must have  $\Lambda \sim 100$  TeV. The origin of the  $\mu$  and  $B$ -parameters is quite model-dependent, and lies somewhat outside the ansatz of gauge-mediated supersymmetry breaking. The simplest models of this type are even more restrictive than mSUGRA, with two fewer degrees of freedom. Benchmark reference points for GMSB models have been proposed in Ref. [92] to facilitate collider studies. However, minimal GMSB is not a fully realized model. The sector of supersymmetry-breaking dynamics can be very complex, and no complete model of gauge-mediated supersymmetry yet exists that is both simple and compelling.

It was noted in Section I.2 that the gravitino is the LSP in GMSB models. Thus, in such models, the next-to-lightest supersymmetric particle (NLSP) plays a crucial role in the phenomenology of supersymmetric particle production and decay. Note that unlike the LSP, the NLSP can be charged. In GMSB models, the most likely candidates for the NLSP are  $\tilde{\chi}_1^0$  and  $\tilde{\tau}_R^\pm$ . The NLSP will decay into its superpartner plus a gravitino (*e.g.*,  $\tilde{\chi}_1^0 \rightarrow \gamma \tilde{g}_{3/2}$ ,  $\tilde{\chi}_1^0 \rightarrow Z \tilde{g}_{3/2}$  or  $\tilde{\tau}_R^\pm \rightarrow \tau^\pm \tilde{g}_{3/2}$ ), with lifetimes and branching ratios that depend on the model parameters.

Different choices for the identity of the NLSP and its decay rate lead to a variety of distinctive supersymmetric phenomenologies [47,94]. For example, a long-lived  $\tilde{\chi}_1^0$ -NLSP that decays outside collider detectors leads to supersymmetric decay chains with missing energy in association with leptons and/or hadronic jets (this case is indistinguishable from the canonical phenomenology of the  $\tilde{\chi}_1^0$ -LSP). On the other hand, if  $\tilde{\chi}_1^0 \rightarrow \gamma \tilde{g}_{3/2}$  is the dominant decay mode, and the decay occurs inside the detector, then nearly *all* supersymmetric particle

decay chains would contain a photon. In contrast, the case of a  $\tilde{\tau}_R^\pm$ -NLSP would lead either to a new long-lived charged particle (*i.e.*, the  $\tilde{\tau}_R^\pm$ ) or to supersymmetric particle decay chains with  $\tau$  leptons.

**I.7.3. Supersymmetric grand unification:** Finally, grand unification [95] can impose additional constraints on the MSSM parameters. As emphasized in Section I.1, it is striking that the  $SU(3)\times SU(2)\times U(1)$  gauge couplings unify in models of supersymmetric grand unified theories (SGUTs) [7,15,96,97] with (some of) the supersymmetry-breaking parameters of order 1 TeV or below. Gauge coupling unification, which takes place at an energy scale of order  $10^{16}$  GeV, is quite robust [98]. For example, successful unification depends weakly on the details of the theory at the unification scale. In particular, given the low-energy values of the electroweak couplings  $g(m_Z)$  and  $g'(m_Z)$ , one can predict  $\alpha_s(m_Z)$  by using the MSSM renormalization group equations to extrapolate to higher energies, and by imposing the unification condition on the three gauge couplings at some high-energy scale,  $M_X$ . This procedure, which fixes  $M_X$ , can be successful (*i.e.*, three running couplings will meet at a single point) only for a unique value of  $\alpha_s(m_Z)$ . The extrapolation depends somewhat on the low-energy supersymmetric spectrum (so-called low-energy “threshold effects”), and on the SGUT spectrum (high-energy threshold effects), which can somewhat alter the evolution of couplings. Ref. [99] summarizes the comparison of data with the expectations of SGUTs, and shows that the measured value of  $\alpha_s(m_Z)$  is in good agreement with the predictions of supersymmetric grand unification for a reasonable choice of supersymmetric threshold corrections.

Additional SGUT predictions arise through the unification of the Higgs-fermion Yukawa couplings ( $\lambda_f$ ). There is some



evidence that  $\lambda_b = \lambda_\tau$  is consistent with observed low-energy data [100], and an intriguing possibility that  $\lambda_b = \lambda_\tau = \lambda_t$  may be phenomenologically viable [89,101] in the parameter regime where  $\tan\beta \simeq m_t/m_b$ . Finally, grand unification imposes constraints on the soft-supersymmetry-breaking parameters. For example, gaugino-mass unification leads to the relations given by Eq. (13). Diagonal squark and slepton soft-supersymmetry-breaking scalar masses may also be unified, which is analogous to the unification of Higgs-fermion Yukawa couplings.

In the absence of a fundamental theory of supersymmetry breaking, further progress will require a detailed knowledge of the supersymmetric-particle spectrum in order to determine the nature of the high-energy parameters. Of course, any of the theoretical assumptions described in this section could be wrong and must eventually be tested experimentally.

### ***1.8. Massive neutrinos in low-energy supersymmetry:***

With the overwhelming evidence for neutrino masses and mixing [102], it is clear that any viable supersymmetric model of fundamental particles must incorporate some form of  $L$  violation in the low-energy theory [103]. This requires an extension of the MSSM, which (as in the case of the minimal Standard Model) contains three generations of massless neutrinos. To construct a supersymmetric model with massive neutrinos, one can follow one of two different approaches.

***1.8.1. The supersymmetric seesaw:*** In the first approach, one starts with a modified Standard Model which incorporates new structure that yields nonzero neutrino masses. Following the procedures of Sections I.2 and I.3, one then formulates the supersymmetric extension of the modified Standard Model. For example, neutrino masses can be incorporated into

the Standard Model by introducing an  $SU(3) \times SU(2) \times U(1)$  singlet right-handed neutrino ( $\nu_R$ ) and a super-heavy Majorana mass (typically of order a grand unified mass) for the  $\nu_R$ . In addition, one must also include a standard Yukawa coupling between the lepton doublet, the Higgs doublet and  $\nu_R$ . The Higgs vacuum expectation value then induces an off-diagonal  $\nu_L$ - $\nu_R$  mass of order the electroweak scale. Diagonalizing the neutrino mass matrix (in the three-generation model) yields three superheavy neutrino states and three very light neutrino states that are identified as the light neutrino states observed in nature. This is the seesaw mechanism [104]. The supersymmetric generalization of the seesaw model of neutrino masses is now easily constructed [105,106].

***1.8.2. R-parity-violating supersymmetry:*** Another approach to incorporating massive neutrinos in supersymmetric models is to retain the minimal particle content of the MSSM but remove the assumption of R-parity invariance [107]. The most general R-parity-violating (RPV) theory involving the MSSM spectrum introduces many new parameters to both the supersymmetry-conserving and the supersymmetry-breaking sectors. Each new interaction term violates either  $B$  or  $L$  conservation. For example, consider new scalar-fermion Yukawa couplings derived from the following interactions:

$$(\lambda_L)_{pmn} \widehat{L}_p \widehat{L}_m \widehat{E}_n^c + (\lambda'_L)_{pmn} \widehat{L}_p \widehat{Q}_m \widehat{D}_n^c + (\lambda_B)_{pmn} \widehat{U}_p^c \widehat{D}_m^c \widehat{D}_n^c, \quad (17)$$

where  $p$ ,  $m$ , and  $n$  are generation indices, and gauge group indices are suppressed. In the notation above,  $\widehat{Q}$ ,  $\widehat{U}^c$ ,  $\widehat{D}^c$ ,  $\widehat{L}$ , and  $\widehat{E}^c$  respectively represent  $(u, d)_L$ ,  $u_L^c$ ,  $d_L^c$ ,  $(\nu, e^-)_L$ , and  $e_L^c$  and the corresponding superpartners. The Yukawa interactions are obtained from Eq. (17) by taking all possible combinations involving two fermions and one scalar superpartner. Note that

the term in Eq. (17) proportional to  $\lambda_B$  violates  $B$ , while the other two terms violate  $L$ . Even if all the terms of Eq. (17) are absent, there is one more possible supersymmetric source of R-parity violation. In the notation of Eq. (17), one can add a term of the form  $(\mu_L)_p \widehat{H}_u \widehat{L}_p$ , where  $\widehat{H}_u$  represents the  $Y = 1$  Higgs doublet and its higgsino superpartner. This term is the RPV generalization of the supersymmetry-conserving Higgs mass parameter  $\mu$  of the MSSM, in which the  $Y = -1$  Higgs/higgsino super-multiplet  $\widehat{H}_d$  is replaced by the slepton/lepton super-multiplet  $\widehat{L}_p$ . The RPV-parameters  $(\mu_L)_p$  also violate  $L$ .

Phenomenological constraints derived from data on various low-energy  $B$ - and  $L$ -violating processes can be used to establish limits on each of the coefficients  $(\lambda_L)_{pmn}$ ,  $(\lambda'_L)_{pmn}$ , and  $(\lambda_B)_{pmn}$  taken one at a time [107,108]. If more than one coefficient is simultaneously non-zero, then the limits are, in general, more complicated [109]. All possible RPV terms cannot be simultaneously present and unsuppressed; otherwise the proton decay rate would be many orders of magnitude larger than the present experimental bound. One way to avoid proton decay is to impose  $B$  or  $L$  invariance (either one alone would suffice). Otherwise, one must accept the requirement that certain RPV coefficients must be extremely suppressed.

One particularly interesting class of RPV models is one in which  $B$  is conserved, but  $L$  is violated. It is possible to enforce baryon number conservation, while allowing for lepton number violating interactions by imposing a discrete  $\mathbf{Z}_3$  baryon *triality* symmetry on the low-energy theory [110], in place of the standard  $\mathbf{Z}_2$  R-parity. Since the distinction between the Higgs and matter super-multiplets is lost in RPV models, R-parity violation permits the mixing of sleptons and Higgs bosons, the mixing of neutrinos and neutralinos, and the mixing of

charged leptons and charginos, leading to more complicated mass matrices and mass eigenstates than in the MSSM.

The supersymmetric phenomenology of the RPV models exhibits features that are quite distinct from that of the MSSM [107]. The LSP is no longer stable, which implies that not all supersymmetric decay chains must yield missing-energy events at colliders. Nevertheless, the loss of the missing-energy signature is often compensated by other striking signals (which depend on which R-parity-violating parameters are dominant). For example, supersymmetric particles in RPV models can be singly produced (in contrast to R-parity-conserving models where supersymmetric particles must be produced in pairs). The phenomenology of pair-produced supersymmetric particles in RPV models can also differ significantly from expectations due to new decay chains not present in R-parity-conserving supersymmetry [107].

In RPV models with lepton number violation (these include low-energy supersymmetry models with baryon triality mentioned above), both  $\Delta L = 1$  and  $\Delta L = 2$  phenomena are allowed, leading to neutrino masses and mixing [111], neutrinoless double-beta decay [112], sneutrino-antisneutrino mixing [106,113,114], and  $s$ -channel resonant production of sneutrinos in  $e^+e^-$  collisions [115] and charged sleptons in  $p\bar{p}$  and  $pp$  collisions [116]. For example, Ref. [117] demonstrates how one can fit both the solar and atmospheric neutrino data in an RPV supersymmetric model where  $\mu_L$  provides the dominant source of R-parity violation.

### ***I.9. Other non-minimal extensions of the MSSM:***

There are additional motivations for extending the supersymmetric model beyond the MSSM. Here we mention just a few. The  $\mu$  parameter of the MSSM is a supersymmetric-preserving parameter; nevertheless it must be of order the

supersymmetry-breaking scale to yield a consistent supersymmetric phenomenology. In the MSSM, one must devise a theoretical mechanism to guarantee that the magnitude of  $\mu$  is not larger than the TeV-scale (*e.g.*, in gravity-mediated supersymmetry, the Giudice-Masiero mechanism of Ref. [118] is the most cited explanation).

In extensions of the MSSM, new compelling solutions to the so-called  $\mu$ -problem are possible. For example, one can replace  $\mu$  by the vacuum expectation value of a new  $SU(3) \times SU(2) \times U(1)$  singlet scalar field. In such a model, the Higgs sector of the MSSM is enlarged (and the corresponding fermionic higgsino superpartner is added). This is the so-called NMSSM (here, NM stands for non-minimal) [119].

Non-minimal extensions of the MSSM involving additional matter super-multiplets can also yield a less restrictive bound on the mass of the lightest Higgs boson (as compared to the MSSM Higgs mass bound quoted in Section I.5.2). For example, by imposing gauge coupling unification, the upper limit on the lightest Higgs boson mass can be as high as 200—300 GeV [120] (a similar relaxation of the Higgs mass bound has been observed in split supersymmetry [121] and in extra-dimensional scenarios [122]). Note that these less restrictive Higgs mass upper bounds are comparable to the (experimentally determined) upper bound for the Higgs boson mass based on the Standard Model global fits to precision electroweak data [26,123].

Other MSSM extensions considered in the literature include an enlarged electroweak gauge group beyond  $SU(2) \times U(1)$  [124]; and/or the addition of new, possibly exotic, matter super-multiplets (*e.g.*, a vector-like color triplet with electric charge  $\frac{1}{3}e$ ; such states sometimes occur as low-energy remnants in  $E_6$  grand unification models). A possible theoretical motivation for

such new structures arises from the study of phenomenologically viable string theory ground states [125].

## References

1. R. Haag, J. T. Lopuszanski and M. Sohnius, Nucl. Phys. **B88**, 257 (1975) S.R. Coleman and J. Mandula, Phys. Rev. **159** (1967) 1251.
2. H.P. Nilles, Phys. Reports **110**, 1 (1984).
3. P. Nath, R. Arnowitt, and A.H. Chamseddine, *Applied N = 1 Supergravity* (World Scientific, Singapore, 1984).
4. S.P. Martin, in *Perspectives on Supersymmetry*, edited by G.L. Kane (World Scientific, Singapore, 1998) pp. 1–98; and a longer archive version in [hep-ph/9709356](http://arxiv.org/abs/hep-ph/9709356); I.J.R. Aitchison, [hep-ph/0505105](http://arxiv.org/abs/hep-ph/0505105).
5. S. Weinberg, *The Quantum Theory of Fields, Volume III: Supersymmetry* (Cambridge University Press, Cambridge, UK, 2000).
6. E. Witten, Nucl. Phys. **B188**, 513 (1981).
7. S. Dimopoulos and H. Georgi, Nucl. Phys. **B193**, 150 (1981).
8. N. Sakai, Z. Phys. **C11**, 153 (1981); R.K. Kaul, Phys. Lett. **109B**, 19 (1982).
9. L. Susskind, Phys. Reports **104**, 181 (1984).
10. L. Girardello and M. Grisaru, Nucl. Phys. **B194**, 65 (1982); L.J. Hall and L. Randall, Phys. Rev. Lett. **65**, 2939 (1990); I. Jack and D.R.T. Jones, Phys. Lett. **B457**, 101 (1999).
11. For a review, see N. Polonsky, *Supersymmetry: Structure and phenomena. Extensions of the standard model*, Lect. Notes Phys. **M68**, 1 (2001).
12. G. Bertone, D. Hooper and J. Silk, Phys. Reports **405**, 279 (2005).
13. G. Jungman, M. Kamionkowski, and K. Griest, Phys. Reports **267**, 195 (1996).

14. V. Agrawal, S.M. Barr, J.F. Donoghue and D. Seckel, Phys. Rev. **D57**, 5480 (1998).
15. N. Arkani-Hamed and S. Dimopoulos, JHEP **0506**, 073 (2005);  
G.F. Giudice and A. Romanino, Nucl. Phys. **B699**, 65 (2004) [erratum: **B706**, 65 (2005)].
16. N. Arkani-Hamed, S. Dimopoulos, G.F. Giudice and A. Romanino, Nucl. Phys. **B709**, 3 (2005);  
W. Kilian, T. Plehn, P. Richardson and E. Schmidt, Eur. Phys. J. **C39**, 229 (2005).
17. H.E. Haber and G.L. Kane, Phys. Reports **117**, 75 (1985);  
M. Drees, R. Godbole and P. Roy, *Theory and Phenomenology of Sparticles* (World Scientific, Singapore, 2005).
18. P. Fayet, Nucl. Phys. **B78**, 14 (1974); **B90**, 104 (1975).
19. K. Inoue *et al.*, Prog. Theor. Phys. **67**, 1889 (1982) [erratum: **70**, 330 (1983)]; **71**, 413 (1984);  
R. Flores and M. Sher, Ann. Phys. (NY) **148**, 95 (1983).
20. J.F. Gunion and H.E. Haber, Nucl. Phys. **B272**, 1 (1986) [erratum: **B402**, 567 (1993)].
21. See, *e.g.*, R. Barbieri and G.F. Giudice, Nucl. Phys. **B305**, 63 (1988);  
G.W. Anderson and D.J. Castano, Phys. Lett. **B347**, 300 (1995); Phys. Rev. **D52**, 1693 (1995); Phys. Rev. **D53**, 2403 (1996);  
J.L. Feng, K.T. Matchev, and T. Moroi, Phys. Rev. **D61**, 075005 (2000).
22. S. Dimopoulos and G.F. Giudice, Phys. Lett. **B357**, 573 (1995);  
A. Pomarol and D. Tommasini, Nucl. Phys. **B466**, 3 (1996);  
A.G. Cohen, D.B. Kaplan, and A.E. Nelson, Phys. Lett. **B388**, 588 (1996);  
J.L. Feng, K.T. Matchev, and T. Moroi, Phys. Rev. Lett. **84**, 2322 (2000).

23. M. Schmitt, “Supersymmetry Part II (Experiment),” immediately following, in the section on Reviews, Tables, and Plots in this *Review*. See also *Particle Listings: Other Searches—Supersymmetric Particles* in this *Review*.
24. See, *e.g.*, F. Gabbiani *et al.*, Nucl. Phys. **B477**, 321 (1996).
25. For a recent review and references to the original literature, see: A. Masiero, and O. Vives, New J. Phys. **4**, 4.1 (2002).
26. J. Erler and P. Langacker, “Electroweak Model and Constraints on New Physics,” in the section on Reviews, Tables, and Plots in this *Review*.
27. P.H. Chankowski and S. Pokorski, in *Perspectives on Supersymmetry*, edited by G.L. Kane (World Scientific, Singapore, 1998) pp. 402–422.
28. A. Dobado, M.J. Herrero, and S. Penaranda, Eur. Phys. J. **C7**, 313 (1999); **C12**, 673 (2000); **C17**, 487 (2000).
29. J. Erler and D.M. Pierce, Nucl. Phys. **B526**, 53 (1998); G. Altarelli, F. Caravaglios, G.F. Giudice, P. Gambino and G. Ridolfi, JHEP **0106**, 018 (2001); J.R. Ellis, S. Heinemeyer, K.A. Olive and G. Weiglein, JHEP **0502**, 013 (2005); J. Haestier, S. Heinemeyer, D. Stockinger and G. Weiglein, JHEP **0512**, 027 (2005).
30. M. Davier and W.J. Marciano, Ann. Rev. Nucl. Part. Sci. **54** (2004) 115; M. Passera, J. Phys. **G31**, R75 (2005); M. Davier, A. Hocker and Z. Zhang, [hep-ph/0507078](http://arxiv.org/abs/hep-ph/0507078).
31. For a recent review and references to the literature, see T. Hurth, Rev. Mod. Phys. **75**, 1159 (2003).
32. See, *e.g.*, M. Ciuchini, E. Franco, A. Masiero and L. Silvestrini, Phys. Rev. **D67**, 075016 (2003).
33. U. Chattopadhyay and P. Nath, Phys. Rev. **D66**, 093001 (2002);



- S.P. Martin and J.D. Wells, Phys. Rev. **D67**, 015002 (2003).
34. P. Fayet, Phys. Lett. **69B**, 489 (1977);  
G. Farrar and P. Fayet, Phys. Lett. **76B**, 575 (1978).
  35. J. Ellis, J.S. Hagelin, D.V. Nanopoulos, K.A. Olive and M. Srednicki, Nucl. Phys. **B238**, 453 (1984).
  36. S. Raby, Phys. Lett. **B422**, 158 (1998);  
S. Raby and K. Tobe, Nucl. Phys. **B539**, 3 (1999);  
A. Mafi and S. Raby, Phys. Rev. **D62**, 035003 (2000).
  37. A.R. Liddle and D.H. Lyth, Phys. Reports **213**, 1 (1993).
  38. M. Drees and G. Gerbier, “Dark Matter,” in the section on Reviews, Tables, and Plots in this *Review*.
  39. P. Fayet, Phys. Lett. **84B**, 421 (1979); Phys. Lett. **86B**, 272 (1979).
  40. P. van Nieuwenhuizen, Phys. Reports **68**, 189 (1981).
  41. S. Deser and B. Zumino, Phys. Rev. Lett. **38**, 1433 (1977).
  42. A.H. Chamseddine, R. Arnowitt and P. Nath, Phys. Rev. Lett. **49**, 970 (1982);  
R. Barbieri, S. Ferrara and C.A. Savoy, Phys. Lett. **119B**, 343 (1982);  
H.-P. Nilles, M. Srednicki and D. Wyler, Phys. Lett. **120B**, 346 (1983); **124B**, 337 (1983) 337;  
E. Cremmer, P. Fayet and L. Girardello, Phys. Lett. **122B**, 41 (1983);  
L. Ibáñez, Nucl. Phys. **B218**, 514 (1982);  
L. Alvarez-Gaumé, J. Polchinski and M.B. Wise, Nucl. Phys. **B221**, 495 (1983).
  43. L.J. Hall, J. Lykken, and S. Weinberg, Phys. Rev. **D27**, 2359 (1983).
  44. S.K. Soni and H.A. Weldon, Phys. Lett. **126B**, 215 (1983);  
Y. Kawamura, H. Murayama, and M. Yamaguchi, Phys. Rev. **D51**, 1337 (1995).
  45. A.B. Lahanas and D.V. Nanopoulos, Phys. Reports **145**, 1 (1987).

46. M. Dine and A.E. Nelson, Phys. Rev. **D48**, 1277 (1993);  
M. Dine, A.E. Nelson, and Y. Shirman, Phys. Rev. **D51**,  
1362 (1995);  
M. Dine *et al.*, Phys. Rev. **D53**, 2658 (1996).
47. G.F. Giudice, and R. Rattazzi, Phys. Reports **322**, 419  
(1999).
48. J. Polchinski, *String Theory, Volumes I and II* (Cam-  
bridge University Press, Cambridge, UK, 1998).
49. Pedagogical lectures describing such mechanisms can be  
found in: M. Quiros, in *Particle Physics and Cosmology:  
The Quest for Physics Beyond the Standard Model(s)*,  
Proceedings of the 2002 Theoretical Advanced Study  
Institute in Elementary Particle Physics (TASI 2002),  
edited by H.E. Haber and A.E. Nelson (World Scientific,  
Singapore, 2004) pp. 549–601;  
C. Csaki, in *ibid.*, pp. 605–698.
50. See, *e.g.*, G.F. Giudice and J.D. Wells, “Extra Dimen-  
sions,” in the section on Reviews, Tables, and Plots in  
this *Review*.
51. These ideas are reviewed in: V.A. Rubakov, Phys. Usp.  
**44**, 871 (2001);  
J. Hewett and M. Spiropulu, Ann. Rev. Nucl. Part. Sci.  
**52**, 397 (2002).
52. L. Randall and R. Sundrum, Nucl. Phys. **B557**, 79  
(1999).
53. Z. Chacko, M.A. Luty, and E. Ponton, JHEP **0007**, 036  
(2000);  
D.E. Kaplan, G.D. Kribs, and M. Schmaltz, Phys. Rev.  
**D62**, 035010 (2000);  
Z. Chacko *et al.*, JHEP **0001**, 003 (2000).
54. J. Scherk and J.H. Schwarz, Phys. Lett. **82B**, 60 (1979);  
Nucl. Phys. **B153**, 61 (1979).
55. See, *e.g.*, R. Barbieri, L.J. Hall, and Y. Nomura, Phys.  
Rev. **D66**, 045025 (2002); Nucl. Phys. **B624**, 63 (2002).
56. K. Cheung and W. Y. Keung, Phys. Rev. **D71**, 015015  
(2005);

- P. Gambino, G.F. Giudice and P. Slavich, Nucl. Phys. **B726**, 35 (2005).
57. H.E. Haber, in *Recent Directions in Particle Theory, Proceedings of the 1992 Theoretical Advanced Study Institute in Particle Physics*, edited by J. Harvey and J. Polchinski (World Scientific, Singapore, 1993) pp. 589–686.
  58. S. Dimopoulos and D. Sutter, Nucl. Phys. **B452**, 496 (1995);  
D.W. Sutter, Stanford Ph. D. thesis, [hep-ph/9704390](http://arxiv.org/abs/hep-ph/9704390).
  59. H.E. Haber, Nucl. Phys. B (Proc. Suppl.) **62A-C**, 469 (1998).
  60. R.M. Barnett, J.F. Gunion and H.E. Haber, Phys. Lett. **B315**, 349 (1993);  
H. Baer, X. Tata and J. Woodside, Phys. Rev. **D41**, 906 (1990).
  61. S.M. Bilenky, E.Kh. Khristova and N.P. Nedelcheva, Phys. Lett. **B161**, 397 (1985); Bulg. J. Phys. **13**, 283 (1986);  
G. Moortgat-Pick and H. Fraas, Eur. Phys. J. **C25**, 189 (2002).
  62. For further details, see *e.g.* Appendix C of Ref. [17] and Appendix A of Ref. [20].
  63. J.L. Kneur and G. Moultaka, Phys. Rev. **D59**, 015005 (1999).
  64. S.Y. Choi, A. Djouadi, M. Guchait, J. Kalinowski, H.S. Song and P.M. Zerwas, Eur. Phys. J. **C14**, 535 (2000).
  65. Roger A. Horn and Charles R. Johnson, *Matrix Analysis*, (Cambridge University Press, Cambridge, UK, 1985).
  66. S.Y. Choi, J. Kalinowski, G. Moortgat-Pick and P.M. Zerwas, Eur. Phys. J. **C22**, 563 (2001); **C23**, 769 (2002).
  67. G.J. Gounaris, C. Le Mouel and P.I. Porfyriadis, Phys. Rev. **D65**, 035002 (2002);  
G.J. Gounaris and C. Le Mouel, Phys. Rev. **D66**, 055007 (2002).
  68. T. Takagi, Japan J. Math. **1**, 83 (1925).

69. M.M. El Kheishen, A.A. Aboshousha and A.A. Shafik, Phys. Rev. **D45**, 4345 (1992);  
M. Guchait, Z. Phys. **C57**, 157 (1993) [Erratum: **C61**, 178 (1994)].
70. J. Ellis and S. Rudaz, Phys. Lett. **128B**, 248 (1983);  
F. Browning, D. Chang and W.Y. Keung, Phys. Rev. **D64**, 015010 (2001);  
A. Bartl, S. Hesselbach, K. Hidaka, T. Kernreiter and W. Porod, Phys. Lett. **B573**, 153 (2003); Phys. Rev. **D70**, 035003 (2004).
71. D.M. Pierce *et al.*, Nucl. Phys. **B491**, 3 (1997).
72. J.A. Aguilar-Saavedra *et al.*, Eur. Phys. J. **C46**, 43 (2006).
73. J.F. Gunion *et al.*, *The Higgs Hunter's Guide* (Perseus Publishing, Cambridge, MA, 1990);  
M. Carena and H.E. Haber, Prog. Part. Nucl. Phys. **50**, 63 (2003).
74. H.E. Haber and R. Hempfling, Phys. Rev. Lett. **66**, 1815 (1991);  
Y. Okada, M. Yamaguchi, and T. Yanagida, Prog. Theor. Phys. **85**, 1 (1991);  
J. Ellis, G. Ridolfi, and F. Zwirner, Phys. Lett. **B257**, 83 (1991).
75. See, *e.g.*, B.C. Allanach, A. Djouadi, J.L. Kneur, W. Porod and P. Slavich, JHEP **0409**, 044 (2004);  
G. Degrassi, S. Heinemeyer, W. Hollik, P. Slavich and G. Weiglein, Eur. Phys. J. **C28**, 133 (2003), and references contained therein.
76. A. Pilaftsis and C.E.M. Wagner, Nucl. Phys. **B553**, 3 (1999);  
D.A. Demir, Phys. Rev. **D60**, 055006 (1999);  
S.Y. Choi, M. Drees and J.S. Lee, Phys. Lett. **B481**, 57 (2000);  
M. Carena *et al.*, Nucl. Phys. **B586**, 92 (2000); Phys. Lett. **B495**, 155 (2000); Nucl. Phys. **B625**, 345 (2002).

77. A summary of MSSM Higgs mass limits can be found in P. Igo-Kemenes, “Higgs Boson Searches,” in the section on Reviews, Tables, and Plots in this *Review*.
78. S. Khalil, Int. J. Mod. Phys. **A18**, 1697 (2003).
79. W. Fischler, S. Paban, and S. Thomas, Phys. Lett. **B289**, 373 (1992);  
S.M. Barr, Int. J. Mod. Phys. **A8**, 209 (1993);  
T. Ibrahim and P. Nath, Phys. Rev. **D58**, 111301 (1998) [erratum: **D60**, 099902 (1999)];  
M. Brhlik, G.J. Good, and G.L. Kane, Phys. Rev. **D59**, 115004 (1999);  
V.D. Barger, T. Falk, T. Han, J. Jiang, T. Li and T. Plehn, Phys. Rev. **D64**, 056007 (2001);  
S. Abel, S. Khalil and O. Lebedev, Nucl. Phys. **B606**, 151 (2001);  
K.A. Olive, M. Pospelov, A. Ritz and Y. Santoso, Phys. Rev. **D72**, 075001 (2005);  
G.F. Giudice and A. Romanino, Phys. Lett. **B634**, 307 (2006).
80. A. Masiero and L. Silvestrini, in *Perspectives on Supersymmetry*, edited by G.L. Kane (World Scientific, Singapore, 1998) pp. 423–441.
81. M. Pospelov and A. Ritz, Annals Phys. **318**, 119 (2005).
82. H. Georgi, Phys. Lett. **169B**, 231 (1986);  
L.J. Hall, V.A. Kostelecky, and S. Raby, Nucl. Phys. **B267**, 415 (1986).
83. Y. Nir and N. Seiberg, Phys. Lett. **B309**, 337 (1993);  
S. Dimopoulos, G.F. Giudice, and N. Tetradis, Nucl. Phys. **B454**, 59 (1995);  
G.F. Giudice *et al.*, JHEP **12**, 027 (1998);  
J.L. Feng and T. Moroi, Phys. Rev. **D61**, 095004 (2000).
84. L.E. Ibáñez and G.G. Ross, Phys. Lett. **B110**, 215 (1982).
85. J.F. Gunion and H.E. Haber, Phys. Rev. **D37**, 2515 (1988).
86. J.L. Feng *et al.*, Phys. Rev. Lett. **83**, 1731 (1999);  
T. Gherghetta, G.F. Giudice, and J.D. Wells, Nucl. Phys.

- B559**, 27 (1999);  
 J.F. Gunion and S. Mrenna, Phys. Rev. **D62**, 015002 (2000).
87. For a number of recent attempts to resolve the tachyonic slepton problem, see *e.g.*, I. Jack, D.R.T. Jones and R. Wild, Phys. Lett. **B535**, 193 (2002);  
 B. Murakami and J.D. Wells, Phys. Rev. **D68**, 035006 (2003);  
 R. Kitano, G.D. Kribs and H. Murayama, Phys. Rev. **D70**, 035001 (2004);  
 R. Hodgson, I. Jack, D.R.T. Jones and G.G. Ross, Nucl. Phys. **B728**, 192 (2005);  
 Q. Shafi and Z. Tavartkiladze, [hep-ph/0408156](http://arxiv.org/abs/hep-ph/0408156).
88. M. Drees and S.P. Martin, in *Electroweak Symmetry Breaking and New Physics at the TeV Scale*, edited by T. Barklow *et al.* (World Scientific, Singapore, 1996) pp. 146–215.
89. M. Carena *et al.*, Nucl. Phys. **B426**, 269 (1994).
90. J.R. Ellis, K.A. Olive, Y. Santoso and V.C. Spanos, Phys. Lett. **B565**, 176 (2003);  
 H. Baer and C. Balazs, JCAP **0305**, 006 (2003).
91. L.E. Ibáñez and D. Lüst, Nucl. Phys. **B382**, 305 (1992);  
 B. de Carlos, J.A. Casas, and C. Muñoz, Phys. Lett. **B299**, 234 (1993);  
 V. Kaplunovsky and J. Louis, Phys. Lett. **B306**, 269 (1993);  
 A. Brignole, L.E. Ibáñez, and C. Muñoz, Nucl. Phys. **B422**, 125 (1994) [erratum: **B436**, 747 (1995)].
92. B.C. Allanach *et al.*, Eur. Phys. J. **C25**, 113 (2002).
93. M. Battaglia *et al.*, Eur. Phys. J. **C33**, 273 (2004).
94. For a review and guide to the literature, see J.F. Gunion and H.E. Haber, in *Perspectives on Supersymmetry*, edited by G.L. Kane (World Scientific, Singapore, 1998) pp. 235–255.
95. S. Raby, “Grand Unified Theories,” in the section on Reviews, Tables, and Plots in this *Review*.

96. M.B. Einhorn and D.R.T. Jones, Nucl. Phys. **B196**, 475 (1982).
97. For a review, see R.N. Mohapatra, in *Particle Physics 1999*, ICTP Summer School in Particle Physics, Trieste, Italy, 21 June—9 July, 1999, edited by G. Senjanovic and A.Yu. Smirnov (World Scientific, Singapore, 2000) pp. 336–394;  
W.J. Marciano and G. Senjanovic, Phys. Rev. **D25**, 3092 (1982).
98. D.M. Ghilencea and G.G. Ross, Nucl. Phys. **B606**, 101 (2001).
99. S. Pokorski, Acta Phys. Polon. **B30**, 1759 (1999).
100. H. Arason *et al.*, Phys. Rev. Lett. **67**, 2933 (1991);  
Phys. Rev. **D46**, 3945 (1992);  
V. Barger, M.S. Berger, and P. Ohmann, Phys. Rev. **D47**, 1093 (1993);  
M. Carena, S. Pokorski, and C.E.M. Wagner, Nucl. Phys. **B406**, 59 (1993);  
P. Langacker and N. Polonsky, Phys. Rev. **D49**, 1454 (1994).
101. M. Olechowski and S. Pokorski, Phys. Lett. **B214**, 393 (1988);  
B. Ananthanarayan, G. Lazarides, and Q. Shafi, Phys. Rev. **D44**, 1613 (1991);  
S. Dimopoulos, L.J. Hall, and S. Raby, Phys. Rev. Lett. **68**, 1984 (1992);  
L.J. Hall, R. Rattazzi, and U. Sarid, Phys. Rev. **D50**, 7048 (1994);  
R. Rattazzi and U. Sarid, Phys. Rev. **D53**, 1553 (1996).
102. See the section on neutrinos in “Particle Listings—Leptons” in this *Review*.
103. For a recent review of neutrino masses in supersymmetry, see B. Mukhopadhyaya, Proc. Indian National Science Academy **A70**, 239 (2004).
104. P. Minkowski, Phys. Lett. **67B**, 421 (1977);  
M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, edited by D. Freedman and P. van Nieuwenhuizen

- (North Holland, Amsterdam, 1979) p. 315;  
T. Yanagida, in *Proceedings of the Workshop on Unified Theory and Baryon Number in the Universe*, edited by O. Sawada and A. Sugamoto (KEK, Tsukuba, Japan, 1979);  
R. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980); Phys. Rev. **D23**, 165 (1981).
105. J. Hisano, T. Moroi, K. Tobe, M. Yamaguchi and T. Yanagida, Phys. Lett. **B357**, 579 (1995);  
J. Hisano, T. Moroi, K. Tobe and M. Yamaguchi, Phys. Rev. **D53**, 2442 (1996);  
J. Ellis, J. Hisano, M. Raidal and Y. Shimizu, Phys. Rev. **D66**, 115013 (2002);  
A. Masiero, S.K. Vempati and O. Vives, New J. Phys. **6**, 202 (2004).
106. Y. Grossman and H.E. Haber, Phys. Rev. Lett. **78**, 3438 (1997).
107. For a recent review and references to the original literature, see M. Chemtob, Prog. Part. Nucl. Phys. **54**, 71 (2005);  
R. Barbier *et al.*, Phys. Reports **420**, 1 (2005).
108. H. Dreiner, in *Perspectives on Supersymmetry*, edited by G.L. Kane (World Scientific, Singapore, 1998) pp. 462–479.
109. B.C. Allanach, A. Dedes and H.K. Dreiner, Phys. Rev. **D60**, 075014 (1999).
110. L.E. Ibáñez and G.G. Ross, Nucl. Phys. **B368**, 3 (1992);  
L.E. Ibáñez, Nucl. Phys. **B398**, 301 (1993).
111. For a review, see J.C. Romao, Nucl. Phys. Proc. Suppl. **81**, 231 (2000).
112. R.N. Mohapatra, Phys. Rev. **D34**, 3457 (1986);  
K.S. Babu and R.N. Mohapatra, Phys. Rev. Lett. **75**, 2276 (1995);  
M. Hirsch, H.V. Klapdor-Kleingrothaus, and S.G. Kovalenko, Phys. Rev. Lett. **75**, 17 (1995); Phys. Rev. **D53**, 1329 (1996).



113. M. Hirsch, H.V. Klapdor-Kleingrothaus, and S.G. Kovalenko, Phys. Lett. **B398**, 311 (1997).
114. Y. Grossman and H.E. Haber, Phys. Rev. **D59**, 093008 (1999).
115. S. Dimopoulos and L.J. Hall, Phys. Lett. **B207**, 210 (1988);  
J. Kalinowski *et al.*, Phys. Lett. **B406**, 314 (1997);  
J. Erler, J.L. Feng, and N. Polonsky, Phys. Rev. Lett. **78**, 3063 (1997).
116. H.K. Dreiner, P. Richardson and M.H. Seymour, Phys. Rev. **D63**, 055008 (2001).
117. See, *e.g.*, M. Hirsch *et al.*, Phys. Rev. **D62**, 113008 (2000) [erratum: **D65**, 119901 (2002)];  
A. Abada, G. Bhattacharyya and M. Losada, Phys. Rev. **D66**, 071701 (2002);  
M.A. Diaz, *et al.*, Phys. Rev. **D68**, 013009 (2003);  
M. Hirsch and J.W.F. Valle, New J. Phys. **6**, 76 (2004).
118. G.F. Giudice and A. Masiero, Phys. Lett. **B206**, 480 (1988).
119. See, *e.g.*, U. Ellwanger, M. Rausch de Traubenberg, and C.A. Savoy, Nucl. Phys. **B492**, 21 (1997);  
U. Ellwanger and C. Hugonie, Eur. J. Phys. **C25**, 297 (2002), and references contained therein.
120. J.R. Espinosa and M. Quiros, Phys. Rev. Lett. **81**, 516 (1998);  
K.S. Babu, I. Gogoladze and C. Kolda, [hep-ph/0410085](http://arxiv.org/abs/hep-ph/0410085).
121. R. Mahbubani, [hep-ph/0408096](http://arxiv.org/abs/hep-ph/0408096).
122. A. Birkedal, Z. Chacko and Y. Nomura, Phys. Rev. **D71**, 015006 (2005).
123. J. Alcaraz *et al.* [The LEP Collaborations ALEPH, DELPHI, L3, OPAL, and the LEP Electroweak Working Group], [hep-ex/0511027](http://arxiv.org/abs/hep-ex/0511027).
124. J.L. Hewett and T.G. Rizzo, Phys. Reports **183**, 193 (1989).
125. K.R. Dienes, Phys. Reports **287**, 447 (1997).

## SUPERSYMMETRY, PART II (EXPERIMENT)

Revised September, 2003 by M. Schmitt (Northwestern University)

**II.1. Introduction:** The theoretical strong points of supersymmetry (SUSY) have motivated many searches for supersymmetric particles. Many of these have been based on the canonical missing-energy signature caused by the escape of weakly-interacting LSP's ('lightest supersymmetric particles'). Other scenarios also have been investigated, widening the range of topologies and experimental signatures in which new physics might be found. Unfortunately, no convincing evidence for the production of supersymmetric particles has been found.

Theoretical aspects of supersymmetry have been covered in Part I of this review by H.E. Haber (see also Ref. 1, 2); we use his notations and terminology.

**II.2. Common supersymmetry scenarios:** In the 'canonical' scenario [1], supersymmetric particles are pair-produced and decay directly or via cascades to the LSP. It follows that there are always at least two LSP's per event. If  $R$ -parity, the quantum number which distinguishes SM and SUSY particles, is conserved, the LSP is stable. For most typical choices of model parameters, the lightest neutralino is the LSP. Since the neutralino is neutral and colorless, interacting only weakly with matter, it will escape detection, giving signal events the characteristic appearance of "missing energy." In  $e^+e^-$  machines, the total visible energy and total visible momentum can be well measured. Since the electron beam energy has a very small spread, the missing energy ( $E^{\text{miss}} = \sqrt{s} - E^{\text{vis}}$ ) and the missing momentum ( $\vec{p}^{\text{miss}} = -\vec{p}^{\text{vis}}$ ) are well correlated with the net energy and momentum of the LSP's. In

proton colliders, the distribution of the energy and longitudinal momentum of the partons (quarks and gluons inside the (anti-)protons) is very broad, so in practice only the transverse momentum is useful. It is calculated from the vector sum of energy deposits registered in the calorimetry and is called “missing transverse energy” ( $\cancel{E}_T$ ). Collimated jets, isolated leptons or photons, and appropriate kinematic and topological cuts provide additional handles for reducing backgrounds.

The conservation of  $R$ -parity is not required in supersymmetry, however, and in some searches it is assumed that supersymmetric particles decay via interactions which violate  $R$ -parity (RPV). For the most part the production of superpartners is unchanged, but the missing-energy signature is lost. Depending on the choice of the  $R$ -parity-violating interaction, SUSY events are characterized by an excess of leptons or hadronic jets, and in many cases it is relatively easy to suppress SM backgrounds [3]. A distinction is made between “indirect” RPV, in which the LSP decays close to the interaction point but no other decays are modified, and “direct” RPV, in which the supersymmetric particles decay to SM particles, producing no LSP’s. The LSP’s themselves provide a visible signal by virtue of their decay to ordinary fermions. Note that the cosmological constraint which requires stable LSP’s to be charge and color neutral no longer applies when there  $R$ -parity is violated.

In models assuming gauge-mediated supersymmetry breaking (GMSB) [4], the gravitino,  $\tilde{g}_{3/2}$ , is a weakly-interacting fermion with a mass so small that it can be neglected when considering the event kinematics. It is the LSP, and the lightest neutralino,  $\tilde{\chi}_1^0$ , decays to it radiatively, possibly with a long lifetime. With few exceptions the decays and production of

other superpartners are the same as in the canonical scenario, so when the neutralino lifetime is not too long, the event topologies are augmented by the presence of energetic and isolated photons. If the lifetime is so long that the neutralino decays outside the detector, the event topologies are the same as in the canonical scenario. In some variants of this theory the right-sleptons are lighter than the lightest neutralino, and they decay to a lepton and a gravitino. The most important case of this type is the channel  $\tilde{\tau}_R \rightarrow \tau \tilde{G}$ . The lifetime of the  $\tilde{\tau}_R$  can vary over a wide range depending on model parameters, leading to new exotic signatures, including quasi-stable, heavily ionizing charged particles.

Finally, there is another phenomenologically important scenario in which the gluino  $\tilde{g}$  is assumed to be relatively light ( $M_{\tilde{g}} < 5 \text{ GeV}/c^2$ ). Experimental evidence does not support the hypothesis, however, as discussed further in the review by H. Murayama.

**II.3. Experimental issues:** When given no signal for supersymmetric particles, experimenters are obliged to derive limits on their production. The most general formulation of supersymmetry is so flexible that few universal bounds can be obtained. Often more restricted forms of the theory are evoked for which predictions are more definite. The most popular of these is minimal supergravity ('mSUGRA'). As explained in Part I of this review, parameter freedom is drastically reduced by requiring related parameters to be equal at the unification scale,  $M_X$ . Thus, the gaugino masses are equal with value  $m_{1/2}$ , and the slepton, squark, and Higgs masses depend on a *common* scalar mass parameter,  $m_0$ . In the individual experimental analyses, only some of these assumptions are necessary.

For example, the gluino and squark searches at proton machines constrain mainly  $M_3$  and a scalar mass parameter  $m_0$  for the squark masses, while the chargino, neutralino, and slepton searches at  $e^+e^-$  colliders constrain  $M_2$  and a scalar mass parameter  $m_0$  for the slepton masses. In addition, results from the Higgs searches can be used to constrain  $m_{1/2}$  and  $m_0$  as a function of  $\tan\beta$ . (The full analysis involves large radiative corrections coming from squark mixing, which is where the dependence on  $m_{1/2}$  and  $m_0$  enter.) In the mSUGRA framework, all the scalar mass parameters  $m_0$  are the same and the three gaugino mass parameters are proportional to  $m_{1/2}$ , so limits from squarks, sleptons, charginos, gluinos, and Higgs all can be used together to constrain the parameter space. A slightly less constrained model allows the Higgs sector to be independent of the sfermion sector, while still requiring that the scalar mass parameter  $m_0$  is the same for sleptons and squarks and that the gaugino mass parameter  $m_{1/2}$  is the same for charginos, neutralinos and gluinos. This model is called the ‘constrained MSSM’ (cMSSM) [5,6].

While the mSUGRA framework is convenient, it is based on several highly specific theoretical assumptions, so limits presented in this framework cannot easily be applied to other supersymmetric models. It has been possible in some instances to reduce the model dependence of experimental results by combining several searches. When model-independent results are impossible, the underlying assumptions and their consequences are (or should be) carefully delineated.

In the analysis of data from hadron collider experiments, the experimenter considers several supersymmetric processes simultaneously. In contrast to experiments at  $e^+e^-$  colliders, it does not make sense to talk about one process at a time due to

the very broad mass range spanned. This makes the utilization of some sort of organizing device, such as a constrained version of the MSSM, practically unavoidable.

#### ***II.4. Supersymmetry searches at $e^+e^-$ colliders:***

The large electron-positron collider (LEP) at CERN ran at energies ranging from the  $Z$  peak up to  $\sqrt{s} = 209 \text{ GeV}/c^2$ . Each experiment (ALEPH, DELPHI, L3, OPAL) accumulated large data sets at a series of energies, as detailed in [7]. For the limits discussed here, the most relevant data samples include  $180 \text{ pb}^{-1}$  at  $189 \text{ GeV}/c^2$ , and  $220 \text{ pb}^{-1}$  at higher energies, of which  $140 \text{ pb}^{-1}$  was delivered above  $206 \text{ GeV}/c^2$ . Since the last edition of this review, several of the searches at the highest energies have been finalized.

Running at the  $Z$  pole, the LEP experiments and SLD at SLAC excluded many supersymmetric particles up to about half the  $Z$  mass. These limits come mainly from the comparison of the measured  $Z$  widths to SM expectations, and are relatively insensitive to the details of SUSY particle decays [8]. The data taken at higher energies allow much stronger limits to be set, although the complex interplay of masses, cross sections, and branching ratios allow for a few exceptions to simple general limits.

The main signals come from SUSY particles with charge, weak isospin, or large Yukawa couplings. The gauge fermions (charginos and neutralinos) generally are produced with large cross sections, while the scalar particles (sleptons and squarks) are suppressed near threshold by kinematic factors.

The various SUSY particles considered at LEP typically decay directly to SM particles and LSP's, so signatures consist of some combination of jets, leptons, possibly photons, and missing energy. Consequently the search criteria are geared toward a

few distinct topologies. Although they may be optimized for one specific signal, they are often efficient for others. For example, acoplanar jets are expected in both  $\tilde{t}_1\tilde{t}_1^*$  and  $\tilde{\chi}_1^0\tilde{\chi}_2^0$  production, and acoplanar leptons for both  $\tilde{\ell}^+\tilde{\ell}^-$  and  $\tilde{\chi}^+\tilde{\chi}^-$ .

Backgrounds come mainly from three sources. First, there are the so-called ‘two-photon interactions,’ in which the beam electrons emit photons which combine to produce a low mass hadronic or leptonic system leaving little visible energy in the detector. Since the electrons are seldom deflected through large angles,  $p_T^{\text{miss}}$  is low. Second, there is difermion production, usually accompanied by large initial-state radiation induced by the  $Z$  pole, which gives events that are well balanced with respect to the beam direction. Finally, there is four-fermion production through states with one or two resonating bosons ( $W^+W^-$ ,  $ZZ$ ,  $We\nu$ ,  $Ze^+e^-$ , etc.) which can give events with large  $E^{\text{miss}}$  and  $p_T^{\text{miss}}$  due to neutrinos and electrons lost down the beam pipe.

In the canonical case,  $E^{\text{miss}}$  and  $p_T^{\text{miss}}$  are large enough to eliminate most of these backgrounds. The  $e^+e^-$  initial state is well defined so searches utilize both transverse and longitudinal momentum components. It is possible to measure the missing mass ( $M_{\text{miss}} = \{(\sqrt{s} - E_{\text{vis}})^2 - \vec{p}_{\text{vis}}^2\}^{1/2}$ ) which is small if  $p_T^{\text{miss}}$  is caused by a single neutrino or an undetected electron or photon, and large when there are two massive LSP’s. The four-fermion processes cannot be entirely eliminated, however, and a non-negligible irreducible background is expected. Fortunately, the uncertainties for these backgrounds are not large.

High efficiencies are easily achieved when the mass of the LSP ( $M_{\text{LSP}}$ ) is less than the parent particle ( $M_{\text{parent}}$ ) by at least 10 GeV/ $c^2$  and greater than about 10 GeV/ $c^2$ . Difficulties arise when the mass difference  $\Delta M = M_{\text{parent}} - M_{\text{LSP}}$  is smaller

than  $10 \text{ GeV}/c^2$  as the signal resembles background from two-photon interactions. A very light LSP is challenging also since, kinematically speaking, it plays a role similar to a neutrino, so that, for example, a signal for charginos of mass  $\sim 80 \text{ GeV}/c^2$  is difficult to distinguish from the production of  $W^+W^-$  pairs. The lower signal efficiency obtained in these two extreme cases has been offset by the large integrated luminosities delivered, so mass limits are not degraded.

***Charginos and Neutralinos:*** The phenomenology of charginos and neutralinos depends on their field content: they tend to be ‘gaugino-like’ (for  $M_2 \ll |\mu|$ ) or ‘higgsino-like’ ( $|\mu| \ll M_2$ ), with a ‘mixed’ field content available only for a relatively small region of parameter space. The cross section for gauginos varies with the masses of sleptons exchanged in the  $t$ -channel. In particular, chargino production can be suppressed by more than an order of magnitude for particular values of  $M_{\tilde{\nu}_e}$ . The gaugino branching ratios also depend on the sfermion sector. When the sfermion masses are larger than  $\sim 200 \text{ GeV}/c^2$ , the chargino and neutralino branching ratios are close to those of the  $W$  and  $Z$  bosons. Enhancements of leptonic branching ratios are important when sleptons are light. Light squarks are excluded by hadron collider experiments and are not considered. Cross sections and branching ratios for higgsinos are, in contrast, insensitive to the masses of the sfermions.

In the gaugino-like region, the lightest chargino mass is driven by  $M_2$  and the lightest neutralino mass by  $M_1$ . For many popular models (such as ‘supergravity’),  $M_1$  and  $M_2$  unify at a GUT scale, with  $M_1 \approx M_2/2$  at the electroweak scale. Consequently, the mass difference  $\Delta M = M_{\tilde{\chi}^\pm} - M_{\tilde{\chi}_1^0}$  is not very small and selection efficiencies are high. However, as explained




in the theoretical section of this review, this unification scheme is not required by Supersymmetry, and it is important to consider both  $M_1 \approx M_2$  and  $M_1 \ll M_2$ . In the higgsino-like region, chargino and neutralino masses are all close to  $|\mu|$ , and hence, small mass differences of order  $5 \text{ GeV}/c^2$  are typical. In the mixed region of moderate, negative  $\mu$ ,  $\Delta M \approx M_W$ , and cuts designed to reject  $W$  background lead to lower efficiencies.

Chargino masses have been excluded up to  $103 \text{ GeV}/c^2$ . However, this limit can be degraded when the sneutrino is lighter than  $\sim 200 \text{ GeV}/c^2$ . Thanks to the large integrated luminosity and the combination of four experiments [7], the impact for  $M_{\tilde{\nu}_e} \gtrsim 100 \text{ GeV}/c^2$  is less than a  $\text{GeV}/c^2$ . The limit is also weakened when the mass difference is small ( $\Delta M = M_{\tilde{\chi}^\pm} - M_{\tilde{\chi}_1^0} \lesssim 3 \text{ GeV}/c^2$ ), as in the higgsino region; however, in this case the associated production of neutralino pairs  $\tilde{\chi}_1^0 \tilde{\chi}_2^0$  is large and the problem of small mass differences ( $M_{\tilde{\chi}_2^0} - M_{\tilde{\chi}_1^0}$ ) less severe. Experimental sensitivity now extends down to mass differences of  $3 \text{ GeV}/c^2$ , corresponding to  $M_2$  above  $2 \text{ TeV}/c^2$ .

For a summary of the interplay of chargino field content and sfermion masses, see Fig. 1.

The possibility of extremely small mass differences has been raised in several theoretical papers which propose models rather different from supergravity [9]. The DELPHI Collaboration was the first to engineer searches to cover this scenario [10], and other collaborations have followed suit [11]. For  $\Delta M \sim 1 \text{ GeV}/c^2$ , the signal can be distinguished from two-photon background on the basis of isolated photons detected at low angles: hard initial-state radiation sometimes accompanies the signal process but is absent for the background. For  $\Delta M \sim 0.2 \text{ GeV}/c^2$ , the chargino acquires a non-negligible



**Figure 1:** Heuristic diagram of the interplay of chargino field content and sfermion masses. See full-color version on color pages at end of book.

lifetime and decays at a significant distance from the interaction point, producing tracks which do not extrapolate back to the interaction point. When  $\Delta M < m_\pi$ , the lifetime is so long that the chargino appears as a heavily ionizing particle which exits the tracking detector before decaying. The bounds on the chargino mass vary from 68 to 88  $\text{GeV}/c^2$  depending on the assumed sneutrino mass; the limit is 92  $\text{GeV}/c^2$  from the combination of the four LEP experiments when  $M_{\tilde{\nu}_e} > 500 \text{ GeV}/c^2$  [7].

The limits from chargino and neutralino production are most often used to constrain  $M_2$  and  $\mu$  for fixed  $\tan\beta$ . For large  $|\mu|$  (the gaugino case), chargino bounds limit  $M_2$ , and vice versa (the Higgsino case). When  $\tan\beta$  is not large, the region of parameter space with  $\mu < 0$  and  $|\mu| \sim M_2$  corresponds to ‘mixed’ field content, and the limits on  $M_2$  and  $|\mu|$  are relatively modest, especially when electron sneutrinos are light. This is the weak point when inferring an indirect limit on the LSP mass [12].

When the sleptons are light, branching ratios to leptons are enhanced, especially to  $\tau$ ’s via  $\tilde{\tau}$ ’s when there is non-negligible mixing of  $\tilde{\tau}_R$  and  $\tilde{\tau}_L$ . These effects are greatest when the chargino has a large gaugino component. The weakest bounds are found for small negative  $\mu$  and small  $\tan\beta$ , as the cross section is reduced with respect to larger  $|\mu|$ , the impact of  $\tilde{\tau}$  mixing can be large, and the efficiency is not optimal because  $\Delta M$  is large. If sneutrinos are lighter than the chargino, then two-body decays  $\tilde{\chi}^+ \rightarrow \ell^+ \tilde{\nu}$  dominate, and in the ‘corridor’  $0 < M_{\tilde{\chi}^\pm} - M_{\tilde{\nu}} \lesssim 3 \text{ GeV}/c^2$  the acceptance is so low that no direct exclusion is possible [13]. However, in the context of the cMSSM it is possible to cover this region with slepton and neutralino searches.

**Sleptons:** Sleptons and squarks are produced via  $\gamma^*$  and  $Z^*$  exchange. For selectrons there is an important contribution from  $t$ -channel neutralino exchange which generally increases the cross section. Even though the cross section is suppressed near threshold, the large luminosity at LEP has allowed mass limits to be placed close to the kinematic threshold [14]. For equal masses, the cross section for the  $R$  state is smaller than for the  $L$  state, so limits are set conservatively for the production of  $R$ -sleptons only. In grand unified theories the masses of the

$R$  and  $L$  states are linked, and usually the  $R$  state is lighter, especially when  $\tan\beta$  is large. For  $\tilde{\tau}$  sleptons, mixing can be important.

The simplest slepton topology results from  $\tilde{\ell} \rightarrow \ell\tilde{\chi}_1^0$ , though for some particular parameter choices, branching ratios for decays to  $\tilde{\chi}_2^0$  reach a few percent. Combined mass limits have been obtained by the LEP SUSY working group [7]. For  $\tilde{\mu}_R$ , the limit is  $95 \text{ GeV}/c^2$ . The limit for  $\tilde{e}_R$  is  $4 \text{ GeV}/c^2$  higher due to the higher cross section coming from  $\tilde{\chi}^0$  exchange. Since the selection of  $\tau$ 's is relatively difficult, the limit is expected to be lower, and the actual limit is  $86 \text{ GeV}/c^2$ . These limits hold provided the slepton is at least  $10 \text{ GeV}/c^2$  heavier than the neutralino.

Assuming a common scalar mass term  $m_0$ , as in the cMSSM, the masses of the  $R$  and  $L$ -sleptons can be related as a function of  $\tan\beta$ , and one finds  $m_{\tilde{\ell}_L} > m_{\tilde{\ell}_R}$  by a few  $\text{GeV}/c^2$ . Consequently, in associated  $\tilde{e}_L\tilde{e}_R$  production, the special case of a neutralino close in mass to the right-selectron still results in a viable signature: a single energetic electron. ALEPH and L3 have used this to close the gap  $M_{\tilde{e}_R} - M_{\tilde{\chi}} \rightarrow 0$ , and place an absolute limit  $M_{\tilde{e}_R} > 73 \text{ GeV}/c^2$  [15,16].

**Squarks:** Although the Tevatron experiments had placed general limits on squark masses far beyond the reach of LEP, a light top squark ('stop') could still have been found since the interaction eigenstates can mix to give a large splitting between the mass eigenstates. While theoretically less natural, light sbottoms also have been considered. LEP limits on stop and sbottom masses vary with the mixing angle because the cross section does: for  $\theta_{\tilde{t}} = 56^\circ$  and  $\theta_{\tilde{b}} = 67^\circ$  the contribution from  $Z$  exchange is "turned off." In fact the variation in mass limits

is only a couple of  $\text{GeV}/c^2$  due to the large luminosity used for these searches [7].

The stop decay  $\tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$  proceeds through loops, giving a lifetime long enough to allow the top squark to form supersymmetric hadrons which provide a pair of jets and missing energy. The conservative limit is  $M_{\tilde{t}_1} > 95 \text{ GeV}/c^2$ , valid for  $\Delta M > 5 \text{ GeV}/c^2$ . If sneutrinos are light, the decay  $\tilde{t}_1 \rightarrow bl\tilde{\nu}$  dominates, giving two leptons in addition to jets, and the limit is  $96 \text{ GeV}/c^2$ . The same signature obtains when sleptons are light. A somewhat more difficult case comes when  $\tilde{\tau}$ 's are light [17,18,16]. Four-fermion final states ( $b f \bar{f}' \tilde{\chi}_1^0$ ) dominate when charginos are light, a topology covered by ALEPH [18]. Access to very small  $\Delta M$  is possible due to the visibility of the decay products of the  $c$  and  $b$  hadrons [19], in which case conservative limit is  $M_{\tilde{t}_1} > 59 \text{ GeV}/c^2$  is obtained. A comparison to results from the Tevatron is given below.

The electric charge of the sbottoms is smaller than that of stops, so the cross section is considerably lower. The only decay channel considered is  $\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$ . Use of  $b$ -jet tagging helps retain sensitivity: the bound is  $M_{\tilde{b}} > 96 \text{ GeV}/c^2$ . It has been pointed out that very light bottoms squarks ( $M_{\tilde{b}} < 5 \text{ GeV}/c^2$ ) which are decoupled from the  $Z$  are not generally excluded by LEP searches. There is, however, a constraint from a CLEO analysis [20] applicable when the sbottoms always decay semileptonically.

The results from the search for acoplanar jets and missing energy has been interpreted as a limit on the production of generic squarks [21,16,7]. A comparison with Tevatron results is given below.

***The Lightest Neutralino:*** In canonical SUSY scenarios the lightest neutralino leaves no signal in the detector. Nonetheless,

the tight correspondences among the neutralino and chargino masses allow an indirect limit on  $M_{\tilde{\chi}_1^0}$  to be derived [12,22]. The key assumption is that the gaugino mass parameters  $M_1$  and  $M_2$  unify at the GUT scale, which leads to a definite relation between them at the electroweak scale:  $M_1 = \frac{5}{3} \tan^2 \theta_W M_2$ . Assuming slepton masses to be high, the bound on  $M_{\tilde{\chi}_1^0}$  is derived from the results of chargino and neutralino searches, and the limit is  $M_{\tilde{\chi}_1^0} > 39 \text{ GeV}/c^2$  [23,11].

When sleptons are lighter than  $\sim 200 \text{ GeV}/c^2$ , all the effects of light sneutrinos on both the production and decay of charginos and heavier neutralinos must be taken into account. Although the bounds from charginos are weakened, useful additional constraints from slepton and higher-mass neutralino searches rule out the possibility of a light neutralino. A combined limit has been obtained in the cMSSM for any  $\tan \beta$ :  $M_{\tilde{\chi}_1^0} > 37 \text{ GeV}/c^2$  [23]. The results of Higgs searches can be brought into play on the basis of mSUGRA mass relations, to very good effect. They exclude large regions at low  $m_0$  and  $m_{1/2}$  for low  $\tan \beta$ , and strengthen the neutralino bound to  $M_{\tilde{\chi}_1^0} > 45 \text{ GeV}/c^2$  [7].

There is a special case for light neutralinos not excluded by collider experiments: when the  $\tilde{\chi}_1^0$  is a pure bino, the constraints from the invisible  $Z$  width and from the cross section for  $\gamma$ +invisible are ineffective [24]. If one does not assume any relation between  $M_1$  and  $M_2$  then the constraints from chargino searches can be evaded also. Thus a bino of mass  $\mathcal{O}(0.1 \text{ MeV}/c^2)$  is not excluded by collider experiments.

***Gauge-Mediated Scenarios:*** All of the limits above obtain in supergravity models. In models with gauge-mediated supersymmetry breaking (GMSB), however, the phenomenology is rather different, and several interesting new topologies are

expected. They can be classified on the basis of the ‘next-to-lightest supersymmetric particle’ (NLSP) which can be either the lightest neutralino or charged sleptons, in particular,  $\tilde{\tau}_R$ . The gravitino is the LSP, with mass well below a keV.

In the case in which  $\tilde{\chi}_1^0$  is the NLSP, high energy photons are present from the decay  $\tilde{\chi}_1^0 \rightarrow \gamma \tilde{g}_{3/2}$ . They facilitate the separation of signal and background, so for gauginos and sfermions, the resulting limits are very similar to the canonical case. The pair production of  $\tilde{\chi}_1^0$ 's provides an additional search channel consisting of two acollinear photons and missing energy. The mass limit derived is  $99 \text{ GeV}/c^2$ , from ALEPH, assuming the neutralino lifetime is negligible [25]. A more general limit of  $54 \text{ GeV}/c^2$  is set by combining searches for photons which do not point back to the interaction point with indirect limits derived from slepton and chargino searches [26]. Also, single-photon production has been used to constrain the processes  $e^+e^- \rightarrow \tilde{g}_{3/2}\tilde{\chi}_1^0$  and  $e^+e^- \rightarrow \tilde{g}_{3/2}\tilde{g}_{3/2}$ .

When sleptons are the NLSP, there are two possibilities: all three flavors enter more or less equally, or, due to significant mixing, the lightest stau dominates. Considering first three flavors of sleptons, the topology depends strongly on the slepton lifetime which is determined by the scale parameter  $\sqrt{F}$ . For very short lifetimes, the decay  $\tilde{\ell}_R \rightarrow \ell \tilde{g}_{3/2}$  corresponds to the searches described above with a very light neutralino. When the sleptons have some lifetime, the leptons will have impact parameters which help to reject backgrounds. For even longer lifetimes, the apparatus can actually resolve the decay vertex, consisting of an incoming slepton and an outgoing lepton – a track with a ‘kink’ in the tracking volume. Finally, if the lifetime is long, the experimental signature is a pair of collinear, heavily ionizing tracks. By combining searches for all of these

signatures, limits of approximately  $82 \text{ GeV}/c^2$  for staus can be placed independent of the slepton lifetime [27,26].

When, due to mixing, the lightest stau is significantly lighter than the other sleptons, special topologies may result. For example,  $4\tau$  final states result from neutralino pair production. No evidence for a signal was found [27,28].

***R-parity Violation:*** If  $R$ -parity is not conserved, searches based on missing energy are not viable. The three possible RPV interaction terms ( $LL\bar{E}$ ,  $LQ\bar{D}$ ,  $\bar{U}\bar{D}\bar{D}$ ) violate lepton or baryon number, consequently precisely measured SM processes constrain products of dissimilar terms. Collider searches assume only one of the many possible terms dominates; given this assumption, searches for charginos and neutralinos, sleptons and squarks have been performed. At LEP all sets of generational indices ( $\lambda_{ijk}$ ,  $\lambda'_{ijk}$ ,  $\lambda''_{ijk}$ ) have been considered. Signatures of indirect and also direct RPV have been utilized. Rather exotic topologies can occur, such as six-lepton final states in slepton production with  $LL\bar{E}$  dominating, or ten-jet final states in chargino production with  $\bar{U}\bar{D}\bar{D}$  dominating; entirely new search criteria keyed to an excess of leptons and/or jets have been devised [29]. Searches with a wide scope have found no evidence for supersymmetry with  $R$ -parity violation, and limits are as constraining as in the canonical scenario. In fact, the direct exclusion of pair-produced  $\tilde{\chi}_1^0$ 's rules out some parameter space not accessible in the canonical case.

### ***II.5. Supersymmetry searches at hadron machines:***

While the LEP experiments can investigate a wide range of scenarios and cover corners of theoretical parameter space, they cannot match the mass reach of the Tevatron experiments (CDF and DØ). Although the full  $p\bar{p}$  energy is never available for annihilation, the cross sections for supersymmetric particle



production are large due to color factors and strong coupling. Each experiment has analyzed approximately  $110 \text{ pb}^{-1}$  of data at  $\sqrt{s} = 1.8 \text{ TeV}$  during Run I, which ended in 1996. Now Run IIa is underway, with an expected  $2 \text{ fb}^{-1}$  to be logged by 2006.

The main source of signals for supersymmetry are squarks and gluinos, in contradistinction to LEP. Pairs of squarks or gluinos are produced in  $s$ ,  $t$  and  $u$ -channel processes. These particles decay directly or via cascades to at least two  $\tilde{\chi}_1^0$ 's. The number of observed hadronic jets depends on whether the gluino or the squark is heavier, with the latter occurring naturally in mSUGRA models. The possibility of cascade decays through charginos or heavier neutralinos also enriches the possibilities of the search. The  $u$ ,  $d$ ,  $s$ ,  $c$ , and (usually)  $b$  squarks are assumed to have similar masses; the search results are reported in terms of their average mass  $M_{\tilde{q}}$  and the gluino mass  $M_{\tilde{g}}$ .

The spread of partonic energies in hadron machines is very large, so one has to consider the possible presence of several SUSY signals in one data set. A search in a given topology, such as  $\geq 3 \text{ jets} + \cancel{E}_T$ , can capture events from  $\tilde{q}$ 's,  $\tilde{g}$ 's and even  $\tilde{\chi}^{(\pm,0)}$ , with or without cascade decays. Applying experimental bounds on one production mechanism while ignoring the rest would be invalid, so the experimenters must find a relatively simple way of organizing the full phenomenology. Traditionally, they have turned to mSUGRA, in part because the fundamental parameters  $m_0$  and  $m_{1/2}$  can be fairly easily related to the squark, gluino and gaugino masses which determine the event kinematics and hence the signal acceptance.

Backgrounds at the Tevatron are relatively much higher than at LEP. There are essentially two types. First, ordinary multijet events can appear to have missing energy due to measurement errors. While large mismeasurements are rare,

there are very many di-jet and tri-jet ‘QCD’ events. This background must be estimated directly from control samples. Second, much rarer processes yield energetic neutrinos which produce a genuine missing energy signature. Examples include the production of  $W$  and  $Z$  bosons with initial-state jets, of boson pairs, and of the top quark. Estimates for these backgrounds commonly are based on theoretical cross sections, although in some analyses direct measurements are used to reduce uncertainties.


***Squarks and Gluinos:*** The classic searches [30] rely on large missing transverse energy  $\cancel{E}_T$  caused by the escaping neutralinos. Jets with high transverse energy are also required as evidence of a hard interaction; care is taken to distinguish genuine  $\cancel{E}_T$  from fluctuations in the jet energy measurement. Backgrounds from  $W$ ,  $Z$  and top production can be reduced by rejecting events with identified leptons. Uncertainties in the rates of these processes can be reduced by normalizing related samples, such as events with two jets and one or more leptons. The tails of more ordinary hard-scattering processes accompanied by multiple gluon emission are estimated directly using simulations normalized using the data.

The bounds traditionally are derived for the  $(M_{\tilde{g}}, M_{\tilde{q}})$  plane. The most recent analysis by the CDF Collaboration places significantly stronger bounds than previous analyses [31]. The removal of instrumental backgrounds is keyed more directly to the detector, which, together with specific topological cuts against poorly reconstructed multijet backgrounds, leaves gauge boson and  $t\bar{t}$  backgrounds dominant. The estimates for these are tied directly to CDF measurements, which greatly reduces systematic uncertainties. The signal region is loosely specified by demanding high  $\cancel{E}_T$  and  $H_T$ , the scalar sum of the  $\cancel{E}_T$  of the

second and third jets, and  $\cancel{E}_T$ . The number of isolated tracks allows the experimentalist to switch between a background-dominated sample and one which could contain SUSY events. As a measure of analysis rigor, the region expected to be potentially rich in SUSY events is ignored as the event counts in background-dominated samples are examined. No excess is observed, and the cuts on  $\cancel{E}_T$  and  $H_T$  are tuned to obtain the exclusion shown in Fig. 2.

If squarks are heavier than gluinos, then  $M_{\tilde{g}} \gtrsim 195 \text{ GeV}/c^2$ . If they all have the same mass, then that mass is at least  $300 \text{ GeV}/c^2$ . If the squarks are much lighter than the gluino (in which case they decay via  $\tilde{q} \rightarrow q\tilde{\chi}_1^0$ ), the bound on the gluino mass is generally high, much more than  $300 \text{ GeV}/c^2$ . A small region in which the neutralino-squark mass difference is small, is covered by the LEP experiments (see Fig. 2).

Since these results are expressed in terms of the physical masses relevant to the production process and experimental signature, the excluded region depends primarily on the assumption of nearly equal squark masses with only a small dependence on other parameters such as  $\mu$  and  $\tan\beta$ . Direct constraints on the theoretical parameters  $m_0$  and  $m_{1/2} \approx 0.34 M_3$  have been obtained by DØ assuming the mass relations of the mSUGRA model (see the first paper in [30]). These bounds do not carry significantly more information than contained in the region above the diagonal of Fig. 2. It is interesting to note that, if the LEP limits on chargino production are interpreted in this context as an indirect limit on gluinos, then roughly one obtains  $M_{\tilde{g}} > 310 \text{ GeV}/c^2$  [6].



**Figure 2:** Regions in the  $M_{\tilde{g}}-M_{\tilde{q}}$  plane excluded by searches for jets and missing energy at CDF,  $D\bar{0}$ , and LEP. See full-color version on color pages at end of book.

**Gauginos:** In the context of the mSUGRA model, which fixes  $|\mu|$  by the requirement of radiative electroweak symmetry breaking, the lightest chargino and neutralinos are dominantly gaugino. They may be produced directly by annihilation ( $q\bar{q} \rightarrow \tilde{\chi}_i^\pm \tilde{\chi}_j^0$ ) or in the decays of heavier squarks ( $\tilde{q} \rightarrow q' \tilde{\chi}_i^\pm, q \tilde{\chi}_j^0$ ).

They decay to energetic leptons ( $\tilde{\chi}^\pm \rightarrow \ell^\pm \nu^{(*)} \tilde{\chi}_1^0$  and  $\tilde{\chi}_2^0 \rightarrow \ell^+ \ell^- \tilde{\chi}_1^0$ ) and the branching ratio can be high for some parameter choices. The presence of energetic leptons has been exploited in two ways: the ‘trilepton’ signature and the ‘dilepton’ signature.

The search for trileptons is most effective for the associated production of  $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$  [32]. The requirement of three energetic leptons ( $e$  or  $\mu$ ), augmented by simple angular cuts against Drell-Yan production and cosmic rays, isolation requirements against semileptonic decays of heavy mesons, and significant  $\cancel{E}_T$  reduce backgrounds to a very small level. The bounds have been derived in the context of mSUGRA models, which generally predict modest leptonic branching ratios for charginos and neutralinos. Consequently, in this framework, the results are not competitive with the LEP bounds. When  $\tan\beta$  is large, final states with  $\tau$ 's are enhanced, and existing searches are inefficient. Nonetheless the search is completely independent of the jets+ $\cancel{E}_T$  search and could be more effective in particular models with light sleptons, for example.

The dilepton signal is geared more for the production of gauginos in gluino and squark cascades [33]. Jets are required as expected from the rest of the decay chain; the leptons should be well separated from the jets in order to avoid backgrounds from heavy quark decays. Drell-Yan events are rejected with simple cuts on the relative azimuthal angle of the leptons and their transverse momentum and by a cut on  $\cancel{E}_T$ . The Majorana nature of the gluino can be exploited by requiring two leptons with the same charge, thereby greatly reducing the background. In this scenario limits on squarks and gluinos are comparable to those from the jets+  $\cancel{E}_T$  when couched in an mSUGRA context.

DØ tried to find squarks tagged by  $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 \gamma$ , where the  $\tilde{\chi}_2^0$  appear in cascade decays [34]. The branching ratio

can be large for a selected set of model parameters leading to a Higgsino-like  $\tilde{\chi}_1^0$  and a gaugino-like  $\tilde{\chi}_2^0$ . DØ assumed a branching ratio of 100% to place the limits  $M_{\tilde{g}} > 240 \text{ GeV}/c^2$  for heavy squarks, and  $M_{\tilde{g}} > 310 \text{ GeV}/c^2$  for squarks of the same mass as the gluino.

***Stops and Sbottoms:*** The top squark is unique among the squarks because its SM partner is so massive: large off-diagonal terms in the squared-mass matrix lead to large mixing effects and a mass eigenstate possibly much lighter than all the others. This can also happen for bottom squarks for rather special parameter choices. Hence, special analyses have been developed for  $\tilde{t}_1$ 's and  $\tilde{b}_1$ 's among all the squarks.

Top squarks are pair-produced with no dependence on the mixing angle, in contrast to LEP. The searches are based on two final states:  $c\cancel{E}_T$  and  $b\ell\cancel{E}_T$ , and it is assumed that one or the other dominates. Theoretical calculations show that if chargino and slepton masses are well above  $M_{\tilde{t}_1}$ , then the loop-induced FCNC decay  $\tilde{t}_1 \rightarrow c\tilde{\chi}^0$  dominates. If  $M_{\tilde{\chi}^\pm} < M_{\tilde{t}_1}$ , then  $\tilde{t}_1 \rightarrow b\tilde{\chi}^\pm$  is the main decay mode, and the experimenters assume  $BR(\tilde{\chi}^\pm \rightarrow \ell\nu\tilde{\chi}^0) = BR(W \rightarrow \ell\nu)$ . When charginos are heavy but  $M_{\tilde{\nu}} < M_{\tilde{t}_1}$ , leptonic final states again are favored via  $\tilde{t}_1 \rightarrow b\ell\tilde{\nu}$ . In this case the branching ratio is assumed to be 1/3 for each lepton flavor. In fact, all these channels compete, and the assumption of a 100% branching ratio is not general. Furthermore, four-body decays to  $b\ell\nu\tilde{\chi}$  should not be neglected, for which limits would be reported in the  $(M_{\tilde{t}}, M_{\tilde{\chi}})$  plane [36].


CDF have obtained a result for the  $c\cancel{E}_T$  final state [37]. They employed their vertex detector to select charm jets. After a lepton veto and  $\cancel{E}_T$  requirement, this result surpasses the prior result from DØ [38]. The vertex detector was also used to tag  $b$ -quark jets for the final state  $b\ell\cancel{E}_T$ . In this case, CDF

went beyond simple event counting and applied a likelihood test to the shapes of kinematic distributions. Like the first  $D\bar{O}$  result, however, this search did not exclude any signal in the channel  $\tilde{t}_1 \rightarrow b\tilde{\chi}^\pm$ , and covered a small region for  $\tilde{t}_1 \rightarrow b\ell\tilde{\nu}$ . A new result from  $D\bar{O}$  is much more performant [39] and significantly extends the parameter space excluded by LEP searches. Finally, CDF considered the possibility  $t \rightarrow \tilde{t}_1\tilde{\chi}$  followed by  $\tilde{t}_1 \rightarrow b\tilde{\chi}^+$  [40]. Such events would remain in the top event sample and can be discriminated using a multivariate technique. No events were found compatible with the kinematics of SUSY decays, and limits on  $BR(t \rightarrow \tilde{t}_1\tilde{\chi})$  were derived in a fairly limited range of stop and chargino masses.

The search for light  $\tilde{b}_1 \rightarrow b\tilde{\chi}$  follows the  $\tilde{t}_1$  search in the charm channel [37]. The CDF search tightens the requirements for a jet with heavy flavor to good effect. An earlier  $D\bar{O}$  result tagged  $b$ -jets through semileptonic decays to muons [41].

A summary of the searches for stops is shown in Fig. 3. Given the modest luminosity and small detection efficiencies, the mass reach of the Tevatron searches is impressive. New data would likely extend this reach (as would the combination of results from the two experiments). Unfortunately, the region with  $M_{\tilde{\chi}_0} > M_{\tilde{t}_1} + 20 \text{ GeV}/c^2$  will remain inaccessible in Run 2, due to the necessity of requiring a minimum missing energy in the experimental trigger.

***R-Parity Violation:*** The CDF and  $D\bar{O}$  collaborations have searched for supersymmetry in certain RPV scenarios [42] in which the lightest neutralino decays to a lepton and two quarks.  $D\bar{O}$  considered all possible production processes as a function of mSUGRA parameters. Their trilepton search amounts to strong bounds on these parameters, stronger than



**Figure 3:** Regions excluded in the  $(M_{\tilde{t}_1}, M_{\tilde{\chi}_1^0})$  plane. The results for the  $c\tilde{\chi}_1^0$  decay mode are displayed from LEP and CDF. A DELPHI result for stable stops is indicated for  $M_{\tilde{t}_1} < M_{\tilde{\chi}_1^0}$ . Finally, the indirect limit on  $M_{\tilde{\chi}_1^0}$  is also shown. There is effectively no exclusion in the region where  $\tilde{t}_1 \rightarrow bW\tilde{\chi}_1^0$ . See full-color version on color pages at end of book.

the limits from their search for two electrons and jets. CDF used their same-sign dielectron and jets topology to look for gluino and squark (including stop) production and obtained some specific upper limits on cross sections corresponding to



$M_{\tilde{q}} > 200 \text{ GeV}/c^2$  and  $M_{\tilde{t}_1} > 120 \text{ GeV}/c^2$ . They also completed a search for  $R$ -parity violating stop decays,  $\tilde{t}_1 \rightarrow b\tau$  in which one tau decays leptonically and the other hadronically, giving the limit  $M_{\tilde{t}_1} > 122 \text{ GeV}/c^2$  [43].

**Gauge-Mediated Models:** Interest in GMSB models was spurred by an anomalous ‘ $ee\gamma\cancel{E}_T$ ’ event found by the CDF Collaboration [44]. Some of these models predict large inclusive signals for  $p\bar{p} \rightarrow \gamma\gamma + X$  given kinematic constraints derived from the properties of the CDF event. The photons arise from the decay  $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{g}_{3/2}$  and the ‘superlight’ gravitino has a mass much smaller than the charged fermions. DØ examined their sample of  $\gamma\gamma\cancel{E}_T$  events and reported limits on neutralino and chargino production corresponding to  $M_{\tilde{\chi}_1^0} > 75 \text{ GeV}/c^2$  [45]. CDF experimenters carried out a systematic survey of events with photons and SM particles (leptons, jets, missing energy) and found no signal confirming the interpretation of the original anomalous event [44,46]. They also looked for evidence of light gravitino pairs without additional SUSY particles. The invisible gravitinos are tagged by a high- $E_T$  jet from the initial state; this is the so-called ‘monojet’ signature [47]. The limit  $\sqrt{F} > 215 \text{ GeV}/c^2$  is placed on the fundamental parameter of this model.

DØ also reported limits on  $\tilde{q}$  and  $\tilde{g}$  production in this same scenario [35]. If  $\tilde{q}$  and  $\tilde{g}$  have similar masses, then that mass is great than  $310 \text{ GeV}/c^2$ .

In GMSB models, a heavy ‘sGoldstino’ is possible, which may have sizable branching ratios to photon pairs. CDF looked for narrow diphoton resonances and placed a limit  $\sqrt{F} > 1 \text{ TeV}/c^2$ , depending on assumed mass of the sGoldstino [48].

***The Search for  $B_s \rightarrow \mu^+\mu^-$ :*** Indirect evidence for SUSY could come from measurements of rare processes, especially those which are highly suppressed in the Standard Model. For example, the branching fraction for the flavor-changing neutral decay  $B_s \rightarrow \mu^+\mu^-$  is only  $3 \times 10^{-9}$  [49]. In the MSSM, however, it can be greatly enhanced due to Higgsino and possibly gluino contributions, and in fact,  $\mathcal{B}(B_s \rightarrow \mu^+\mu^-) \propto \tan^6 \beta$  [50]. The exact value for the branching fraction is highly model dependent, but in mSUGRA values as high as  $0.5 \times 10^{-7}$  can be obtained for  $\tan \beta = 55$ .

CDF found no evidence for  $B_s \rightarrow \mu^+\mu^-$  in their Run I data, and placed the upper limit  $\mathcal{B}(B_s \rightarrow \mu^+\mu^-) < 20 \times 10^{-7}$  at 90% C.L. [51]. The sensitivity will be substantially improved for Run II due to a much higher trigger acceptance and better vertex reconstruction. Recent preliminary results from Run II have strengthened the bound to  $9.5 \times 10^{-7}$  (CDF,  $113 \text{ pb}^{-1}$ ) and  $16 \times 10^{-7}$  (DØ,  $\sim 100 \text{ pb}^{-1}$ ), both at 90% C.L. [52]. The sensitivity for an integrated luminosity of  $4 \text{ fb}^{-1}$  could reach, optimistically,  $0.5 \times 10^{-7}$  [53].

If the decay  $B_s \rightarrow \mu^+\mu^-$  is observed, then a general lower bound on  $\tan \beta$  can be derived [54]. It is also worth noting that, if a signal is observed at the Tevatron, then models based on anomaly-mediated or gauge-mediated supersymmetry breaking would not be favored [50,54].

***II.7. Searches at HERA:*** The initial state for collisions at HERA includes an electron (or positron) and a proton, which provides a special opportunity to probe RPV scenarios with a dominant  $\lambda'_{1jk}$  coupling [55]. The H1 and ZEUS experiments have searched for the resonant production of squarks. The most up-to-date results include the search by H1 based on

**Table 1:** Table 1: Lower limits on supersymmetric particle masses. ‘GMSB’ refers to models with gauge-mediated supersymmetry breaking, and ‘RPV’ refers to models allowing  $R$ -parity violation.

particle		Condition	Lower limit (GeV/ $c^2$ )	Source
$\tilde{\chi}_1^\pm$	gaugino	$M_{\tilde{\nu}} > 200 \text{ GeV}/c^2$	103	LEP 2
		$M_{\tilde{\nu}} > M_{\tilde{\chi}^\pm}$	85	LEP 2
		any $M_{\tilde{\nu}}$	45	$Z$ width
	Higgsino	$M_2 < 1 \text{ TeV}/c^2$	99	LEP 2
	GMSB		150	$D\bar{O}$ isolated photons
	RPV	$LL\bar{E}$ worst case	87	LEP 2
		$LQ\bar{D}$ $m_0 > 500 \text{ GeV}/c^2$	88	LEP 2
$\tilde{\chi}_1^0$	indirect	any $\tan\beta$ , $M_{\tilde{\nu}} > 500 \text{ GeV}/c^2$	39	LEP 2
		any $\tan\beta$ , any $m_0$	36	LEP 2
		any $\tan\beta$ , any $m_0$ , SUGRA Higgs	59	LEP 2 combined
	GMSB		93	LEP 2 combined
	RPV	$LL\bar{E}$ worst case	23	LEP 2
$\tilde{e}_R$	$e\tilde{\chi}_1^0$	$\Delta M > 10 \text{ GeV}/c^2$	99	LEP 2 combined
$\tilde{\mu}_R$	$\mu\tilde{\chi}_1^0$	$\Delta M > 10 \text{ GeV}/c^2$	95	LEP 2 combined
$\tilde{\tau}_R$	$\tau\tilde{\chi}_1^0$	$M_{\tilde{\chi}_1^0} < 20 \text{ GeV}/c^2$	80	LEP 2 combined
$\tilde{\nu}$			43	$Z$ width
$\tilde{\mu}_R, \tilde{\tau}_R$		stable	86	LEP 2 combined
$\tilde{t}_1$	$c\tilde{\chi}_1^0$	any $\theta_{\text{mix}}$ , $\Delta M > 10 \text{ GeV}/c^2$	95	LEP 2 combined
		any $\theta_{\text{mix}}$ , $M_{\tilde{\chi}_1^0} \sim \frac{1}{2}M_{\tilde{t}}$	115	CDF
		any $\theta_{\text{mix}}$ and any $\Delta M$	59	ALEPH
	$b\ell\tilde{\nu}$	any $\theta_{\text{mix}}$ , $\Delta M > 7 \text{ GeV}/c^2$	96	LEP 2 combined
$\tilde{g}$	any $M_{\tilde{q}}$		195	CDF jets+ $\cancel{E}_T$
$\tilde{q}$	$M_{\tilde{q}} = M_{\tilde{g}}$		300	CDF jets+ $\cancel{E}_T$

37 pb<sup>-1</sup> of  $e^+p$  data [56]. Both  $R_p$ -violating and conserving decays of the squark were covered by a combination of seven different topologies. Bounds are placed on the  $R_p$ -violating coupling as a function of the squark mass. Completely general limits on the squark mass are impossible. However, in the constrained MSSM, and assuming  $M_{\tilde{\chi}_1^0} > 30$  GeV/ $c^2$ , the limit  $M_{\tilde{u}_L} > 160$  GeV/ $c^2$  can be placed (235 GeV/ $c^2$  for the third generation). See Ref. [56] for more details, and the Particle Listings for a list of previous results from both H1 and ZEUS.

**II.8. Conclusions:** A huge variety of searches for supersymmetry have been carried out at LEP, the Tevatron, and in fixed-target experiments. Despite all the effort, no inarguable signal has been found, forcing the experimenters to derive limits. We have tried to summarize the interesting cases in Table 1. At the present time there is little room for SUSY particles lighter than  $M_Z$ . The LEP collaborations have analyzed all their data, so prospects for the immediate future pass to the Tevatron collaborations. If still no sign of supersymmetry is found, definitive tests will be made at the LHC.

## References

1. H.E. Haber and G. Kane, Phys. Reports **117**, 75 (1985);  
H.P. Nilles, Phys. Reports **110**, 1 (1984);  
M. Chen, C. Dionisi, M. Martinez, and X. Tata, Phys. Reports **159**, 201 (1988).
2. H.E. Haber, Nucl. Phys. (Proc. Supp.) **B62**, 469 (1998);  
S. Dawson, *SUSY and Such*, hep-ph/9612229.
3. H. Dreiner, *An Introduction to Explicit R-parity Violation*, in **Perspectives on Supersymmetry**, ed. by G.L. Kane, World Scientific, 1997, p.462;  
G. Bhattacharyya, Nucl. Phys. Proc. Suppl. **A52**, 83 (1997);  
V. Barger, G.F. Giudice, and T. Han, Phys. Rev. **D40**,

- 1987 (1989);  
S. Dawson, Nucl. Phys. **B261**, 297 (1985).
4. M. Dine, Nucl. Phys. Proc. Suppl. **52A**, 201(1997);  
K.S. Babu, C. Kolda, and F. Wilczek, Phys. Rev. Lett. **77**, 3070 (1996);  
S. Dimopoulos *et al.*, Phys. Rev. Lett. **76**, 3494 (1996);  
S. Dimopoulos, S. Thomas, J.D. Wells, Phys. Rev. **D54**, 3283 (1996), and Nucl. Phys. **B488**, 39 (1997);  
D.R. Stump, M. Wiest, C.P. Yuan, Phys. Rev. **D54**, 1936 (1996);  
M. Dine, A. Nelson, and Y. Shirman Phys. Rev. **D51**, 1362 (1995);  
D.A. Dicus, S. Nandi, and J. Woodside, Phys. Rev. **D41**, 2347 (1990) and Phys. Rev. **D43**, 2951 (1990);  
P. Fayet, Phys. Lett. **B175**, 471 (1986);  
J. Ellis, K. Enqvist, and D.V. Nanopoulos, Phys. Lett. **B151**, 357 (1985), and Phys. Lett. **B147**, 99 (1984);  
P. Fayet, Phys. Lett. **B69**, 489 (1977) and Phys. Lett. **B70**, 461 (1977).
  5. F. Gianotti, New Jour. Phys. **4**,63(2002).
  6. A. Lipniacka, hep-ph/0112280.
  7. **LEPSUSYWG, ALEPH, DELPHI, L3** and **OPAL** Collab., Preliminary results from the combination of LEP data, prepared by the LEP SUSY Working Group. LEPSUSYWG/02-01.1, 02-02.1, 02-04.1, 02-05.1, 02-06.2, 02-07.1, 02-08.1, 02-09.2, 02-10.1, 01-03.1, 01-07.1 See also <http://www.cern.ch/lepsusy/>.
  8. J.-F. Grivaz, *Supersymmetric Particle Searches at LEP*, in **Perspectives on Supersymmetry**, *ibid.*, p.179;  
M. Drees and X. Tata, Phys. Rev. **D43**, 2971 (1991).
  9. J. L. Feng and T. Moroi, Phys. Rev. **D61**, 095004 (2000);  
L. Randall and R. Sundrum, Nucl. Phys. **B557**, 79 (1999).
  10. **DELPHI**: Eur. Phys. J. **C11**, 1 (1999).
  11. **ALEPH**: Phys. Lett. **B533**, 223 (2002);  
**OPAL**: hep-ex/0210043;  
**L3**: Phys. Lett. **B482**, 31 (2000).

12. **ALEPH**: Z. Phys. **C72**, 549 (1996) and Eur. Phys. J. **C11**, 193 (1999).
13. **ALEPH**: Eur. Phys. J. **C2**, 417 (1998).
14. **ALEPH**: Phys. Lett. **B526**, 206 (2002);  
**OPAL**: hep-ex/0309014;  
**DELPHI**: Eur. Phys. J. **C19**, 29 (2001);  
**L3**: Phys. Lett. **B471**, 280 (1999).
15. **ALEPH**: Phys. Lett. **B544**, 73 (2002).
16. **L3**: hep-ex/0310007.
17. **OPAL**: Phys. Lett. **B545**, 272 (2002) Err. *ibid.* **B548**, 258 (2002).
18. **ALEPH**: Phys. Lett. **B537**, 5 (2002).
19. **ALEPH**: Phys. Lett. **B488**, 234 (2000).
20. **CLEO**: Phys. Rev. **D63**, 051101 (2001).
21. **ALEPH**: Phys. Lett. **B469**, 303 (1999).
22. **OPAL**: Eur. Phys. J. **C8**, 255 (1999);  
**L3**: Eur. Phys. J. **C4**, 207 (1998).
23. **ALEPH**: Phys. Lett. **B499**, 67 (2001).
24. H. K. Dreiner *et al.*, hep-ph/0304289;  
D. Choudhury, *et al.*, Phys. Rev. **D61**, 095009 (2000).
25. **ALEPH**: Eur. Phys. J. **C28**, 1 (2003).
26. **ALEPH**: Eur. Phys. J. **C25**, 339 (2002).
27. **DELPHI**: Eur. Phys. J. **C27**, 153 (2003);  
**DELPHI**: Phys. Lett. **B503**, 34 (2001);  
**ALEPH**: Eur. Phys. J. **C16**, 71 (1999).
28. **DELPHI**: Eur. Phys. J. **C7**, 595 (1999).
29. **ALEPH**: Eur. Phys. J. **C19**, 415 (2001) and Eur. Phys. J. **C13**, 29 (2000);  
**OPAL**: Eur. Phys. J. **C12**, 1 (2000) and Eur. Phys. J. **C11**, 619 (1999);  
**DELPHI**: Phys. Lett. **B502**, 24 (2001);  
**L3**: Eur. Phys. J. **C19**, 397 (2001) and Phys. Lett. **B524**, 65 (2002).
30. **DØ**: Phys. Rev. Lett. **83**, 4937 (1999) and Phys. Rev. Lett. **75**, 618 (1995);

- CDF**: Phys. Rev. **D56**, 1357 (1997) and Phys. Rev. Lett. **76**, 2006 (1996).
31. **CDF**: Phys. Rev. Lett. **88**, 041801 (2002).
  32. **DØ**: Phys. Rev. Lett. **80**, 1591 (1998);  
**CDF**: Phys. Rev. Lett. **80**, 5275 (1998).
  33. **DØ**: Phys. Rev. **D63**, 091102 (2001);  
**CDF**: Phys. Rev. Lett. **76**, 2006 (1996) and Phys. Rev. Lett. **87**, 251803 (2001).
  34. **DØ**: Phys. Rev. Lett. **82**, 29 (1999).
  35. **DØ**: Phys. Rev. Lett. **78**, 2070 (1997).
  36. A. Djouadi *et al.*, Phys. Rev. **D71**, 095006 (2000) and Phys. Rev. **D63**, 115005 (2001).
  37. **CDF**: Phys. Rev. Lett. **84**, 5704 (2000).
  38. **DØ**: Phys. Rev. Lett. **76**, 2222 (1996).
  39. **DØ**: Phys. Rev. Lett. **88**, 171802 (2002).
  40. **CDF**: Phys. Rev. **D63**, 091101 (2001).
  41. **DØ**: Phys. Rev. **D60**, 031101 (1999).
  42. **CDF**: Phys. Rev. Lett. **83**, 2133 (1999) and Phys. Rev. Lett. **87**, 251803 (2001);  
**DØ**: Phys. Rev. **D62**, 071701 (2000) and Phys. Rev. Lett. **83**, 4476 (1999).
  43. **CDF**: [hep-ex/0305010](http://arxiv.org/abs/hep-ex/0305010).
  44. **CDF**: Phys. Rev. **D59**, 092002 (1999).
  45. **DØ**: Phys. Rev. Lett. **78**, 2070 (1997).
  46. **CDF**: Phys. Rev. Lett. **81**, 1791 (1998).
  47. **CDF**: Phys. Rev. Lett. **85**, 1378 (2000).
  48. **CDF**: Phys. Rev. Lett. **81**, 1791 (1998).
  49. Andrzej J. Buras, [hep-ph/9806471](http://arxiv.org/abs/hep-ph/9806471) and references therein.
  50. A. Dedes, H.K. Dreiner, U. Nierste and P. Richardson, [hep-ph/0207026](http://arxiv.org/abs/hep-ph/0207026).
  51. **CDF**: Phys. Rev. Lett. **57**, 3811 (1998).
  52. Prelim. results from **CDF** and **DØ** were reported by M.Schmitt at Lepton-Photon, 2003, at FNAL.

- 53. R. Arnowitt, B.Dutta, T.Kamon and M.Tanaka, Phys. Lett. **B538**, 121 (2002).
- 54. G.L.Kane, C.Kolda and J.E.Lennon, [hep-ph/0310042](http://arxiv.org/abs/hep-ph/0310042).
- 55. M. Kuze and Y. Sirois, Prog. in Part. Nucl. Phys. **50**, 1 (2003).
- 56. **H1**: Eur. Phys. J. **C20**, 639 (2001);  
**H1**: Phys. Lett. **B568**, 35 (2003).

## SUPERSYMMETRIC MODEL ASSUMPTIONS

The exclusion of particle masses within a mass range ( $m_1, m_2$ ) will be denoted with the notation "none  $m_1 - m_2$ " in the VALUE column of the following Listings.

Most of the results shown below, unless stated otherwise, are based on the Minimal Supersymmetric Standard Model (MSSM), as described in the Note on Supersymmetry. Unless otherwise indicated, this includes the assumption of common gaugino and scalar masses at the scale of Grand Unification (GUT), and use of the resulting relations in the spectrum and decay branching ratios. It is also assumed that  $R$ -parity ( $R$ ) is conserved. Unless otherwise indicated, the results also assume that:

- 1) The  $\tilde{\chi}_1^0$  is the lightest supersymmetric particle (LSP)
- 2)  $m_{\tilde{f}_L} = m_{\tilde{f}_R}$ , where  $\tilde{f}_{L,R}$  refer to the scalar partners of left- and right-handed fermions.

Limits involving different assumptions are identified in the Comments or in the Footnotes. We summarize here the notations used in this Chapter to characterize some of the most common deviations from the MSSM (for further details, see the Note on Supersymmetry).

Theories with  $R$ -parity violation ( $\mathcal{R}$ ) are characterized by a superpotential of the form:  $\lambda_{ijk} L_i L_j e_k^c + \lambda'_{ijk} L_i Q_j d_k^c + \lambda''_{ijk} u_i^c d_j^c d_k^c$ , where  $i, j, k$  are generation indices. The presence of any of these couplings is often identified in the following



by the symbols  $L\bar{L}\bar{E}$ ,  $LQ\bar{D}$ , and  $\bar{U}\bar{D}\bar{D}$ . Mass limits in the presence of  $\mathcal{R}$  will often refer to “direct” and “indirect” decays. Direct refers to  $\mathcal{R}$  decays of the particle in consideration. Indirect refers to cases where  $\mathcal{R}$  appears in the decays of the LSP.

In several models, most notably in theories with so-called Gauge Mediated Supersymmetry Breaking (GMSB), the gravitino ( $\tilde{G}$ ) is the LSP. It is usually much lighter than any other massive particle in the spectrum, and  $m_{\tilde{G}}$  is then neglected in all decay processes involving gravitinos. In these scenarios, particles other than the neutralino are sometimes considered as the next-to-lightest supersymmetric particle (NLSP), and are assumed to decay to their even- $R$  partner plus  $\tilde{G}$ . If the lifetime is short enough for the decay to take place within the detector,  $\tilde{G}$  is assumed to be undetected and to give rise to missing energy ( $\cancel{E}$ ) or missing transverse energy ( $\cancel{E}_T$ ) signatures.

When needed, specific assumptions on the eigenstate content of  $\tilde{\chi}^0$  and  $\tilde{\chi}^\pm$  states are indicated, using the notation  $\tilde{\gamma}$  (photino),  $\tilde{H}$  (higgsino),  $\tilde{W}$  (wino), and  $\tilde{Z}$  (zino) to signal that the limit of pure states was used. The terms gaugino is also used, to generically indicate wino-like charginos and zino-like neutralinos.

### $\tilde{\chi}_1^0$ (Lightest Neutralino) MASS LIMIT

$\tilde{\chi}_1^0$  is often assumed to be the lightest supersymmetric particle (LSP). See also the  $\tilde{\chi}_2^0$ ,  $\tilde{\chi}_3^0$ ,  $\tilde{\chi}_4^0$  section below.

We have divided the  $\tilde{\chi}_1^0$  listings below into five sections:

- 1) Accelerator limits for stable  $\tilde{\chi}_1^0$ ,
- 2) Bounds on  $\tilde{\chi}_1^0$  from dark matter searches,
- 3) Bounds on  $\tilde{\chi}_1^0$  elastic cross sections from dark matter searches,
- 4) Other bounds on  $\tilde{\chi}_1^0$  from astrophysics and cosmology, and
- 5) Bounds on unstable  $\tilde{\chi}_1^0$ .

#### ———— Accelerator limits for stable $\tilde{\chi}_1^0$ ————

Unless otherwise stated, results in this section assume spectra, production rates, decay modes, and branching ratios as evaluated in the MSSM, with gaugino and sfermion mass unification at the GUT scale. These papers generally study production of  $\tilde{\chi}_i^0 \tilde{\chi}_j^0$  ( $i \geq 1, j \geq 2$ ),  $\tilde{\chi}_1^+ \tilde{\chi}_1^-$ , and (in the case of hadronic collisions)  $\tilde{\chi}_1^+ \tilde{\chi}_2^0$  pairs. The mass limits on  $\tilde{\chi}_1^0$  are either direct, or follow indirectly from the constraints set by the non-observation of  $\tilde{\chi}_1^\pm$  and  $\tilde{\chi}_2^0$  states on the gaugino and higgsino MSSM parameters  $M_2$  and  $\mu$ . In some cases, information is used from the nonobservation of slepton decays.

Obsolete limits obtained from  $e^+ e^-$  collisions up to  $\sqrt{s}=184$  GeV have been removed from this compilation and can be found in the 2000 Edition (The European Physical Journal **C15** 1 (2000)) of this Review.

$$\Delta m_0 = m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}.$$

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>40	95	<sup>1</sup> ABBIENDI	04H OPAL	all $\tan\beta$ , $\Delta m_0 > 5$ GeV, $m_0 > 500$ GeV, $A_0 = 0$
>42.4	95	<sup>2</sup> HEISTER	04 ALEP	all $\tan\beta$ , all $\Delta m_0$ , all $m_0$
>39.2	95	<sup>3</sup> ABDALLAH	03M DLPH	all $\tan\beta$ , $m_{\tilde{\nu}} > 500$ GeV
<b>&gt;46</b>	95	<sup>4</sup> ABDALLAH	03M DLPH	all $\tan\beta$ , all $\Delta m_0$ , all $m_0$
>32.5	95	<sup>5</sup> ACCIARRI	00D L3	$\tan\beta > 0.7$ , $\Delta m_0 > 3$ GeV, all $m_0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		<sup>6</sup> ABBOTT	98C D0	$p\bar{p} \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_2^0$
>41	95	<sup>7</sup> ABE	98J CDF	$p\bar{p} \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_2^0$

<sup>1</sup> ABBIENDI 04H search for charginos and neutralinos in events with acoplanar leptons+jets and multi-jet final states in the 192–209 GeV data, combined with the results on leptonic final states from ABBIENDI 04. The results hold for a scan over the parameter space covering the region  $0 < M_2 < 5000$  GeV,  $-1000 < \mu < 1000$  GeV and  $\tan\beta$  from 1 to 40. This limit supersedes ABBIENDI 00H.

<sup>2</sup> HEISTER 04 data collected up to 209 GeV. Updates earlier analysis of selectrons from HEISTER 02E, includes a new analysis of charginos and neutralinos decaying into stau and uses results on charginos with initial state radiation from HEISTER 02J. The limit is based on the direct search for charginos and neutralinos, the constraints from the slepton search and the Higgs mass limits from HEISTER 02 using a top mass of 175 GeV, interpreted in a framework with universal gaugino and sfermion masses. Assuming the mixing in the stau sector to be negligible, the limit improves to 43.1 GeV. Under the assumption of MSUGRA with unification of the Higgs and sfermion masses, the limit improves to 50 GeV, and reaches 53 GeV for  $A_0 = 0$ . These limits include and update the results of BARATE 01.

<sup>3</sup> ABDALLAH 03M uses data from  $\sqrt{s} = 192\text{--}208$  GeV. A limit on the mass of  $\tilde{\chi}_1^0$  is derived from direct searches for neutralinos combined with the chargino search. Neutralinos are searched in the production of  $\tilde{\chi}_1^0 \tilde{\chi}_2^0$ ,  $\tilde{\chi}_1^0 \tilde{\chi}_3^0$ , as well as  $\tilde{\chi}_2^0 \tilde{\chi}_3^0$  and  $\tilde{\chi}_2^0 \tilde{\chi}_4^0$  giving rise to cascade decays, and  $\tilde{\chi}_1^0 \tilde{\chi}_2^0$  and  $\tilde{\chi}_1^0 \tilde{\chi}_2^0$ , followed by the decay  $\tilde{\chi}_2^0 \rightarrow \tilde{\tau} \tau$ . The results hold for the parameter space defined by values of  $M_2 < 1$  TeV,  $|\mu| \leq 2$  TeV with the  $\tilde{\chi}_1^0$  as LSP. The limit is obtained for  $\tan\beta = 1$  and large  $m_0$ , where  $\tilde{\chi}_2^0 \tilde{\chi}_4^0$  and chargino pair production are important. If the constraint from Higgs searches is also imposed, the limit improves to 49.0 GeV in the  $M_h^{max}$  scenario with  $m_t=174.3$  GeV. These limits update the results of ABREU 00J.

<sup>4</sup> ABDALLAH 03M uses data from  $\sqrt{s} = 192\text{--}208$  GeV. An indirect limit on the mass of  $\tilde{\chi}_1^0$  is derived by constraining the MSSM parameter space by the results from direct

searches for neutralinos (including cascade decays and  $\tilde{\tau}\tau$  final states), for charginos (for all  $\Delta m_+$ ) and for sleptons, stop and sbottom. The results hold for the full parameter space defined by values of  $M_2 < 1$  TeV,  $|\mu| \leq 2$  TeV with the  $\tilde{\chi}_1^0$  as LSP. Constraints from the Higgs search in the  $M_h^{max}$  scenario assuming  $m_t=174.3$  GeV are included. The limit is obtained for  $\tan\beta \geq 5$  when stau mixing leads to mass degeneracy between  $\tilde{\tau}_1$  and  $\tilde{\chi}_1^0$  and the limit is based on  $\tilde{\chi}_2^0$  production followed by its decay to  $\tilde{\tau}_1\tau$ . In the pathological scenario where  $m_0$  and  $|\mu|$  are large, so that the  $\tilde{\chi}_2^0$  production cross section is negligible, and where there is mixing in the stau sector but not in stop nor sbottom, the limit is based on charginos with soft decay products and an ISR photon. The limit then degrades to 39 GeV. See Figs 40–42 for the dependence of the limit on  $\tan\beta$  and  $m_{\tilde{\nu}}$ . These limits update the results of ABREU 00W.

<sup>5</sup> ACCIARRI 00D data collected at  $\sqrt{s}=189$  GeV. The results hold over the full parameter space defined by  $0.7 \leq \tan\beta \leq 60$ ,  $0 \leq M_2 \leq 2$  TeV,  $m_0 \leq 500$  GeV,  $|\mu| \leq 2$  TeV. The minimum mass limit is reached for  $\tan\beta=1$  and large  $m_0$ . The results of slepton searches from ACCIARRI 99W are used to help set constraints in the region of small  $m_0$ . The limit improves to 48 GeV for  $m_0 \gtrsim 200$  GeV and  $\tan\beta \gtrsim 10$ . See their Figs. 6–8 for the  $\tan\beta$  and  $m_0$  dependence of the limits. Updates ACCIARRI 98F.

<sup>6</sup> ABBOTT 98C searches for trilepton final states ( $\ell=e,\mu$ ). See footnote to ABBOTT 98C in the Chargino Section for details on the assumptions. Assuming a negligible decay rate of  $\tilde{\chi}_1^\pm$  and  $\tilde{\chi}_2^0$  to quarks, they obtain  $m_{\tilde{\chi}_2^0} \gtrsim 51$  GeV.

<sup>7</sup> ABE 98J searches for trilepton final states ( $\ell=e,\mu$ ). See footnote to ABE 98J in the Chargino Section for details on the assumptions. The quoted result corresponds to the best limit within the selected range of parameters, obtained for  $m_{\tilde{q}} > m_{\tilde{g}}$ ,  $\tan\beta=2$ , and  $\mu=-600$  GeV.

### ————— Bounds on $\tilde{\chi}_1^0$ from dark matter searches —————

These papers generally exclude regions in the  $M_2 - \mu$  parameter plane assuming that  $\tilde{\chi}_1^0$  is the dominant form of dark matter in the galactic halo. These limits are based on the lack of detection in laboratory experiments or by the absence of a signal in underground neutrino detectors. The latter signal is expected if  $\tilde{\chi}_1^0$  accumulates in the Sun or the Earth and annihilates into high-energy  $\nu$ 's.

VALUE	DOCUMENT ID	TECN
-------	-------------	------

• • • We do not use the following data for averages, fits, limits, etc. • • •

	8 DESAI	04	SKAM
	8 AMBROSIO	99	MCRO
	9 LOSECCO	95	RVUE
	10 MORI	93	KAMI
	11 BOTTINO	92	COSM
	12 BOTTINO	91	RVUE
	13 GELMINI	91	COSM
	14 KAMIONKOW.	91	RVUE
	15 MORI	91B	KAMI
none 4–15 GeV	16 OLIVE	88	COSM

- <sup>8</sup> AMBROSIO 99 and DESAI 04 set new neutrino flux limits which can be used to limit the parameter space in supersymmetric models based on neutralino annihilation in the Sun and the Earth.
- <sup>9</sup> LOSECCO 95 reanalyzed the IMB data and places lower limit on  $m_{\tilde{\chi}_1^0}$  of 18 GeV if the LSP is a photino and 10 GeV if the LSP is a higgsino based on LSP annihilation in the sun producing high-energy neutrinos and the limits on neutrino fluxes from the IMB detector.
- <sup>10</sup> MORI 93 excludes some region in  $M_2-\mu$  parameter space depending on  $\tan\beta$  and lightest scalar Higgs mass for neutralino dark matter  $m_{\tilde{\chi}_0} > m_W$ , using limits on upgoing muons produced by energetic neutrinos from neutralino annihilation in the Sun and the Earth.
- <sup>11</sup> BOTTINO 92 excludes some region  $M_2-\mu$  parameter space assuming that the lightest neutralino is the dark matter, using upgoing muons at Kamiokande, direct searches by Ge detectors, and by LEP experiments. The analysis includes top radiative corrections on Higgs parameters and employs two different hypotheses for nucleon-Higgs coupling. Effects of rescaling in the local neutralino density according to the neutralino relic abundance are taken into account.
- <sup>12</sup> BOTTINO 91 excluded a region in  $M_2-\mu$  plane using upgoing muon data from Kamioka experiment, assuming that the dark matter surrounding us is composed of neutralinos and that the Higgs boson is not too heavy.
- <sup>13</sup> GELMINI 91 exclude a region in  $M_2-\mu$  plane using dark matter searches.
- <sup>14</sup> KAMIONKOWSKI 91 excludes a region in the  $M_2-\mu$  plane using IMB limit on upgoing muons originated by energetic neutrinos from neutralino annihilation in the sun, assuming that the dark matter is composed of neutralinos and that  $m_{H_1^0} \lesssim 50$  GeV. See Fig. 8 in the paper.
- <sup>15</sup> MORI 91B exclude a part of the region in the  $M_2-\mu$  plane with  $m_{\tilde{\chi}_1^0} \lesssim 80$  GeV using a limit on upgoing muons originated by energetic neutrinos from neutralino annihilation in the earth, assuming that the dark matter surrounding us is composed of neutralinos and that  $m_{H_1^0} \lesssim 80$  GeV.
- <sup>16</sup> OLIVE 88 result assumes that photinos make up the dark matter in the galactic halo. Limit is based on annihilations in the sun and is due to an absence of high energy neutrinos detected in underground experiments. The limit is model dependent.

### ———— $\tilde{\chi}_1^0$ - $p$ elastic cross section ————

Experimental results on the  $\tilde{\chi}_1^0$ - $p$  elastic cross section are evaluated at  $m_{\tilde{\chi}_1^0}=100$  GeV. The experimental results on the cross section are often mass dependent. Therefore, the mass and cross section results are also given where the limit is strongest, when appropriate. Results are quoted separately for spin-dependent interactions (based on an effective 4-Fermi Lagrangian of the form  $\bar{\chi}\gamma^\mu\gamma^5\chi\bar{q}\gamma_\mu\gamma^5q$ ) and spin-independent interactions ( $\bar{\chi}\chi\bar{q}q$ ). For calculational details see GRIEST 88B, ELLIS 88D, BARBIERI 89C, DREES 93B, ARNOWITT 96, BERGSTROM 96, and BAER 97 in addition to the theory papers listed in the Tables. For a description of the theoretical assumptions and experimental techniques underlying most of the listed papers, see the review on "Dark matter" in this "Review of Particle Physics," and references therein. Most of the following papers use galactic halo and nuclear interaction assumptions from (LEWIN 96).

### Spin-dependent interactions

VALUE (pb)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

< 5	17 AKERIB	06 CDMS Ge
< 0.4	18 ALNER	05 NAIA NaI Spin Dep.
< 2	19 BARNABE-HE..05	PICA C
< 1.4	20 GIRARD	05 SMPL F, Cl
< 4	21 KLAPDOR-K...05	HDMS Ge
$2 \times 10^{-11}$ to $1 \times 10^{-4}$	22 ELLIS	04 THEO $\mu > 0$
< 16	23 GIULIANI	04 SIMP F
< 0.8	24 AHMED	03 NAIA NaI Spin Dep.
< 40	25 TAKEDA	03 BOLO NaF Spin Dep.
< 10	26 ANGLOHER	02 CRES Sapphire
$8 \times 10^{-7}$ to $2 \times 10^{-5}$	27 ELLIS	01C THEO $\tan\beta \leq 10$
< 3.8	28 BERNABEI	00D DAMA Xe
< 15	29 COLLAR	00 SMPL F
< 0.8	SPOONER	00 UKDM NaI
< 4.8	30 BELLI	99C DAMA F
<100	31 OOTANI	99 BOLO LiF
< 0.6	BERNABEI	98C DAMA Xe
< 5	30 BERNABEI	97 DAMA F

17 The strongest upper limit is 4 pb and occurs at  $m_\chi \simeq 60$  GeV. The limit on the neutron spin-dependent elastic cross section is 0.07 pb.

18 The strongest upper limit is 0.35 pb and occurs at  $m_\chi \simeq 60$  GeV.

19 The strongest upper limit is 1.2 pb and occurs  $m_\chi \simeq 30$  GeV.

20 The strongest upper limit is 1.2 pb and occurs  $m_\chi \simeq 40$  GeV.

21 Limit applies to neutron elastic cross section.

22 ELLIS 04 calculates the  $\chi$ - $p$  elastic scattering cross section in the framework of  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry, but without universal scalar masses. In the case of universal squark and slepton masses, but non-universal Higgs masses, the limit becomes  $2 \times 10^{-4}$ , see ELLIS 03E.

23 The strongest upper limit is 10 pb and occurs at  $m_\chi \simeq 30$  GeV.

24 The strongest upper limit is 0.75 pb and occurs at  $m_\chi \approx 70$  GeV.

25 The strongest upper limit is 30 pb and occurs at  $m_\chi \approx 20$  GeV.

26 The strongest upper limit is 8 pb and occurs at  $m_\chi \simeq 30$  GeV.

27 ELLIS 01C calculates the  $\chi$ - $p$  elastic scattering cross section in the framework of  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry. In models with nonuniversal Higgs masses, the upper limit to the cross section is  $6 \times 10^{-4}$ .

28 The strongest upper limit is 3 pb and occurs at  $m_\chi \simeq 60$  GeV. The limits are for inelastic scattering  $\chi^0 + {}^{129}\text{Xe} \rightarrow \chi^0 + {}^{129}\text{Xe}^*$  (39.58 keV).

29 The strongest upper limit is 9 pb and occurs at  $m_\chi \simeq 30$  GeV.

30 The strongest upper limit is 4.4 pb and occurs at  $m_\chi \simeq 60$  GeV.

31 The strongest upper limit is about 35 pb and occurs at  $m_\chi \simeq 15$  GeV.

### Spin-independent interactions

VALUE (pb)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
< $2 \times 10^{-7}$	32 AKERIB	06A CDMS Ge	
< $5 \times 10^{-7}$	33 AKERIB	05 CDMS Ge	
< $9 \times 10^{-6}$	ALNER	05 NAIA NaI Spin Indep.	
< $1.2 \times 10^{-6}$	34 ALNER	05A ZEPL	

$< 2 \times 10^{-6}$	35	ANGLOHER	05	CRES	CaWO <sub>4</sub>
$< 1.4 \times 10^{-6}$		SANGLARD	05	EDEL	Ge
$< 4 \times 10^{-7}$	36	AKERIB	04	CDMS	Ge
$2 \times 10^{-11}$ to $8 \times 10^{-6}$	37,38	ELLIS	04	THEO	$\mu > 0$
$< 5 \times 10^{-8}$	39	PIERCE	04A	THEO	
$< 2 \times 10^{-5}$	40	AHMED	03	NAIA	NaI Spin Indep.
$< 3 \times 10^{-6}$	41	AKERIB	03	CDMS	Ge
$2 \times 10^{-13}$ to $2 \times 10^{-7}$	42	BAER	03A	THEO	
$< 1.4 \times 10^{-5}$	43	KLAPDOR-K...	03	HDMS	Ge
$< 6 \times 10^{-6}$	44	ABRAMS	02	CDMS	Ge
$< 1.4 \times 10^{-6}$	45	BENOIT	02	EDEL	Ge
$10^{-12}$ to $7 \times 10^{-6}$	37	KIM	02B	THEO	
$< 3 \times 10^{-5}$	46	MORALES	02B	CSME	Ge
$< 10^{-5}$	47	MORALES	02C	IGEX	Ge
$< 10^{-6}$		BALTZ	01	THEO	
$< 3 \times 10^{-5}$	48	BAUDIS	01	HDMS	Ge
$< 4.5 \times 10^{-6}$		BENOIT	01	EDEL	Ge
$< 7 \times 10^{-6}$	49	BOTTINO	01	THEO	
$< 10^{-8}$	50	CORSETTI	01	THEO	$\tan\beta \leq 25$
$5 \times 10^{-10}$ to $1.5 \times 10^{-8}$	51	ELLIS	01C	THEO	$\tan\beta \leq 10$
$< 4 \times 10^{-6}$	50	GOMEZ	01	THEO	
$2 \times 10^{-10}$ to $10^{-7}$	50	LAHANAS	01	THEO	
$< 3 \times 10^{-6}$		ABUSAIDI	00	CDMS	Ge, Si
$< 6 \times 10^{-7}$	52	ACCOMANDO	00	THEO	
	53	BERNABEI	00	DAMA	NaI
$2.5 \times 10^{-9}$ to $3.5 \times 10^{-8}$	54	FENG	00	THEO	$\tan\beta=10$
$< 1.5 \times 10^{-5}$		MORALES	00	IGEX	Ge
$< 4 \times 10^{-5}$		SPOONER	00	UKDM	NaI
$< 7 \times 10^{-6}$		BAUDIS	99	HDMS	<sup>76</sup> Ge
	55	BERNABEI	99	DAMA	NaI
	56	BERNABEI	98	DAMA	NaI
$< 7 \times 10^{-6}$		BERNABEI	98C	DAMA	Xe

32 AKERIB 06A updates the results of AKERIB 05. The strongest upper limit is  $1.6 \times 10^{-7}$  pb and occurs at  $m_\chi \approx 60$  GeV.

33 AKERIB 05 is incompatible with the DAMA most likely value. The strongest upper limit is  $4 \times 10^{-7}$  pb and occurs at  $m_\chi \approx 60$  GeV.

34 The strongest upper limit is also close to  $1.0 \times 10^{-6}$  pb and occurs at  $m_\chi \approx 70$  GeV.

35 The strongest upper limit is also close to  $1.4 \times 10^{-6}$  pb and occurs at  $m_\chi \approx 70$  GeV.

36 AKERIB 04 is incompatible with BERNABEI 00 most likely value, under the assumption of standard WIMP-halo interactions. The strongest upper limit is  $4 \times 10^{-7}$  pb and occurs at  $m_\chi \approx 60$  GeV.

37 KIM 02 and ELLIS 04 calculate the  $\chi - p$  elastic scattering cross section in the framework of  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry, but without universal scalar masses.

38 In the case of universal squark and slepton masses, but non-universal Higgs masses, the limit becomes  $2 \times 10^{-6}$  ( $2 \times 10^{-11}$  when constraint from the BNL  $g-2$  experiment are included), see ELLIS 03E. ELLIS 05 display the sensitivity of the elastic scattering cross section to the  $\pi$ -Nucleon  $\Sigma$  term.

- 39 PIERCE 04A calculates the  $\chi-p$  elastic scattering cross section in the framework of models with very heavy scalar masses. See Fig. 2 of the paper.
- 40 The strongest upper limit is  $1.8 \times 10^{-5}$  pb and occurs at  $m_\chi \approx 80$  GeV.
- 41 Under the assumption of standard WIMP-halo interactions, Akerib 03 is incompatible with BERNABEI 00 most likely value at the 99.98% CL. See Fig. 4.
- 42 BAER 03A calculates the  $\chi-p$  elastic scattering cross section in several models including the framework of  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry.
- 43 The strongest upper limit is  $7 \times 10^{-6}$  pb and occurs at  $m_\chi \simeq 30$  GeV.
- 44 ABRAMS 02 is incompatible with the DAMA most likely value at the 99.9% CL. The strongest upper limit is  $3 \times 10^{-6}$  pb and occurs at  $m_\chi \simeq 30$  GeV.
- 45 BENOIT 02 excludes the central result of DAMA at the 99.8%CL.
- 46 The strongest upper limit is  $2 \times 10^{-5}$  pb and occurs at  $m_\chi \simeq 40$  GeV.
- 47 The strongest upper limit is  $7 \times 10^{-6}$  pb and occurs at  $m_\chi \simeq 46$  GeV.
- 48 The strongest upper limit is  $1.8 \times 10^{-5}$  pb and occurs at  $m_\chi \simeq 32$  GeV
- 49 BOTTINO 01 calculates the  $\chi-p$  elastic scattering cross section in the framework of the following supersymmetric models:  $N=1$  supergravity with the radiative breaking of the electroweak gauge symmetry,  $N=1$  supergravity with nonuniversal scalar masses and an effective MSSM model at the electroweak scale.
- 50 Calculates the  $\chi-p$  elastic scattering cross section in the framework of  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry.
- 51 ELLIS 01C calculates the  $\chi-p$  elastic scattering cross section in the framework of  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry. ELLIS 02B find a range  $2 \times 10^{-8}$ – $1.5 \times 10^{-7}$  at  $\tan\beta=50$ . In models with nonuniversal Higgs masses, the upper limit to the cross section is  $4 \times 10^{-7}$ .
- 52 ACCOMANDO 00 calculate the  $\chi-p$  elastic scattering cross section in the framework of minimal  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry. The limit is relaxed by at least an order of magnitude when models with nonuniversal scalar masses are considered. A subset of the authors in ARNOWITT 02 updated the limit to  $< 9 \times 10^{-8}$  ( $\tan\beta < 55$ ).
- 53 BERNABEI 00 search for annual modulation of the WIMP signal. The data favor the hypothesis of annual modulation at  $4\sigma$  and are consistent, for a particular model framework quoted there, with  $m_{\chi^0} = 44^{+12}_{-9}$  GeV and a spin-independent  $\chi^0$ -proton cross section of  $(5.4 \pm 1.0) \times 10^{-6}$  pb. See also BERNABEI 01 and BERNABEI 00c.
- 54 FENG 00 calculate the  $\chi-p$  elastic scattering cross section in the framework of  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry with a particular emphasis on focus point models. At  $\tan\beta=50$ , the range is  $8 \times 10^{-8}$ – $4 \times 10^{-7}$ .
- 55 BERNABEI 99 search for annual modulation of the WIMP signal. The data favor the hypothesis of annual modulation at 99.6%CL and are consistent, for the particular model framework considered there, with  $m_{\chi^0} = 59^{+17}_{-14}$  GeV and spin-independent  $\chi^0$ -proton cross section of  $(7.0^{+0.4}_{-1.2}) \times 10^{-6}$  pb ( $1\sigma$  errors).
- 56 BERNABEI 98 search for annual modulation of the WIMP signal. The data are consistent, for the particular model framework considered there, with  $m_{\chi^0} = 59^{+36}_{-19}$  GeV and spin-independent  $\chi^0$ -proton cross section of  $(1.0^{+0.1}_{-0.4}) \times 10^{-5}$  pb ( $1\sigma$  errors).

**Other bounds on  $\tilde{\chi}_1^0$  from astrophysics and cosmology**

Most of these papers generally exclude regions in the  $M_2 - \mu$  parameter plane by requiring that the  $\tilde{\chi}_1^0$  contribution to the overall cosmological density is less than some maximal value to avoid overclosure of the Universe. Those not based on the cosmological density are indicated. Many of these papers also include LEP and/or other bounds.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&gt;46 GeV</b>	57 ELLIS	00	RVUE
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
> 6 GeV	58,59 BELANGER	04	THEO
	60 ELLIS	04B	COSM
	61 PIERCE	04A	COSM
	62 BAER	03	COSM
> 6 GeV	58 BOTTINO	03	COSM
	62 CHATTOPAD...03		COSM
	63 ELLIS	03	COSM
	64 ELLIS	03B	COSM
	62 ELLIS	03C	COSM
> 18 GeV	58 HOOPER	03	COSM $\Omega_\chi = 0.05-0.3$
	62 LAHANAS	03	COSM
	65 BAER	02	COSM
	66 ELLIS	02	COSM
	67 LAHANAS	02	COSM
	68 BARGER	01C	COSM
	65 DJOUADI	01	COSM
	69 ELLIS	01B	COSM
	65 ROSZKOWSKI	01	COSM
	63 BOEHM	00B	COSM
	70 FENG	00	COSM
	71 LAHANAS	00	COSM
< 600 GeV	72 ELLIS	98B	COSM
	73 EDSJO	97	COSM Co-annihilation
	74 BAER	96	COSM
	75 BEREZINSKY	95	COSM
	76 FALK	95	COSM CP-violating phases
	77 DREES	93	COSM Minimal supergravity
	78 FALK	93	COSM Sfermion mixing
	77 KELLEY	93	COSM Minimal supergravity
	79 MIZUTA	93	COSM Co-annihilation
	80 LOPEZ	92	COSM Minimal supergravity, $m_0=A=0$
	81 MCDONALD	92	COSM
	82 GRIEST	91	COSM
	83 NOJIRI	91	COSM Minimal supergravity
	84 OLIVE	91	COSM
	85 ROSZKOWSKI	91	COSM
	86 GRIEST	90	COSM
	84 OLIVE	89	COSM
none 100 eV – 15 GeV	SREDNICKI	88	COSM $\tilde{\gamma}; m_{\tilde{f}}=100$ GeV



none	100 eV–5 GeV	ELLIS	84	COSM	$\tilde{\gamma}$ ; for $m_{\tilde{f}}=100$ GeV
		GOLDBERG	83	COSM	$\tilde{\gamma}$
		87 KRAUSS	83	COSM	$\tilde{\gamma}$
		VYSOTSKII	83	COSM	$\tilde{\gamma}$

- 57 ELLIS 00 updates ELLIS 98. Uses LEP  $e^+e^-$  data at  $\sqrt{s}=202$  and 204 GeV to improve bound on neutralino mass to 51 GeV when scalar mass universality is assumed and 46 GeV when Higgs mass universality is relaxed. Limits on  $\tan\beta$  improve to  $> 2.7$  ( $\mu > 0$ ),  $> 2.2$  ( $\mu < 0$ ) when scalar mass universality is assumed and  $> 1.9$  (both signs of  $\mu$ ) when Higgs mass universality is relaxed.
- 58 HOOPER 03, BOTTINO 03 (see also BOTTINO 03A and BOTTINO 04) , and BELANGER 04 do not assume gaugino or scalar mass unification.
- 59 Limit assumes a pseudo scalar mass  $< 200$  GeV. For larger pseudo scalar masses,  $m_{\chi} > 18(29)$  GeV for  $\tan\beta = 50(10)$ . Bounds from WMAP,  $(g-2)_{\mu}$ ,  $b \rightarrow s\gamma$ , LEP.
- 60 ELLIS 04B places constraints on the SUSY parameter space in the framework of  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry including supersymmetry breaking relations between A and B parameters. See also ELLIS 03D.
- 61 PIERCE 04A places constraints on the SUSY parameter space in the framework of models with very heavy scalar masses.
- 62 BAER 03, CHATTOPADHYAY 03, ELLIS 03C and LAHANAS 03 place constraints on the SUSY parameter space in the framework of  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry based on WMAP results for the cold dark matter density.
- 63 BOEHM 00B and ELLIS 03 place constraints on the SUSY parameter space in the framework of minimal  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry. Includes the effect of  $\chi-\tilde{t}$  co-annihilations.
- 64 BEREZINSKY 95 and ELLIS 03B places constraints on the SUSY parameter space in the framework of  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry but non-Universal Higgs masses.
- 65 DJOUADI 01, ROSZKOWSKI 01, and BAER 02 place constraints on the SUSY parameter space in the framework of minimal  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry.
- 66 ELLIS 02 places constraints on the soft supersymmetry breaking masses in the framework of minimal  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry.
- 67 LAHANAS 02 places constraints on the SUSY parameter space in the framework of minimal  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry. Focuses on the role of pseudo-scalar Higgs exchange.
- 68 BARGER 01C use the cosmic relic density inferred from recent CMB measurements to constrain the parameter space in the framework of minimal  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry.
- 69 ELLIS 01B places constraints on the SUSY parameter space in the framework of minimal  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry. Focuses on models with large  $\tan\beta$ .
- 70 FENG 00 explores cosmologically allowed regions of MSSM parameter space with multi-TeV masses.
- 71 LAHANAS 00 use the new cosmological data which favor a cosmological constant and its implications on the relic density to constrain the parameter space in the framework of minimal  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry.
- 72 ELLIS 98B assumes a universal scalar mass and radiative supersymmetry breaking with universal gaugino masses. The upper limit to the LSP mass is increased due to the inclusion of  $\chi - \tilde{\tau}_R$  coannihilations.
- 73 EDSJO 97 included all coannihilation processes between neutralinos and charginos for any neutralino mass and composition.

- 74 Notes the location of the neutralino  $Z$  resonance and  $h$  resonance annihilation corridors in minimal supergravity models with radiative electroweak breaking.
- 75 BEREZINSKY 95 and ELLIS 02C places constraints on the SUSY parameter space in the framework of  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry but non-Universal Higgs masses.
- 76 Mass of the bino (=LSP) is limited to  $m_{\tilde{B}} \lesssim 350$  GeV for  $m_t = 174$  GeV.
- 77 DREES 93, KELLEY 93 compute the cosmic relic density of the LSP in the framework of minimal  $N=1$  supergravity models with radiative breaking of the electroweak gauge symmetry.
- 78 FALK 93 relax the upper limit to the LSP mass by considering sfermion mixing in the MSSM.
- 79 MIZUTA 93 include coannihilations to compute the relic density of Higgsino dark matter.
- 80 LOPEZ 92 calculate the relic LSP density in a minimal SUSY GUT model.
- 81 MCDONALD 92 calculate the relic LSP density in the MSSM including exact tree-level annihilation cross sections for all two-body final states.
- 82 GRIEST 91 improve relic density calculations to account for coannihilations, pole effects, and threshold effects.
- 83 NOJIRI 91 uses minimal supergravity mass relations between squarks and sleptons to narrow cosmologically allowed parameter space.
- 84 Mass of the bino (=LSP) is limited to  $m_{\tilde{B}} \lesssim 350$  GeV for  $m_t \leq 200$  GeV. Mass of the higgsino (=LSP) is limited to  $m_{\tilde{H}} \lesssim 1$  TeV for  $m_t \leq 200$  GeV.
- 85 ROSZKOWSKI 91 calculates LSP relic density in mixed gaugino/higgsino region.
- 86 Mass of the bino (=LSP) is limited to  $m_{\tilde{B}} \lesssim 550$  GeV. Mass of the higgsino (=LSP) is limited to  $m_{\tilde{H}} \lesssim 3.2$  TeV.
- 87 KRAUSS 83 finds  $m_{\tilde{\gamma}}$  not 30 eV to 2.5 GeV. KRAUSS 83 takes into account the gravitino decay. Find that limits depend strongly on reheated temperature. For example a new allowed region  $m_{\tilde{\gamma}} = 4\text{--}20$  MeV exists if  $m_{\text{gravitino}} < 40$  TeV. See figure 2.

### ———— Unstable $\tilde{\chi}_1^0$ (Lightest Neutralino) MASS LIMIT ————

Unless otherwise stated, results in this section assume spectra and production rates as evaluated in the MSSM. Unless otherwise stated, the goldstino or gravitino mass  $m_{\tilde{G}}$  is assumed to be negligible relative to all other masses. In the following,  $\tilde{G}$  is assumed to be undetected and to give rise to a missing energy ( $\cancel{E}$ ) signature.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>108	95	88 ABAZOV	05A D0	$p\bar{p} \rightarrow \tilde{\chi}\tilde{\chi}, \tilde{\chi}=\tilde{\chi}_2^0, \tilde{\chi}_1^\pm, \tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}, \text{GMSB}$
		89 ABDALLAH	05B DLPH	$e^+e^- \rightarrow \tilde{G}\tilde{\chi}_1^0, (\tilde{\chi}_1^0 \rightarrow \tilde{G}\gamma)$
> 96	95	90 ABDALLAH	05B DLPH	$e^+e^- \rightarrow \tilde{B}\tilde{B}, (\tilde{B} \rightarrow \tilde{G}\gamma)$
> 93	95	91 ACOSTA	05E CDF	$p\bar{p} \rightarrow \tilde{\chi}\tilde{\chi}, \tilde{\chi}=\tilde{\chi}_2^0, \tilde{\chi}_1^\pm, \tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}, \text{GMSB}$
		92 AKTAS	05 H1	$e^\pm p \rightarrow q\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}, \text{GMSB}+\cancel{R} LQ\bar{D}$
		93 ABBIENDI	04N OPAL	$e^+e^- \rightarrow \gamma\gamma\cancel{E}$
> 66	95	94,95 ABDALLAH	04H DLPH	AMSB, $\mu > 0$
> 38.0	95	96,97 ABDALLAH	04M DLPH	$\cancel{R}(\bar{U}\bar{D}\bar{D})$

		98	ACHARD	04E L3	$e^+e^- \rightarrow \tilde{G}\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \tilde{G}\gamma$
> 99.5	95	99	ACHARD	04E L3	$e^+e^- \rightarrow \tilde{B}\tilde{B}, (\tilde{B} \rightarrow \tilde{G}\gamma)$
> 89	95	100	ABDALLAH	03D DLPH	$e^+e^- \rightarrow \tilde{\chi}_1^0\tilde{\chi}_1^0, \text{GMSB}, m(\tilde{G}) < 1 \text{ eV}$
		101	HEISTER	03C ALEP	$e^+e^- \rightarrow \tilde{B}\tilde{B}, (\tilde{B} \rightarrow \gamma\tilde{G})$
		102	HEISTER	03C ALEP	$e^+e^- \rightarrow \tilde{G}\tilde{\chi}_1^0, (\tilde{\chi}_1^0 \rightarrow \tilde{G}\gamma)$
> 39.9	95	103	ACHARD	02 L3	$\cancel{E}_T, \text{MSUGRA}$
> 92	95	104	HEISTER	02R ALEP	short lifetime
> 54	95	104	HEISTER	02R ALEP	any lifetime
> 85	95	105	ABBIENDI	01 OPAL	$e^+e^- \rightarrow \tilde{\chi}_1^0\tilde{\chi}_1^0, \text{GMSB}, \tan\beta=2$
> 76	95	105	ABBIENDI	01 OPAL	$e^+e^- \rightarrow \tilde{\chi}_1^0\tilde{\chi}_1^0, \text{GMSB}, \tan\beta=20$
> 32.5	95	106	ACCIARRI	01 L3	$\cancel{E}_T, \text{all } m_0, 0.7 \leq \tan\beta \leq 40$
		107	ADAMS	01 NTEV	$\tilde{\chi}^0 \rightarrow \mu\mu\nu, \cancel{E}_T, LL\bar{E}$
> 29	95	108	ABBIENDI	99T OPAL	$e^+e^- \rightarrow \tilde{\chi}_1^0\tilde{\chi}_1^0, \cancel{E}_T, m_0=500 \text{ GeV}, \tan\beta > 1.2$
		109	ACCIARRI	99R L3	Superseded by ACHARD 04E
> 88.2	95	110	ACCIARRI	99R L3	Superseded by ACHARD 04E
> 29	95	111	BARATE	99E ALEP	$\cancel{E}_T, LQ\bar{D}, \tan\beta=1.41, m_0=500 \text{ GeV}$
		112	ABREU	98 DLPH	$e^+e^- \rightarrow \tilde{\chi}_1^0\tilde{\chi}_1^0 (\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G})$
> 23	95	113	BARATE	98S ALEP	$\cancel{E}_T, LL\bar{E}$
		114	ELLIS	97 THEO	$e^+e^- \rightarrow \tilde{\chi}_1^0\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$
		115	CABIBBO	81 COSM	

88 ABAZOV 05A looked in  $263 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96 \text{ TeV}$  for diphoton events with large  $\cancel{E}_T$ . They may originate from the production of  $\tilde{\chi}^\pm$  in pairs or associated to a  $\tilde{\chi}_2^0$ , decaying to a  $\tilde{\chi}_1^0$  which itself decays promptly in GMSB to  $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$ . No significant excess was found at large  $\cancel{E}_T$  compared to the background expectation. A limit is derived on the masses of SUSY particles in the GMSB framework for  $M = 2 \Lambda$ ,  $N = 1$ ,  $\tan\beta = 15$  and  $\mu > 0$ , see Figure 2. It also excludes  $\Lambda < 79.6 \text{ TeV}$ . Very similar results are obtained for different choices of parameters, see their Table 2. Supersedes the results of ABBOTT 98.

89 ABDALLAH 05B use data from  $\sqrt{s} = 180\text{--}209 \text{ GeV}$ . They look for events with single photons +  $\cancel{E}_T$  final states. Limits are computed in the plane  $(m(\tilde{G}), m(\tilde{\chi}_1^0))$ , shown in their Fig. 9b for a pure Bino state in the GMSB framework and in Fig. 9c for a no-scale supergravity model. Supersedes the results of ABREU 00Z.

90 ABDALLAH 05B use data from  $\sqrt{s} = 130\text{--}209 \text{ GeV}$ . They look for events with diphotons +  $\cancel{E}_T$  final states and single photons not pointing to the vertex, expected in GMSB when the  $\tilde{\chi}_1^0$  is the NLSP. Limits are computed in the plane  $(m(\tilde{G}), m(\tilde{\chi}_1^0))$ , see their Fig. 10. The lower limit is derived on the  $\tilde{\chi}_1^0$  mass for a pure Bino state assuming a prompt decay and  $m_{\tilde{e}_R} = m_{\tilde{e}_L} = 2 m_{\tilde{\chi}_1^0}$ . It improves to  $100 \text{ GeV}$  for  $m_{\tilde{e}_R} = m_{\tilde{e}_L} = 1.1 m_{\tilde{\chi}_1^0}$  and the limit in the plane  $(m(\tilde{\chi}_1^0), m(\tilde{e}_R))$  is shown in Fig. 10b. For long-lived neutralinos, cross-section limits are displayed in their Fig 11. Supersedes the results of ABREU 00Z.

91 ACOSTA 05E looked in  $202 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.96 \text{ TeV}$  for diphoton events with large  $\cancel{E}_T$ . They may originate from the production of  $\tilde{\chi}^\pm$  in pairs or associated to a  $\tilde{\chi}_2^0$ , decaying to a  $\tilde{\chi}_1^0$  which itself decays promptly in GMSB to  $\gamma\tilde{G}$ . No events are selected at large  $\cancel{E}_T$  compared to the background expectation. A limit is derived on the masses of SUSY particles in the GMSB framework for  $M = 2 \Lambda$ ,  $N = 1$ ,  $\tan\beta = 15$  and  $\mu > 0$ , see Figure 2. It also excludes  $\Lambda < 69 \text{ TeV}$ . Supersedes the results of ABE 99I.

- 92 AKTAS 05 data collected at 319 GeV with  $64.3 \text{ pb}^{-1}$  of  $e^+p$  and  $13.5 \text{ pb}^{-1}$  of  $e^-p$ . They look for  $\cancel{R}$  resonant  $\tilde{\chi}_1^0$  production via  $t$ -channel exchange of a  $\tilde{e}$ , followed by prompt GMSB decay of the  $\tilde{\chi}_1^0$  to  $\gamma\tilde{G}$ . Upper limits at 95% on the cross section are derived, see their Figure 4, and compared to two example scenarios. In Figure 5, they display 95% exclusion limits in the plane of  $M(\tilde{\chi}_1^0)$  versus  $M(\tilde{e}_L) - M(\tilde{\chi}_1^0)$  for the two scenarios and several values of the  $\lambda'$  Yukawa coupling.
- 93 ABBIENDI 04N use data from  $\sqrt{s} = 189\text{--}209$  GeV, setting limits on  $\sigma(e^+e^- \rightarrow XX) \times B^2(X \rightarrow Y\gamma)$ , with  $Y$  invisible (see their Fig. 4). Limits on  $\tilde{\chi}_1^0$  masses for a specific model are given. Supersedes the results of ABBIENDI, G 00D.
- 94 ABDALLAH 04H use data from LEP 1 and  $\sqrt{s} = 192\text{--}208$  GeV. They re-use results or re-analyze the data from ABDALLAH 03M to put limits on the parameter space of anomaly-mediated supersymmetry breaking (AMSB), which is scanned in the region  $1 < m_{3/2} < 50$  TeV,  $0 < m_0 < 1000$  GeV,  $1.5 < \tan\beta < 35$ , both signs of  $\mu$ . The constraints are obtained from the searches for mass degenerate chargino and neutralino, for SM-like and invisible Higgs, for leptonically decaying charginos and from the limit on non-SM  $Z$  width of 3.2 MeV. The limit is for  $m_t = 174.3$  GeV (see Table 2 for other  $m_t$  values).
- 95 The limit improves to 73 GeV for  $\mu < 0$ .
- 96 ABDALLAH 04M use data from  $\sqrt{s} = 192\text{--}208$  GeV to derive limits on sparticle masses under the assumption of  $\cancel{R}$  with  $LL\bar{E}$  or  $\overline{UD}\bar{D}$  couplings. The results are valid in the ranges  $90 < m_0 < 500$  GeV,  $0.7 < \tan\beta < 30$ ,  $-200 < \mu < 200$  GeV,  $0 < M_2 < 400$  GeV. Supersedes the result of ABREU 01D and ABREU 00U.
- 97 The limit improves to 39.5 GeV for  $LL\bar{E}$  couplings.
- 98 ACHARD 04E use data from  $\sqrt{s} = 189\text{--}209$  GeV. They look for events with single photons +  $\cancel{R}$  final states. Limits are computed in the plane ( $m(\tilde{G})$ ,  $m(\tilde{\chi}_1^0)$ ), shown in their Fig. 8c for a no-scale supergravity model, excluding, e.g., Gravitino masses below  $10^{-5}$  eV for neutralino masses below 172 GeV. Supersedes the results of ACCIARRI 99R.
- 99 ACHARD 04E use data from  $\sqrt{s} = 189\text{--}209$  GeV. They look for events with diphotons +  $\cancel{R}$  final states. Limits are computed in the plane ( $m(\tilde{\chi}_1^0)$ ,  $m(\tilde{e}_R)$ ), see their Fig. 8d. The limit on the  $\tilde{\chi}_1^0$  mass is for a pure Bino state assuming a prompt decay, with  $m_{\tilde{e}_L} = 1.1 m_{\tilde{\chi}_1^0}$  and  $m_{\tilde{e}_R} = 2.5 m_{\tilde{\chi}_1^0}$ . Supersedes the results of ACCIARRI 99R.
- 100 ABDALLAH 03D use data from  $\sqrt{s} = 161\text{--}208$  GeV. They look for 4-tau +  $\cancel{R}$  final states, expected in GMSB when the  $\tilde{\tau}_1$  is the NLSP, and 4-lepton +  $\cancel{R}$  final states, expected in the co-NLSP scenario, and assuming a short-lived  $\tilde{\chi}_1^0$  ( $m(\tilde{G}) < 1$  eV). Limits are computed in the plane ( $m(\tilde{\tau}_1)$ ,  $m(\tilde{\chi}_1^0)$ ) from a scan of the GMSB parameters space, after combining these results with the search for slepton pair production from the same paper to cover prompt decays and for the case of  $\tilde{\chi}_1^0$  NLSP from ABREU 00Z. The limit above is reached for a single generation of messengers and when the  $\tilde{\tau}_1$  is the NLSP. Stronger limits are obtained when more messenger generations are assumed or when the other sleptons are co-NLSP, see their Fig. 10. Supersedes the results of ABREU 01G.
- 101 HEISTER 03C use the data from  $\sqrt{s} = 189\text{--}209$  GeV to search for  $\gamma\cancel{R}_T$  final states with non-pointing photons and  $\gamma\cancel{R}_T$  events. Interpreted in the framework of Minimal GMSB, a lower bound on the  $\tilde{\chi}_1^0$  mass is obtained as function of its lifetime. For a laboratory lifetime of less than 3 ns, the limit at 95% CL is 98.8 GeV. For other lifetimes, see their Fig. 5. These results are interpreted in a more general GMSB framework in HEISTER 02R.
- 102 HEISTER 03C use the data from  $\sqrt{s} = 189\text{--}209$  GeV to search for  $\gamma\cancel{R}_T$  final states. They obtained an upper bound on the cross section for the process  $e^+e^- \rightarrow \tilde{G}\tilde{\chi}_1^0$ , followed by the prompt decay  $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$ , shown in their Fig. 4. These results supersede BARATE 98H.
- 103 ACHARD 02 searches for the production of sparticles in the case of  $\cancel{R}$  prompt decays with  $LL\bar{E}$  or  $\overline{UDD}$  couplings at  $\sqrt{s} = 189\text{--}208$  GeV. The search is performed for direct and

- indirect decays, assuming one coupling at the time to be nonzero. The MSUGRA limit results from a scan over the MSSM parameter space with the assumption of gaugino and scalar mass unification at the GUT scale, imposing simultaneously the exclusions from neutralino, chargino, sleptons, and squarks analyses. The limit holds for  $\overline{UDD}$  couplings and increases to 40.2 GeV for  $LL\overline{E}$  couplings. For L3 limits from  $LQ\overline{D}$  couplings, see ACCIARRI 01.
- 104 HEISTER 02R search for signals of GMSB in the 189–209 GeV data. For the  $\tilde{\chi}_1^0$  NLSP scenario, they looked for topologies consisting of  $\gamma\gamma\cancel{E}$  or a single  $\gamma$  not pointing to the interaction vertex. For the  $\tilde{\ell}$  NLSP case, the topologies consist of  $\ell\ell\cancel{E}$  or  $4\ell\cancel{E}$  (from  $\tilde{\chi}_1^0\tilde{\chi}_1^0$  production), including leptons with large impact parameters, kinks, or stable particles. Limits are derived from a scan over the GMSB parameters (see their Table 5 for the ranges). The limits are valid whichever is the NLSP. The absolute mass bound on the  $\tilde{\chi}_1^0$  for any lifetime includes indirect limits from the chargino search, and from the slepton search HEISTER 02E performed within the MSUGRA framework. A bound for any NLSP and any lifetime of 77 GeV has also been derived by using the constraints from the neutral Higgs search in HEISTER 02. Limits on the universal SUSY mass scale  $\Lambda$  are also derived in the paper. Supersedes the results from BARATE 00G.
- 105 ABBIENDI 01 looked for final states with  $\gamma\gamma\cancel{E}$ ,  $\ell\ell\cancel{E}$ , with possibly additional activity and four leptons +  $\cancel{E}$  to search for prompt decays of  $\tilde{\chi}_1^0$  or  $\tilde{\ell}_1$  in GMSB. They derive limits in the plane  $(m_{\tilde{\chi}_1^0}, m_{\tilde{\tau}_1})$ , see Fig. 6, allowing either the  $\tilde{\chi}_1^0$  or a  $\tilde{\ell}_1$  to be the NLSP. Two scenarios are considered:  $\tan\beta=2$  with the 3 sleptons degenerate in mass and  $\tan\beta=20$  where the  $\tilde{\tau}_1$  is lighter than the other sleptons. Data taken at  $\sqrt{s}=189$  GeV.
- 106 ACCIARRI 01 searches for multi-lepton and/or multi-jet final states from  $\cancel{R}$  prompt decays with  $LL\overline{E}$ ,  $LQ\overline{D}$ , or  $\overline{UDD}$  couplings at  $\sqrt{s}=189$  GeV. The search is performed for direct and indirect decays of neutralinos, charginos, and scalar leptons, with the  $\tilde{\chi}_1^0$  or a  $\tilde{\ell}$  as LSP and assuming one coupling to be nonzero at a time. Mass limits are derived using simultaneously the constraints from the neutralino, chargino, and slepton analyses; and the  $Z^0$  width measurements from ACCIARRI 00C in a scan of the parameter space assuming MSUGRA with gaugino and scalar mass universality. Updates and supersedes the results from ACCIARRI 99I.
- 107 ADAMS 01 looked for neutral particles with mass  $> 2.2$  GeV, produced by 900 GeV protons incident on a Beryllium oxide target and decaying through weak interactions into  $\mu\mu$ ,  $\mu e$ , or  $\mu\pi$  final states in the decay channel of the NuTeV detector (E815) at Fermilab. The number of observed events is  $3\mu\mu$ ,  $0\mu e$ , and  $0\mu\pi$  with an expected background of  $0.069 \pm 0.010$ ,  $0.13 \pm 0.02$ , and  $0.14 \pm 0.02$ , respectively. The  $\mu\mu$  events are consistent with the  $\cancel{R}$  decay of a neutralino with mass around 5 GeV. However, they share several aspects with  $\nu$ -interaction backgrounds. An upper limit on the differential production cross section of neutralinos in  $p\bar{p}$  interactions as function of the decay length is given in Fig. 3.
- 108 ABBIENDI 99T searches for the production of neutralinos in the case of  $R$ -parity violation with  $LL\overline{E}$ ,  $LQ\overline{D}$ , or  $\overline{UDD}$  couplings using data from  $\sqrt{s}=183$  GeV. They investigate topologies with multiple leptons, jets plus leptons, or multiple jets, assuming one coupling at the time to be non-zero and giving rise to direct or indirect decays. Mixed decays (where one particle has a direct, the other an indirect decay) are also considered for the  $\overline{UDD}$  couplings. Upper limits on the cross section are derived which, combined with the constraint from the  $Z^0$  width, allow to exclude regions in the  $M_2$  versus  $\mu$  plane for any coupling. Limits on the neutralino mass are obtained for non-zero  $LL\overline{E}$  couplings  $> 10^{-5}$ . The limit disappears for  $\tan\beta < 1.2$  and it improves to 50 GeV for  $\tan\beta > 20$ .
- 109 ACCIARRI 99R searches for  $\gamma\cancel{E}$  final states using data from  $\sqrt{s}=189$  GeV. From limits on cross section times branching ratio, mass limits are derived in a no-scale SUGRA model, see their Fig. 5. Supersedes the results of ACCIARRI 98v.
- 110 ACCIARRI 99R searches for  $\gamma\cancel{E}$  final states using data from  $\sqrt{s}=189$  GeV. From a scan over the GMSB parameter space, a limit on the mass is derived under the assumption that the neutralino is the NLSP. Supersedes the results of ACCIARRI 98v.

- 111 BARATE 99E looked for the decay of gauginos via  $R$ -violating couplings  $LQ\bar{D}$ . The bound is significantly reduced for smaller values of  $m_0$ . Data collected at  $\sqrt{s}=130\text{--}172$  GeV.
- 112 ABREU 98 uses data at  $\sqrt{s}=161$  and 172 GeV. Upper bounds on  $\gamma\gamma\cancel{E}$  cross section are obtained. Similar limits on  $\gamma\cancel{E}$  are also given, relevant for  $e^+e^- \rightarrow \tilde{\chi}_1^0\tilde{G}$  production.
- 113 BARATE 98S looked for the decay of gauginos via  $R$ -violating coupling  $LL\bar{E}$ . The bound improves to 25 GeV if the chargino decays into neutralino which further decays into lepton pairs. Data collected at  $\sqrt{s}=130\text{--}172$  GeV.
- 114 ELLIS 97 reanalyzed the LEP2 ( $\sqrt{s}=161$  GeV) limits of  $\sigma(\gamma\gamma+E_{\text{miss}}) < 0.2$  pb to exclude  $m_{\tilde{\chi}_1^0} < 63$  GeV if  $m_{\tilde{e}_L} = m_{\tilde{e}_R} < 150$  GeV and  $\tilde{\chi}_1^0$  decays to  $\gamma\tilde{G}$  inside detector.
- 115 CABIBBO 81 consider  $\tilde{\gamma} \rightarrow \gamma + \text{goldstino}$ . Photino must be either light enough ( $< 30$  eV) to satisfy cosmology bound, or heavy enough ( $> 0.3$  MeV) to have disappeared at early universe.

## $\tilde{\chi}_2^0, \tilde{\chi}_3^0, \tilde{\chi}_4^0$ (Neutralinos) MASS LIMITS

Neutralinos are unknown mixtures of photinos, z-inos, and neutral higgsinos (the supersymmetric partners of photons and of  $Z$  and Higgs bosons). The limits here apply only to  $\tilde{\chi}_2^0, \tilde{\chi}_3^0,$  and  $\tilde{\chi}_4^0$ .  $\tilde{\chi}_1^0$  is the lightest supersymmetric particle (LSP); see  $\tilde{\chi}_1^0$  Mass Limits. It is not possible to quote rigorous mass limits because they are extremely model dependent; i.e. they depend on branching ratios of various  $\tilde{\chi}^0$  decay modes, on the masses of decay products ( $\tilde{e}, \tilde{\gamma}, \tilde{q}, \tilde{g}$ ), and on the  $\tilde{e}$  mass exchanged in  $e^+e^- \rightarrow \tilde{\chi}_i^0\tilde{\chi}_j^0$ . Limits arise either from direct searches, or from the MSSM constraints set on the gaugino and higgsino mass parameters  $M_2$  and  $\mu$  through searches for lighter charginos and neutralinos. Often limits are given as contour plots in the  $m_{\tilde{\chi}^0} - m_{\tilde{e}}$  plane vs other parameters. When specific assumptions are made, e.g. the neutralino is a pure photino ( $\tilde{\gamma}$ ), pure z-ino ( $\tilde{Z}$ ), or pure neutral higgsino ( $\tilde{H}^0$ ), the neutralinos will be labelled as such.

Limits obtained from  $e^+e^-$  collisions at energies up to 136 GeV, as well as other limits from different techniques, are now superseded and have not been included in this compilation. They can be found in the 1998 Edition (The European Physical Journal **C3** 1 (1998)) of this Review.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
$> 78$	95	116 ABBIENDI	04H OPAL	$\tilde{\chi}_2^0$ , all $\tan\beta, \Delta m_0 > 5$ GeV, $m_0 > 500$ GeV, $A_0 = 0$
$> 62.4$	95	117 ABREU	00W DLPH	$\tilde{\chi}_2^0, 1 \leq \tan\beta \leq 40$ , all $\Delta m_0$ , all $m_0$
$> 99.9$	95	117 ABREU	00W DLPH	$\tilde{\chi}_3^0, 1 \leq \tan\beta \leq 40$ , all $\Delta m_0$ , all $m_0$
$> 116.0$	95	117 ABREU	00W DLPH	$\tilde{\chi}_4^0, 1 \leq \tan\beta \leq 40$ , all $\Delta m_0$ , all $m_0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
		118 ABDALLAH	05B DLPH	$e^+e^- \rightarrow \tilde{\chi}_2^0\tilde{\chi}_2^0, (\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0\gamma)$
		119 ACHARD	04E L3	$e^+e^- \rightarrow \tilde{\chi}_2^0\tilde{\chi}_2^0, (\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0\gamma)$
$> 80.0$	95	120 ACHARD	02 L3	$\tilde{\chi}_2^0, \mathcal{R}, \text{MSUGRA}$
$> 107.2$	95	120 ACHARD	02 L3	$\tilde{\chi}_3^0, \mathcal{R}, \text{MSUGRA}$
		121 ABREU	01B DLPH	$e^+e^- \rightarrow \tilde{\chi}_i^0\tilde{\chi}_j^0$
$> 68.0$	95	122 ACCIARRI	01 L3	$\tilde{\chi}_2^0, \mathcal{R}$ , all $m_0, 0.7 \leq \tan\beta \leq 40$

> 99.0	95	122 ACCIARRI	01 L3	$\tilde{\chi}_3^0$ , $\mathcal{R}$ , all $m_0$ , $0.7 \leq \tan\beta \leq 40$
> 50	95	123 ABREU	00U DLPH	$\tilde{\chi}_2^0$ , $\mathcal{R}$ ( $LL\bar{E}$ ), all $\Delta m_0$ , $1 \leq \tan\beta \leq 30$
		124 ABBIENDI	99F OPAL	$e^+e^- \rightarrow \tilde{\chi}_2^0\tilde{\chi}_1^0$ ( $\tilde{\chi}_2^0 \rightarrow \gamma\tilde{\chi}_1^0$ )
		125 ABBIENDI	99F OPAL	$e^+e^- \rightarrow \tilde{\chi}_2^0\tilde{\chi}_2^0$ ( $\tilde{\chi}_2^0 \rightarrow \gamma\tilde{\chi}_1^0$ )
		126 ABBOTT	98C D0	$p\bar{p} \rightarrow \tilde{\chi}_1^\pm\tilde{\chi}_2^0$
> 82.2	95	127 ABE	98J CDF	$p\bar{p} \rightarrow \tilde{\chi}_1^\pm\tilde{\chi}_2^0$
> 92	95	128 ACCIARRI	98F L3	$\tilde{H}_2^0$ , $\tan\beta=1.41$ , $M_2 < 500$ GeV
		129 ACCIARRI	98V L3	$e^+e^- \rightarrow \tilde{\chi}_2^0\tilde{\chi}_{1,2}^0$ ( $\tilde{\chi}_2^0 \rightarrow \gamma\tilde{\chi}_1^0$ )
> 53	95	130 BARATE	98H ALEP	$e^+e^- \rightarrow \tilde{\gamma}\tilde{\gamma}$ ( $\tilde{\gamma} \rightarrow \gamma\tilde{H}^0$ )
> 74	95	131 BARATE	98J ALEP	$e^+e^- \rightarrow \tilde{\gamma}\tilde{\gamma}$ ( $\tilde{\gamma} \rightarrow \gamma\tilde{H}^0$ )
		132 ABACHI	96 D0	$p\bar{p} \rightarrow \tilde{\chi}_1^\pm\tilde{\chi}_2^0$
		133 ABE	96K CDF	$p\bar{p} \rightarrow \tilde{\chi}_1^\pm\tilde{\chi}_2^0$

- 116 ABBIENDI 04H search for charginos and neutralinos in events with acoplanar leptons+jets and multi-jet final states in the 192–209 GeV data, combined with the results on leptonic final states from ABBIENDI 04. The results hold for a scan over the parameter space covering the region  $0 < M_2 < 5000$  GeV,  $-1000 < \mu < 1000$  GeV and  $\tan\beta$  from 1 to 40. This limit supersedes ABBIENDI 00H.
- 117 ABREU 00W combines data collected at  $\sqrt{s}=189$  GeV with results from lower energies. The mass limit is obtained by constraining the MSSM parameter space with gaugino and sfermion mass universality at the GUT scale, using the results of negative direct searches for neutralinos (including cascade decays and  $\tilde{\tau}\tau$  final states) from ABREU 01, for charginos from ABREU 00J and ABREU 00T (for all  $\Delta m_+$ ), and for charged sleptons from ABREU 01B. The results hold for the full parameter space defined by all values of  $M_2$  and  $|\mu| \leq 2$  TeV with the  $\tilde{\chi}_1^0$  as LSP.
- 118 ABDALLAH 05B use data from  $\sqrt{s} = 130$ –209 GeV, looking for events with diphotons +  $\cancel{E}$ . Limits on the cross-section are computed in the plane  $(m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0))$ , see Fig. 12. Supersedes the results of ABREU 00Z.
- 119 ACHARD 04E use data from  $\sqrt{s} = 189$ –209 GeV, looking for events with diphotons +  $\cancel{E}$ . Limits are computed in the plane  $(m(\tilde{\chi}_2^0), m(\tilde{e}_R))$ , for  $\Delta m_0 > 10$  GeV, see Fig. 7. Supersedes the results of ACCIARRI 99R.
- 120 ACHARD 02 searches for the production of sparticles in the case of  $\mathcal{R}$  prompt decays with  $LL\bar{E}$  or  $UDD$  couplings at  $\sqrt{s}=189$ –208 GeV. The search is performed for direct and indirect decays, assuming one coupling at the time to be nonzero. The MSUGRA limit results from a scan over the MSSM parameter space with the assumption of gaugino and scalar mass unification at the GUT scale, imposing simultaneously the exclusions from neutralino, chargino, sleptons, and squarks analyses. The limit of  $\tilde{\chi}_2^0$  holds for  $UDD$  couplings and increases to 84.0 GeV for  $LL\bar{E}$  couplings. The same  $\tilde{\chi}_3^0$  limit holds for both  $LL\bar{E}$  and  $UDD$  couplings. For L3 limits from  $LQ\bar{D}$  couplings, see ACCIARRI 01.
- 121 ABREU 01B used data from  $\sqrt{s}=189$  GeV to search for the production of  $\tilde{\chi}_i^0\tilde{\chi}_j^0$ . They looked for di-jet and di-lepton pairs with  $\cancel{E}$  for events from  $\tilde{\chi}_i^0\tilde{\chi}_j^0$  with the decay  $\tilde{\chi}_j^0 \rightarrow f\bar{f}\tilde{\chi}_1^0$ ; multi-jet and multi-lepton pairs with or without additional photons to cover the cascade decays  $\tilde{\chi}_j^0 \rightarrow f\bar{f}\tilde{\chi}_2^0$ , followed by  $\tilde{\chi}_2^0 \rightarrow f\bar{f}\tilde{\chi}_1^0$  or  $\tilde{\chi}_2^0 \rightarrow \gamma\tilde{\chi}_1^0$ ; multi-tau final states from  $\tilde{\chi}_2^0 \rightarrow \tilde{\tau}\tau$  with  $\tilde{\tau} \rightarrow \tau\tilde{\chi}_1^0$ . See Figs. 9 and 10 for limits on the  $(\mu, M_2)$  plane for  $\tan\beta=1.0$  and different values of  $m_0$ .
- 122 ACCIARRI 01 searches for multi-lepton and/or multi-jet final states from  $\mathcal{R}$  prompt decays with  $LL\bar{E}$ ,  $LQ\bar{D}$ , or  $UDD$  couplings at  $\sqrt{s}=189$  GeV. The search is performed for

- direct and indirect decays of neutralinos, charginos, and scalar leptons, with the  $\tilde{\chi}_1^0$  or a  $\tilde{\ell}$  as LSP and assuming one coupling to be nonzero at a time. Mass limits are derived using simultaneously the constraints from the neutralino, chargino, and slepton analyses; and the  $Z^0$  width measurements from ACCIARRI 00C in a scan of the parameter space assuming MSUGRA with gaugino and scalar mass universality. Updates and supersedes the results from ACCIARRI 99I.
- 123 ABREU 00U searches for the production of charginos and neutralinos in the case of  $R$ -parity violation with  $LL\bar{E}$  couplings, using data from  $\sqrt{s}=189$  GeV. They investigate topologies with multiple leptons or jets plus leptons, assuming one coupling to be nonzero at the time and giving rise to direct or indirect decays. Limits are obtained in the  $M_2$  versus  $\mu$  plane and a limit on the neutralino mass is derived from a scan over the parameters  $m_0$  and  $\tan\beta$ .
- 124 ABBIENDI 99F looked for  $\gamma\cancel{E}$  final states at  $\sqrt{s}=183$  GeV. They obtained an upper bound on the cross section for the production  $e^+e^- \rightarrow \tilde{\chi}_2^0\tilde{\chi}_1^0$  followed by the prompt decay  $\tilde{\chi}_2^0 \rightarrow \gamma\tilde{\chi}_1^0$  of 0.075–0.80 pb in the region  $m_{\tilde{\chi}_2^0} + m_{\tilde{\chi}_1^0} > m_Z$ ,  $m_{\tilde{\chi}_2^0} = 91$ –183 GeV, and  $\Delta m_0 > 5$  GeV. See Fig. 7 for explicit limits in the  $(m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_1^0})$  plane.
- 125 ABBIENDI 99F looked for  $\gamma\gamma\cancel{E}$  final states at  $\sqrt{s}=183$  GeV. They obtained an upper bound on the cross section for the production  $e^+e^- \rightarrow \tilde{\chi}_2^0\tilde{\chi}_2^0$  followed by the prompt decay  $\tilde{\chi}_2^0 \rightarrow \gamma\tilde{\chi}_1^0$  of 0.08–0.37 pb for  $m_{\tilde{\chi}_2^0} = 45$ –81.5 GeV, and  $\Delta m_0 > 5$  GeV. See Fig. 11 for explicit limits in the  $(m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_1^0})$  plane.
- 126 ABBOTT 98C searches for trilepton final states ( $\ell=e,\mu$ ). See footnote to ABBOTT 98C in the Chargino Section for details on the assumptions. Assuming a negligible decay rate of  $\tilde{\chi}_1^\pm$  and  $\tilde{\chi}_2^0$  to quarks, they obtain  $m_{\tilde{\chi}_2^0} \gtrsim 103$  GeV.
- 127 ABE 98J searches for trilepton final states ( $\ell=e,\mu$ ). See footnote to ABE 98J in the Chargino Section for details on the assumptions. The quoted result for  $m_{\tilde{\chi}_2^0}$  corresponds to the best limit within the selected range of parameters, obtained for  $m_{\tilde{q}} > m_{\tilde{g}}$ ,  $\tan\beta=2$ , and  $\mu=-600$  GeV.
- 128 ACCIARRI 98F is obtained from direct searches in the  $e^+e^- \rightarrow \tilde{\chi}_{1,2}^0\tilde{\chi}_2^0$  production channels, and indirectly from  $\tilde{\chi}_1^\pm$  and  $\tilde{\chi}_1^0$  searches within the MSSM. See footnote to ACCIARRI 98F in the chargino Section for further details on the assumptions. Data taken at  $\sqrt{s} = 130$ –172 GeV.
- 129 ACCIARRI 98V looked for  $\gamma(\gamma)\cancel{E}$  final states at  $\sqrt{s}=183$  GeV. They obtained an upper bound on the cross section for the production  $e^+e^- \rightarrow \tilde{\chi}_2^0\tilde{\chi}_{1,2}^0$  followed by the prompt decay  $\tilde{\chi}_2^0 \rightarrow \gamma\tilde{\chi}_1^0$ . See Figs. 4a and 6a for explicit limits in the  $(m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_1^0})$  plane.
- 130 BARATE 98H looked for  $\gamma\gamma\cancel{E}$  final states at  $\sqrt{s} = 161, 172$  GeV. They obtained an upper bound on the cross section for the production  $e^+e^- \rightarrow \tilde{\chi}_2^0\tilde{\chi}_2^0$  followed by the prompt decay  $\tilde{\chi}_2^0 \rightarrow \gamma\tilde{\chi}_1^0$  of 0.4–0.8 pb for  $m_{\tilde{\chi}_2^0} = 10$ –80 GeV. The bound above is for the specific case of  $\tilde{\chi}_1^0 = \tilde{H}^0$  and  $\tilde{\chi}_2^0 = \tilde{\gamma}$  and  $m_{\tilde{e}_R} = 100$  GeV. See Fig. 6 and 7 for explicit limits in the  $(\tilde{\chi}_2^0, \tilde{\chi}_1^0)$  plane and in the  $(\tilde{\chi}_2^0, \tilde{e}_R)$  plane.
- 131 BARATE 98J looked for  $\gamma\gamma\cancel{E}$  final states at  $\sqrt{s} = 161$ –183 GeV. They obtained an upper bound on the cross section for the production  $e^+e^- \rightarrow \tilde{\chi}_2^0\tilde{\chi}_2^0$  followed by the prompt decay  $\tilde{\chi}_2^0 \rightarrow \gamma\tilde{\chi}_1^0$  of 0.08–0.24 pb for  $m_{\tilde{\chi}_2^0} < 91$  GeV. The bound above is for the specific case of  $\tilde{\chi}_1^0 = \tilde{H}^0$  and  $\tilde{\chi}_2^0 = \tilde{\gamma}$  and  $m_{\tilde{e}_R} = 100$  GeV.



- 132 ABACHI 96 searches for 3-lepton final states. Efficiencies are calculated using mass relations and branching ratios in the Minimal Supergravity scenario. Results are presented as lower bounds on  $\sigma(\tilde{\chi}_1^\pm \tilde{\chi}_2^0) \times B(\tilde{\chi}_1^\pm \rightarrow \ell \nu_\ell \tilde{\chi}_1^0) \times B(\tilde{\chi}_2^0 \rightarrow \ell^+ \ell^- \tilde{\chi}_1^0)$  as a function of  $m_{\tilde{\chi}_1^0}$ . Limits range from 3.1 pb ( $m_{\tilde{\chi}_1^0} = 45$  GeV) to 0.6 pb ( $m_{\tilde{\chi}_1^0} = 100$  GeV).
- 133 ABE 96K looked for trilepton events from chargino-neutralino production. They obtained lower bounds on  $m_{\tilde{\chi}_2^0}$  as a function of  $\mu$ . The lower bounds are in the 45–50 GeV range for gaugino-dominant  $\tilde{\chi}_2^0$  with negative  $\mu$ , if  $\tan\beta < 10$ . See paper for more details of the assumptions.

## $\tilde{\chi}_1^\pm, \tilde{\chi}_2^\pm$ (Charginos) MASS LIMITS

Charginos are unknown mixtures of  $w$ -inos and charged higgsinos (the supersymmetric partners of  $W$  and Higgs bosons). A lower mass limit for the lightest chargino ( $\tilde{\chi}_1^\pm$ ) of approximately 45 GeV, independent of the field composition and of the decay mode, has been obtained by the LEP experiments from the analysis of the  $Z$  width and decays. These results, as well as other now superseded limits from  $e^+e^-$  collisions at energies below 136 GeV, and from hadronic collisions, can be found in the 1998 Edition (The European Physical Journal **C3** 1 (1998)) of this Review.

Unless otherwise stated, results in this section assume spectra, production rates, decay modes and branching ratios as evaluated in the MSSM, with gaugino and sfermion mass unification at the GUT scale. These papers generally study production of  $\tilde{\chi}_1^0 \tilde{\chi}_2^0$ ,  $\tilde{\chi}_1^+ \tilde{\chi}_1^-$  and (in the case of hadronic collisions)  $\tilde{\chi}_1^+ \tilde{\chi}_2^0$  pairs, including the effects of cascade decays. The mass limits on  $\tilde{\chi}_1^\pm$  are either direct, or follow indirectly from the constraints set by the non-observation of  $\tilde{\chi}_2^0$  states on the gaugino and higgsino MSSM parameters  $M_2$  and  $\mu$ . For generic values of the MSSM parameters, limits from high-energy  $e^+e^-$  collisions coincide with the highest value of the mass allowed by phase-space, namely  $m_{\tilde{\chi}_1^\pm} \lesssim \sqrt{s}/2$ . At the time of this writing, preliminary and unpublished results from the 2000 run of LEP2 at  $\sqrt{s}$  up to  $\simeq 209$  GeV give therefore a lower mass limit of approximately 104 GeV valid for general MSSM models. The limits become however weaker in special regions of the MSSM parameter space where the detection efficiencies or production cross sections are suppressed. For example, this may happen when: (i) the mass differences  $\Delta m_{\pm} = m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0}$  or  $\Delta m_{\nu} = m_{\tilde{\chi}_1^\pm} - m_{\tilde{\nu}}$  are very small, and the detection efficiency is reduced; (ii) the electron sneutrino mass is small, and the  $\tilde{\chi}_1^\pm$  production rate is suppressed due to a destructive interference between  $s$  and  $t$  channel exchange diagrams. The regions of MSSM parameter space where the following limits are valid are indicated in the comment lines or in the footnotes.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>101	95	134 ABBIENDI	04H OPAL	all $\tan\beta, \Delta m_0 > 5$ GeV, $m_0 > 500$ GeV, $A_0 = 0$
> 89	95	135 ABBIENDI	03H OPAL	$0.5 \leq \Delta m_{\pm} \leq 5$ GeV, higgsino-like, $\tan\beta=1.5$
> 97.1	95	136 ABDALLAH	03M DLPH	$\tilde{\chi}_1^\pm, \Delta m_{\pm} \geq 3$ GeV, $m_{\tilde{\nu}} > m_{\tilde{\chi}_1^\pm}$
> 75	95	136 ABDALLAH	03M DLPH	$\tilde{\chi}_1^\pm$ , higgsino, all $\Delta m_{\pm}, m_{\tilde{f}} > m_{\tilde{\chi}_1^\pm}$

> 70	95	136	ABDALLAH	03M DLPH	$\tilde{\chi}_1^\pm$ , all $\Delta m_+$ , $m_{\tilde{\nu}} > 500$ GeV, $M_2 \leq 2M_1 \leq 10M_2$
<b>&gt; 94</b>	95	137	ABDALLAH	03M DLPH	$\tilde{\chi}_1^\pm$ , $\tan\beta \leq 40$ , $\Delta m_+ > 3$ GeV, all $m_0$
> 88	95	138	HEISTER	02J ALEP	$\tilde{\chi}_1^\pm$ , all $\Delta m_+$ , large $m_0$
> 67.7	95	139	ACCIARRI	00D L3	$\tan\beta > 0.7$ , all $\Delta m_+$ , all $m_0$
> 69.4	95	140	ACCIARRI	00K L3	$e^+e^- \rightarrow \tilde{\chi}^\pm \tilde{\chi}^\mp$ , all $\Delta m_+$ , heavy scalars
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>195	95	141	ABAZOV	05A D0	$\rho\bar{\rho} \rightarrow \tilde{\chi}\tilde{\chi}$ , $\tilde{\chi} = \tilde{\chi}_2^0$ , $\tilde{\chi}_1^\pm, \tilde{\chi}_1^0 \rightarrow$ $\gamma\tilde{G}$ , GMSB
>117	95	142	ABAZOV	05U D0	$\rho\bar{\rho} \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_2^0$
>167	95	143	ACOSTA	05E CDF	$\rho\bar{\rho} \rightarrow \tilde{\chi}\tilde{\chi}$ , $\tilde{\chi} = \tilde{\chi}_2^0$ , $\tilde{\chi}_1^\pm, \tilde{\chi}_1^0 \rightarrow$ $\gamma\tilde{G}$ , GMSB
> 66	95	144,145	ABDALLAH	04H DLPH	AMSB, $\mu > 0$
>102.5	95	146,147	ABDALLAH	04M DLPH	$\mathcal{R}(\overline{U}\overline{D}\overline{D})$
>100	95	148	ABDALLAH	03D DLPH	$e^+e^- \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_1^\mp$ ( $\tilde{\chi}_1^\pm \rightarrow$ $\tilde{\tau}_1\nu_\tau, \tilde{\tau}_1 \rightarrow \tau\tilde{G}$ )
>103	95	149	HEISTER	03G ALEP	$\mathcal{R}$ decays, $m_0 > 500$ GeV
>102.7	95	150	ACHARD	02 L3	$\mathcal{R}$ , MSUGRA
		151	GHODBANE	02 THEO	
> 94.3	95	152	ABREU	01C DLPH	$\tilde{\chi}^\pm \rightarrow \tau J$
> 93.8	95	153	ACCIARRI	01 L3	$\mathcal{R}$ , all $m_0$ , $0.7 \leq \tan\beta \leq 40$
>100	95	154	BARATE	01B ALEP	$\mathcal{R}$ decays, $m_0 > 500$ GeV
> 91.8	95	155	ABREU	00V DLPH	$e^+e^- \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_1^\pm$ ( $\tilde{\chi}_1^\pm \rightarrow \tilde{\tau}_1\nu_\tau$ , $\tilde{\tau}_1 \rightarrow \tau\tilde{G}$ )
		156	CHO	00B THEO	EW analysis
> 76	95	157	ABBIENDI	99T OPAL	$\mathcal{R}$ , $m_0=500$ GeV
> 51	95	158	MALTONI	99B THEO	EW analysis, $\Delta m_+ \sim 1$ GeV
> 81.5	95	159	ABE	98J CDF	$\rho\bar{\rho} \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_2^0$
		160	ACKERSTAFF	98K OPAL	$\tilde{\chi}^+ \rightarrow \ell^+ \cancel{E}$
> 65.7	95	161	ACKERSTAFF	98L OPAL	$\Delta m_+ > 3$ GeV, $\Delta m_\nu > 2$ GeV
		162	ACKERSTAFF	98V OPAL	light gluino
		163	CARENA	97 THEO	$g_\mu - 2$
		164	KALINOWSKI	97 THEO	$W \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_1^0$
		165	ABE	96K CDF	$\rho\bar{\rho} \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_2^0$

<sup>134</sup> ABBIENDI 04H search for charginos and neutralinos in events with acoplanar leptons+jets and multi-jet final states in the 192–209 GeV data, combined with the results on leptonic final states from ABBIENDI 04. The results hold for a scan over the parameter space covering the region  $0 < M_2 < 5000$  GeV,  $-1000 < \mu < 1000$  GeV and  $\tan\beta$  from 1 to 40. This limit supersedes ABBIENDI 00H.

<sup>135</sup> ABBIENDI 03H used  $e^+e^-$  data at  $\sqrt{s} = 188$ –209 GeV to search for chargino pair production in the case of small  $\Delta m_+$ . They select events with an energetic photon, large  $\cancel{E}$  and little hadronic or leptonic activity. The bound applies to higgsino-like charginos with zero lifetime and a 100% branching ratio  $\tilde{\chi}_1^\pm \rightarrow \tilde{\chi}_1^0 W^*$ . The mass limit for gaugino-like charginos, in case of non-universal gaugino masses, is of 92 GeV for  $m_{\tilde{\nu}} = 1000$  GeV and is lowered to 74 GeV for  $m_{\tilde{\nu}} \geq 100$  GeV. Limits in the plane  $(m_{\tilde{\nu}}, \Delta m_+)$

are shown in Fig. 7. Exclusion regions are also derived for the AMSB scenario in the  $(m_{3/2}, \tan\beta)$  plane, see their Fig. 9.

- 136 ABDALLAH 03M searches for the production of charginos using data from  $\sqrt{s}=192$  to 208 GeV to investigate topologies with multiple leptons, jets plus leptons, multi-jets, or isolated photons. The first limit holds for  $\tan\beta \geq 1$  and is obtained at  $\Delta m_+ = 3$  GeV in the higgsino region. For  $\Delta m_+ \geq 10$  (5) GeV and large  $m_0$ , the limit improves to 102.7 (101.7) GeV. For the region of small  $\Delta m_+$ , all data from  $\sqrt{s}=130$  to 208 GeV are used to investigate final states with heavy stable charged particles, decay vertices inside the detector and soft topologies with a photon from initial state radiation. The second limit is obtained in the higgsino region, assuming gaugino mass universality at the GUT scale and  $1 < \tan\beta < 50$ . For the case of non-universality of gaugino masses, the parameter space is scanned in the domain  $1 < \tan\beta < 50$  and, for  $\Delta m_+ < 3$  GeV, for values of  $M_1$ ,  $M_2$  and  $\mu$  such that  $M_2 \leq 2M_1 \leq 10M_2$  and  $|\mu| \geq M_2$ . The third limit is obtained in the gaugino region. See Fig. 36 for the dependence of the low  $\Delta m_+$  limits on  $\Delta m_+$ . These limits include and update the results of ABREU 00J and ABREU 00T.
- 137 ABDALLAH 03M uses data from  $\sqrt{s} = 192$ –208 GeV to obtain limits in the framework of the MSSM with gaugino and sfermion mass universality at the GUT scale. An indirect limit on the mass of charginos is derived by constraining the MSSM parameter space by the results from direct searches for neutralinos (including cascade decays), for charginos and for sleptons. These limits are valid for values of  $M_2 < 1$  TeV,  $|\mu| \leq 2$  TeV with the  $\tilde{\chi}_1^0$  as LSP. Constraints from the Higgs search in the  $M_h^{max}$  scenario assuming  $m_t=174.3$  GeV are included. The quoted limit applies if there is no mixing in the third family or when  $m_{\tilde{\tau}_1} - m_{\tilde{\chi}_1^0} > 6$  GeV. If mixing is included the limit degrades to 90 GeV. See Fig. 43 for the mass limits as a function of  $\tan\beta$ . These limits update the results of ABREU 00W.
- 138 HEISTER 02J search for chargino production with small  $\Delta m_+$  in final states with a hard isolated initial state radiation photon and few low-momentum particles, using 189–208 GeV data. This search is sensitive in the intermediate  $\Delta m_+$  region. Combined with searches for  $\cancel{E}$  topologies and for stable charged particles, the above bound is obtained for  $m_0$  larger than few hundred GeV,  $1 < \tan\beta < 300$  and holds for any chargino field contents. For light scalars, the general limit reduces to the one from the  $Z^0$ , but under the assumption of gaugino and sfermion mass unification the above bound is recovered. See Figs. 4–6 for the more general dependence of the limits on  $\Delta m_+$ . Updates BARATE 98X.
- 139 ACCIARRI 00D data collected at  $\sqrt{s}=189$  GeV. The results hold over the full parameter space defined by  $0.7 \leq \tan\beta \leq 60$ ,  $0 \leq M_2 \leq 2$  TeV,  $|\mu| \leq 2$  TeV  $m_0 \leq 500$  GeV. The results of slepton searches from ACCIARRI 99W are used to help set constraints in the region of small  $m_0$ . See their Figs. 5 for the  $\tan\beta$  and  $M_2$  dependence on the limits. See the text for the impact of a large  $B(\tilde{\chi}^\pm \rightarrow \tau \tilde{\nu}_\tau)$  on the result. The region of small  $\Delta m_+$  is excluded by the analysis of ACCIARRI 00K. Updates ACCIARRI 98F.
- 140 ACCIARRI 00K searches for the production of charginos with small  $\Delta m_+$  using data from  $\sqrt{s}=189$  GeV. They investigate soft final states with a photon from initial state radiation. The results are combined with the limits on prompt decays from ACCIARRI 00D and from heavy stable charged particles from ACCIARRI 99L (see Heavy Charged Lepton Searches). The production and decay branching ratios are evaluated within the MSSM, assuming heavy sfermions. The parameter space is scanned in the domain  $1 < \tan\beta < 50$ ,  $0.3 < M_1/M_2 < 50$ , and  $0 < |\mu| < 2$  TeV. The limit is obtained in the higgsino region and improves to 78.6 GeV for gaugino-like charginos. The limit is unchanged for light scalar quarks. For light  $\tilde{\tau}$  or  $\tilde{\nu}_\tau$ , the limit is unchanged in the gaugino-like region and is lowered by 0.8 GeV in the higgsino-like case. For light  $\tilde{\mu}$  or  $\tilde{\nu}_\mu$ , the limit is unchanged in the higgsino-like region and is lowered by 0.9 GeV in the gaugino-like region. No direct mass limits are obtained for light  $\tilde{e}$  or  $\tilde{\nu}_e$ .

- 141 ABAZOV 05A looked in  $263 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96 \text{ TeV}$  for diphoton events with large  $\cancel{E}_T$ . They may originate from the production of  $\tilde{\chi}^\pm$  in pairs or associated to a  $\tilde{\chi}_2^0$ , decaying to a  $\tilde{\chi}_1^0$  which itself decays promptly in GMSB to  $\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$ . No significant excess was found at large  $\cancel{E}_T$  compared to the background expectation. A limit is derived on the masses of SUSY particles in the GMSB framework for  $M = 2 \Lambda$ ,  $N = 1$ ,  $\tan\beta = 15$  and  $\mu > 0$ , see Figure 2. It also excludes  $\Lambda < 79.6 \text{ TeV}$ . Very similar results are obtained for different choices of parameters, see their Table 2. Supersedes the results of ABBOTT 98.
- 142 ABAZOV 05U looked in  $320 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96 \text{ TeV}$  for events with large  $\cancel{E}_T$ , no jets and three leptons ( $e, \mu, \tau$ ) of which at least two are  $e$  or  $\mu$ . No significant excess was found at large  $\cancel{E}_T$  compared to the background expectation. A limit is derived on the cross section times branching ratio to 3 leptons, see their Figures 2 and 3. The mass limit assumes gaugino mass universality, three degenerate sleptons and “maximally enhanced” leptonic branching fraction, i.e. a decay dominated by a slepton rather than  $W/Z$ . If, in addition, squarks are heavy, the limit improves to  $132 \text{ GeV}$ . Supersedes the results of ABBOTT 98C.
- 143 ACOSTA 05E looked in  $202 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.96 \text{ TeV}$  for diphoton events with large  $\cancel{E}_T$ . They may originate from the production of  $\tilde{\chi}^\pm$  in pairs or associated to a  $\tilde{\chi}_2^0$ , decaying to a  $\tilde{\chi}_1^0$  which itself decays promptly in GMSB to  $\gamma \tilde{G}$ . No events are selected at large  $\cancel{E}_T$  compared to the background expectation. A limit is derived on the masses of SUSY particles in the GMSB framework for  $M = 2 \Lambda$ ,  $N = 1$ ,  $\tan\beta = 15$  and  $\mu > 0$ , see Figure 2. It also excludes  $\Lambda < 69 \text{ TeV}$ . Supersedes the results of ABE 99I.
- 144 ABDALLAH 04H use data from LEP 1 and  $\sqrt{s} = 192\text{--}208 \text{ GeV}$ . They re-use results or re-analyze the data from ABDALLAH 03M to put limits on the parameter space of anomaly-mediated supersymmetry breaking (AMSB), which is scanned in the region  $1 < m_{3/2} < 50 \text{ TeV}$ ,  $0 < m_0 < 1000 \text{ GeV}$ ,  $1.5 < \tan\beta < 35$ , both signs of  $\mu$ . The constraints are obtained from the searches for mass degenerate chargino and neutralino, for SM-like and invisible Higgs, for leptonically decaying charginos and from the limit on non-SM  $Z$  width of  $3.2 \text{ MeV}$ . The limit is for  $m_t = 174.3 \text{ GeV}$  (see Table 2 for other  $m_t$  values).
- 145 The limit improves to  $73 \text{ GeV}$  for  $\mu < 0$ .
- 146 ABDALLAH 04M use data from  $\sqrt{s} = 192\text{--}208 \text{ GeV}$  to derive limits on sparticle masses under the assumption of  $\cancel{R}$  with  $LL\bar{E}$  or  $\bar{U}DD$  couplings. The results are valid in the ranges  $90 < m_0 < 500 \text{ GeV}$ ,  $0.7 < \tan\beta < 30$ ,  $-200 < \mu < 200 \text{ GeV}$ ,  $0 < M_2 < 400 \text{ GeV}$ . Supersedes the result of ABREU 01D and ABREU 00U.
- 147 The limit improves to  $103 \text{ GeV}$  for  $LL\bar{E}$  couplings.
- 148 ABDALLAH 03D use data from  $\sqrt{s}= 183\text{--}208 \text{ GeV}$ . They look for final states with two acoplanar leptons, expected in GMSB when the  $\tilde{\tau}_1$  is the NLSP and assuming a short-lived  $\tilde{\chi}_1^\pm$ . Limits are obtained in the plane  $(m(\tilde{\tau}), m(\tilde{\chi}_1^\pm))$  for different domains of  $m(\tilde{G})$ , after combining these results with the search for slepton pair production from the same paper. The limit above is valid if the  $\tilde{\tau}_1$  is the NLSP for all values of  $m(\tilde{G})$  provided  $m(\tilde{\chi}_1^\pm) - m(\tilde{\tau}_1) \geq 0.3 \text{ GeV}$ . For larger  $m(\tilde{G}) > 100 \text{ eV}$  the limit improves to  $102 \text{ GeV}$ , see their Fig. 11. In the co-NLSP scenario, the limits are  $96$  and  $102 \text{ GeV}$  for all  $m(\tilde{G})$  and  $m(\tilde{G}) > 100 \text{ eV}$ , respectively. Supersedes the results of ABREU 01G.
- 149 HEISTER 03G searches for the production of charginos prompt decays. in the case of  $\cancel{R}$  prompt decays with  $LL\bar{E}$ ,  $LQ\bar{D}$  or  $\bar{U}DD$  couplings at  $\sqrt{s}=189\text{--}209 \text{ GeV}$ . The search is performed for indirect decays, assuming one coupling at a time to be non-zero. The limit holds for  $\tan\beta = 1.41$ . Excluded regions in the  $(\mu, M_2)$  plane are shown in their Fig. 3.
- 150 ACHARD 02 searches for the production of sparticles in the case of  $\cancel{R}$  prompt decays with  $LL\bar{E}$  or  $\bar{U}DD$  couplings at  $\sqrt{s}=189\text{--}208 \text{ GeV}$ . The search is performed for direct and indirect decays, assuming one coupling at the time to be nonzero. The MSUGRA limit results from a scan over the MSSM parameter space with the assumption of gaugino and scalar mass unification at the GUT scale, imposing simultaneously the exclusions from neutralino, chargino, sleptons, and squarks analyses. The limit of  $\tilde{\chi}_1^\pm$  holds for

- $\overline{UDD}$  couplings and increases to 103.0 GeV for  $LL\overline{E}$  couplings. For L3 limits from  $LQ\overline{D}$  couplings, see ACCIARRI 01.
- 151 GHODBANE 02 reanalyzes DELPHI data at  $\sqrt{s}=189$  GeV in the presence of complex phases for the MSSM parameters.
- 152 ABREU 01C looked for  $\tau$  pairs with  $\cancel{E}$  at  $\sqrt{s}=183\text{--}189$  GeV to search for the associated production of charginos, followed by the decay  $\tilde{\chi}^{\pm} \rightarrow \tau J$ ,  $J$  being an invisible massless particle. See Fig. 6 for the regions excluded in the  $(\mu, M_2)$  plane.
- 153 ACCIARRI 01 searches for multi-lepton and/or multi-jet final states from  $\cancel{R}$  prompt decays with  $LL\overline{E}$ ,  $LQ\overline{D}$ , or  $\overline{UDD}$  couplings at  $\sqrt{s}=189$  GeV. The search is performed for direct and indirect decays of neutralinos, charginos, and scalar leptons, with the  $\tilde{\chi}_1^0$  or a  $\tilde{\ell}$  as LSP and assuming one coupling to be nonzero at a time. Mass limits are derived using simultaneously the constraints from the neutralino, chargino, and slepton analyses; and the  $Z^0$  width measurements from ACCIARRI 00C in a scan of the parameter space assuming MSUGRA with gaugino and scalar mass universality. Updates and supersedes the results from ACCIARRI 99I.
- 154 BARATE 01B searches for the production of charginos in the case of  $\cancel{R}$  prompt decays with  $LL\overline{E}$ ,  $LQ\overline{D}$ , or  $\overline{UDD}$  couplings at  $\sqrt{s}=189\text{--}202$  GeV. The search is performed for indirect decays, assuming one coupling at a time to be nonzero. Updates BARATE 00H.
- 155 ABREU 00V use data from  $\sqrt{s}=183\text{--}189$  GeV. They look for final states with two acoplanar leptons, expected in GMSB when the  $\tilde{\tau}_1$  is the NLSP and assuming a short-lived  $\tilde{\chi}_1^{\pm}$ . Limits are obtained in the plane  $(m_{\tilde{\tau}}, m_{\tilde{\chi}_1^{\pm}})$  for different domains of  $m_{\tilde{G}}$ , after combining these results with the search for slepton pair production in the SUGRA framework from ABREU 01 to cover prompt decays and on stable particle searches from ABREU 00Q. The limit above is valid for all values of  $m_{\tilde{G}}$ .
- 156 CHO 00B studied constraints on the MSSM spectrum from precision EW observables. Global fits favour charginos with masses at the lower bounds allowed by direct searches. Allowing for variations of the squark and slepton masses does not improve the fits.
- 157 ABBIENDI 99T searches for the production of neutralinos in the case of  $R$ -parity violation with  $LL\overline{E}$ ,  $LQ\overline{D}$ , or  $\overline{UDD}$  couplings using data from  $\sqrt{s}=183$  GeV. They investigate topologies with multiple leptons, jets plus leptons, or multiple jets, assuming one coupling at the time to be non-zero and giving rise to direct or indirect decays. Mixed decays (where one particle has a direct, the other an indirect decay) are also considered for the  $\overline{UDD}$  couplings. Upper limits on the cross section are derived which, combined with the constraint from the  $Z^0$  width, allow to exclude regions in the  $M_2$  versus  $\mu$  plane for any coupling. Limits on the chargino mass are obtained for non-zero  $LL\overline{E}$  couplings  $> 10^{-5}$  and assuming decays via a  $W^*$ .
- 158 MALTONI 99B studied the effect of light chargino-neutralino to the electroweak precision data with a particular focus on the case where they are nearly degenerate ( $\Delta m_{\pm} \sim 1$  GeV) which is difficult to exclude from direct collider searches. The quoted limit is for higgsino-like case while the bound improves to 56 GeV for wino-like case. The values of the limits presented here are obtained in an update to MALTONI 99B, as described in MALTONI 00.
- 159 ABE 98J searches for trilepton final states ( $\ell=e, \mu$ ). Efficiencies are calculated using mass relations in the Minimal Supergravity scenario, exploring the domain of parameter space defined by  $1.1 < \tan\beta < 8$ ,  $-1000 < \mu(\text{GeV}) < -200$ , and  $m_{\tilde{q}}/m_{\tilde{g}}=1\text{--}2$ . In this region  $m_{\tilde{\chi}_1^{\pm}} \sim m_{\tilde{\chi}_2^0}$  and  $m_{\tilde{\chi}_1^{\pm}} \sim 2m_{\tilde{\chi}_1^0}$ . Results are presented in Fig. 1 as upper bounds on  $\sigma(p\overline{p} \rightarrow \tilde{\chi}_1^{\pm} \tilde{\chi}_2^0) \times B(3\ell)$ . Limits range from 0.8 pb ( $m_{\tilde{\chi}_1^{\pm}}=50$  GeV) to 0.23 pb ( $m_{\tilde{\chi}_1^{\pm}}=100$  GeV) at 95%CL. The gaugino mass unification hypothesis and the assumed mass relation between squarks and gluinos define the value of the leptonic branching ratios. The quoted result corresponds to the best limit within the selected

- range of parameters, obtained for  $m_{\tilde{q}} > m_{\tilde{g}}$ ,  $\tan\beta=2$ , and  $\mu=-600$  GeV. Mass limits for different values of  $\tan\beta$  and  $\mu$  are given in Fig. 2.
- 160 ACKERSTAFF 98K looked for dilepton+ $\cancel{E}_T$  final states at  $\sqrt{s}=130-172$  GeV. Limits on  $\sigma(e^+e^- \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_1^-) \times B^2(\ell)$ , with  $B(\ell)=B(\chi^+ \rightarrow \ell^+ \nu_\ell \chi_1^0)$  ( $B(\ell)=B(\chi^+ \rightarrow \ell^+ \tilde{\nu}_\ell)$ ), are given in Fig. 16 (Fig. 17).
- 161 ACKERSTAFF 98L limit is obtained for  $0 < M_2 < 1500$ ,  $|\mu| < 500$  and  $\tan\beta > 1$ , but remains valid outside this domain. The dependence on the trilinear-coupling parameter  $A$  is studied, and found negligible. The limit holds for the smallest value of  $m_0$  consistent with scalar lepton constraints (ACKERSTAFF 97H) and for all values of  $m_0$  where the condition  $\Delta m_{\tilde{\nu}} > 2.0$  GeV is satisfied.  $\Delta m_{\tilde{\nu}} > 10$  GeV if  $\tilde{\chi}^\pm \rightarrow \ell \tilde{\nu}_\ell$ . The limit improves to 84.5 GeV for  $m_0=1$  TeV. Data taken at  $\sqrt{s}=130-172$  GeV.
- 162 ACKERSTAFF 98V excludes the light gluino with universal gaugino mass where charginos, neutralinos decay as  $\tilde{\chi}_1^\pm, \tilde{\chi}_2^0 \rightarrow q \bar{q} \tilde{g}$  from total hadronic cross sections at  $\sqrt{s}=130-172$  GeV. See paper for the case of nonuniversal gaugino mass.
- 163 CARENA 97 studied the constraints on chargino and sneutrino masses from muon  $g-2$ . The bound can be important for large  $\tan\beta$ .
- 164 KALINOWSKI 97 studies the constraints on the chargino-neutralino parameter space from limits on  $\Gamma(W \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_1^0)$  achievable at LEP2. This is relevant when  $\tilde{\chi}_1^\pm$  is "invisible," i.e., if  $\tilde{\chi}_1^\pm$  dominantly decays into  $\tilde{\nu}_\ell \ell^\pm$  with little energy for the lepton. Small otherwise allowed regions could be excluded.
- 165 ABE 96K looked for trilepton events from chargino-neutralino production. The bound on  $m_{\tilde{\chi}_1^\pm}$  can reach up to 47 GeV for specific choices of parameters. The limits on the combined production cross section times 3-lepton branching ratios range between 1.4 and 0.4 pb, for  $45 < m_{\tilde{\chi}_1^\pm}(\text{GeV}) < 100$ . See the paper for more details on the parameter dependence of the results.

## Long-lived $\tilde{\chi}^\pm$ (Chargino) MASS LIMITS

Limits on charginos which leave the detector before decaying.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>102	95	166 ABBIENDI	03L OPAL	$m_{\tilde{\nu}} > 500$ GeV
none 2-93.0	95	167 ABREU	00T DLPH	$\tilde{H}^\pm$ or $m_{\tilde{\nu}} > m_{\tilde{\chi}^\pm}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

- |        |    |            |          |
|--------|----|------------|----------|
| > 83   | 95 | 168 BARATE | 97K ALEP |
| > 28.2 | 95 | ADACHI     | 90C TOPZ |
- 166 ABBIENDI 03L used  $e^+e^-$  data at  $\sqrt{s} = 130-209$  GeV to select events with two high momentum tracks with anomalous  $dE/dx$ . The excluded cross section is compared to the theoretical expectation as a function of the heavy particle mass in their Fig. 3. The bounds are valid for colorless fermions with lifetime longer than  $10^{-6}$  s. Supersedes the results from ACKERSTAFF 98P.
- 167 ABREU 00T searches for the production of heavy stable charged particles, identified by their ionization or Cherenkov radiation, using data from  $\sqrt{s}=130$  to 189 GeV. These limits include and update the results of ABREU 98P.
- 168 BARATE 97K uses  $e^+e^-$  data collected at  $\sqrt{s} = 130-172$  GeV. Limit valid for  $\tan\beta = \sqrt{2}$  and  $m_{\tilde{\nu}} > 100$  GeV. The limit improves to 86 GeV for  $m_{\tilde{\nu}} > 250$  GeV.

## $\tilde{\nu}$ (Sneutrino) MASS LIMIT

The limits may depend on the number,  $N(\tilde{\nu})$ , of sneutrinos assumed to be degenerate in mass. Only  $\tilde{\nu}_L$  (not  $\tilde{\nu}_R$ ) is assumed to exist. It is possible that  $\tilde{\nu}$  could be the lightest supersymmetric particle (LSP).

We report here, but do not include in the Listings, the limits obtained from the final, but unpublished, fit of the final results obtained by the LEP Collaborations on the invisible width of the Z boson ( $\Delta\Gamma_{\text{inv.}} < 2.0$  MeV, LEP 03):  $m_{\tilde{\nu}} > 43.7$  GeV ( $N(\tilde{\nu})=1$ ) and  $m_{\tilde{\nu}} > 44.7$  GeV ( $N(\tilde{\nu})=3$ ).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt; 94</b>	95	169 ABDALLAH	03M DLPH	$1 \leq \tan\beta \leq 40,$ $m_{\tilde{e}_R} - m_{\tilde{\chi}_1^0} > 10$ GeV
> 84	95	170 HEISTER	02N ALEP	$\tilde{\nu}_e$ , any $\Delta m$
> 37.1	95	171 ADRIANI	93M L3	$\Gamma(Z \rightarrow \text{invisible}); N(\tilde{\nu})=1$
> 41	95	172 DECAMP	92 ALEP	$\Gamma(Z \rightarrow \text{invisible}); N(\tilde{\nu})=3$
> 36	95	ABREU	91F DLPH	$\Gamma(Z \rightarrow \text{invisible}); N(\tilde{\nu})=1$
> 31.2	95	173 ALEXANDER	91F OPAL	$\Gamma(Z \rightarrow \text{invisible}); N(\tilde{\nu})=1$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		174 ABULENCIA	05A CDF	$p\bar{p} \rightarrow \tilde{\nu} \rightarrow ee, \mu\mu, \cancel{R} LQ\bar{D}$
		175 ACOSTA	05R CDF	$p\bar{p} \rightarrow \tilde{\nu} \rightarrow \tau\tau, \cancel{R}, LQ\bar{D}$
		176 ABBIENDI	04F OPAL	$\cancel{R}, \tilde{\nu}_{e,\mu,\tau}$
> 95	95	177,178 ABDALLAH	04H DLPH	AMSB, $\mu > 0$
> 98	95	179 ABDALLAH	04M DLPH	$\cancel{R}(LL\bar{E}), \tilde{\nu}_e$ , indirect, $\Delta m_0 > 5$ GeV
> 85	95	179 ABDALLAH	04M DLPH	$\cancel{R}(LL\bar{E}), \tilde{\nu}_\mu$ , indirect, $\Delta m_0 > 5$ GeV
> 85	95	179 ABDALLAH	04M DLPH	$\cancel{R}(LL\bar{E}), \tilde{\nu}_\tau$ , indirect, $\Delta m_0 > 5$ GeV
		180 ABDALLAH	03F DLPH	$\tilde{\nu}_{\mu,\tau}, \cancel{R} LL\bar{E}$ decays
		181 ACOSTA	03E CDF	$\tilde{\nu}, \cancel{R}, LQ\bar{D}$ production and $LL\bar{E}$ decays
> 88	95	182 HEISTER	03G ALEP	$\tilde{\nu}_e, \cancel{R}$ decays, $\mu = -200$ GeV, $\tan\beta=2$
> 65	95	182 HEISTER	03G ALEP	$\tilde{\nu}_{\mu,\tau}, \cancel{R}$ decays
		183 ABAZOV	02H D0	$\cancel{R}, \lambda_{211}$
> 95	95	184 ACHARD	02 L3	$\tilde{\nu}_e, \cancel{R}$ decays, $\mu = -200$ GeV, $\tan\beta = \sqrt{2}$
> 65	95	184 ACHARD	02 L3	$\tilde{\nu}_{\nu,\tau}, \cancel{R}$ decays
>149	95	184 ACHARD	02 L3	$\tilde{\nu}, \cancel{R}$ decays, MSUGRA
		185 HEISTER	02F ALEP	$e\gamma \rightarrow \tilde{\nu}_{\mu,\tau} \ell_k, \cancel{R} LL\bar{E}$
none 100–264	95	186 ABBIENDI	00R OPAL	$\tilde{\nu}_{\mu,\tau}, \cancel{R}, (s+t)$ -channel
none 100–200	95	187 ABBIENDI	00R OPAL	$\tilde{\nu}_\tau, \cancel{R}, s$ -channel
		188 ABREU	00S DLPH	$\tilde{\nu}_\ell, \cancel{R}, (s+t)$ -channel
none 50–210	95	189 ACCIARRI	00P L3	$\tilde{\nu}_{\mu,\tau}, \cancel{R}, s$ -channel
none 50–210	95	190 BARATE	00I ALEP	$\tilde{\nu}_{\mu,\tau}, \cancel{R}, (s+t)$ -channel
none 90–210	95	191 BARATE	00I ALEP	$\tilde{\nu}_\tau, \cancel{R}, s$ -channel
none 100–160	95	192 ABBIENDI	99 OPAL	$\tilde{\nu}_e, \cancel{R}, t$ -channel
$\neq m_Z$	95	193 ACCIARRI	97U L3	$\tilde{\nu}_\tau, \cancel{R}, s$ -channel
none 125–180	95	193 ACCIARRI	97U L3	$\tilde{\nu}_\tau, \cancel{R}, s$ -channel
		194 CARENA	97 THEO	$g_\mu - 2$
> 46.0	95	195 BUSKULIC	95E ALEP	$N(\tilde{\nu})=1, \tilde{\nu} \rightarrow \nu\nu\ell\bar{\ell}'$

none 20–25000	196	BECK	94	COSM	Stable $\tilde{\nu}$ , dark matter
<600	197	FALK	94	COSM	$\tilde{\nu}$ LSP, cosmic abundance
none 3–90	90	198 SATO	91	KAMI	Stable $\tilde{\nu}_e$ or $\tilde{\nu}_\mu$ , dark matter
none 4–90	90	198 SATO	91	KAMI	Stable $\tilde{\nu}_\tau$ , dark matter

- 169 ABDALLAH 03M uses data from  $\sqrt{s} = 192\text{--}208$  GeV to obtain limits in the framework of the MSSM with gaugino and sfermion mass universality at the GUT scale. An indirect limit on the mass is derived by constraining the MSSM parameter space by the results from direct searches for neutralinos (including cascade decays) and for sleptons. These limits are valid for values of  $M_2 < 1$  TeV,  $|\mu| \leq 1$  TeV with the  $\tilde{\chi}_1^0$  as LSP. The quoted limit is obtained when there is no mixing in the third family. See Fig. 43 for the mass limits as a function of  $\tan\beta$ . These limits update the results of ABREU 00W.
- 170 HEISTER 02N derives a bound on  $m_{\tilde{\nu}_e}$  by exploiting the mass relation between the  $\tilde{\nu}_e$  and  $\tilde{e}$ , based on the assumption of universal GUT scale gaugino and scalar masses  $m_{1/2}$  and  $m_0$  and the search described in the  $\tilde{e}$  section. In the MSUGRA framework with radiative electroweak symmetry breaking, the limit improves to  $m_{\tilde{\nu}_e} > 130$  GeV, assuming a trilinear coupling  $A_0=0$  at the GUT scale. See Figs. 5 and 7 for the dependence of the limits on  $\tan\beta$ .
- 171 ADRIANI 93M limit from  $\Delta\Gamma(Z)(\text{invisible}) < 16.2$  MeV.
- 172 DECAMP 92 limit is from  $\Gamma(\text{invisible})/\Gamma(\ell\ell) = 5.91 \pm 0.15$  ( $N_\nu = 2.97 \pm 0.07$ ).
- 173 ALEXANDER 91F limit is for one species of  $\tilde{\nu}$  and is derived from  $\Gamma(\text{invisible, new})/\Gamma(\ell\ell) < 0.38$ .
- 174 ABULENCIA 05A looked in  $\sim 200$  pb $^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV for dimuon and dielectron events. They may originate from the  $\tilde{\chi}$  production of a sneutrino decaying to dileptons. No significant excess rate was found compared to the background expectation. A limit is derived on the cross section times branching ratio,  $B$ , of  $\tilde{\nu} \rightarrow ee, \mu\mu$  of 25 fb at high mass, see their Figure 2. Sneutrino masses are excluded at 95% CL below 680, 620, 460 GeV ( $ee$  channel) and 665, 590, 450 GeV ( $\mu\mu$  channel) for a  $\lambda'$  coupling and branching ratio such that  $\lambda'^2 B = 0.01, 0.005, 0.001$ , respectively.
- 175 ACOSTA 05R looked in 195 pb $^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV for ditau events with one identified hadronic tau decay and one other tau decay. They may originate from the  $\tilde{\chi}$  production of a sneutrino decaying to  $\tau\tau$ . No significant excess rate was found compared to the background expectation, dominated by Drell-Yan. A limit is derived on the cross section times branching ratio,  $B$ , of  $\tilde{\nu} \rightarrow \tau\tau$ , see their Figure 3. Sneutrino masses below 377 GeV are excluded at 95% CL for a  $\lambda'$  coupling to  $d\bar{d}$  and branching ratio such that  $\lambda'^2 B = 0.01$ .
- 176 ABBIENDI 04F use data from  $\sqrt{s} = 189\text{--}209$  GeV. They derive limits on sparticle masses under the assumption of  $\tilde{\chi}$  with  $LL\bar{E}$  or  $LQ\bar{D}$  couplings. The results are valid for  $\tan\beta = 1.5$ ,  $\mu = -200$  GeV, and a BR for the decay given by CMSSM, assuming no sensitivity to other decays. Limits are quoted for  $m_{\tilde{\chi}_0} = 60$  GeV and degrade for low-mass  $\tilde{\chi}_1^0$ . For  $\tilde{\nu}_e$  the direct (indirect) limits with  $LL\bar{E}$  couplings are 89 (95) GeV and with  $LQ\bar{D}$  they are 89 (88) GeV. For  $\tilde{\nu}_{\mu,\tau}$  the direct (indirect) limits with  $LL\bar{E}$  couplings are 79 (81) GeV and with  $LQ\bar{D}$  they are 74 (no limit) GeV. Supersedes the results of ABBIENDI 00.
- 177 ABDALLAH 04H use data from LEP 1 and  $\sqrt{s} = 192\text{--}208$  GeV. They re-use results or re-analyze the data from ABDALLAH 03M to put limits on the parameter space of anomaly-mediated supersymmetry breaking (AMSB), which is scanned in the region  $1 < m_{3/2} < 50$  TeV,  $0 < m_0 < 1000$  GeV,  $1.5 < \tan\beta < 35$ , both signs of  $\mu$ . The constraints are obtained from the searches for mass degenerate chargino and neutralino, for SM-like and invisible Higgs, for leptonically decaying charginos and from the limit on non-SM  $Z$  width of 3.2 MeV. The limit is for  $m_t = 174.3$  GeV (see Table 2 for other  $m_t$  values).
- 178 The limit improves to 114 GeV for  $\mu < 0$ .



- 179 ABDALLAH 04M use data from  $\sqrt{s} = 189\text{--}208$  GeV. The results are valid for  $\mu = -200$  GeV,  $\tan\beta = 1.5$ ,  $\Delta m_0 > 5$  GeV and assuming a BR of 1 for the given decay. The limit quoted is for indirect decays using the neutralino constraint of 39.5 GeV, also derived in ABDALLAH 04M. For indirect decays the limit on  $\tilde{\nu}_e$  decreases to 96 GeV if the constraint from the neutralino is not used and for direct decays it remains 96 GeV. For indirect decays the limit on  $\tilde{\nu}_\mu$  decreases to 82 GeV if the constraint from the neutralino is not used and to 83 GeV for direct decays. For indirect decays the limit on  $\tilde{\nu}_\tau$  decreases to 82 GeV if the constraint from the neutralino is not used and improves to 91 GeV for direct decays. Supersedes the results of ABREU 00U.
- 180 ABDALLAH 03F looked for events of the type  $e^+e^- \rightarrow \tilde{\nu} \rightarrow \tilde{\chi}^0\nu, \tilde{\chi}^\pm\ell^\mp$  followed by  $\cancel{E}$  decays of the  $\tilde{\chi}^0$  via  $\lambda_{1j1}$  ( $j = 2,3$ ) couplings in the data at  $\sqrt{s} = 183\text{--}208$  GeV. From a scan over the SUGRA parameters, they derive upper limits on the  $\lambda_{1j1}$  couplings as a function of the sneutrino mass, see their Figs. 5-8.
- 181 ACOSTA 03E search for  $e\mu, e\tau$  and  $\mu\tau$  final states, and sets limits on the product of production cross-section and decay branching ratio for a  $\tilde{\nu}$  in RPV models (see Fig. 3).
- 182 HEISTER 03G searches for the production of sneutrinos in the case of  $\cancel{E}$  prompt decays with  $LL\bar{E}$ ,  $LQ\bar{D}$  or  $U\bar{D}\bar{D}$  couplings at  $\sqrt{s} = 189\text{--}209$  GeV. The search is performed for direct and indirect decays, assuming one coupling at a time to be non-zero. The limit holds for indirect  $\bar{\nu}$  decays via  $U\bar{D}\bar{D}$  couplings and  $\Delta m > 10$  GeV. Stronger limits are reached for  $(\bar{\nu}_e, \bar{\nu}_\mu, \bar{\nu}_\tau)$  for  $LL\bar{E}$  direct (100,90) GeV or indirect (98,89) GeV and for  $LQ\bar{D}$  direct (-,79) GeV or indirect (91,78) GeV couplings. For  $LL\bar{E}$  indirect decays, use is made of the bound  $m(\tilde{\chi}_1^0) > 23$  GeV from BARATE 98S. Supersedes the results from BARATE 01B.
- 183 ABAZOV 02H looked in  $94 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV for events with at least 2 muons and 2 jets for  $s$ -channel production of  $\tilde{\mu}$  or  $\tilde{\nu}$  and subsequent decay via  $\cancel{E}$  couplings  $LQ\bar{D}$ . A scan over the MSUGRA parameters is performed to exclude regions of the  $(m_0, m_{1/2})$  plane, examples being shown in Fig. 2.
- 184 ACHARD 02 searches for the associated production of sneutrinos in the case of  $\cancel{E}$  prompt decays with  $LL\bar{E}$  or  $U\bar{D}\bar{D}$  couplings at  $\sqrt{s} = 189\text{--}208$  GeV. The search is performed for direct and indirect decays, assuming one coupling at the time to be nonzero. The limit holds for direct decays via  $LL\bar{E}$  couplings. Stronger limits are reached for  $(\tilde{\nu}_e, \tilde{\nu}_\mu, \tilde{\nu}_\tau)$  for  $LL\bar{E}$  indirect (99,78) GeV and for  $U\bar{D}\bar{D}$  direct or indirect (99,70) GeV decays. The MSUGRA limit results from a scan over the MSSM parameter space with the assumption of gaugino and scalar mass unification at the GUT scale, imposing simultaneously the exclusions from neutralino, chargino, sleptons, and squarks analyses. The limit holds for  $U\bar{D}\bar{D}$  couplings and increases to 152.7 GeV for  $LL\bar{E}$  couplings.
- 185 HEISTER 02F searched for single sneutrino production via  $e\gamma \rightarrow \tilde{\nu}; \ell_k$  mediated by  $\cancel{E} LL\bar{E}$  couplings, decaying directly or indirectly via a  $\tilde{\chi}_1^0$  and assuming a single coupling to be nonzero at a time. Final states with three leptons and possible  $\cancel{E}_T$  due to neutrinos were selected in the 189–209 GeV data. Limits on the couplings  $\lambda_{1jk}$  as function of the sneutrino mass are shown in Figs. 10–14. The couplings  $\lambda_{232}$  and  $\lambda_{233}$  are not accessible and  $\lambda_{121}$  and  $\lambda_{131}$  are measured with better accuracy in sneutrino resonant production. For all tested couplings, except  $\lambda_{133}$ , the limits are significantly improved compared to the low-energy limits.
- 186 ABBIENDI 00R studied the effect of  $s$ - and  $t$ -channel  $\tau$  or  $\mu$  sneutrino exchange in  $e^+e^- \rightarrow e^+e^-$  at  $\sqrt{s} = 130\text{--}189$  GeV, via the  $R$ -parity violating coupling  $\lambda_{1i1} L_1 L_i e_1$  ( $i=2$  or  $3$ ). The limits quoted here hold for  $\lambda_{1j1} > 0.13$ , and supersede the results of ABBIENDI 99. See Fig. 11 for limits on  $m_{\tilde{\nu}}$  versus coupling.
- 187 ABBIENDI 00R studied the effect of  $s$ -channel  $\tau$  sneutrino exchange in  $e^+e^- \rightarrow \mu^+\mu^-$  at  $\sqrt{s} = 130\text{--}189$  GeV, in presence of the  $R$ -parity violating couplings  $\lambda_{i3i} L_i L_3 e_i$  ( $i=1$  and  $2$ ), with  $\lambda_{131} = \lambda_{232}$ . The limits quoted here hold for  $\lambda_{131} > 0.09$ , and supersede the results of ABBIENDI 99. See Fig. 12 for limits on  $m_{\tilde{\nu}}$  versus coupling.
- 188 ABREU 00S searches for anomalies in the production cross sections and forward-backward asymmetries of the  $\ell^+\ell^-(\gamma)$  final states ( $\ell=e,\mu,\tau$ ) from  $e^+e^-$  collisions

- at  $\sqrt{s}=130\text{--}189$  GeV. Limits are set on the  $s$ - and  $t$ -channel exchange of sneutrinos in the presence of  $\tilde{R}$  with  $\lambda L\tilde{L}\tilde{E}$  couplings. For points between the energies at which data were taken, information is obtained from events in which a photon was radiated. Exclusion limits in the  $(\lambda, m_{\tilde{\nu}}$ ) plane are given in Fig. 5. These limits include and update the results of ABREU 99A.
- 189 ACCIARRI 00P use the dilepton total cross sections and asymmetries at  $\sqrt{s}=m_Z$  and  $\sqrt{s}=130\text{--}189$  GeV data to set limits on the effect of  $\tilde{R}$   $L\tilde{L}\tilde{E}$  couplings giving rise to  $\mu$  or  $\tau$  sneutrino exchange. See their Fig. 5 for limits on the sneutrino mass versus couplings.
- 190 BARATE 00I studied the effect of  $s$ -channel and  $t$ -channel  $\tau$  or  $\mu$  sneutrino exchange in  $e^+e^- \rightarrow e^+e^-$  at  $\sqrt{s}=130\text{--}183$  GeV, via the  $R$ -parity violating coupling  $\lambda_{1j1}L_1L_je_1^c$  ( $i=2$  or  $3$ ). The limits quoted here hold for  $\lambda_{1j1} > 0.1$ . See their Fig. 15 for limits as a function of the coupling.
- 191 BARATE 00I studied the effect of  $s$ -channel  $\tau$  sneutrino exchange in  $e^+e^- \rightarrow \mu^+\mu^-$  at  $\sqrt{s}=130\text{--}183$  GeV, in presence of the  $R$ -parity violating coupling  $\lambda_{i3j}L_iL_3e_j^c$  ( $i=1$  and  $2$ ). The limits quoted here hold for  $\sqrt{|\lambda_{131}\lambda_{232}|} > 0.2$ . See their Fig. 16 for limits as a function of the coupling.
- 192 ABBIENDI 99 studied the effect of  $t$ -channel electron sneutrino exchange in  $e^+e^- \rightarrow \tau^+\tau^-$  at  $\sqrt{s}=130\text{--}183$  GeV, in presence of the  $R$ -parity violating couplings  $\lambda_{131}L_1L_3e_1^c$ . The limits quoted here hold for  $\lambda_{131} > 0.6$ .
- 193 ACCIARRI 97U studied the effect of the  $s$ -channel tau-sneutrino exchange in  $e^+e^- \rightarrow e^+e^-$  at  $\sqrt{s}=m_Z$  and  $\sqrt{s}=130\text{--}172$  GeV, via the  $R$ -parity violating coupling  $\lambda_{131}L_1L_3e_1^c$ . The limits quoted here hold for  $\lambda_{131} > 0.05$ . Similar limits were studied in  $e^+e^- \rightarrow \mu^+\mu^-$  together with  $\lambda_{232}L_2L_3e_2^c$  coupling.
- 194 CARENA 97 studied the constraints on chargino and sneutrino masses from muon  $g-2$ . The bound can be important for large  $\tan\beta$ .
- 195 BUSKULIC 95E looked for  $Z \rightarrow \tilde{\nu}\tilde{\nu}$ , where  $\tilde{\nu} \rightarrow \nu\chi_1^0$  and  $\chi_1^0$  decays via  $R$ -parity violating interactions into two leptons and a neutrino.
- 196 BECK 94 limit can be inferred from limit on Dirac neutrino using  $\sigma(\tilde{\nu}) = 4\sigma(\nu)$ . Also private communication with H.V. Klapdor-Kleingrothaus.
- 197 FALK 94 puts an upper bound on  $m_{\tilde{\nu}}$  when  $\tilde{\nu}$  is LSP by requiring its relic density does not overclose the Universe.
- 198 SATO 91 search for high-energy neutrinos from the sun produced by annihilation of sneutrinos in the sun. Sneutrinos are assumed to be stable and to constitute dark matter in our galaxy. SATO 91 follow the analysis of NG 87, OLIVE 88, and GAISSER 86.

## CHARGED SLEPTONS

This section contains limits on charged scalar leptons ( $\tilde{\ell}$ , with  $\ell=e,\mu,\tau$ ). Studies of width and decays of the  $Z$  boson (use is made here of  $\Delta\Gamma_{\text{inv}} < 2.0$  MeV, LEP 00) conclusively rule out  $m_{\tilde{\ell}_R} < 40$  GeV (41

GeV for  $\tilde{\ell}_L$ ), independently of decay modes, for each individual slepton. The limits improve to 43 GeV (43.5 GeV for  $\tilde{\ell}_L$ ) assuming all 3 flavors to be degenerate. Limits on higher mass sleptons depend on model assumptions and on the mass splitting  $\Delta m = m_{\tilde{\ell}} - m_{\tilde{\chi}_1^0}$ . The mass and composition

of  $\tilde{\chi}_1^0$  may affect the selectron production rate in  $e^+e^-$  collisions through  $t$ -channel exchange diagrams. Production rates are also affected by the potentially large mixing angle of the lightest mass eigenstate  $\tilde{\ell}_1 = \tilde{\ell}_R \sin\theta_\ell + \tilde{\ell}_L \cos\theta_\ell$ . It is generally assumed that only  $\tilde{\tau}$  may have significant mixing. The coupling to the  $Z$  vanishes for  $\theta_\ell=0.82$ . In the high-energy limit of  $e^+e^-$  collisions the interference between  $\gamma$  and  $Z$  exchange leads to a minimal cross section for  $\theta_\ell=0.91$ , a value which is sometimes used in the

following entries relative to data taken at LEP2. When limits on  $m_{\tilde{\ell}_R}$  are quoted, it is understood that limits on  $m_{\tilde{\ell}_L}$  are usually at least as strong.

Possibly open decays involving gauginos other than  $\tilde{\chi}_1^0$  will affect the detection efficiencies. Unless otherwise stated, the limits presented here result from the study of  $\tilde{\ell}^+\tilde{\ell}^-$  production, with production rates and decay properties derived from the MSSM. Limits made obsolete by the recent analyses of  $e^+e^-$  collisions at high energies can be found in previous Editions of this Review.

For decays with final state gravitinos ( $\tilde{G}$ ),  $m_{\tilde{G}}$  is assumed to be negligible relative to all other masses.

### $\tilde{e}$ (Selectron) MASS LIMIT

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
> 97.5	95	199 ABBIENDI	04 OPAL	$\tilde{e}_R, \Delta m > 11$ GeV, $ \mu  > 100$ GeV, $\tan\beta=1.5$
> 94.4	95	200 ACHARD	04 L3	$\tilde{e}_R, \Delta m > 10$ GeV, $ \mu  > 200$ GeV, $\tan\beta \geq 2$
> 71.3	95	200 ACHARD	04 L3	$\tilde{e}_R$ , all $\Delta m$
none 30–94	95	201 ABDALLAH	03M DLPH	$\Delta m > 15$ GeV, $\tilde{e}_R^+\tilde{e}_R^-$
> 94	95	202 ABDALLAH	03M DLPH	$\tilde{e}_R, 1 \leq \tan\beta \leq 40$ , $\Delta m > 10$ GeV
> 95	95	203 HEISTER	02E ALEP	$\Delta m > 15$ GeV, $\tilde{e}_R^+\tilde{e}_R^-$
<b>&gt; 73</b>	95	204 HEISTER	02N ALEP	$\tilde{e}_R$ , any $\Delta m$
<b>&gt;107</b>	95	204 HEISTER	02N ALEP	$\tilde{e}_L$ , any $\Delta m$

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 89	95	205 ABBIENDI	04F OPAL	$\tilde{\mu}, \tilde{e}_L$
> 92	95	206 ABDALLAH	04M DLPH	$\tilde{\mu}, \tilde{e}_R$ , indirect, $\Delta m > 5$ GeV
> 93	95	207 HEISTER	03G ALEP	$\tilde{e}_R, \tilde{\mu}$ decays, $\mu=-200$ GeV, $\tan\beta=2$
> 69	95	208 ACHARD	02 L3	$\tilde{e}_R, \tilde{\mu}$ decays, $\mu=-200$ GeV, $\tan\beta=\sqrt{2}$
> 92	95	209 BARATE	01 ALEP	$\Delta m > 10$ GeV, $\tilde{e}_R^+\tilde{e}_R^-$
> 77	95	210 ABBIENDI	00J OPAL	$\Delta m > 5$ GeV, $\tilde{e}_R^+\tilde{e}_R^-$
> 83	95	211 ABREU	00U DLPH	Superseded by ABDALLAH 04M
> 67	95	212 ABREU	00V DLPH	$\tilde{e}_R\tilde{e}_R$ ( $\tilde{e}_R \rightarrow e\tilde{G}$ ), $m_{\tilde{G}} > 10$ eV
> 85	95	213 BARATE	00G ALEP	$\tilde{\ell}_R \rightarrow \ell\tilde{G}$ , any $\tau(\tilde{\ell}_R)$
> 29.5	95	214 ACCIARRI	99I L3	$\tilde{e}_R, \tilde{\mu}$ , $\tan\beta \geq 2$
> 56	95	215 ACCIARRI	98F L3	$\Delta m > 5$ GeV, $\tilde{e}_R^+\tilde{e}_R^-$ , $\tan\beta \geq 1.41$
> 77	95	216 BARATE	98K ALEP	Any $\Delta m$ , $\tilde{e}_R^+\tilde{e}_R^-$ , $\tilde{e}_R \rightarrow e\gamma\tilde{G}$
> 77	95	217 BREITWEG	98 ZEUS	$m_{\tilde{q}}=m_{\tilde{e}}$ , $m(\tilde{\chi}_1^0)=40$ GeV
> 63	95	218 AID	96C H1	$m_{\tilde{q}}=m_{\tilde{e}}$ , $m_{\tilde{\chi}_1^0}=35$ GeV

<sup>199</sup> ABBIENDI 04 search for  $\tilde{e}_R\tilde{e}_R$  production in acoplanar di-electron final states in the 183–208 GeV data. See Fig. 13 for the dependence of the limits on  $m_{\tilde{\chi}_1^0}$  and for the limit at  $\tan\beta=35$ . This limit supersedes ABBIENDI 00G.

<sup>200</sup> ACHARD 04 search for  $\tilde{e}_R\tilde{e}_L$  and  $\tilde{e}_R\tilde{e}_R$  production in single- and acoplanar di-electron final states in the 192–209 GeV data. Absolute limits on  $m_{\tilde{e}_R}$  are derived from a scan over the MSSM parameter space with universal GUT scale gaugino and scalar masses

- $m_{1/2}$  and  $m_0$ ,  $1 \leq \tan\beta \leq 60$  and  $-2 \leq \mu \leq 2$  TeV. See Fig. 4 for the dependence of the limits on  $m_{\tilde{\chi}_1^0}$ . This limit supersedes ACCIARRI 99W.
- 201 ABDALLAH 03M looked for acoplanar dielectron +  $\cancel{B}$  final states at  $\sqrt{s} = 189\text{--}208$  GeV. The limit assumes  $\mu = -200$  GeV and  $\tan\beta = 1.5$  in the calculation of the production cross section and  $B(\tilde{e} \rightarrow e\tilde{\chi}_1^0)$ . See Fig. 15 for limits in the  $(m_{\tilde{e}_R}, m_{\tilde{\chi}_1^0})$  plane. These limits include and update the results of ABREU 01
- 202 ABDALLAH 03M uses data from  $\sqrt{s} = 192\text{--}208$  GeV to obtain limits in the framework of the MSSM with gaugino and sfermion mass universality at the GUT scale. An indirect limit on the mass is derived by constraining the MSSM parameter space by the results from direct searches for neutralinos (including cascade decays) and for sleptons. These limits are valid for values of  $M_2 < 1$  TeV,  $|\mu| \leq 1$  TeV with the  $\tilde{\chi}_1^0$  as LSP. The quoted limit is obtained when there is no mixing in the third family. See Fig. 43 for the mass limits as a function of  $\tan\beta$ . These limits update the results of ABREU 00W.
- 203 HEISTER 02E looked for acoplanar dielectron +  $\cancel{B}_T$  final states from  $e^+e^-$  interactions between 183 and 209 GeV. The mass limit assumes  $\mu < -200$  GeV and  $\tan\beta = 2$  for the production cross section and  $B(\tilde{e} \rightarrow e\tilde{\chi}_1^0) = 1$ . See their Fig. 4 for the dependence of the limit on  $\Delta m$ . These limits include and update the results of BARATE 01.
- 204 HEISTER 02N search for  $\tilde{e}_R\tilde{e}_L$  and  $\tilde{e}_R\tilde{e}_R$  production in single- and acoplanar di-electron final states in the 183–208 GeV data. Absolute limits on  $m_{\tilde{e}_R}$  are derived from a scan over the MSSM parameter space with universal GUT scale gaugino and scalar masses  $m_{1/2}$  and  $m_0$ ,  $1 \leq \tan\beta \leq 50$  and  $-10 \leq \mu \leq 10$  TeV. The region of small  $|\mu|$ , where cascade decays are important, is covered by a search for  $\tilde{\chi}_1^0\tilde{\chi}_3^0$  in final states with leptons and possibly photons. Limits on  $m_{\tilde{e}_L}$  are derived by exploiting the mass relation between the  $\tilde{e}_L$  and  $\tilde{e}_R$ , based on universal  $m_0$  and  $m_{1/2}$ . When the constraint from the mass limit of the lightest Higgs from HEISTER 02 is included, the bounds improve to  $m_{\tilde{e}_R} > 77(75)$  GeV and  $m_{\tilde{e}_L} > 115(115)$  GeV for a top mass of 175(180) GeV. In the MSUGRA framework with radiative electroweak symmetry breaking, the limits improve further to  $m_{\tilde{e}_R} > 95$  GeV and  $m_{\tilde{e}_L} > 152$  GeV, assuming a trilinear coupling  $A_0 = 0$  at the GUT scale. See Figs. 4, 5, 7 for the dependence of the limits on  $\tan\beta$ .
- 205 ABBIENDI 04F use data from  $\sqrt{s} = 189\text{--}209$  GeV. They derive limits on sparticle masses under the assumption of  $\cancel{B}$  with  $LL\bar{E}$  or  $LQ\bar{D}$  couplings. The results are valid for  $\tan\beta = 1.5$ ,  $\mu = -200$  GeV, with, in addition,  $\Delta m > 5$  GeV for indirect decays via  $LQ\bar{D}$ . The limit quoted applies to direct decays via  $LL\bar{E}$  or  $LQ\bar{D}$  couplings. For indirect decays, the limits on the  $\tilde{e}_R$  mass are respectively 99 and 92 GeV for  $LL\bar{E}$  and  $LQ\bar{D}$  couplings and  $m_{\tilde{\chi}_1^0} = 10$  GeV and degrade slightly for larger  $\tilde{\chi}_1^0$  mass. Supersedes the results of ABBIENDI 00.
- 206 ABDALLAH 04M use data from  $\sqrt{s} = 192\text{--}208$  GeV to derive limits on sparticle masses under the assumption of  $\cancel{B}$  with  $LL\bar{E}$  or  $UD\bar{D}$  couplings. The results are valid for  $\mu = -200$  GeV,  $\tan\beta = 1.5$ ,  $\Delta m > 5$  GeV and assuming a BR of 1 for the given decay. The limit quoted is for indirect  $UD\bar{D}$  decays using the neutralino constraint of 39.5 GeV for  $LL\bar{E}$  and of 38.0 GeV for  $UD\bar{D}$  couplings, also derived in ABDALLAH 04M. For indirect decays via  $LL\bar{E}$  the limit improves to 95 GeV if the constraint from the neutralino is used and to 94 GeV if it is not used. For indirect decays via  $UD\bar{D}$  couplings it remains unchanged when the neutralino constraint is not used. Supersedes the result of ABREU 00U.
- 207 HEISTER 03G searches for the production of selectrons in the case of  $\cancel{B}$  prompt decays with  $LL\bar{E}$ ,  $LQ\bar{D}$  or  $UD\bar{D}$  couplings at  $\sqrt{s} = 189\text{--}209$  GeV. The search is performed for direct and indirect decays, assuming one coupling at a time to be non-zero. The limit holds for indirect decays mediated by  $LQ\bar{D}$  couplings with  $\Delta m > 10$  GeV. Limits are also given for  $LL\bar{E}$  direct ( $m_{\tilde{e}_R} > 96$  GeV) and indirect decays ( $m_{\tilde{e}_R} > 96$  GeV for  $m(\tilde{\chi}_1^0) > 23$  GeV from BARATE 98s) and for  $UD\bar{D}$  indirect decays ( $m_{\tilde{e}_R} > 94$  GeV with  $\Delta m > 10$  GeV). Supersedes the results from BARATE 01B.

- 208 ACHARD 02 searches for the production of selectrons in the case of  $\tilde{R}$  prompt decays with  $LL\bar{E}$  or  $UDD$  couplings at  $\sqrt{s}=189\text{--}208$  GeV. The search is performed for direct and indirect decays, assuming one coupling at the time to be nonzero. The limit holds for direct decays via  $LL\bar{E}$  couplings. Stronger limits are reached for  $LL\bar{E}$  indirect (79 GeV) and for  $UDD$  direct or indirect (96 GeV) decays.
- 209 BARATE 01 looked for acoplanar dielectron +  $\cancel{E}_T$  final states at 189 to 202 GeV. The limit assumes  $\mu=-200$  GeV and  $\tan\beta=2$  for the production cross section and 100% branching ratio for  $\tilde{e} \rightarrow e\tilde{\chi}_1^0$ . See their Fig. 1 for the dependence of the limit on  $\Delta m$ . These limits include and update the results of BARATE 99Q.
- 210 ABBIENDI 00J looked for acoplanar dielectron +  $\cancel{E}_T$  final states at  $\sqrt{s}=161\text{--}183$  GeV. The limit assumes  $\mu < -100$  GeV and  $\tan\beta=1.5$  for the production cross section and decay branching ratios, evaluated within the MSSM, and zero efficiency for decays other than  $\tilde{e} \rightarrow e\tilde{\chi}_1^0$ . See their Fig. 12 for the dependence of the limit on  $\Delta m$  and  $\tan\beta$ .
- 211 ABREU 00U studies decays induced by  $R$ -parity violating  $LL\bar{E}$  couplings, using data from  $\sqrt{s}=189$  GeV. They investigate topologies with multiple leptons, assuming one coupling at the time to be nonzero and giving rise to indirect decays. The limits assume a neutralino mass limit of 30 GeV, also derived in ABREU 00U. Updates ABREU 00I.
- 212 ABREU 00V use data from  $\sqrt{s}=130\text{--}189$  GeV to search for tracks with large impact parameter or visible decay vertices. Limits are obtained as a function of  $m_{\tilde{G}}$ , from a scan of the GMSB parameters space, after combining these results with the search for slepton pair production in the SUGRA framework from ABREU 01 to cover prompt decays and on stable particle searches from ABREU 00Q. For limits at different  $m_{\tilde{G}}$ , see their Fig. 12.
- 213 BARATE 00G combines the search for acoplanar dileptons, leptons with large impact parameters, kinks, and stable heavy-charged tracks, assuming 3 flavors of degenerate sleptons, produced in the  $s$  channel. Data collected at  $\sqrt{s}=189$  GeV.
- 214 ACCIARRI 99I establish indirect limits on  $m_{\tilde{e}_R}$  from the regions excluded in the  $M_2$  versus  $m_0$  plane by their chargino and neutralino searches at  $\sqrt{s}=130\text{--}183$  GeV. The situations where the  $\tilde{\chi}_1^0$  is the LSP (indirect decays) and where a  $\tilde{\ell}$  is the LSP (direct decays) were both considered. The weakest limit, quoted above, comes from direct decays with  $UDD$  couplings;  $LL\bar{E}$  couplings or indirect decays lead to a stronger limit.
- 215 ACCIARRI 98F looked for acoplanar dielectron+ $\cancel{E}_T$  final states at  $\sqrt{s}=130\text{--}172$  GeV. The limit assumes  $\mu=-200$  GeV, and zero efficiency for decays other than  $\tilde{e}_R \rightarrow e\tilde{\chi}_1^0$ . See their Fig. 6 for the dependence of the limit on  $\Delta m$ .
- 216 BARATE 98K looked for  $e^+e^-\gamma\gamma + \cancel{E}$  final states at  $\sqrt{s}=161\text{--}184$  GeV. The limit assumes  $\mu=-200$  GeV and  $\tan\beta=2$  for the evaluation of the production cross section. See Fig. 4 for limits on the  $(m_{\tilde{e}_R}, m_{\tilde{\chi}_1^0})$  plane and for the effect of cascade decays.
- 217 BREITWEG 98 used positron+jet events with missing energy and momentum to look for  $e^+q \rightarrow \tilde{e}\tilde{q}$  via gaugino-like neutralino exchange with decays into  $(e\tilde{\chi}_1^0)(q\tilde{\chi}_1^0)$ . See paper for dependences in  $m(\tilde{q})$ ,  $m(\tilde{\chi}_1^0)$ .
- 218 AID 96C used positron+jet events with missing energy and momentum to look for  $e^+q \rightarrow \tilde{e}\tilde{q}$  via neutralino exchange with decays into  $(e\tilde{\chi}_1^0)(q\tilde{\chi}_1^0)$ . See the paper for dependences on  $m_{\tilde{q}}$ ,  $m_{\tilde{\chi}_1^0}$ .

### $\tilde{\mu}$ (Smuon) MASS LIMIT

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>91.0	95	219 ABBIENDI	04 OPAL	$\Delta m > 3$ GeV, $\tilde{\mu}_R^+\tilde{\mu}_R^-$ , $ \mu  > 100$ GeV, $\tan\beta=1.5$
>86.7	95	220 ACHARD	04 L3	$\Delta m > 10$ GeV, $\tilde{\mu}_R^+\tilde{\mu}_R^-$ , $ \mu  > 200$ GeV, $\tan\beta \geq 2$
none 30–88	95	221 ABDALLAH	03M DLPH	$\Delta m > 5$ GeV, $\tilde{\mu}_R^+\tilde{\mu}_R^-$

- |     |    |     |          |          |                                                                            |
|-----|----|-----|----------|----------|----------------------------------------------------------------------------|
| >94 | 95 | 222 | ABDALLAH | 03M DLPH | $\tilde{\mu}_{R,1} \leq \tan\beta \leq 40,$<br>$\Delta m > 10 \text{ GeV}$ |
| >88 | 95 | 223 | HEISTER  | 02E ALEP | $\Delta m > 15 \text{ GeV}, \tilde{\mu}_R^+ \tilde{\mu}_R^-$               |
- • • We do not use the following data for averages, fits, limits, etc. • • •
- |     |    |     |          |          |                                                                                                  |
|-----|----|-----|----------|----------|--------------------------------------------------------------------------------------------------|
| >74 | 95 | 224 | ABBIENDI | 04F OPAL | $\tilde{R}, \tilde{\mu}_L$                                                                       |
| >87 | 95 | 225 | ABDALLAH | 04M DLPH | $\tilde{R}, \tilde{\mu}_R$ , indirect, $\Delta m > 5 \text{ GeV}$                                |
| >81 | 95 | 226 | HEISTER  | 03G ALEP | $\tilde{\mu}_L, \tilde{R}$ decays                                                                |
|     |    | 227 | ABAZOV   | 02H D0   | $\tilde{R}, \lambda'_{211}$                                                                      |
| >61 | 95 | 228 | ACHARD   | 02 L3    | $\tilde{\mu}_R, \tilde{R}$ decays                                                                |
| >85 | 95 | 229 | BARATE   | 01 ALEP  | $\Delta m > 10 \text{ GeV}, \tilde{\mu}_R^+ \tilde{\mu}_R^-$                                     |
| >65 | 95 | 230 | ABBIENDI | 00J OPAL | $\Delta m > 2 \text{ GeV}, \tilde{\mu}_R^+ \tilde{\mu}_R^-$                                      |
| >80 | 95 | 231 | ABREU    | 00V DLPH | $\tilde{\mu}_R \tilde{\mu}_R (\tilde{\mu}_R \rightarrow \mu \tilde{G}), m_{\tilde{G}} > 8$<br>eV |
| >77 | 95 | 232 | BARATE   | 98K ALEP | Any $\Delta m, \tilde{\mu}_R^+ \tilde{\mu}_R^-, \tilde{\mu}_R \rightarrow \mu \gamma \tilde{G}$  |
- 219 ABBIENDI 04 search for  $\tilde{\mu}_R \tilde{\mu}_R$  production in acoplanar di-muon final states in the 183–208 GeV data. See Fig. 14 for the dependence of the limits on  $m_{\tilde{\chi}_1^0}$  and for the limit at  $\tan\beta=35$ . Under the assumption of 100% branching ratio for  $\tilde{\mu}_R \rightarrow \mu \tilde{\chi}_1^0$ , the limit improves to 94.0 GeV for  $\Delta m > 4 \text{ GeV}$ . See Fig. 11 for the dependence of the limits on  $m_{\tilde{\chi}_1^0}$  at several values of the branching ratio. This limit supersedes ABBIENDI 00G.
- 220 ACHARD 04 search for  $\tilde{\mu}_R \tilde{\mu}_R$  production in acoplanar di-muon final states in the 192–209 GeV data. Limits on  $m_{\tilde{\mu}_R}$  are derived from a scan over the MSSM parameter space with universal GUT scale gaugino and scalar masses  $m_{1/2}$  and  $m_0$ ,  $1 \leq \tan\beta \leq 60$  and  $-2 \leq \mu \leq 2 \text{ TeV}$ . See Fig. 4 for the dependence of the limits on  $m_{\tilde{\chi}_1^0}$ . This limit supersedes ACCIARRI 99W.
- 221 ABDALLAH 03M looked for acoplanar dimuon  $+ \cancel{E}$  final states at  $\sqrt{s} = 189\text{--}208 \text{ GeV}$ . The limit assumes  $B(\tilde{\mu} \rightarrow \mu \tilde{\chi}_1^0) = 100\%$ . See Fig. 16 for limits on the  $(m_{\tilde{\mu}_R}, m_{\tilde{\chi}_1^0})$  plane. These limits include and update the results of ABREU 01.
- 222 ABDALLAH 03M uses data from  $\sqrt{s} = 192\text{--}208 \text{ GeV}$  to obtain limits in the framework of the MSSM with gaugino and sfermion mass universality at the GUT scale. An indirect limit on the mass is derived by constraining the MSSM parameter space by the results from direct searches for neutralinos (including cascade decays) and for sleptons. These limits are valid for values of  $M_2 < 1 \text{ TeV}$ ,  $|\mu| \leq 1 \text{ TeV}$  with the  $\tilde{\chi}_1^0$  as LSP. The quoted limit is obtained when there is no mixing in the third family. See Fig. 43 for the mass limits as a function of  $\tan\beta$ . These limits update the results of ABREU 00W.
- 223 HEISTER 02E looked for acoplanar dimuon  $+ \cancel{E}_T$  final states from  $e^+ e^-$  interactions between 183 and 209 GeV. The mass limit assumes  $B(\tilde{\mu} \rightarrow \mu \tilde{\chi}_1^0) = 1$ . See their Fig. 4 for the dependence of the limit on  $\Delta m$ . These limits include and update the results of BARATE 01.
- 224 ABBIENDI 04F use data from  $\sqrt{s} = 189\text{--}209 \text{ GeV}$ . They derive limits on sparticle masses under the assumption of  $\tilde{R}$  with  $LL\bar{E}$  or  $LQ\bar{D}$  couplings. The results are valid for  $\tan\beta = 1.5$ ,  $\mu = -200 \text{ GeV}$ , with, in addition,  $\Delta m > 5 \text{ GeV}$  for indirect decays via  $LQ\bar{D}$ . The limit quoted applies to direct decays with  $LL\bar{E}$  couplings and improves to 75 GeV for  $LQ\bar{D}$  couplings. The limits on the  $\tilde{\mu}_R$  mass for indirect decays are respectively 94 and 87 GeV for  $LL\bar{E}$  and  $LQ\bar{D}$  couplings and  $m_{\tilde{\chi}_1^0} = 10 \text{ GeV}$ . Supersedes the results of ABBIENDI 00.

- 225 ABDALLAH 04M use data from  $\sqrt{s} = 192\text{--}208$  GeV to derive limits on sparticle masses under the assumption of  $\tilde{R}$  with  $LL\bar{E}$  or  $\overline{UDD}$  couplings. The results are valid for  $\mu = -200$  GeV,  $\tan\beta = 1.5$ ,  $\Delta m > 5$  GeV and assuming a BR of 1 for the given decay. The limit quoted is for indirect  $\overline{UDD}$  decays using the neutralino constraint of 39.5 GeV for  $LL\bar{E}$  and of 38.0 GeV for  $\overline{UDD}$  couplings, also derived in ABDALLAH 04M. For indirect decays via  $LL\bar{E}$  the limit improves to 90 GeV if the constraint from the neutralino is used and remains at 87 GeV if it is not used. For indirect decays via  $\overline{UDD}$  couplings it degrades to 85 GeV when the neutralino constraint is not used. Supersedes the result of ABREU 00U.
- 226 HEISTER 03G searches for the production of smuons in the case of  $\tilde{R}$  prompt decays with  $LL\bar{E}$ ,  $LQ\bar{D}$  or  $\overline{UDD}$  couplings at  $\sqrt{s} = 189\text{--}209$  GeV. The search is performed for direct and indirect decays, assuming one coupling at a time to be non-zero. The limit holds for direct decays mediated by  $\tilde{R}$   $LQ\bar{D}$  couplings and improves to 90 GeV for indirect decays (for  $\Delta m > 10$  GeV). Limits are also given for  $LL\bar{E}$  direct ( $m_{\tilde{\mu}R} > 87$  GeV) and indirect decays ( $m_{\tilde{\mu}R} > 96$  GeV for  $m(\tilde{\chi}_1^0) > 23$  GeV from BARATE 98S) and for  $\overline{UDD}$  indirect decays ( $m_{\tilde{\mu}R} > 85$  GeV for  $\Delta m > 10$  GeV). Supersedes the results from BARATE 01B.
- 227 ABAZOV 02H looked in  $94\text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.8$  TeV for events with at least 2 muons and 2 jets for  $s$ -channel production of  $\tilde{\mu}$  or  $\tilde{\nu}$  and subsequent decay via  $\tilde{R}$  couplings  $LQ\bar{D}$ . A scan over the MSUGRA parameters is performed to exclude regions of the  $(m_0, m_{1/2})$  plane, examples being shown in Fig. 2.
- 228 ACHARD 02 searches for the production of smuons in the case of  $\tilde{R}$  prompt decays with  $LL\bar{E}$  or  $\overline{UDD}$  couplings at  $\sqrt{s}=189\text{--}208$  GeV. The search is performed for direct and indirect decays, assuming one coupling at the time to be nonzero. The limit holds for direct decays via  $LL\bar{E}$  couplings. Stronger limits are reached for  $LL\bar{E}$  indirect (87 GeV) and for  $\overline{UDD}$  direct or indirect (86 GeV) decays.
- 229 BARATE 01 looked for acoplanar dimuon +  $\cancel{E}_T$  final states at 189 to 202 GeV. The limit assumes 100% branching ratio for  $\tilde{\mu} \rightarrow \mu\tilde{\chi}_1^0$ . See their Fig. 1 for the dependence of the limit on  $\Delta m$ . These limits include and update the results of BARATE 99Q.
- 230 ABBIENDI 00J looked for acoplanar dimuon +  $\cancel{E}_T$  final states at  $\sqrt{s}= 161\text{--}183$  GeV. The limit assumes  $B(\tilde{\mu} \rightarrow \mu\tilde{\chi}_1^0)=1$ . Using decay branching ratios derived from the MSSM, a lower limit of 65 GeV is obtained for  $\mu < -100$  GeV and  $\tan\beta=1.5$ . See their Figs. 10 and 13 for the dependence of the limit on the branching ratio and on  $\Delta m$ .
- 231 ABREU 00V use data from  $\sqrt{s}= 130\text{--}189$  GeV to search for tracks with large impact parameter or visible decay vertices. Limits are obtained as function of  $m_{\tilde{G}}$ , after combining these results with the search for slepton pair production in the SUGRA framework from ABREU 01 to cover prompt decays and on stable particle searches from ABREU 00Q. For limits at different  $m_{\tilde{G}}$ , see their Fig. 12.
- 232 BARATE 98K looked for  $\mu^+\mu^-\gamma\gamma + \cancel{E}$  final states at  $\sqrt{s}= 161\text{--}184$  GeV. See Fig. 4 for limits on the  $(m_{\tilde{\mu}R}, m_{\tilde{\chi}_1^0})$  plane and for the effect of cascade decays.

### $\tilde{\tau}$ (Stau) MASS LIMIT

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>85.2	95	233 ABBIENDI	04 OPAL	$\Delta m > 6$ GeV, $\theta_\tau=\pi/2$ , $ \mu  > 100$ GeV, $\tan\beta=1.5$
>78.3	95	234 ACHARD	04 L3	$\Delta m > 15$ GeV, $\theta_\tau=\pi/2$ , $ \mu  > 200$ GeV, $\tan\beta \geq 2$
<b>&gt;81.9</b>	95	235 ABDALLAH	03M DLPH	$\Delta m > 15$ GeV, all $\theta_\tau$
none $m_\tau - 26.3$	95	235 ABDALLAH	03M DLPH	$\Delta m > m_\tau$ , all $\theta_\tau$
>79	95	236 HEISTER	02E ALEP	$\Delta m > 15$ GeV, $\theta_\tau=\pi/2$
>76	95	236 HEISTER	02E ALEP	$\Delta m > 15$ GeV, $\theta_\tau=0.91$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>74	95	237	ABBIENDI	04F	OPAL	$\tilde{R}, \tilde{\tau}_L$
>68	95	238,239	ABDALLAH	04H	DLPH	AMSB, $\mu > 0$
>90	95	240	ABDALLAH	04M	DLPH	$\tilde{R}, \tilde{\tau}_R$ , indirect, $\Delta m > 5$ GeV
>82.5	95	241	ABDALLAH	03D	DLPH	$\tilde{\tau}_R \rightarrow \tau \tilde{G}$ , all $\tau(\tilde{\tau}_R)$
>70	95	242	HEISTER	03G	ALEP	$\tilde{\tau}_R, \tilde{R}$ decay
>61	95	243	ACHARD	02	L3	$\tilde{\tau}_R, \tilde{R}$ decays
>77	95	244	HEISTER	02R	ALEP	$\tau_1$ , any lifetime
>70	95	245	BARATE	01	ALEP	$\Delta m > 10$ GeV, $\theta_\tau = \pi/2$
>68	95	245	BARATE	01	ALEP	$\Delta m > 10$ GeV, $\theta_\tau = 0.91$
>64	95	246	ABBIENDI	00J	OPAL	$\Delta m > 10$ GeV, $\tilde{\tau}_R^+ \tilde{\tau}_R^-$
>84	95	247	ABREU	00V	DLPH	$\tilde{\ell}_R \tilde{\ell}_R (\tilde{\ell}_R \rightarrow \ell \tilde{G}), m_{\tilde{G}} > 9$ eV
>73	95	248	ABREU	00V	DLPH	$\tilde{\tau}_1 \tilde{\tau}_1 (\tilde{\tau}_1 \rightarrow \tau \tilde{G})$ , all $\tau(\tilde{\tau}_1)$
>52	95	249	BARATE	98K	ALEP	Any $\Delta m, \theta_\tau = \pi/2, \tilde{\tau}_R \rightarrow \tau \gamma \tilde{G}$

233 ABBIENDI 04 search for  $\tilde{\tau}\tilde{\tau}$  production in acoplanar di-tau final states in the 183–208 GeV data. See Fig. 15 for the dependence of the limits on  $m_{\tilde{\chi}_1^0}$  and for

the limit at  $\tan\beta=35$ . Under the assumption of 100% branching ratio for  $\tilde{\tau}_R \rightarrow \tau \tilde{\chi}_1^0$ , the limit improves to 89.8 GeV for  $\Delta m > 8$  GeV. See Fig. 12 for the dependence of the limits on  $m_{\tilde{\chi}_1^0}$  at several values of the branching ratio and for their dependence on  $\theta_\tau$ .

This limit supersedes ABBIENDI 00G.

234 ACHARD 04 search for  $\tilde{\tau}\tilde{\tau}$  production in acoplanar di-tau final states in the 192–209 GeV data. Limits on  $m_{\tilde{\tau}_R}$  are derived from a scan over the MSSM parameter space with universal GUT scale gaugino and scalar masses  $m_{1/2}$  and  $m_0$ ,  $1 \leq \tan\beta \leq 60$  and  $-2 \leq \mu \leq 2$  TeV. See Fig. 4 for the dependence of the limits on  $m_{\tilde{\chi}_1^0}$ .

235 ABDALLAH 03M looked for acoplanar ditau +  $\cancel{E}$  final states at  $\sqrt{s} = 130$ –208 GeV. A dedicated search was made for low mass  $\tilde{\tau}$ s decoupling from the  $Z^0$ . The limit assumes  $B(\tilde{\tau} \rightarrow \tau \tilde{\chi}_1^0) = 100\%$ . See Fig. 20 for limits on the  $(m_{\tilde{\tau}}, m_{\tilde{\chi}_1^0})$  plane and as function

of the  $\tilde{\chi}_1^0$  mass and of the branching ratio. The limit in the low-mass region improves to 29.6 and 31.1 GeV for  $\tilde{\tau}_R$  and  $\tilde{\tau}_L$ , respectively, at  $\Delta m > m_\tau$ . The limit in the high-mass region improves to 84.7 GeV for  $\tilde{\tau}_R$  and  $\Delta m > 15$  GeV. These limits include and update the results of ABREU 01.

236 HEISTER 02E looked for acoplanar ditau +  $\cancel{E}_T$  final states from  $e^+e^-$  interactions between 183 and 209 GeV. The mass limit assumes  $B(\tilde{\tau} \rightarrow \tau \tilde{\chi}_1^0) = 1$ . See their Fig. 4 for the dependence of the limit on  $\Delta m$ . These limits include and update the results of BARATE 01.

237 ABBIENDI 04F use data from  $\sqrt{s} = 189$ –209 GeV. They derive limits on sparticle masses under the assumption of  $\tilde{R}$  with  $LL\bar{E}$  or  $LQ\bar{D}$  couplings. The results are valid for  $\tan\beta = 1.5$ ,  $\mu = -200$  GeV, with, in addition,  $\Delta m > 5$  GeV for indirect decays via  $LQ\bar{D}$ . The limit quoted applies to direct decays with  $LL\bar{E}$  couplings and improves to 75 GeV for  $LQ\bar{D}$  couplings. The limit on the  $\tilde{\tau}_R$  mass for indirect decays is 92 GeV for  $LL\bar{E}$  couplings at  $m_{\tilde{\chi}_1^0} = 10$  GeV and no exclusion is obtained for  $LQ\bar{D}$  couplings. Supersedes the results of ABBIENDI 00.

238 ABDALLAH 04H use data from LEP 1 and  $\sqrt{s} = 192$ –208 GeV. They re-use results or re-analyze the data from ABDALLAH 03M to put limits on the parameter space of anomaly-mediated supersymmetry breaking (AMSB), which is scanned in the region  $1 < m_{3/2} < 50$  TeV,  $0 < m_0 < 1000$  GeV,  $1.5 < \tan\beta < 35$ , both signs of  $\mu$ . The constraints are obtained from the searches for mass degenerate chargino and neutralino, for SM-like and invisible Higgs, for leptonically decaying charginos and from the limit on non-SM  $Z$  width of 3.2 MeV. The limit is for  $m_t = 174.3$  GeV (see Table 2 for other  $m_t$  values).



- 239 The limit improves to 75 GeV for  $\mu < 0$ .
- 240 ABDALLAH 04M use data from  $\sqrt{s} = 192\text{--}208$  GeV to derive limits on sparticle masses under the assumption of  $\cancel{R}$  with  $LL\bar{E}$  couplings. The results are valid for  $\mu = -200$  GeV,  $\tan\beta = 1.5$ ,  $\Delta m > 5$  GeV and assuming a BR of 1 for the given decay. The limit quoted is for indirect decays using the neutralino constraint of 39.5 GeV, also derived in ABDALLAH 04M. For indirect decays via  $LL\bar{E}$  the limit decreases to 86 GeV if the constraint from the neutralino is not used. Supersedes the result of ABREU 00U.
- 241 ABDALLAH 03D use data from  $\sqrt{s} = 130\text{--}208$  GeV to search for tracks with large impact parameter or visible decay vertices and for heavy charged stable particles. Limits are obtained as function of  $m(\tilde{G})$ , after combining these results with the search for slepton pair production in the SUGRA framework from ABDALLAH 03M to cover prompt decays. The above limit is reached for the stau decaying promptly,  $m(\tilde{G}) < 6$  eV, and is computed for stau mixing yielding the minimal cross section. Stronger limits are obtained for longer lifetimes. See their Fig. 9. Supersedes the results of ABREU 01G.
- 242 HEISTER 03G searches for the production of stau in the case of  $\cancel{R}$  prompt decays with  $LL\bar{E}$ ,  $LQ\bar{D}$  or  $UDD$  couplings at  $\sqrt{s} = 189\text{--}209$  GeV. The search is performed for direct and indirect decays, assuming one coupling at a time to be non-zero. The limit holds for indirect decays mediated by  $\cancel{R}$   $UDD$  couplings with  $\Delta m > 10$  GeV. Limits are also given for  $LL\bar{E}$  direct ( $m_{\tilde{\tau}_R} > 87$  GeV) and indirect decays ( $m_{\tilde{\tau}_R} > 95$  GeV for  $m(\tilde{\chi}_1^0) > 23$  GeV from BARATE 98S) and for  $LQ\bar{D}$  indirect decays ( $m_{\tilde{\tau}_R} > 76$  GeV). Supersedes the results from BARATE 01B.
- 243 ACHARD 02 searches for the production of staus in the case of  $\cancel{R}$  prompt decays with  $LL\bar{E}$  or  $UDD$  couplings at  $\sqrt{s} = 189\text{--}208$  GeV. The search is performed for direct and indirect decays, assuming one coupling at a time to be nonzero. The limit holds for direct decays via  $LL\bar{E}$  couplings. Stronger limits are reached for  $LL\bar{E}$  indirect (86 GeV) and for  $UDD$  direct or indirect (75 GeV) decays.
- 244 HEISTER 02R search for signals of GMSB in the 189–209 GeV data. For the  $\tilde{\chi}_1^0$  NLSP scenario, they looked for topologies consisting of  $\gamma\gamma\cancel{E}$  or a single  $\gamma$  not pointing to the interaction vertex. For the  $\tilde{\ell}$  NLSP case, the topologies consist of  $\ell\ell\cancel{E}$ , including leptons with large impact parameters, kinks, or stable particles. Limits are derived from a scan over the GMSB parameters (see their Table 5 for the ranges). The limit remains valid whichever is the NLSP. The absolute mass bound on the  $\tilde{\chi}_1^0$  for any lifetime includes indirect limits from the slepton search HEISTER 02E performed within the MSUGRA framework. A bound for any NLSP and any lifetime of 77 GeV has also been derived by using the constraints from the neutral Higgs search in HEISTER 02. In the co-NLSP scenario, limits  $m_{\tilde{e}_R} > 83$  GeV (neglecting  $t$ -channel exchange) and  $m_{\tilde{\mu}_R} > 88$  GeV are obtained independent of the lifetime. Supersedes the results from BARATE 00G.
- 245 BARATE 01 looked for acoplanar ditau +  $\cancel{E}_T$  final states at 189 to 202 GeV. A slight excess (with 1.2% probability) of events is observed relative to the expected SM background. The limit assumes 100% branching ratio for  $\tilde{\tau} \rightarrow \tau\tilde{\chi}_1^0$ . See their Fig. 1 for the dependence of the limit on  $\Delta m$ . These limits include and update the results of BARATE 99Q.
- 246 ABBIENDI 00J looked for acoplanar ditau +  $\cancel{E}_T$  final states at  $\sqrt{s} = 161\text{--}183$  GeV. The limit assumes  $B(\tilde{\tau} \rightarrow \tau\tilde{\chi}_1^0) = 1$ . Using decay branching ratios derived from the MSSM, a lower limit of 60 GeV at  $\Delta m > 9$  GeV is obtained for  $\mu < -100$  GeV and  $\tan\beta = 1.5$ . See their Figs. 11 and 14 for the dependence of the limit on the branching ratio and on  $\Delta m$ .
- 247 ABREU 00V use data from  $\sqrt{s} = 130\text{--}189$  GeV to search for tracks with large impact parameter or visible decay vertices. Limits are obtained as function of  $m_{\tilde{G}}$ , after combining these results with the search for slepton pair production in the SUGRA framework from ABREU 01 to cover prompt decays and on stable particle searches from ABREU 00Q. The above limit assumes the degeneracy of stau and smuon. For limits at different  $m_{\tilde{G}}$ , see their Fig. 12.
- 248 ABREU 00V use data from  $\sqrt{s} = 130\text{--}189$  GeV to search for tracks with large impact parameter or visible decay vertices. Limits are obtained as function of  $m_{\tilde{G}}$ , after combining

these results with the search for slepton pair production in the SUGRA framework from ABREU 01 to cover prompt decays and on stable particle searches from ABREU 00Q. The above limit is reached for the stau mixing yielding the minimal cross section and decaying promptly. Stronger limits are obtained for longer lifetimes or for  $\tilde{\tau}_R$ ; see their Fig. 11. For  $10 \leq m_{\tilde{G}} \leq 310$  eV, the whole range  $2 \leq m_{\tilde{\tau}_1} \leq 80$  GeV is excluded.

Supersedes the results of ABREU 99C and ABREU 99F.

- 249 BARATE 98K looked for  $\tau^+ \tau^- \gamma \gamma + \cancel{E}$  final states at  $\sqrt{s}=161\text{--}184$  GeV. See Fig. 4 for limits on the  $(m_{\tilde{\tau}_R}, m_{\tilde{\chi}_1^0})$  plane and for the effect of cascade decays.

### Degenerate Charged Sleptons

Unless stated otherwise in the comment lines or in the footnotes, the following limits assume 3 families of degenerate charged sleptons.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>93	95	250 BARATE	01 ALEP	$\Delta m > 10$ GeV, $\tilde{\ell}_R^+ \tilde{\ell}_R^-$
>70	95	250 BARATE	01 ALEP	all $\Delta m$ , $\tilde{\ell}_R^+ \tilde{\ell}_R^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>88	95	251 ABDALLAH	03D DLPH	$\tilde{\ell}_R \rightarrow \ell \tilde{G}$ , all $\tau(\tilde{\ell}_R)$
>82.7	95	252 ACHARD	02 L3	$\tilde{\ell}_R, \cancel{R}$ decays, MSUGRA
>83	95	253 ABBIENDI	01 OPAL	$e^+ e^- \rightarrow \tilde{\ell}_1 \tilde{\ell}_1$ , GMSB, $\tan\beta=2$
		254 ABREU	01 DLPH	$\tilde{\ell} \rightarrow \ell \tilde{\chi}_2^0, \tilde{\chi}_2^0 \rightarrow \gamma \tilde{\chi}_1^0$ , $\ell=e,\mu$
>68.8	95	255 ACCIARRI	01 L3	$\tilde{\ell}_R, \cancel{R}$ , $0.7 \leq \tan\beta \leq 40$
>84	95	256,257 ABREU	00V DLPH	$\tilde{\ell}_R \tilde{\ell}_R (\tilde{\ell}_R \rightarrow \ell \tilde{G})$ , $m_{\tilde{G}} > 9$ eV

- 250 BARATE 01 looked for acoplanar dilepton +  $\cancel{E}_T$  and single electron (for  $\tilde{e}_R \tilde{e}_L$ ) final states at 189 to 202 GeV. The limit assumes  $\mu=-200$  GeV and  $\tan\beta=2$  for the production cross section and decay branching ratios, evaluated within the MSSM, and zero efficiency for decays other than  $\tilde{\ell} \rightarrow \ell \tilde{\chi}_1^0$ . The slepton masses are determined from the GUT relations without stau mixing. See their Fig. 1 for the dependence of the limit on  $\Delta m$ .

- 251 ABDALLAH 03D use data from  $\sqrt{s}=130\text{--}208$  GeV to search for tracks with large impact parameter or visible decay vertices and for heavy charged stable particles. Limits are obtained as function of  $m(\tilde{G})$ , after combining these results with the search for slepton pair production in the SUGRA framework from ABDALLAH 03M to cover prompt decays. The above limit is reached for prompt decays and assumes the degeneracy of the sleptons. For limits at different  $m(\tilde{G})$ , see their Fig. 9. Supersedes the results of ABREU 01G.

- 252 ACHARD 02 searches for the production of sparticles in the case of  $\cancel{R}$  prompt decays with  $L\bar{L}E$  or  $U\bar{D}D$  couplings at  $\sqrt{s}=189\text{--}208$  GeV. The search is performed for direct and indirect decays, assuming one coupling at the time to be nonzero. The MSUGRA limit results from a scan over the MSSM parameter space with the assumption of gaugino and scalar mass unification at the GUT scale and no mixing in the slepton sector, imposing simultaneously the exclusions from neutralino, chargino, sleptons, and squarks analyses. The limit holds for  $L\bar{L}E$  couplings and increases to 88.7 GeV for  $U\bar{D}D$  couplings. For L3 limits from  $LQ\bar{D}$  couplings, see ACCIARRI 01.

- 253 ABBIENDI 01 looked for final states with  $\gamma\gamma\cancel{E}$ ,  $\ell\ell\cancel{E}$ , with possibly additional activity and four leptons +  $\cancel{E}$  to search for prompt decays of  $\tilde{\chi}_1^0$  or  $\tilde{\ell}_1$  in GMSB. They derive limits in the plane  $(m_{\tilde{\chi}_1^0}, m_{\tilde{\tau}_1})$ , see Fig. 6, allowing either the  $\tilde{\chi}_1^0$  or a  $\tilde{\ell}_1$  to be the NLSP.

Two scenarios are considered:  $\tan\beta=2$  with the 3 sleptons degenerate in mass and  $\tan\beta=20$  where the  $\tilde{\tau}_1$  is lighter than the other sleptons. Data taken at  $\sqrt{s}=189$  GeV. For  $\tan\beta=20$ , the obtained limits are  $m_{\tilde{\tau}_1} > 69$  GeV and  $m_{\tilde{e}_1, \tilde{\mu}_1} > 88$  GeV.

- 254 ABREU 01 looked for acoplanar dilepton + diphoton +  $\cancel{E}$  final states from  $\tilde{\ell}$  cascade decays at  $\sqrt{s}=130\text{--}189$  GeV. See Fig. 9 for limits on the  $(\mu, M_2)$  plane for  $m_{\tilde{\ell}}=80$  GeV,  $\tan\beta=1.0$ , and assuming degeneracy of  $\tilde{\mu}$  and  $\tilde{e}$ .
- 255 ACCIARRI 01 searches for multi-lepton and/or multi-jet final states from  $\cancel{R}$  prompt decays with  $LL\bar{E}$ ,  $LQ\bar{D}$ , or  $UD\bar{D}$  couplings at  $\sqrt{s}=189$  GeV. The search is performed for direct and indirect decays of neutralinos, charginos, and scalar leptons, with the  $\tilde{\chi}_1^0$  or a  $\tilde{\ell}$  as LSP and assuming one coupling to be nonzero at a time. Mass limits are derived using simultaneously the constraints from the neutralino, chargino, and slepton analyses; and the  $Z^0$  width measurements from ACCIARRI 00C in a scan of the parameter space assuming MSUGRA with gaugino and scalar mass universality. Updates and supersedes the results from ACCIARRI 99i.
- 256 ABREU 00V use data from  $\sqrt{s}=130\text{--}189$  GeV to search for tracks with large impact parameter or visible decay vertices. Limits are obtained as function of  $m_{\tilde{G}}$ , after combining these results with the search for slepton pair production in the SUGRA framework from ABREU 01 to cover prompt decays and on stable particle searches from ABREU 00Q. For limits at different  $m_{\tilde{G}}$ , see their Fig. 12.
- 257 The above limit assumes the degeneracy of stau and smuon.

### Long-lived $\tilde{\ell}$ (Slepton) MASS LIMIT

Limits on scalar leptons which leave detector before decaying. Limits from  $Z$  decays are independent of lepton flavor. Limits from continuum  $e^+e^-$  annihilation are also independent of flavor for smuons and staus. Selectron limits from  $e^+e^-$  collisions in the continuum depend on MSSM parameters because of the additional neutralino exchange contribution.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>98	95	258 ABBIENDI	03L OPAL	$\tilde{\mu}_R, \tilde{\tau}_R$
<b>none 2–87.5</b>	95	259 ABREU	00Q DLPH	$\tilde{\mu}_R, \tilde{\tau}_R$
>81.2	95	260 ACCIARRI	99H L3	$\tilde{\mu}_R, \tilde{\tau}_R$
>81	95	261 BARATE	98K ALEP	$\tilde{\mu}_R, \tilde{\tau}_R$

- 258 ABBIENDI 03L used  $e^+e^-$  data at  $\sqrt{s}=130\text{--}209$  GeV to select events with two high momentum tracks with anomalous  $dE/dx$ . The excluded cross section is compared to the theoretical expectation as a function of the heavy particle mass in their Fig. 3. The limit improves to 98.5 GeV for  $\tilde{\mu}_L$  and  $\tilde{\tau}_L$ . The bounds are valid for colorless spin 0 particles with lifetimes longer than  $10^{-6}$  s. Supersedes the results from ACKERSTAFF 98P.
- 259 ABREU 00Q searches for the production of pairs of heavy, charged stable particles in  $e^+e^-$  annihilation at  $\sqrt{s}=130\text{--}189$  GeV. The upper bound improves to 88 GeV for  $\tilde{\mu}_L, \tilde{\tau}_L$ . These limits include and update the results of ABREU 98P.
- 260 ACCIARRI 99H searched for production of pairs of back-to-back heavy charged particles at  $\sqrt{s}=130\text{--}183$  GeV. The upper bound improves to 82.2 GeV for  $\tilde{\mu}_L, \tilde{\tau}_L$ .
- 261 The BARATE 98K mass limit improves to 82 GeV for  $\tilde{\mu}_L, \tilde{\tau}_L$ . Data collected at  $\sqrt{s}=161\text{--}184$  GeV.

### $\tilde{q}$ (Squark) MASS LIMIT

For  $m_{\tilde{q}} > 60\text{--}70$  GeV, it is expected that squarks would undergo a cascade decay via a number of neutralinos and/or charginos rather than undergo a direct decay to photinos as assumed by some papers. Limits obtained when direct decay is assumed are usually higher than limits when cascade decays are included.

Limits from  $e^+e^-$  collisions depend on the mixing angle of the lightest mass eigenstate  $\tilde{q}_1 = \tilde{q}_R \sin\theta_q + \tilde{q}_L \cos\theta_q$ . It is usually assumed that only the sbottom and stop squarks have non-trivial mixing angles (see the stop and sbottom sections). Here, unless

otherwise noted, squarks are always taken to be either left/right degenerate, or purely of left or right type. Data from  $Z$  decays have set squark mass limits above 40 GeV, in the case of  $\tilde{q} \rightarrow q\tilde{\chi}_1^0$  decays if  $\Delta m = m_{\tilde{q}} - m_{\tilde{\chi}_1^0} \gtrsim 5$  GeV. For smaller values of  $\Delta m$ , current constraints on the invisible width of the  $Z$  ( $\Delta\Gamma_{\text{inv}} < 2.0$  MeV, LEP 00) exclude  $m_{\tilde{u}_{L,R}} < 44$  GeV,  $m_{\tilde{d}_R} < 33$  GeV,  $m_{\tilde{d}_L} < 44$  GeV and, assuming all squarks degenerate,  $m_{\tilde{q}} < 45$  GeV.

Limits made obsolete by the most recent analyses of  $e^+e^-$ ,  $p\bar{p}$ , and  $ep$  collisions can be found in previous Editions of this Review.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
> 99.5	95	262 ACHARD	04 L3	$\Delta m > 10$ GeV, $e^+e^- \rightarrow \tilde{q}_{L,R} \tilde{q}_{L,R}$
> 97	95	262 ACHARD	04 L3	$\Delta m \geq 10$ GeV, $e^+e^- \rightarrow \tilde{q}_R \tilde{q}_R$
>138	95	263 ABBOTT	01D D0	$ll+\text{jets}+\cancel{E}_T$ , $\tan\beta < 10$ , $m_0 < 300$ GeV, $\mu < 0$ , $A_0=0$
>255	95	263 ABBOTT	01D D0	$\tan\beta=2$ , $m_{\tilde{g}}=m_{\tilde{q}}$ , $\mu < 0$ , $A_0=0$ , $ll+\text{jets}+\cancel{E}_T$
> 97	95	264 BARATE	01 ALEP	$e^+e^- \rightarrow \tilde{q}\tilde{q}$ , $\Delta m > 6$ GeV
<b>&gt;250</b>	95	265 ABBOTT	99L D0	$\tan\beta=2$ , $\mu < 0$ , $A=0$ , $\text{jets}+\cancel{E}_T$
>224	95	266 ABE	96D CDF	$m_{\tilde{g}} \leq m_{\tilde{q}}$ ; with cascade decays, $ll+\text{jets}+\cancel{E}_T$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>275		267 AKTAS	04D H1	$e^\pm p \rightarrow \tilde{u}_L, \tilde{R}, LQ\bar{D}$
>280		267 AKTAS	04D H1	$e^\pm p \rightarrow \tilde{D}_R, \tilde{R}, LQ\bar{D}$
		268 ADLOFF	03 H1	$e^\pm p \rightarrow \tilde{q}, \tilde{R}, LQ\bar{D}$
>276	95	269 CHEKANOV	03B ZEUS	$\tilde{d} \rightarrow e^- u, \nu d, \tilde{R}, LQ\bar{D}, \lambda > 0.1$
>260	95	269 CHEKANOV	03B ZEUS	$\tilde{u} \rightarrow e^+ d, \tilde{R}, LQ\bar{D}, \lambda > 0.1$
> 82.5	95	270 HEISTER	03G ALEP	$\tilde{u}_R, \tilde{R}$ decay
> 77	95	270 HEISTER	03G ALEP	$\tilde{d}_R, \tilde{R}$ decay
>240	95	271 ABAZOV	02F D0	$\tilde{q}, \tilde{R} \lambda'_{2jk}$ indirect decays, $\tan\beta=2$ , any $m_{\tilde{g}}$
>265	95	271 ABAZOV	02F D0	$\tilde{q}, \tilde{R} \lambda'_{2jk}$ indirect decays, $\tan\beta=2$ , $m_{\tilde{q}}=m_{\tilde{g}}$
		272 ABAZOV	02G D0	$p\bar{p} \rightarrow \tilde{g}\tilde{g}, \tilde{g}\tilde{q}$
none 80–121	95	273 ABBIENDI	02 OPAL	$e\gamma \rightarrow \tilde{u}_L, \tilde{R} LQ\bar{D}, \lambda=0.3$
none 80–158	95	273 ABBIENDI	02 OPAL	$e\gamma \rightarrow \tilde{d}_R, \tilde{R} LQ\bar{D}, \lambda=0.3$
none 80–185	95	274 ABBIENDI	02B OPAL	$e\gamma \rightarrow \tilde{u}_L, \tilde{R} LQ\bar{D}, \lambda=0.3$
none 80–196	95	274 ABBIENDI	02B OPAL	$e\gamma \rightarrow \tilde{d}_R, \tilde{R} LQ\bar{D}, \lambda=0.3$
> 79	95	275 ACHARD	02 L3	$\tilde{u}_R, \tilde{R}$ decays
> 55	95	275 ACHARD	02 L3	$\tilde{d}_R, \tilde{R}$ decays
>263	95	276 CHEKANOV	02 ZEUS	$\tilde{u}_L \rightarrow \mu q, \tilde{R}, LQ\bar{D}, \lambda=0.3$
>258	95	276 CHEKANOV	02 ZEUS	$\tilde{u}_L \rightarrow \tau q, \tilde{R}, LQ\bar{D}, \lambda=0.3$
> 82	95	277 BARATE	01B ALEP	$\tilde{u}_R, \tilde{R}$ decays
> 68	95	277 BARATE	01B ALEP	$\tilde{d}_R, \tilde{R}$ decays
none 150–204	95	278 BREITWEG	01 ZEUS	$e^+ p \rightarrow \tilde{d}_R, \tilde{R} LQ\bar{D}, \lambda=0.3$
>200	95	279 ABBOTT	00C D0	$\tilde{u}_L, \tilde{R}, \lambda'_{2jk}$ decays

>180	95	279	ABBOTT	00C D0	$\tilde{d}_R, \mathcal{R}, \lambda'_{2jk}$ decays
>390	95	280	ACCIARRI	00P L3	$e^+ e^- \rightarrow q\bar{q}, \mathcal{R}, \lambda=0.3$
>148	95	281	AFFOLDER	00K CDF	$\tilde{d}_L, \mathcal{R}, \lambda'_{ij3}$ decays
>200	95	282	BARATE	00I ALEP	$e^+ e^- \rightarrow q\bar{q}, \mathcal{R}, \lambda=0.3$
none 150–269	95	283	BREITWEG	00E ZEUS	$e^+ p \rightarrow \tilde{u}_L, \mathcal{R}, LQ\bar{D}, \lambda=0.3$
>240	95	284	ABBOTT	99 D0	$\tilde{q} \rightarrow \tilde{\chi}_2^0 X \rightarrow \tilde{\chi}_1^0 \gamma X, m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} > 20 \text{ GeV}$
>320	95	284	ABBOTT	99 D0	$\tilde{q} \rightarrow \tilde{\chi}_1^0 X \rightarrow \tilde{G} \gamma X$
>243	95	285	ABBOTT	99K D0	any $m_{\tilde{g}}, \mathcal{R}, \tan\beta=2, \mu < 0$
>200	95	286	ABE	99M CDF	$p\bar{p} \rightarrow \tilde{q}\tilde{q}, \mathcal{R}$
none 80–134	95	287	ABREU	99G DLPH	$e\gamma \rightarrow \tilde{u}_L, \mathcal{R}, LQ\bar{D}, \lambda=0.3$
none 80–161	95	287	ABREU	99G DLPH	$e\gamma \rightarrow \tilde{d}_R, \mathcal{R}, LQ\bar{D}, \lambda=0.3$
>225	95	288	ABBOTT	98E D0	$\tilde{u}_L, \mathcal{R}, \lambda'_{1jk}$ decays
>204	95	288	ABBOTT	98E D0	$\tilde{d}_R, \mathcal{R}, \lambda'_{1jk}$ decays
> 79	95	288	ABBOTT	98E D0	$\tilde{d}_L, \mathcal{R}, \lambda'_{ijk}$ decays
>202	95	289	ABE	98S CDF	$\tilde{u}_L, \mathcal{R}, \lambda'_{2jk}$ decays
>160	95	289	ABE	98S CDF	$\tilde{d}_R, \mathcal{R}, \lambda'_{2jk}$ decays
>140	95	290	ACKERSTAFF	98V OPAL	$e^+ e^- \rightarrow q\bar{q}, \mathcal{R}, \lambda=0.3$
> 77	95	291	BREITWEG	98 ZEUS	$m_{\tilde{q}}=m_{\tilde{e}}, m(\tilde{\chi}_1^0)=40 \text{ GeV}$
		292	DATTA	97 THEO	$\tilde{\nu}$ 's lighter than $\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$
>216	95	293	DERRICK	97 ZEUS	$e p \rightarrow \tilde{q}, \tilde{q} \rightarrow \mu j \text{ or } \tau j, \mathcal{R}$
none 130–573	95	294	HEWETT	97 THEO	$q\tilde{g} \rightarrow \tilde{q}, \tilde{q} \rightarrow q\tilde{g}$ , with a light gluino
none 190–650	95	295	TEREKHOV	97 THEO	$q\tilde{g} \rightarrow \tilde{q}\tilde{g}, \tilde{q} \rightarrow q\tilde{g}$ , with a light gluino
> 63	95	296	AID	96C H1	$m_{\tilde{q}}=m_{\tilde{e}}, m_{\tilde{\chi}_1^0}=35 \text{ GeV}$
none 330–400	95	297	TEREKHOV	96 THEO	$u g \rightarrow \tilde{u}\tilde{g}, \tilde{u} \rightarrow u\tilde{g}$ with a light gluino
>176	95	298	ABACHI	95C D0	Any $m_{\tilde{g}} < 300 \text{ GeV}$ ; with cascade decays
		299	ABE	95T CDF	$\tilde{q} \rightarrow \tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 \gamma$
> 90	90	300	ABE	92L CDF	Any $m_{\tilde{g}} < 410 \text{ GeV}$ ; with cascade decay
>100		301	ROY	92 RVUE	$p\bar{p} \rightarrow \tilde{q}\tilde{q}, \mathcal{R}$
		302	NOJIRI	91 COSM	

262 ACHARD 04 search for the production of  $\tilde{q}\tilde{q}$  of the first two generations in acoplanar di-jet final states in the 192–209 GeV data. Degeneracy of the squark masses is assumed either for both left and right squarks or for right squarks only, as well as  $B(\tilde{q} \rightarrow q\tilde{\chi}_1^0) = 1$ . See Fig. 7 for the dependence of the limits on  $m_{\tilde{\chi}_1^0}$ . This limit supersedes ACCIARRI 99v.

263 ABBOTT 01D looked in  $\sim 108 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.8 \text{ TeV}$  for events with  $e e, \mu\mu,$  or  $e\mu$  accompanied by at least 2 jets and  $\cancel{E}_T$ . Excluded regions are obtained in the MSUGRA framework from a scan over the parameters  $0 < m_0 < 300 \text{ GeV}, 10 < m_{1/2} < 110 \text{ GeV},$  and  $1.2 < \tan\beta < 10$ .

264 BARATE 01 looked for acoplanar dijets +  $\cancel{E}_T$  final states at 189 to 202 GeV. The limit assumes  $B(\tilde{q} \rightarrow q\tilde{\chi}_1^0)=1,$  with  $\Delta m = m_{\tilde{q}} - m_{\tilde{\chi}_1^0}$ . It applies to  $\tan\beta=4, \mu=-400 \text{ GeV}$ .

- See their Fig. 2 for the exclusion in the  $(m_{\tilde{q}}, m_{\tilde{g}})$  plane. These limits include and update the results of BARATE 99Q.
- 265 ABBOTT 99L consider events with three or more jets and large  $\cancel{E}_T$ . Spectra and decay rates are evaluated in the framework of minimal Supergravity, assuming five flavors of degenerate squarks, and scanning the space of the universal gaugino ( $m_{1/2}$ ) and scalar ( $m_0$ ) masses. See their Figs. 2–3 for the dependence of the limit on the relative value of  $m_{\tilde{q}}$  and  $m_{\tilde{g}}$ .
- 266 ABE 96D searched for production of gluinos and five degenerate squarks in final states containing a pair of leptons, two jets, and missing  $E_T$ . The two leptons arise from the semileptonic decays of charginos produced in the cascade decays. The limit is derived for fixed  $\tan\beta = 4.0$ ,  $\mu = -400$  GeV, and  $m_{H^\pm} = 500$  GeV, and with the cascade decays of the squarks and gluinos calculated within the framework of the Minimal Supergravity scenario.
- 267 AKTAS 04D looked in  $77.8 \text{ pb}^{-1}$  of  $e^\pm p$  collisions at  $\sqrt{s} = 319$  GeV for resonant production of  $\tilde{q}$  by R-parity violating  $LQ\bar{D}$  couplings assuming that one of the  $\lambda'$  couplings dominates over all others. They consider final states with or without leptons and/or jets and/or  $\cancel{p}_T$  resulting from direct and indirect decays. They combine the channels to derive limits on  $\lambda'_{1j1}$  and  $\lambda'_{11k}$  as a function of the squark mass, see their Figs. 8 and 9, from a scan over the parameters  $70 < M_2 < 350$  GeV,  $-300 < \mu < 300$  GeV,  $\tan\beta = 6$ , for a fixed mass of 90 GeV for degenerate sleptons and an LSP mass  $> 30$  GeV. The quoted limits refer to  $\lambda' = 0.3$ , with  $U=u,c,t$  and  $D=d,s,b$ . Supersedes the results of ADLOFF 01B.
- 268 ADLOFF 03 looked for the s-channel production of squarks via  $\cancel{R} LQ\bar{D}$  couplings in  $117.2 \text{ pb}^{-1}$  of  $e^+p$  data at  $\sqrt{s} = 301$  and 319 GeV and of  $e^-p$  data at  $\sqrt{s} = 319$  GeV. The comparison of the data with the SM differential cross section allows limits to be set on couplings for processes mediated through contact interactions. They obtain lower bounds on the value of  $m_{\tilde{q}}/\lambda'$  of 710 GeV for the process  $e^+\bar{u} \rightarrow \tilde{d}^k$  (and charge conjugate), mediated by  $\lambda'_{11k}$ , and of 430 GeV for the process  $e^+d \rightarrow \tilde{u}^j$  (and charge conjugate), mediated by  $\lambda'_{1j1}$ .
- 269 CHEKANOV 03B used  $131.5 \text{ pb}^{-1}$  of  $e^+p$  and  $e^-p$  data taken at 300 and 318 GeV to look for narrow resonances in the  $eq$  or  $\nu q$  final states. Such final states may originate from  $LQ\bar{D}$  couplings with non-zero  $\lambda'_{1j1}$  (leading to  $\tilde{u}_j$ ) or  $\lambda'_{11k}$  (leading to  $\tilde{d}_k$ ). See their Fig. 8 and explanations in the text for limits. The quoted mass bound assumes that only direct squark decays contribute.
- 270 HEISTER 03G searches for the production of squarks in the case of  $\cancel{R}$  prompt decays with  $\cancel{UDD}$  direct couplings at  $\sqrt{s} = 189\text{--}209$  GeV.
- 271 ABAZOV 02F looked in  $77.5 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at 1.8 TeV for events with  $\geq 2\mu + \geq 4$  jets, originating from associated production of squarks followed by an indirect  $\cancel{R}$  decay (of the  $\tilde{\chi}_1^0$ ) via  $LQ\bar{D}$  couplings of the type  $\lambda'_{2jk}$  where  $j=1,2$  and  $k=1,2,3$ . Bounds are obtained in the MSUGRA scenario by a scan in the range  $0 \leq M_0 \leq 400$  GeV,  $60 \leq m_{1/2} \leq 120$  GeV for fixed values  $A_0=0$ ,  $\mu < 0$ , and  $\tan\beta=2$  or 6. The bounds are weaker for  $\tan\beta=6$ . See Figs. 2,3 for the exclusion contours in  $m_{1/2}$  versus  $m_0$  for  $\tan\beta=2$  and 6, respectively.
- 272 ABAZOV 02G search for associated production of gluinos and squarks in  $92.7 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.8$  TeV, using events with one electron,  $\geq 4$  jets, and large  $\cancel{E}_T$ . The results are compared to a MSUGRA scenario with  $\mu < 0$ ,  $A_0=0$ , and  $\tan\beta=3$  and allow to exclude a region of the  $(m_0, m_{1/2})$  shown in Fig. 11.
- 273 ABBIENDI 02 looked for events with an electron or neutrino and a jet in  $e^+e^-$  at 189 GeV. Squarks (or leptoquarks) could originate from a  $LQ\bar{D}$  coupling of an electron with a quark from the fluctuation of a virtual photon. Limits on the couplings  $\lambda'_{1jk}$  as a

- function of the squark mass are shown in Figs. 8–9, assuming that only direct squark decays contribute.
- 274 ABBIENDI 02B looked for events with an electron or neutrino and a jet in  $e^+e^-$  at 189–209 GeV. Squarks (or leptoquarks) could originate from a  $LQ\bar{D}$  coupling of an electron with a quark from the fluctuation of a virtual photon. Limits on the couplings  $\lambda'_{1jk}$  as a function of the squark mass are shown in Fig. 4, assuming that only direct squark decays contribute. The quoted limits are read off from Fig. 4. Supersedes the results of ABBIENDI 02.
- 275 ACHARD 02 searches for the production of squarks in the case of  $\tilde{R}$  prompt decays with  $\overline{UDD}$  couplings at  $\sqrt{s}=189\text{--}208$  GeV. The search is performed for direct and indirect decays, assuming one coupling at the time to be nonzero. The limit holds for indirect decays. Stronger limits are reached for  $(\tilde{u}_R, \tilde{d}_R)$  direct (80,56) GeV and  $(\tilde{u}_L, \tilde{d}_L)$  direct or indirect (87,86) GeV decays.
- 276 CHEKANOV 02 search for lepton flavor violating processes  $e^+p \rightarrow \ell X$ , where  $\ell = \mu$  or  $\tau$  with high  $p_T$ , in  $47.7 \text{ pb}^{-1}$  of  $e^+p$  collisions at 300 GeV. Such final states may originate from  $LQ\bar{D}$  couplings with simultaneously nonzero  $\lambda'_{1jk}$  and  $\lambda'_{ijk}$  ( $i=2$  or  $3$ ). The quoted mass bound assumes that only direct squark decays contribute.
- 277 BARATE 01B searches for the production of squarks in the case of  $\tilde{R}$  prompt decays with  $LL\bar{E}$  indirect or  $\overline{UDD}$  direct couplings at  $\sqrt{s}=189\text{--}202$  GeV. The limit holds for direct decays mediated by  $\tilde{R}$   $\overline{UDD}$  couplings. Limits are also given for  $LL\bar{E}$  indirect decays ( $m_{\tilde{u}_R} > 90$  GeV and  $m_{\tilde{d}_R} > 89$  GeV). Supersedes the results from BARATE 00H.
- 278 BREITWEG 01 searches for squark production in  $47.7 \text{ pb}^{-1}$  of  $e^+p$  collisions, mediated by  $\tilde{R}$  couplings  $LQ\bar{D}$  and leading to final states with  $\tilde{\nu}$  and  $\geq 1$  jet, complementing the  $e^+X$  final states of BREITWEG 00E. Limits are derived on  $\lambda'\sqrt{\beta}$ , where  $\beta$  is the branching fraction of the squarks into  $e^+q+\bar{\nu}q$ , as function of the squark mass, see their Fig. 15. The quoted mass limit assumes that only direct squark decays contribute.
- 279 ABBOTT 00C searched in  $\sim 94 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions for events with  $\mu\mu$ +jets, originating from associated production of leptoquarks. The results can be interpreted as limits on production of squarks followed by direct  $\tilde{R}$  decay via  $\lambda'_{2jk}L_2Q_jd_k^c$  couplings. Bounds are obtained on the cross section for branching ratios of 1 and of 1/2, see their Fig. 4. The former yields the limit on the  $\tilde{u}_L$ . The latter is combined with the bound of ABBOTT 99J from the  $\mu\nu$ +jets channel and of ABBOTT 98E and ABBOTT 98J from the  $\nu\nu$ +jets channel to yield the limit on  $\tilde{d}_R$ .
- 280 ACCIARRI 00P studied the effect on hadronic cross sections of  $t$ -channel down-type squark exchange via  $R$ -parity violating coupling  $\lambda'_{1jk}L_1Q_jd_k^c$ . The limit here refers to the case  $j=1,2$ , and holds for  $\lambda'_{1jk}=0.3$ . Data collected at  $\sqrt{s}=130\text{--}189$  GeV, superseding the results of ACCIARRI 98J.
- 281 AFFOLDER 00K searched in  $\sim 88 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions for events with 2–3 jets, at least one being  $b$ -tagged, large  $\cancel{E}_T$  and no high  $p_T$  leptons. Such  $\nu\nu+b$ -jets events would originate from associated production of squarks followed by direct  $\tilde{R}$  decay via  $\lambda'_{ij3}L_iQ_jd_3^c$  couplings. Bounds are obtained on the production cross section assuming zero branching ratio to charged leptons.
- 282 BARATE 00I studied the effect on hadronic cross sections and charge asymmetries of  $t$ -channel down-type squark exchange via  $R$ -parity violating coupling  $\lambda'_{1jk}L_1Q_jd_k^c$ . The limit here refers to the case  $j=1,2$ , and holds for  $\lambda'_{1jk}=0.3$ . A 50 GeV limit is found for up-type squarks with  $k=3$ . Data collected at  $\sqrt{s}=130\text{--}183$  GeV.
- 283 BREITWEG 00E searches for squark exchange in  $e^+p$  collisions, mediated by  $\tilde{R}$  couplings  $LQ\bar{D}$  and leading to final states with an identified  $e^+$  and  $\geq 1$  jet. The limit applies to up-type squarks of all generations, and assumes  $B(\tilde{q} \rightarrow qe)=1$ .

- 284 ABBOTT 99 searched for  $\gamma \cancel{E}_T + \geq 2$  jet final states, and set limits on  $\sigma(p\bar{p} \rightarrow \tilde{q} + X) \cdot B(\tilde{q} \rightarrow \gamma \cancel{E}_T X)$ . The quoted limits correspond to  $m_{\tilde{g}} \geq m_{\tilde{q}}$ , with  $B(\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 \gamma) = 1$  and  $B(\tilde{\chi}_1^0 \rightarrow \tilde{G} \gamma) = 1$ , respectively. They improve to 310 GeV (360 GeV in the case of  $\gamma \tilde{G}$  decay) for  $m_{\tilde{g}} = m_{\tilde{q}}$ .
- 285 ABBOTT 99K uses events with an electron pair and four jets to search for the decay of the  $\tilde{\chi}_1^0$  LSP via  $\cancel{R} LQ\bar{D}$  couplings. The particle spectrum and decay branching ratios are taken in the framework of minimal supergravity. An excluded region at 95% CL is obtained in the  $(m_0, m_{1/2})$  plane under the assumption that  $A_0 = 0$ ,  $\mu < 0$ ,  $\tan\beta = 2$  and any one of the couplings  $\lambda'_{1jk} > 10^{-3}$  ( $j=1,2$  and  $k=1,2,3$ ) and from which the above limit is computed. For equal mass squarks and gluinos, the corresponding limit is 277 GeV. The results are essentially independent of  $A_0$ , but the limit deteriorates rapidly with increasing  $\tan\beta$  or  $\mu > 0$ .
- 286 ABE 99M looked in  $107 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV for events with like sign dielectrons and two or more jets from the sequential decays  $\tilde{q} \rightarrow q\tilde{\chi}_1^0$  and  $\tilde{\chi}_1^0 \rightarrow e q \tilde{q}'$ , assuming  $\cancel{R}$  coupling  $L_1 Q_j D_k^c$ , with  $j=2,3$  and  $k=1,2,3$ . They assume five degenerate squark flavors,  $B(\tilde{q} \rightarrow q\tilde{\chi}_1^0) = 1$ ,  $B(\tilde{\chi}_1^0 \rightarrow e q \tilde{q}') = 0.25$  for both  $e^+$  and  $e^-$ , and  $m_{\tilde{g}} \geq 200$  GeV. The limit is obtained for  $m_{\tilde{\chi}_1^0} \geq m_{\tilde{q}}/2$  and improves for heavier gluinos or heavier  $\tilde{\chi}_1^0$ .
- 287 ABREU 99G looked for events with an electron or neutrino and a jet in  $e^+ e^-$  at 183 GeV. Squarks (or leptoquarks) could originate from a  $LQ\bar{D}$  coupling of an electron with a quark from the fluctuation of a virtual photon. Limits on the couplings  $\lambda'_{1jk}$  as a function of the squark mass are shown in Fig. 4, assuming that only direct squark decays contribute.
- 288 ABBOTT 98E searched in  $\sim 115 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions for events with  $e\nu$ +jets, originating from associated production of squarks followed by direct  $\cancel{R}$  decay via  $\lambda'_{1jk} L_1 Q_j d_k^c$  couplings. Bounds are obtained by combining these results with the previous bound of ABBOTT 97B from the  $ee$ +jets channel and with a reinterpretation of ABACHI 96B  $\nu\nu$ +jets channel.
- 289 ABE 98S looked in  $\sim 110 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV for events with  $\mu\mu$ +jets originating from associated production of squarks followed by direct  $\cancel{R}$  decay via  $\lambda'_{2jk} L_2 Q_j d_k^c$  couplings. Bounds are obtained on the production cross section times the square of the branching ratio, see Fig. 2. Mass limits result from the comparison with theoretical cross sections and branching ratio equal to 1 for  $\tilde{u}_L$  and 1/2 for  $\tilde{d}_R$ .
- 290 ACKERSTAFF 98V and ACCIARRI 98J studied the interference of  $t$ -channel squark ( $\tilde{d}_R$ ) exchange via  $R$ -parity violating  $\lambda'_{1jk} L_1 Q_j d_k^c$  coupling in  $e^+ e^- \rightarrow q\bar{q}$ . The limit is for  $\lambda'_{1jk} = 0.3$ . See paper for related limits on  $\tilde{u}_L$  exchange. Data collected at  $\sqrt{s} = 130\text{--}172$  GeV.
- 291 BREITWEG 98 used positron+jet events with missing energy and momentum to look for  $e^+ q \rightarrow \tilde{e}\tilde{q}$  via gaugino-like neutralino exchange with decays into  $(e\tilde{\chi}_1^0)(q\tilde{\chi}_1^0)$ . See paper for dependences in  $m_{\tilde{e}}, m_{\tilde{\chi}_1^0}$ .
- 292 DATTA 97 argues that the squark mass bound by ABACHI 95C can be weakened by 10–20 GeV if one relaxes the assumption of the universal scalar mass at the GUT-scale so that the  $\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$  in the squark cascade decays have dominant and invisible decays to  $\tilde{\nu}$ .
- 293 DERRICK 97 looked for lepton-number violating final states via  $R$ -parity violating couplings  $\lambda'_{ijk} L_i Q_j d_k$ . When  $\lambda'_{11k} \lambda'_{ijk} \neq 0$ , the process  $e u \rightarrow \tilde{d}_k^* \rightarrow \ell_i u_j$  is possible.



- When  $\lambda'_{1j1}\lambda'_{ijk} \neq 0$ , the process  $e\bar{d} \rightarrow \tilde{u}_j^* \rightarrow \ell_i\bar{d}_k$  is possible. 100% branching fraction  $\tilde{q} \rightarrow \ell j$  is assumed. The limit quoted here corresponds to  $\tilde{t} \rightarrow \tau q$  decay, with  $\lambda'=0.3$ . For different channels, limits are slightly better. See Table 6 in their paper.
- 294 HEWETT 97 reanalyzed the limits on possible resonances in di-jet mode ( $\tilde{q} \rightarrow q\tilde{g}$ ) from ALITTI 93 quoted in "Limits for Excited  $q$  ( $q^*$ ) from Single Production," ABE 96 in "SCALE LIMITS for Contact Interactions:  $\Lambda(qqqq)$ ," and unpublished CDF, DØ bounds. The bound applies to the gluino mass of 5 GeV, and improves for lighter gluino. The analysis has gluinos in parton distribution function.
- 295 TEREKHOV 97 improved the analysis of TEREKHOV 96 by including di-jet angular distributions in the analysis.
- 296 AID 96C used positron+jet events with missing energy and momentum to look for  $e^+ q \rightarrow \tilde{e}\tilde{q}$  via neutralino exchange with decays into  $(e\tilde{\chi}_1^0)(q\tilde{\chi}_1^0)$ . See the paper for dependences on  $m_{\tilde{e}}, m_{\tilde{\chi}_1^0}$ .
- 297 TEREKHOV 96 reanalyzed the limits on possible resonances in di-jet mode ( $\tilde{u} \rightarrow u\tilde{g}$ ) from ABE 95N quoted in "MASS LIMITS for  $g_A$  (axigluon)." The bound applies only to the case with a light gluino.
- 298 ABACHI 95C assume five degenerate squark flavors with  $m_{\tilde{q}_L} = m_{\tilde{q}_R}$ . Sleptons are assumed to be heavier than squarks. The limits are derived for fixed  $\tan\beta = 2.0$ ,  $\mu = -250$  GeV, and  $m_{H^+}=500$  GeV, and with the cascade decays of the squarks and gluinos calculated within the framework of the Minimal Supergravity scenario. The bounds are weakly sensitive to the three fixed parameters for a large fraction of parameter space. No limit is given for  $m_{\text{gluino}} > 547$  GeV.
- 299 ABE 95T looked for a cascade decay of five degenerate squarks into  $\tilde{\chi}_2^0$  which further decays into  $\tilde{\chi}_1^0$  and a photon. No signal is observed. Limits vary widely depending on the choice of parameters. For  $\mu = -40$  GeV,  $\tan\beta = 1.5$ , and heavy gluinos, the range  $50 < m_{\tilde{q}} \text{ (GeV)} < 110$  is excluded at 90% CL. See the paper for details.
- 300 ABE 92L assume five degenerate squark flavors and  $m_{\tilde{q}_L} = m_{\tilde{q}_R}$ . ABE 92L includes the effect of cascade decay, for a particular choice of parameters,  $\mu = -250$  GeV,  $\tan\beta = 2$ . Results are weakly sensitive to these parameters over much of parameter space. No limit for  $m_{\tilde{q}} \leq 50$  GeV (but other experiments rule out that region). Limits are 10–20 GeV higher if  $B(\tilde{q} \rightarrow q\tilde{\gamma}) = 1$ . Limit assumes GUT relations between gaugino masses and the gauge coupling; in particular that for  $|\mu|$  not small,  $m_{\tilde{\chi}_1^0} \approx m_{\tilde{g}}/6$ . This last relation implies that as  $m_{\tilde{g}}$  increases, the mass of  $\tilde{\chi}_1^0$  will eventually exceed  $m_{\tilde{q}}$  so that no decay is possible. Even before that occurs, the signal will disappear; in particular no bounds can be obtained for  $m_{\tilde{g}} > 410$  GeV.  $m_{H^+}=500$  GeV.
- 301 ROY 92 reanalyzed CDF limits on di-lepton events to obtain limits on squark production in  $R$ -parity violating models. The 100% decay  $\tilde{q} \rightarrow q\tilde{\chi}$  where  $\tilde{\chi}$  is the LSP, and the LSP decays either into  $\ell q\bar{d}$  or  $\ell\ell\bar{e}$  is assumed.
- 302 NOJIRI 91 argues that a heavy squark should be nearly degenerate with the gluino in minimal supergravity not to overclose the universe.

## Long-lived $\tilde{q}$ (Squark) MASS LIMIT

The following are bounds on long-lived scalar quarks, assumed to hadronise into hadrons with lifetime long enough to escape the detector prior to a possible decay. Limits may depend on the mixing angle of mass eigenstates:  $\tilde{q}_1 = \tilde{q}_L \cos\theta_q + \tilde{q}_R \sin\theta_q$ .

The coupling to the  $Z^0$  boson vanishes for up-type squarks when  $\theta_u=0.98$ , and for down type squarks when  $\theta_d=1.17$ .

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
-------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>95	95	303 HEISTER	03H ALEP	$\tilde{u}$
>92	95	303 HEISTER	03H ALEP	$\tilde{d}$
none 2–85	95	304 ABREU	98P DLPH	$\tilde{u}_L$
none 2–81	95	304 ABREU	98P DLPH	$\tilde{u}_R$
none 2–80	95	304 ABREU	98P DLPH	$\tilde{u}, \theta_u=0.98$
none 2–83	95	304 ABREU	98P DLPH	$\tilde{d}_L$
none 5–40	95	304 ABREU	98P DLPH	$\tilde{d}_R$
none 5–38	95	304 ABREU	98P DLPH	$\tilde{d}, \theta_d=1.17$

303 HEISTER 03H use  $e^+e^-$  data at and around the  $Z^0$  peak to look for hadronizing stable squarks. Combining their results on searches for charged and neutral R-hadrons with JANOT 03, a lower limit of 15.7 GeV on the mass is obtained. Combining this further with the results of searches for tracks with anomalous ionization in data from 183 to 208 GeV yields the quoted bounds.

304 ABREU 98P assumes that 40% of the squarks will hadronise into a charged hadron, and 60% into a neutral hadron which deposits most of its energy in hadron calorimeter. Data collected at  $\sqrt{s}=130\text{--}183$  GeV.

## $\tilde{b}$ (Sbottom) MASS LIMIT

Limits in  $e^+e^-$  depend on the mixing angle of the mass eigenstate  $\tilde{b}_1 = \tilde{b}_L \cos\theta_b + \tilde{b}_R \sin\theta_b$ . Coupling to the  $Z$  vanishes for  $\theta_b \sim 1.17$ . As a consequence, no absolute constraint in the mass region  $\lesssim 40$  GeV is available in the literature at this time from  $e^+e^-$  collisions. In the Listings below, we use  $\Delta m = m_{\tilde{b}_1} - m_{\tilde{\chi}_1^0}$ .

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>95	95	305 ACHARD	04 L3	$\tilde{b} \rightarrow b\tilde{\chi}_1^0, \theta_b=0, \Delta m > 15\text{--}25$ GeV
>81	95	305 ACHARD	04 L3	$\tilde{b} \rightarrow b\tilde{\chi}_1^0$ , all $\theta_b$ , $\Delta m > 15\text{--}25$ GeV
> 7.5	95	306 JANOT	04 THEO	unstable $\tilde{b}_1, e^+e^- \rightarrow$ hadrons
>93	95	307 ABDALLAH	03M DLPH	$\tilde{b} \rightarrow b\tilde{\chi}_1^0, \theta_b=0, \Delta m > 7$ GeV
>76	95	307 ABDALLAH	03M DLPH	$\tilde{b} \rightarrow b\tilde{\chi}_1^0$ , all $\theta_b, \Delta m > 7$ GeV
>85.1	95	308 ABBIENDI	02H OPAL	$\tilde{b} \rightarrow b\tilde{\chi}_1^0$ , all $\theta_b, \Delta m > 10$ GeV,
<b>&gt;89</b>	95	309 HEISTER	02K ALEP	CDF $\tilde{b} \rightarrow b\tilde{\chi}_1^0$ , all $\theta_b, \Delta m > 8$ GeV,
none 3.5–4.5	95	310 SAVINOV	01 CLEO	CDF $\tilde{B}$ meson
none 80–145		311 AFFOLDER	00D CDF	$\tilde{b} \rightarrow b\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} < 50$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>78	95	312 ABDALLAH	04M DLPH	$\tilde{R}, \tilde{b}_L$ , indirect, $\Delta m > 5$ GeV
none 50–82	95	313 ABDALLAH	03C DLPH	$\tilde{b} \rightarrow b\tilde{g}$ , stable $\tilde{g}$ , all $\theta_b$ , $\Delta m > 10$ GeV
		314 BERGER	03 THEO	
>71.5	95	315 HEISTER	03G ALEP	$\tilde{b}_L, \tilde{R}$ decay
>27.4	95	316 HEISTER	03H ALEP	$\tilde{b} \rightarrow b\tilde{g}$ , stable $\tilde{g}$ or $\tilde{b}$
>48	95	317 ACHARD	02 L3	$\tilde{b}_1, \tilde{R}$ decays
		318 BAEK	02 THEO	
		319 BECHER	02 THEO	
		320 CHEUNG	02B THEO	
		321 CHO	02 THEO	
		322 BERGER	01 THEO	$p\bar{p} \rightarrow X+b\text{-quark}$
none 52–115	95	323 ABBOTT	99F D0	$\tilde{b} \rightarrow b\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} < 20$ GeV

- 305 ACHARD 04 search for the production of  $\tilde{b}\tilde{b}$  in acoplanar b-tagged di-jet final states in the 192–209 GeV data. See Fig. 6 for the dependence of the limits on  $m_{\tilde{\chi}_1^0}$ . This limit supersedes ACCIARRI 99v.
- 306 JANOT 04 reanalyzes  $e^+e^- \rightarrow$  hadrons total cross section data with  $\sqrt{s} \in [20, 209]$  GeV from PEP, PETRA, TRISTAN, SLC, and LEP and constrains the mass of  $\tilde{b}_1$  assuming it decays quickly to hadrons.
- 307 ABDALLAH 03M looked for  $\tilde{b}$  pair production in events with acoplanar jets and  $\cancel{E}_T$  at  $\sqrt{s} = 189$ –208 GeV. The limit improves to 87 (98) GeV for all  $\theta_b$  ( $\theta_b = 0$ ) for  $\Delta m > 10$  GeV. See Fig. 24 and Table 11 for other choices of  $\Delta m$ . These limits include and update the results of ABREU,P 00D.
- 308 ABBIENDI 02H search for events with two acoplanar jets and  $\cancel{p}_T$  in the 161–209 GeV data. The limit assumes 100% branching ratio and uses the exclusion at large  $\Delta m$  from CDF (AFFOLDER 00D). For  $\theta_b=0$ , the bound improves to  $> 96.9$  GeV. See Fig. 4 and Table 6 for the more general dependence on the limits on  $\Delta m$ . These results supersede ABBIENDI 99M.
- 309 HEISTER 02K search for bottom squarks in final states with acoplanar jets with  $b$  tagging, using 183–209 GeV data. The mass bound uses the CDF results from AFFOLDER 00D. See Fig. 5 for the more general dependence of the limits on  $\Delta m$ . Updates BARATE 01.
- 310 SAVINOV 01 use data taken at  $\sqrt{s}=10.52$  GeV, below the  $B\bar{B}$  threshold. They look for events with a pair of leptons with opposite charge and a fully reconstructed hadronic  $D$  or  $D^*$  decay. These could originate from production of a light-sbottom hadron followed by  $\tilde{B} \rightarrow D^{(*)} \ell^- \tilde{\nu}$ , in case the  $\tilde{\nu}$  is the LSP, or  $\tilde{B} \rightarrow D^{(*)} \pi \ell^-$ , in case of  $\cancel{R}$ . The mass range  $3.5 \leq M(\tilde{B}) \leq 4.5$  GeV was explored, assuming 100% branching ratio for either of the decays. In the  $\tilde{\nu}$  LSP scenario, the limit holds only for  $M(\tilde{\nu})$  less than about 1 GeV and for the  $D^*$  decays it is reduced to the range 3.9–4.5 GeV. For the  $\cancel{R}$  decay, the whole range is excluded.
- 311 AFFOLDER 00D search for final states with 2 or 3 jets and  $\cancel{E}_T$ , one jet with a  $b$  tag. See their Fig. 3 for the mass exclusion in the  $m_{\tilde{t}}, m_{\tilde{\chi}_1^0}$  plane.
- 312 ABDALLAH 04M use data from  $\sqrt{s} = 192$ –208 GeV to derive limits on sparticle masses under the assumption of  $\cancel{R}$  with  $\overline{UDD}$  couplings. The results are valid for  $\mu = -200$  GeV,  $\tan\beta = 1.5$ ,  $\Delta m > 5$  GeV and assuming a BR of 1 for the given decay. The limit quoted is for indirect  $\overline{UDD}$  decays using the neutralino constraint of 38.0 GeV, also derived in ABDALLAH 04M, and assumes no mixing. For indirect decays it remains at 78 GeV when the neutralino constraint is not used. Supersedes the result of ABREU 01D.
- 313 ABDALLAH 03C looked for events of the type  $q\bar{q}R^\pm R^\pm$ ,  $q\bar{q}R^\pm R^0$  or  $q\bar{q}R^0 R^0$  in  $e^+e^-$  interactions at  $\sqrt{s} = 189$ –208 GeV. The  $R^\pm$  bound states are identified by anomalous  $dE/dx$  in the tracking chambers and the  $R^0$  by missing energy due to their reduced energy loss in the calorimeters. Excluded mass regions in the  $(m(\tilde{b}), m(\tilde{g}))$  plane for  $m(\tilde{g}) > 2$  GeV are obtained for several values of the probability for the gluino to fragment into  $R^\pm$  or  $R^0$ , as shown in their Fig. 19. The limit improves to 94 GeV for  $\theta_b = 0$ .
- 314 BERGER 03 studies the constraints on a  $\tilde{b}_1$  with mass in the 2.2–5.5 GeV region coming from radiative decays of  $\Upsilon(nS)$  into sbottomonium. The constraints apply only if  $\tilde{b}_1$  lives long enough to permit formation of the sbottomonium bound state. A small region of mass in the  $m_{\tilde{b}_1} - m_{\tilde{g}}$  plane survives current experimental constraints from CLEO.
- 315 HEISTER 03G searches for the production of  $\tilde{b}$  pairs in the case of  $\cancel{R}$  prompt decays with  $L\bar{L}\bar{E}$ ,  $LQ\bar{D}$  or  $\overline{UDD}$  couplings at  $\sqrt{s} = 189$ –209 GeV. The limit holds for indirect decays mediated by  $\cancel{R}$   $\overline{UDD}$  couplings. It improves to 90 GeV for indirect decays mediated by  $\cancel{R}$   $L\bar{L}\bar{E}$  couplings and to 80 GeV for indirect decays mediated by  $\cancel{R}$   $LQ\bar{D}$  couplings. Supersedes the results from BARATE 01B.
- 316 HEISTER 03H use their results on bounds on stable squarks, on stable gluinos and on squarks decaying to a stable gluino from the same paper to derive a mass limit on  $\tilde{b}$ , see their Fig. 13. The limit for a long-lived  $\tilde{b}_1$  is 92 GeV.

- 317 **ACHARD 02** searches for the production of squarks in the case of  $\tilde{R}$  prompt decays with  $UDD$  couplings at  $\sqrt{s}=189\text{--}208$  GeV. The search is performed for direct and indirect decays, assuming one coupling at the time to be nonzero. The limit is computed for the minimal cross section and holds for indirect decays and reaches 55 GeV for direct decays.
- 318 **BAEK 02** studies the constraints on a  $\tilde{b}_1$  with mass in the 2.2–5.5 GeV region coming from precision measurements of  $Z^0$  decays. It is noted that  $CP$ -violating couplings in the MSSM parameters relax the strong constraints otherwise derived from  $CP$  conservation.
- 319 **BECHER 02** studies the constraints on a  $\tilde{b}_1$  with mass in the 2.2–5.5 GeV region coming from radiative  $B$  meson decays, and sets limits on the off-diagonal flavor-changing couplings  $q\tilde{b}\tilde{g}$  ( $q=d,s$ ).
- 320 **CHEUNG 02B** studies the constraints on a  $\tilde{b}_1$  with mass in the 2.2–5.5 GeV region and a gluino in the mass range 12–16 GeV, using precision measurements of  $Z^0$  decays and  $e^+e^-$  annihilations at LEP2. Few detectable events are predicted in the LEP2 data for the model proposed by BERGER 01.
- 321 **CHO 02** studies the constraints on a  $\tilde{b}_1$  with mass in the 2.2–5.5 GeV region coming from precision measurements of  $Z^0$  decays. Strong constraints are obtained for  $CP$ -conserving MSSM couplings.
- 322 **BERGER 01** reanalyzed interpretation of Tevatron data on bottom-quark production. Argues that pair production of light gluinos ( $m\sim 12\text{--}16$  GeV) with subsequent 2-body decay into a light sbottom ( $m\sim 2\text{--}5.5$  GeV) and bottom can reconcile Tevatron data with predictions of perturbative QCD for the bottom production rate. The sbottom must either decay hadronically via a  $R$ -parity- and  $B$ -violating interaction, or be long-lived. Constraints on the mass spectrum are derived from the measurements of time-averaged  $B^0\text{--}\bar{B}^0$  mixing.
- 323 **ABBOTT 99F** looked for events with two jets, with or without an associated muon from  $b$  decay, and  $\cancel{E}_T$ . See Fig. 2 for the dependence of the limit on  $m_{\tilde{\chi}_1^0}$ . No limit for  $m_{\tilde{\chi}_1^0} > 47$  GeV.

## $\tilde{t}$ (Stop) MASS LIMIT

Limits depend on the decay mode. In  $e^+e^-$  collisions they also depend on the mixing angle of the mass eigenstate  $\tilde{t}_1 = \tilde{t}_L \cos\theta_t + \tilde{t}_R \sin\theta_t$ . The coupling to the  $Z$  vanishes when  $\theta_t = 0.98$ . In the Listings below, we use  $\Delta m \equiv m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0}$  or  $\Delta m \equiv m_{\tilde{t}_1} - m_{\tilde{\nu}}$ , depending on relevant decay mode. See also bounds in “ $\tilde{q}$  (Squark)

MASS LIMIT.” Limits made obsolete by the most recent analyses of  $e^+e^-$  and  $p\bar{p}$  collisions can be found in previous Editions of this *Review*.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
none 80–120	95	324 ABAZOV	04 D0	$\tilde{t} \rightarrow b\ell\nu\tilde{\chi}^0$ , $m_{\tilde{\chi}^0} = 50$ GeV
> 90	95	325 ACHARD	04 L3	$\tilde{t} \rightarrow c\tilde{\chi}_1^0$ , all $\theta_t$ , $\Delta m > 15\text{--}25$ GeV
> 93	95	325 ACHARD	04 L3	$\tilde{b} \rightarrow b\ell\tilde{\nu}$ , all $\theta_t$ , $\Delta m > 15$ GeV
> 88	95	325 ACHARD	04 L3	$\tilde{b} \rightarrow b\tau\tilde{\nu}$ , all $\theta_t$ , $\Delta m > 15$ GeV
> 75	95	326 ABDALLAH	03M DLPH	$\tilde{t} \rightarrow c\tilde{\chi}^0$ , $\theta_t=0$ , $\Delta m > 2$ GeV
> 71	95	326 ABDALLAH	03M DLPH	$\tilde{t} \rightarrow c\tilde{\chi}^0$ , all $\theta_t$ , $\Delta m > 2$ GeV
> 96	95	326 ABDALLAH	03M DLPH	$\tilde{t} \rightarrow c\tilde{\chi}^0$ , $\theta_t=0$ , $\Delta m > 10$ GeV
> 92	95	326 ABDALLAH	03M DLPH	$\tilde{t} \rightarrow c\tilde{\chi}^0$ , all $\theta_t$ , $\Delta m > 10$ GeV
none 80–131	95	327 ACOSTA	03C CDF	$\tilde{t} \rightarrow b\ell\tilde{\nu}$ , $m_{\tilde{\nu}} \leq 63$ GeV

>144	95	328	ABAZOV	02C D0	$\tilde{t} \rightarrow b\ell\tilde{\nu}$ , $m_{\tilde{\nu}}=45$ GeV
<b>&gt; 95.7</b>	95	329	ABBIENDI	02H OPAL	$c\tilde{\chi}_1^0$ , all $\theta_t$ , $\Delta m > 10$ GeV
<b>&gt; 92.6</b>	95	329	ABBIENDI	02H OPAL	$b\ell\tilde{\nu}$ , all $\theta_t$ , $\Delta m > 10$ GeV
> 91.5	95	329	ABBIENDI	02H OPAL	$b\tau\tilde{\nu}$ , all $\theta_t$ , $\Delta m > 10$ GeV
> 63	95	330	HEISTER	02K ALEP	any decay, any lifetime, all $\theta_t$
> 92	95	330	HEISTER	02K ALEP	$\tilde{t} \rightarrow c\tilde{\chi}_1^0$ , all $\theta_t$ , $\Delta m > 8$ GeV,
> 97	95	330	HEISTER	02K ALEP	CDF $\tilde{t} \rightarrow b\ell\tilde{\nu}$ , all $\theta_t$ , $\Delta m > 8$ GeV,
> 78	95	330	HEISTER	02K ALEP	DØ $\tilde{t} \rightarrow b\tilde{\chi}_1^0 W^*$ , all $\theta_t$ , $\Delta m > 8$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
none 80–122	95	331	ABAZOV	04B D0	$\tilde{t} \rightarrow c\tilde{\chi}_1^0$ , $m_{\tilde{\chi}_1^0} < 45$ GeV
> 77	95	332	ABBIENDI	04F OPAL	$\tilde{R}$ , direct, all $\theta_t$
> 77	95	333	ABDALLAH	04M DLPH	$\tilde{R}$ , indirect, all $\theta_t$ , $\Delta m > 5$ GeV
>122	95	334	ACOSTA	04B CDF	$\tilde{R}$ , direct, all $\theta_t$
> 74.5		335	AKTAS	04B H1	$\tilde{R}$ , $\tilde{t}_1$
		336	DAS	04 THEO	$\tilde{t}\tilde{t} \rightarrow b\ell\nu_\ell\chi^0\bar{b}q\bar{q}'\chi^0$ , $m_{\tilde{\chi}_1^0}$ = 15 GeV, no $\tilde{t} \rightarrow c\chi^0$
none 50–87	95	337	ABDALLAH	03C DLPH	$\tilde{t} \rightarrow c\tilde{g}$ , stable $\tilde{g}$ , all $\theta_t$ , $\Delta M > 10$ GeV
> 71.5	95	338	CHAKRAB...	03 THEO	$p\bar{p} \rightarrow \tilde{t}\tilde{t}^*$ , RPV
> 80	95	339	HEISTER	03G ALEP	$\tilde{t}_L, \tilde{R}$ decay
> 77	95	340	HEISTER	03H ALEP	$\tilde{t} \rightarrow c\tilde{g}$ , stable $\tilde{g}$ or $\tilde{t}$ , all $\theta_t$ , all $\Delta M$
> 61	95	341	ACHARD	02 L3	$\tilde{t}_1, \tilde{R}$ decays
		342	AFFOLDER	01B CDF	$t \rightarrow \tilde{t}\chi_1^0$
	95	343	ABREU	00i DLPH	$\tilde{R} (L\bar{L}\bar{E})$ , $\theta_t=0.98$ , $\Delta m > 4$ GeV
none 68–119	95	344	AFFOLDER	00D CDF	$\tilde{t} \rightarrow c\tilde{\chi}_1^0$ , $m_{\tilde{\chi}_1^0} < 40$ GeV
none 84–120	95	345	AFFOLDER	00G CDF	$\tilde{t}_1 \rightarrow b\ell\tilde{\nu}$ , $m_{\tilde{\nu}} < 45$
> 59	95	346	BARATE	00P ALEP	Repl. by HEISTER 02K
>120	95	347	ABE	99M CDF	$p\bar{p} \rightarrow \tilde{t}_1\tilde{t}_1, \tilde{R}$
none 61–91	95	348	ABACHI	96B D0	$\tilde{t} \rightarrow c\tilde{\chi}_1^0$ , $m_{\tilde{\chi}_1^0} < 30$ GeV
none 9–24.4	95	349	AID	96 H1	$e p \rightarrow \tilde{t}\tilde{t}, \tilde{R}$ decays
>138	95	350	AID	96 H1	$e p \rightarrow \tilde{t}, \tilde{R}$ , $\lambda\cos\theta_t > 0.03$
> 45		351	CHO	96 RVUE	$B^0\text{-}\bar{B}^0$ and $\epsilon$ , $\theta_t=0.98$ , $\tan\beta < 2$
none 11–41	95	352	BUSKULIC	95E ALEP	$\tilde{R} (L\bar{L}\bar{E})$ , $\theta_t=0.98$
none 6.0–41.2	95		AKERS	94K OPAL	$\tilde{t} \rightarrow c\tilde{\chi}_1^0$ , $\theta_t=0$ , $\Delta m > 2$ GeV
none 5.0–46.0	95		AKERS	94K OPAL	$\tilde{t} \rightarrow c\tilde{\chi}_1^0$ , $\theta_t=0$ , $\Delta m > 5$ GeV
none 11.2–25.5	95		AKERS	94K OPAL	$\tilde{t} \rightarrow c\tilde{\chi}_1^0$ , $\theta_t=0.98$ , $\Delta m > 2$ GeV
none 7.9–41.2	95		AKERS	94K OPAL	$\tilde{t} \rightarrow c\tilde{\chi}_1^0$ , $\theta_t=0.98$ , $\Delta m > 5$ GeV
none 7.6–28.0	95	353	SHIRAI	94 VNS	$\tilde{t} \rightarrow c\tilde{\chi}_1^0$ , any $\theta_t$ , $\Delta m > 10$ GeV
none 10–20	95	353	SHIRAI	94 VNS	$\tilde{t} \rightarrow c\tilde{\chi}_1^0$ , any $\theta_t$ , $\Delta m > 2.5$ GeV

- 324 ABAZOV 04 looked at  $108.3\text{pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV for events with  $e+\mu+\cancel{E}_T$  as signature for the 3- and 4-body decays of stop into  $b\ell\nu\tilde{\chi}^0$  final states. For the  $b\ell\tilde{\nu}$  channel they use the results from ABAZOV 02C. No significant excess is observed compared to the Standard Model expectation and limits are derived on the mass of  $\tilde{t}_1$  for the 3- and 4-body decays in the  $(m_{\tilde{t}}, m_{\tilde{\chi}^0})$  plane, see their Figure 4.
- 325 ACHARD 04 search in the 192–209 GeV data for the production of  $\tilde{t}\tilde{t}$  in acoplanar di-jet final states and, in case of  $b\ell\tilde{\nu}$  ( $b\tau\tilde{\nu}$ ) final states, two leptons (taus). The limits for  $\theta_t=0$  improve to 95, 96 and 93 GeV, respectively. All limits assume 100% branching ratio for the respective decay modes. See Fig. 6 for the dependence of the limits on  $m_{\tilde{\chi}_1^0}$ . These limits supersede ACCIARRI 99V.
- 326 ABDALLAH 03M looked for  $\tilde{t}$  pair production in events with acoplanar jets and  $\cancel{E}$  at  $\sqrt{s} = 189\text{--}208$  GeV. See Fig. 23 and Table 11 for other choices of  $\Delta m$ . These limits include and update the results of ABREU,P 00D.
- 327 ACOSTA 03C searched in  $107\text{pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.8$  TeV for pair production of  $\tilde{t}$  followed by the decay  $\tilde{t} \rightarrow b\ell\tilde{\nu}$ . They looked for events with two isolated leptons ( $e$  or  $\mu$ ), at least one jet and  $\cancel{E}_T$ . The excluded mass range is reduced for larger  $m_{\tilde{\nu}}$ , and no limit is set for  $m_{\tilde{\nu}} > 88.4$  GeV (see Fig. 2).
- 328 ABAZOV 02C looked in  $108.3\text{pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.8$  TeV for events with  $e\mu\cancel{E}_T$ , originating from associated production  $\tilde{t}\tilde{t}$ . Branching ratios are assumed to be 100%. The bound for the  $b\ell\tilde{\nu}$  decay weakens for large  $\tilde{\nu}$  mass (see Fig. 3), and no limit is set when  $m_{\tilde{\nu}} > 85$  GeV. See Fig. 4 for the limits in case of decays to a real  $\tilde{\chi}_1^\pm$ , followed by  $\tilde{\chi}_1^\pm \rightarrow \ell\tilde{\nu}$ , as a function of  $m_{\tilde{\chi}_1^\pm}$ .
- 329 ABBIENDI 02H looked for events with two acoplanar jets,  $\cancel{p}_T$ , and, in the case of  $b\ell\tilde{\nu}$  final states, two leptons, in the 161–209 GeV data. The bound for  $c\tilde{\chi}_1^0$  applies to the region where  $\Delta m < m_W + m_b$ , else the decay  $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^0 W^+$  becomes dominant. The limit for  $b\ell\tilde{\nu}$  assumes equal branching ratios for the three lepton flavors and for  $b\tau\tilde{\nu}$  100% for this channel. For  $\theta_t=0$ , the bounds improve to  $> 97.6$  GeV ( $c\tilde{\chi}_1^0$ ),  $> 96.0$  GeV ( $b\ell\tilde{\nu}$ ), and  $> 95.5$  ( $b\tau\tilde{\nu}$ ). See Figs. 5–6 and Table 5 for the more general dependence of the limits on  $\Delta m$ . These results supersede ABBIENDI 99M.
- 330 HEISTER 02K search for top squarks in final states with jets (with/without  $b$  tagging or leptons) or long-lived hadrons, using 183–209 GeV data. The absolute mass bound is obtained by varying the branching ratio of  $\tilde{t} \rightarrow c\tilde{\chi}_1^0$  and the lepton fraction in  $\tilde{t} \rightarrow b\tilde{\chi}_1^0 f\bar{f}'$  decays. The mass bound for  $\tilde{t} \rightarrow c\tilde{\chi}_1^0$  uses the CDF results from AFFOLDER 00D and for  $\tilde{t} \rightarrow b\ell\tilde{\nu}$  the DØ results from ABAZOV 02C. See Figs. 2–5 for the more general dependence of the limits on  $\Delta m$ . Updates BARATE 01 and BARATE 00P.
- 331 ABAZOV 04B looked in  $85.2\text{pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV for events with at least two acoplanar jets and  $\cancel{E}_T$ . No significant excess is observed compared to the Standard Model expectation and a limit is derived on the production of  $\tilde{t}_1$ , see their Figure 2 for the limit in the  $(m_{\tilde{t}}, m_{\tilde{\chi}^0})$  plane. No limit can be obtained for  $m_{\tilde{\chi}^0} > 52$  GeV.
- 332 ABBIENDI 04F use data from  $\sqrt{s} = 189\text{--}209$  GeV. They derive limits on the stop mass under the assumption of  $\mathcal{R}$  with  $LQ\bar{D}$  or  $\overline{UD}\bar{D}$  couplings. The limit quoted applies to direct decays with  $\overline{UD}\bar{D}$  couplings when the stop decouples from the  $Z^0$  and improves to 88 GeV for  $\theta_t = 0$ . For  $LQ\bar{D}$  couplings, the limit improves to 98 (100) GeV for  $\lambda'_{13k}$  or  $\lambda'_{23k}$  couplings and all  $\theta_t$  ( $\theta_t = 0$ ). For  $\lambda'_{33k}$  couplings it is 96 (98) GeV for all  $\theta_t$  ( $\theta_t = 0$ ). Supersedes the results of ABBIENDI 00.

- 333 ABDALLAH 04M use data from  $\sqrt{s} = 192\text{--}208$  GeV to derive limits on sparticle masses under the assumption of  $\tilde{R}$  with  $LL\bar{E}$  or  $\overline{UDD}$  couplings. The results are valid for  $\mu = -200$  GeV,  $\tan\beta = 1.5$ ,  $\Delta m > 5$  GeV and assuming a BR of 1 for the given decay. The limit quoted is for decoupling of the stop from the  $Z^0$  and indirect  $\overline{UDD}$  decays using the neutralino constraint of 39.5 GeV for  $LL\bar{E}$  and of 38.0 GeV for  $\overline{UDD}$  couplings, also derived in ABDALLAH 04M. For no mixing (decoupling) and indirect decays via  $LL\bar{E}$  the limit improves to 92 (87) GeV if the constraint from the neutralino is used and to 88 (81) GeV if it is not used. For indirect decays via  $\overline{UDD}$  couplings it improves to 87 GeV for no mixing and using the constraint from the neutralino, whereas it becomes 81 GeV (67) GeV for no mixing (decoupling) if the neutralino constraint is not used. Supersedes the result of ABREU 01D.
- 334 ACOSTA 04B looked in  $106\text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV for R-parity violating decays of  $\tilde{t}_1$  with  $LQ\bar{D}$  couplings. They search for events of the type  $\tilde{t}_1\bar{\tilde{t}}_1 \rightarrow \ell\tau_h jj$  where  $\ell = e, \mu$  originates from a leptonic  $\tau$  decay and  $\tau_h$  represents a hadronic decay of  $\tau$ . They derive limits on the stop mass for direct decays after combining the results from  $e$  and  $\mu$  and under the assumption that BR = 1 for the decay to  $\tau$ .
- 335 AKTAS 04B looked in  $106\text{ pb}^{-1}$  of  $e^\pm p$  collisions at  $\sqrt{s} = 319$  GeV and 301 GeV for resonant production of  $\tilde{t}_1$  by R-parity violating  $LQ\bar{D}$  couplings with  $\lambda'_{131}$ , others being zero. They consider the decays  $\tilde{t}_1 \rightarrow e^+ d$  and  $\tilde{t}_1 \rightarrow W\bar{b}$  followed by  $\bar{b} \rightarrow \bar{\nu}_e d$  and assume gauginos too heavy to participate in the decays. They combine the channels  $j\ell\nu\cancel{T}$ ,  $j\mu\nu\cancel{T}$ ,  $jjj\nu\cancel{T}$  to derive limits in the plane  $(m_{\tilde{t}}, \lambda'_{131})$ , see their Fig. 5.
- 336 DAS 04 reanalyzes AFFOLDER 00G data and obtains constraints on  $m_{\tilde{t}_1}$  as a function of  $B(\tilde{t} \rightarrow b\ell\nu\chi^0) \times B(\tilde{t} \rightarrow b\bar{q}q'\chi^0)$ ,  $B(\tilde{t} \rightarrow c\chi^0)$  and  $m_{\chi^0}$ . Bound weakens for larger  $B(\tilde{t} \rightarrow c\chi^0)$  and  $m_{\chi^0}$ .
- 337 ABDALLAH 03C looked for events of the type  $q\bar{q}R^\pm R^\pm$ ,  $q\bar{q}R^\pm R^0$  or  $q\bar{q}R^0 R^0$  in  $e^+e^-$  interactions at  $\sqrt{s} = 189\text{--}208$  GeV. The  $R^\pm$  bound states are identified by anomalous  $dE/dx$  in the tracking chambers and the  $R^0$  by missing energy, due to their reduced energy loss in the calorimeters. Excluded mass regions in the  $(m(\tilde{t}), m(\tilde{g}))$  plane for  $m(\tilde{g}) > 2$  GeV are obtained for several values of the probability for the gluino to fragment into  $R^\pm$  or  $R^0$ , as shown in their Fig. 18. The limit improves to 90 GeV for  $\theta_t = 0$ .
- 338 Theoretical analysis of  $e^+e^- + 2$  jet final states from the RPV decay of  $\tilde{t}\tilde{t}^*$  pairs produced in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8$  TeV. 95%CL limits of 220 (165) GeV are derived for  $B(\tilde{t} \rightarrow eq) = 1$  (0.5).
- 339 HEISTER 03G searches for the production of  $\tilde{t}$  pairs in the case of  $\tilde{R}$  prompt decays with  $LL\bar{E}$ ,  $LQ\bar{D}$  or  $\overline{UDD}$  couplings at  $\sqrt{s} = 189\text{--}209$  GeV. The limit holds for indirect decays mediated by  $\tilde{R}$   $\overline{UDD}$  couplings. It improves to 91 GeV for indirect decays mediated by  $\tilde{R}$   $LL\bar{E}$  couplings, to 97 GeV for direct (assuming  $B(\tilde{t}_L \rightarrow q\tau) = 100\%$ ) and to 85 GeV for indirect decays mediated by  $\tilde{R}$   $LQ\bar{D}$  couplings. Supersedes the results from BARATE 01B.
- 340 HEISTER 03H use  $e^+e^-$  data from 183–208 GeV to look for the production of stop decaying into a c quark and a stable gluino hadronizing into charged or neutral R-hadrons. Combining these results with bounds on stable squarks and on a stable gluino LSP from the same paper yields the quoted limit. See their Fig. 13 for the dependence of the mass limit on the gluino mass and on  $\theta_t$ .
- 341 ACHARD 02 searches for the production of squarks in the case of  $\tilde{R}$  prompt decays with  $\overline{UDD}$  couplings at  $\sqrt{s} = 189\text{--}208$  GeV. The search is performed for direct and indirect decays, assuming one coupling at the time to be nonzero. The limit is computed for the minimal cross section and holds for both direct and indirect decays.
- 342 AFFOLDER 01B searches for decays of the top quark into stop and LSP, in  $t\bar{t}$  events. Limits on the stop mass as a function of the LSP mass and of the decay branching ratio are shown in Fig. 3. They exclude branching ratios in excess of 45% for SLP masses up to 40 GeV.

- 343 ABREU 00I searches for the production of stop in the case of  $R$ -parity violation with  $LL\bar{E}$  couplings, for which only indirect decays are allowed. They investigate topologies with jets plus leptons in data from  $\sqrt{s}=183$  GeV. The lower bound on the stop mass assumes a neutralino mass limit of 27 GeV, also derived in ABREU 00I.
- 344 AFFOLDER 00D search for final states with 2 or 3 jets and  $\cancel{E}_T$ , one jet with a  $c$  tag. See their Fig. 2 for the mass exclusion in the  $(m_{\tilde{t}}, m_{\tilde{\chi}_1^0})$  plane. The maximum excluded  $m_{\tilde{t}}$  value is 119 GeV, for  $m_{\tilde{\chi}_1^0}=40$  GeV.
- 345 AFFOLDER 00G searches for  $\tilde{t}_1 \tilde{t}_1^*$  production, with  $\tilde{t}_1 \rightarrow b l \tilde{\nu}$ , leading to topologies with  $\geq 1$  isolated lepton ( $e$  or  $\mu$ ),  $\cancel{E}_T$ , and  $\geq 2$  jets with  $\geq 1$  tagged as  $b$  quark by a secondary vertex. See Fig. 4 for the excluded mass range as a function of  $m_{\tilde{\nu}}$ . Cross-section limits for  $\tilde{t}_1 \tilde{t}_1^*$ , with  $\tilde{t}_1 \rightarrow b \chi_1^\pm$  ( $\chi_1^\pm \rightarrow \ell^\pm \nu \tilde{\chi}_1^0$ ), are given in Fig. 2.
- 346 BARATE 00P use data from  $\sqrt{s}=189\text{--}202$  GeV to explore the region of small mass difference between the stop and the neutralino by searching heavy stable charged particles or tracks with large impact parameters. For prompt decays, they make use of acoplanar jets from BARATE 99Q, updated up to 202 GeV. The limit is reached for  $\Delta m=1.6$  GeV and a decay length of 1 cm. If the MSSM relation between the decay width and  $\Delta m$  is used, the limit improves to 63 GeV. It is set for  $\Delta m=1.9$  GeV,  $\tan\beta=2.6$ , and  $\theta_{\tilde{t}}=0.98$ , and large negative  $\mu$ .
- 347 ABE 99M looked in  $107 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.8$  TeV for events with like sign dielectrons and two or more jets from the sequential decays  $\tilde{q} \rightarrow q \tilde{\chi}_1^0$  and  $\tilde{\chi}_1^0 \rightarrow e q \bar{q}'$ , assuming  $\mathcal{R}$  coupling  $L_1 Q_j D_k^c$ , with  $j=2,3$  and  $k=1,2,3$ . They assume  $B(\tilde{t}_1 \rightarrow c \tilde{\chi}_1^0)=1$ ,  $B(\tilde{\chi}_1^0 \rightarrow e q \bar{q}')=0.25$  for both  $e^+$  and  $e^-$ , and  $m_{\tilde{\chi}_1^0} \geq m_{\tilde{t}_1}/2$ . The limit improves for heavier  $\tilde{\chi}_1^0$ .
- 348 ABACHI 96B searches for final states with 2 jets and missing  $E_T$ . Limits on  $m_{\tilde{t}}$  are given as a function of  $m_{\tilde{\chi}_1^0}$ . See Fig. 4 for details.
- 349 AID 96 considers photoproduction of  $\tilde{t}\tilde{t}$  pairs, with 100%  $R$ -parity violating decays of  $\tilde{t}$  to  $e q$ , with  $q=d, s$ , or  $b$  quarks.
- 350 AID 96 considers production and decay of  $\tilde{t}$  via the  $R$ -parity violating coupling  $\lambda' L_1 Q_3 d_1^c$ .
- 351 CHO 96 studied the consistency among the  $B^0\text{--}\bar{B}^0$  mixing,  $\epsilon$  in  $K^0\text{--}\bar{K}^0$  mixing, and the measurements of  $V_{cb}$ ,  $V_{ub}/V_{cb}$ . For the range  $25.5 \text{ GeV} < m_{\tilde{t}_1} < m_Z/2$  left by AKERS 94K for  $\theta_{\tilde{t}}=0.98$ , and within the allowed range in  $M_2\text{--}\mu$  parameter space from chargino, neutralino searches by ACCIARRI 95E, they found the scalar top contribution to  $B^0\text{--}\bar{B}^0$  mixing and  $\epsilon$  to be too large if  $\tan\beta < 2$ . For more on their assumptions, see the paper and their reference 10.
- 352 BUSKULIC 95E looked for  $Z \rightarrow \tilde{t}\tilde{t}$ , where  $\tilde{t} \rightarrow c \chi_1^0$  and  $\chi_1^0$  decays via  $R$ -parity violating interactions into two leptons and a neutrino.
- 353 SHIRAI 94 bound assumes the cross section without the  $s$ -channel  $Z$ -exchange and the QCD correction, underestimating the cross section up to 20% and 30%, respectively. They assume  $m_c=1.5$  GeV.

## Heavy $\tilde{g}$ (Gluino) MASS LIMIT

For  $m_{\tilde{g}} > 60\text{--}70$  GeV, it is expected that gluinos would undergo a cascade decay via a number of neutralinos and/or charginos rather than undergo a direct decay to photinos as assumed by some papers. Limits obtained when direct decay is assumed are usually higher than limits when cascade decays are included. Limits made obsolete by the most recent analyses of  $p\bar{p}$  collisions can be found in previous Editions of this *Review*.



VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>195	95	354 AFFOLDER	02 CDF	Jets+ $\cancel{E}_T$ , any $m_{\tilde{q}}$
>300	95	354 AFFOLDER	02 CDF	Jets+ $\cancel{E}_T$ , $m_{\tilde{q}}=m_{\tilde{g}}$
>129	95	355 ABBOTT	01D D0	$ll$ +jets+ $\cancel{E}_T$ , $\tan\beta < 10$ , $m_0 < 300$ GeV, $\mu < 0$ , $A_0=0$
>175	95	355 ABBOTT	01D D0	$ll$ +jets+ $\cancel{E}_T$ , $\tan\beta=2$ , large $m_0$ , $\mu < 0$ , $A_0=0$
>255	95	355 ABBOTT	01D D0	$ll$ +jets+ $\cancel{E}_T$ , $\tan\beta=2$ , $m_{\tilde{g}}=m_{\tilde{q}}$ , $\mu < 0$ , $A_0=0$
>168	95	356 AFFOLDER	01J CDF	$ll$ +Jets+ $\cancel{E}_T$ , $\tan\beta=2$ , $\mu=-800$ GeV, $m_{\tilde{q}} \gg m_{\tilde{g}}$
>221	95	356 AFFOLDER	01J CDF	$ll$ +Jets+ $\cancel{E}_T$ , $\tan\beta=2$ , $\mu=-800$ GeV, $m_{\tilde{q}}=m_{\tilde{g}}$
>190	95	357 ABBOTT	99L D0	Jets+ $\cancel{E}_T$ , $\tan\beta=2$ , $\mu < 0$ , $A=0$
>260	95	357 ABBOTT	99L D0	Jets+ $\cancel{E}_T$ , $m_{\tilde{g}}=m_{\tilde{q}}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>224	95	358 ABAZOV	02F D0	$\mathcal{R} \lambda'_{2jk}$ indirect decays, $\tan\beta=2$ , any $m_{\tilde{q}}$
>265	95	358 ABAZOV	02F D0	$\mathcal{R} \lambda'_{2jk}$ indirect decays, $\tan\beta=2$ , $m_{\tilde{q}}=m_{\tilde{g}}$
		359 ABAZOV	02G D0	$p\bar{p} \rightarrow \tilde{g}\tilde{g}, \tilde{g}\tilde{q}$
		360 CHEUNG	02B THEO	
		361 BERGER	01 THEO	$p\bar{p} \rightarrow X+b$ -quark
>240	95	362 ABBOTT	99 D0	$\tilde{g} \rightarrow \tilde{\chi}_2^0 X \rightarrow \tilde{\chi}_1^0 \gamma X$ , $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} > 20$ GeV
>320	95	362 ABBOTT	99 D0	$\tilde{g} \rightarrow \tilde{\chi}_1^0 X \rightarrow \tilde{G} \gamma X$
>227	95	363 ABBOTT	99K D0	any $m_{\tilde{q}}$ , $\mathcal{R}$ , $\tan\beta=2$ , $\mu < 0$
>212	95	364 ABACHI	95C D0	$m_{\tilde{g}} \geq m_{\tilde{q}}$ ; with cascade decays
>144	95	364 ABACHI	95C D0	Any $m_{\tilde{q}}$ ; with cascade decays
		365 ABE	95T CDF	$\tilde{g} \rightarrow \tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 \gamma$
		366 HEBBEKER	93 RVUE	$e^+e^-$ jet analyses
>218	90	367 ABE	92L CDF	$m_{\tilde{q}} \leq m_{\tilde{g}}$ ; with cascade decay
>100		368 ROY	92 RVUE	$p\bar{p} \rightarrow \tilde{g}\tilde{g}; \mathcal{R}$
		369 NOJIRI	91 COSM	
none 4–53	90	370 ALBAJAR	87D UA1	Any $m_{\tilde{q}} > m_{\tilde{g}}$
none 4–75	90	370 ALBAJAR	87D UA1	$m_{\tilde{q}} = m_{\tilde{g}}$
none 16–58	90	371 ANSARI	87D UA2	$m_{\tilde{q}} \lesssim 100$ GeV

354 AFFOLDER 02 searched in  $\sim 84 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions for events with  $\geq 3$  jets and  $\cancel{E}_T$ , arising from the production of gluinos and/or squarks. Limits are derived by scanning the parameter space, for  $m_{\tilde{q}} \geq m_{\tilde{g}}$  in the framework of minimal Supergravity, assuming five flavors of degenerate squarks, and for  $m_{\tilde{q}} < m_{\tilde{g}}$  in the framework of constrained MSSM, assuming conservatively four flavors of degenerate squarks. See Fig. 3 for the variation of the limit as function of the squark mass. Supersedes the results of ABE 97K.

355 ABBOTT 01D looked in  $\sim 108 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.8$  TeV for events with  $e\bar{e}$ ,  $\mu\bar{\mu}$ , or  $e\mu$  accompanied by at least 2 jets and  $\cancel{E}_T$ . Excluded regions are obtained in the

- MSUGRA framework from a scan over the parameters  $0 < m_0 < 300$  GeV,  $10 < m_{1/2} < 110$  GeV, and  $1.2 < \tan\beta < 10$ .
- 356 AFFOLDER 01J searched in  $\sim 106$  pb $^{-1}$  of  $p\bar{p}$  collisions for events with 2 like-sign leptons ( $e$  or  $\mu$ ),  $\geq 2$  jets and  $\cancel{E}_T$ , expected to arise from the production of gluinos and/or squarks with cascade decays into  $\tilde{\chi}^\pm$  or  $\tilde{\chi}_2^0$ . Spectra and decay rates are evaluated in the framework of minimal Supergravity, assuming five flavors of degenerate squarks and a pseudoscalar Higgs mass  $m_A=500$  GeV. The limits are derived for  $\tan\beta=2$ ,  $\mu=-800$  GeV, and scanning over  $m_{\tilde{g}}$  and  $m_{\tilde{q}}$ . See Fig. 2 for the variation of the limit as function of the squark mass. These limits supersede the results of ABE 96D.
- 357 ABBOTT 99L consider events with three or more jets and large  $\cancel{E}_T$ . Spectra and decay rates are evaluated in the framework of minimal Supergravity, assuming five flavors of degenerate squarks, and scanning the space of the universal gaugino ( $m_{1/2}$ ) and scalar ( $m_0$ ) masses. See their Figs. 2–3 for the dependence of the limit on the relative value of  $m_{\tilde{q}}$  and  $m_{\tilde{g}}$ .
- 358 ABAZOV 02F looked in 77.5 pb $^{-1}$  of  $p\bar{p}$  collisions at 1.8 TeV for events with  $\geq 2\mu + \geq 4$  jets, originating from associated production of squarks followed by an indirect  $\cancel{H}$  decay (of the  $\tilde{\chi}_1^0$ ) via  $LQ\bar{D}$  couplings of the type  $\lambda'_{2jk}$  where  $j=1,2$  and  $k=1,2,3$ . Bounds are obtained in the MSUGRA scenario by a scan in the range  $0 \leq M_0 \leq 400$  GeV,  $60 \leq m_{1/2} \leq 120$  GeV for fixed values  $A_0=0$ ,  $\mu < 0$ , and  $\tan\beta=2$  or 6. The bounds are weaker for  $\tan\beta=6$ . See Figs. 2,3 for the exclusion contours in  $m_{1/2}$  versus  $m_0$  for  $\tan\beta=2$  and 6, respectively.
- 359 ABAZOV 02G search for associated production of gluinos and squarks in 92.7 pb $^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.8$  TeV, using events with one electron,  $\geq 4$  jets, and large  $\cancel{E}_T$ . The results are compared to a MSUGRA scenario with  $\mu < 0$ ,  $A_0=0$ , and  $\tan\beta=3$  and allow to exclude a region of the  $(m_0, m_{1/2})$  shown in Fig. 11.
- 360 CHEUNG 02B studies the constraints on a  $\tilde{b}_1$  with mass in the 2.2–5.5 GeV region and a gluino in the mass range 12–16 GeV, using precision measurements of  $Z^0$  decays and  $e^+e^-$  annihilations at LEP2. Few detectable events are predicted in the LEP2 data for the model proposed by BERGER 01.
- 361 BERGER 01 reanalyzed interpretation of Tevatron data on bottom-quark production. Argues that pair production of light gluinos ( $m \sim 12$ –16 GeV) with subsequent 2-body decay into a light sbottom ( $m \sim 2$ –5.5 GeV) and bottom can reconcile Tevatron data with predictions of perturbative QCD for the bottom production rate. The sbottom must either decay hadronically via a  $R$ -parity- and  $B$ -violating interaction, or be long-lived.
- 362 ABBOTT 99 searched for  $\gamma\cancel{E}_T + \geq 2$  jet final states, and set limits on  $\sigma(p\bar{p} \rightarrow \tilde{g}+X) \cdot B(\tilde{g} \rightarrow \gamma\cancel{E}_T X)$ . The quoted limits correspond to  $m_{\tilde{q}} \geq m_{\tilde{g}}$ , with  $B(\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 \gamma)=1$  and  $B(\tilde{\chi}_1^0 \rightarrow \tilde{G} \gamma)=1$ , respectively. They improve to 310 GeV (360 GeV in the case of  $\gamma\tilde{G}$  decay) for  $m_{\tilde{g}}=m_{\tilde{q}}$ .
- 363 ABBOTT 99K uses events with an electron pair and four jets to search for the decay of the  $\tilde{\chi}_1^0$  LSP via  $\cancel{H} LQ\bar{D}$  couplings. The particle spectrum and decay branching ratios are taken in the framework of minimal supergravity. An excluded region at 95% CL is obtained in the  $(m_0, m_{1/2})$  plane under the assumption that  $A_0=0$ ,  $\mu < 0$ ,  $\tan\beta=2$  and any one of the couplings  $\lambda'_{1jk} > 10^{-3}$  ( $j=1,2$  and  $k=1,2,3$ ) and from which the above limit is computed. For equal mass squarks and gluinos, the corresponding limit is 277 GeV. The results are essentially independent of  $A_0$ , but the limit deteriorates rapidly with increasing  $\tan\beta$  or  $\mu > 0$ .
- 364 ABACHI 95C assume five degenerate squark flavors with  $m_{\tilde{q}_L} = m_{\tilde{q}_R}$ . Sleptons are assumed to be heavier than squarks. The limits are derived for fixed  $\tan\beta = 2.0$ ,  $\mu = -250$  GeV, and  $m_{H^+}=500$  GeV, and with the cascade decays of the squarks and gluinos

- calculated within the framework of the Minimal Supergravity scenario. The bounds are weakly sensitive to the three fixed parameters for a large fraction of parameter space.
- 365 ABE 95T looked for a cascade decay of gluino into  $\tilde{\chi}_2^0$  which further decays into  $\tilde{\chi}_1^0$  and a photon. No signal is observed. Limits vary widely depending on the choice of parameters. For  $\mu = -40$  GeV,  $\tan\beta = 1.5$ , and heavy squarks, the range  $50 < m_{\tilde{g}} \text{ (GeV)} < 140$  is excluded at 90% CL. See the paper for details.
- 366 HEBBEKER 93 combined jet analyses at various  $e^+e^-$  colliders. The 4-jet analyses at TRISTAN/LEP and the measured  $\alpha_s$  at PEP/PETRA/TRISTAN/LEP are used. A constraint on effective number of quarks  $N=6.3 \pm 1.1$  is obtained, which is compared to that with a light gluino,  $N=8$ .
- 367 ABE 92L bounds are based on similar assumptions as ABACHI 95C. Not sensitive to  $m_{\text{gluino}} < 40$  GeV (but other experiments rule out that region).
- 368 ROY 92 reanalyzed CDF limits on di-lepton events to obtain limits on gluino production in  $R$ -parity violating models. The 100% decay  $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}$  where  $\tilde{\chi}$  is the LSP, and the LSP decays either into  $\ell q\bar{d}$  or  $\ell\ell\bar{e}$  is assumed.
- 369 NOJIRI 91 argues that a heavy gluino should be nearly degenerate with squarks in minimal supergravity not to overclose the universe.
- 370 The limits of ALBAJAR 87D are from  $p\bar{p} \rightarrow \tilde{g}\tilde{g}X$  ( $\tilde{g} \rightarrow q\bar{q}\tilde{\gamma}$ ) and assume  $m_{\tilde{q}} > m_{\tilde{g}}$ . These limits apply for  $m_{\tilde{\gamma}} \lesssim 20$  GeV and  $\tau(\tilde{g}) < 10^{-10}$  s.
- 371 The limit of ANSARI 87D assumes  $m_{\tilde{q}} > m_{\tilde{g}}$  and  $m_{\tilde{\gamma}} \approx 0$ .

### Long-lived/light $\tilde{g}$ (Gluino) MASS LIMIT

Limits on light gluinos ( $m_{\tilde{g}} < 5$  GeV), or gluinos which leave the detector before decaying.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
> 12		372 BERGER	05 THEO	hadron scattering data
none 2–18	95	373 ABDALLAH	03C DLPH	$e^+e^- \rightarrow q\bar{q}\tilde{g}\tilde{g}$ , stable $\tilde{g}$
> 5		374 ABDALLAH	03G DLPH	QCD beta function
		375 HEISTER	03 ALEP	Color factors
> 26.9	95	376 HEISTER	03H ALEP	$e^+e^- \rightarrow q\bar{q}\tilde{g}\tilde{g}$
> 6.3		377 JANOT	03 RVUE	$\Delta\Gamma_{had} < 3.9$ MeV
		378 MAFI	00 THEO	$pp \rightarrow \text{jets} + \cancel{p}_T$
		379 ALAVI-HARATI	199E KTEV	$pN \rightarrow R^0$ , with $R^0 \rightarrow \rho^0\tilde{\gamma}$ and $R^0 \rightarrow \pi^0\tilde{\gamma}$
		380 BAER	99 RVUE	Stable $\tilde{g}$ hadrons
		381 FANTI	99 NA48	$p\text{Be} \rightarrow R^0 \rightarrow \eta\tilde{\gamma}$
		382 ACKERSTAFF	98V OPAL	$e^+e^- \rightarrow \tilde{\chi}_1^+\tilde{\chi}_1^-$
		383 ADAMS	97B KTEV	$pN \rightarrow R^0 \rightarrow \rho^0\tilde{\gamma}$
		384 ALBUQUERQ..	97 E761	$R^+(uud\tilde{g}) \rightarrow S^0(uds\tilde{g})\pi^+$ , $X^-(ssd\tilde{g}) \rightarrow S^0\pi^-$
> 6.3	95	385 BARATE	97L ALEP	Color factors
> 5	99	386 CSIKOR	97 RVUE	$\beta$ function, $Z \rightarrow \text{jets}$
> 1.5	90	387 DEGOUVEA	97 THEO	$Z \rightarrow jjjj$
		388 FARRAR	96 RVUE	$R^0 \rightarrow \pi^0\tilde{\gamma}$
none 1.9–13.6	95	389 AKERS	95R OPAL	$Z$ decay into a long-lived $(\tilde{g}q\bar{q})^\pm$
< 0.7		390 CLAVELLI	95 RVUE	quarkonia
none 1.5–3.5		391 CAKIR	94 RVUE	$\Upsilon(1S) \rightarrow \gamma + \text{gluonium}$
not 3–5		392 LOPEZ	93C RVUE	LEP

$\approx 4$		393 CLAVELLI	92 RVUE	$\alpha_s$ running
		394 ANTONIADIS	91 RVUE	$\alpha_s$ running
$> 1$		395 ANTONIADIS	91 RVUE	$pN \rightarrow$ missing energy
		396 NAKAMURA	89 SPEC	$R\text{-}\Delta^{++}$
$> 3.8$	90	397 ARNOLD	87 EMUL	$\pi^-$ (350 GeV). $\sigma \simeq A^1$
$> 3.2$	90	397 ARNOLD	87 EMUL	$\pi^-$ (350 GeV). $\sigma \simeq A^{0.72}$
none 0.6–2.2	90	398 TUTS	87 CUSB	$\Upsilon(1S) \rightarrow \gamma +$ gluinonium
none 1–4.5	90	399 ALBRECHT	86C ARG	$1 \times 10^{-11} \lesssim \tau \lesssim 1 \times 10^{-9} \text{s}$
none 1–4	90	400 BADIER	86 BDMP	$1 \times 10^{-10} < \tau < 1 \times 10^{-7} \text{s}$
none 3–5		401 BARNETT	86 RVUE	$p\bar{p} \rightarrow$ gluino gluino gluon
none		402 VOLOSHIN	86 RVUE	If (quasi) stable; $\tilde{g}uud$
none 0.5–2		403 COOPER-...	85B BDMP	For $m_{\tilde{q}}=300$ GeV
none 0.5–4		403 COOPER-...	85B BDMP	For $m_{\tilde{q}} < 65$ GeV
none 0.5–3		403 COOPER-...	85B BDMP	For $m_{\tilde{q}}=150$ GeV
none 2–4		404 DAWSON	85 RVUE	$\tau > 10^{-7} \text{s}$
none 1–2.5		404 DAWSON	85 RVUE	For $m_{\tilde{q}}=100$ GeV
none 0.5–4.1	90	405 FARRAR	85 RVUE	FNAL beam dump
$> 1$		406 GOLDMAN	85 RVUE	Gluinonium
$> 1-2$		407 HABER	85 RVUE	
		408 BALL	84 CALO	
		409 BRICK	84 RVUE	
		410 FARRAR	84 RVUE	
$> 2$		411 BERGSMA	83C RVUE	For $m_{\tilde{q}} < 100$ GeV
		412 CHANOWITZ	83 RVUE	$\tilde{g}u\bar{d}, \tilde{g}uud$
$> 2-3$		413 KANE	82 RVUE	Beam dump
$> 1.5-2$		FARRAR	78 RVUE	$R$ -hadron

372 BERGER 05 include the light gluino in proton PDF and perform global analysis of hadronic data. Effects on the running of  $\alpha_s$  also included. Strong dependency on  $\alpha_s(m_Z)$ . Bound quoted for  $\alpha_s(m_Z) = 0.118$ .

373 ABDALLAH 03C looked for events of the type  $q\bar{q}R^\pm R^\pm, q\bar{q}R^\pm R^0$  or  $q\bar{q}R^0 R^0$  in  $e^+e^-$  interactions at 91.2 GeV collected in 1994. The  $R^\pm$  bound states are identified by anomalous  $dE/dx$  in the tracking chambers and the  $R^0$  by missing energy, due to their reduced energy loss in the calorimeters. The upper value of the excluded range depends on the probability for the gluino to fragment into  $R^\pm$  or  $R^0$ , see their Fig. 17. It improves to 23 GeV for 100% fragmentation to  $R^\pm$ .

374 ABDALLAH 03G used  $e^+e^-$  data at and around the  $Z^0$  peak, above the  $Z^0$  up to  $\sqrt{s} = 202$  GeV and events from radiative return to cover the low energy region. They perform a direct measurement of the QCD beta-function from the means of fully inclusive event observables. Compared to the energy range, gluinos below 5 GeV can be considered massless and are firmly excluded by the measurement.

375 HEISTER 03 use  $e^+e^-$  data from 1994 and 1995 at and around the  $Z^0$  peak to measure the 4-jet rate and angular correlations. The comparison with QCD NLO calculations allow  $\alpha_S(M_Z)$  and the color factor ratios to be extracted and the results are in agreement with the expectations from QCD. The inclusion of a massless gluino in the beta functions yields  $T_R/C_F = 0.15 \pm 0.06 \pm 0.06$  (expectation is  $T_R/C_F = 3/8$ ), excluding a massless gluino at more than 95% CL. As no NLO calculations are available for massive gluinos, the earlier LO results from BARATE 97L for massive gluinos remain valid.

376 HEISTER 03H use  $e^+e^-$  data at and around the  $Z^0$  peak to look for stable gluinos hadronizing into charged or neutral  $R$ -hadrons with arbitrary branching ratios. Combining these results with bounds on the  $Z^0$  hadronic width from electroweak measurements (JANOT 03) to cover the low mass region the quoted lower limit on the mass of a long-lived gluino is obtained.

- 377 JANOT 03 excludes a light gluino from the upper limit on an additional contribution to the  $Z$  hadronic width. At higher confidence levels,  $m_{\tilde{g}} > 5.3(4.2)$  GeV at  $3\sigma(5\sigma)$  level.
- 378 MAFI 00 reanalyzed CDF data assuming a stable heavy gluino as the LSP, with model for  $R$ -hadron-nucleon scattering. Gluino masses between 35 GeV and 115 GeV are excluded based on the CDF Run I data. Combined with the analysis of BAER 99, this allows a LSP gluino mass between 25 and 35 GeV if the probability of fragmentation into charged  $R$ -hadron  $P > 1/2$ . The cosmological exclusion of such a gluino LSP are assumed to be avoided as in BAER 99. Gluino could be NLSP with  $\tau_{\tilde{g}} \sim 100$  yrs, and decay to gluon gravitino.
- 379 ALAVI-HARATI 99E looked for  $R^0$  bound states, yielding  $\pi^+\pi^-$  or  $\pi^0$  in the final state. The experiment is sensitive to values of  $\Delta m = m_{R^0} - m_{\tilde{\gamma}}$  larger than 280 MeV and 140 MeV for the two decay modes, respectively, and to  $R^0$  mass and lifetime in the ranges 0.8–5 GeV and  $10^{-10}$ – $10^{-3}$  s. The limits obtained depend on  $B(R^0 \rightarrow \pi^+\pi^- \text{ photino})$  and  $B(R^0 \rightarrow \pi^0 \text{ photino})$  on the value of  $m_{R^0}/m_{\tilde{\gamma}}$ , and on the ratio of production rates  $\sigma(R^0)/\sigma(K_L^0)$ . See Figures in the paper for the excluded  $R^0$  production rates as a function of  $\Delta m$ ,  $R^0$  mass and lifetime. Using the production rates expected from perturbative QCD, and assuming dominance of the above decay channels over the suitable phase space,  $R^0$  masses in the range 0.8–5 GeV are excluded at 90%CL for a large fraction of the sensitive lifetime region. ALAVI-HARATI 99E updates and supersedes the results of ADAMS 97B.
- 380 BAER 99 set constraints on the existence of stable  $\tilde{g}$  hadrons, in the mass range  $m_{\tilde{g}} > 3$  GeV. They argue that strong-interaction effects in the low-energy annihilation rates could leave small enough relic densities to evade cosmological constraints up to  $m_{\tilde{g}} < 10$  TeV. They consider  $\text{jet} + \cancel{E}_T$  as well as heavy-ionizing charged-particle signatures from production of stable  $\tilde{g}$  hadrons at LEP and Tevatron, developing modes for the energy loss of  $\tilde{g}$  hadrons inside the detectors. Results are obtained as a function of the fragmentation probability  $P$  of the  $\tilde{g}$  into a charged hadron. For  $P < 1/2$ , and for various energy-loss models, OPAL and CDF data exclude gluinos in the  $3 < m_{\tilde{g}}(\text{GeV}) < 130$  mass range. For  $P > 1/2$ , gluinos are excluded in the mass ranges  $3 < m_{\tilde{g}}(\text{GeV}) < 23$  and  $50 < m_{\tilde{g}}(\text{GeV}) < 200$ .
- 381 FANTI 99 looked for  $R^0$  bound states yielding high  $P_T \eta \rightarrow 3\pi^0$  decays. The experiment is sensitive to a region of  $R^0$  mass and lifetime in the ranges of 1–5 GeV and  $10^{-10}$ – $10^{-3}$  s. The limits obtained depend on  $B(R^0 \rightarrow \eta\tilde{\gamma})$ , on the value of  $m_{R^0}/m_{\tilde{\gamma}}$ , and on the ratio of production rates  $\sigma(R^0)/\sigma(K_L^0)$ . See Fig. 6–7 for the excluded production rates as a function of  $R^0$  mass and lifetime.
- 382 ACKERSTAFF 98V excludes the light gluino with universal gaugino mass where charginos, neutralinos decay as  $\tilde{\chi}_1^\pm, \tilde{\chi}_2^0 \rightarrow q\bar{q}\tilde{g}$  from total hadronic cross sections at  $\sqrt{s}=130$ –172 GeV. See paper for the case of nonuniversal gaugino mass.
- 383 ADAMS 97B looked for  $\rho^0 \rightarrow \pi^+\pi^-$  as a signature of  $R^0 = (\tilde{g}g)$  bound states. The experiment is sensitive to an  $R^0$  mass range of 1.2–4.5 GeV and to a lifetime range of  $10^{-10}$ – $10^{-3}$  sec. Precise limits depend on the assumed value of  $m_{R^0}/m_{\tilde{\gamma}}$ . See Fig. 7 for the excluded mass and lifetime region.
- 384 ALBUQUERQUE 97 looked for weakly decaying baryon-like states which contain a light gluino, following the suggestions in FARRAR 96. See their Table 1 for limits on the production fraction. These limits exclude gluino masses in the range 100–600 MeV for the predicted lifetimes (FARRAR 96) and production rates, which are assumed to be comparable to those of strange or charmed baryons.
- 385 BARATE 97L studied the QCD color factors from four-jet angular correlations and the differential two-jet rate in  $Z$  decay. Limit obtained from the determination of  $n_f = 4.24 \pm 0.29 \pm 1.15$ , assuming  $T_F/C_F=3/8$  and  $C_A/C_F=9/4$ .

- 386 CSIKOR 97 combined the  $\alpha_s$  from  $\sigma(e^+e^- \rightarrow \text{hadron})$ ,  $\tau$  decay, and jet analysis in  $Z$  decay. They exclude a light gluino below 5 GeV at more than 99.7%CL.
- 387 DEGOUVEA 97 reanalyzed AKERS 95A data on  $Z$  decay into four jets to place constraints on a light stable gluino. The mass limit corresponds to the pole mass of 2.8 GeV. The analysis, however, is limited to the leading-order QCD calculation.
- 388 FARRAR 96 studied the possible  $R^0=(\tilde{g}g)$  component in Fermilab E799 experiment and used its bound  $B(K_L^0 \rightarrow \pi^0 \nu \bar{\nu}) \leq 5.8 \times 10^{-5}$  to place constraints on the combination of  $R^0$  production cross section and its lifetime.
- 389 AKERS 95R looked for  $Z$  decay into  $q\bar{q}\tilde{g}\tilde{g}$ , by searching for charged particles with  $dE/dx$  consistent with  $\tilde{g}$  fragmentation into a state  $(\tilde{g}q\bar{q})^\pm$  with lifetime  $\tau > 10^{-7}$  sec. The fragmentation probability into a charged state is assumed to be 25%.
- 390 CLAVELLI 95 updates the analysis of CLAVELLI 93, based on a comparison of the hadronic widths of charmonium and bottomonium  $S$ -wave states. The analysis includes a parametrization of relativistic corrections. Claims that the presence of a light gluino improves agreement with the data by slowing down the running of  $\alpha_s$ .
- 391 CAKIR 94 reanalyzed TUTS 87 and later unpublished data from CUSB to exclude pseudo-scalar gluinonium  $\eta_{\tilde{g}}(\tilde{g}\tilde{g})$  of mass below 7 GeV. It was argued, however, that the perturbative QCD calculation of the branching fraction  $\mathcal{T} \rightarrow \eta_{\tilde{g}}\gamma$  is unreliable for  $m_{\eta_{\tilde{g}}} < 3$  GeV. The gluino mass is defined by  $m_{\tilde{g}}=(m_{\eta_{\tilde{g}}})/2$ . The limit holds for any gluino lifetime.
- 392 LOPEZ 93C uses combined restraint from the radiative symmetry breaking scenario within the minimal supergravity model, and the LEP bounds on the  $(M_2, \mu)$  plane. Claims that the light gluino window is strongly disfavored.
- 393 CLAVELLI 92 claims that a light gluino mass around 4 GeV should exist to explain the discrepancy between  $\alpha_s$  at LEP and at quarkonia ( $\mathcal{T}$ ), since a light gluino slows the running of the QCD coupling.
- 394 ANTONIADIS 91 argue that possible light gluinos ( $< 5$  GeV) contradict the observed running of  $\alpha_s$  between 5 GeV and  $m_Z$ . The significance is less than 2 s.d.
- 395 ANTONIADIS 91 interpret the search for missing energy events in 450 GeV/c  $pN$  collisions, AKESSON 91, in terms of light gluinos.
- 396 NAKAMURA 89 searched for a long-lived ( $\tau \gtrsim 10^{-7}$  s) charge-( $\pm 2$ ) particle with mass  $\lesssim 1.6$  GeV in proton-Pt interactions at 12 GeV and found that the yield is less than  $10^{-8}$  times that of the pion. This excludes  $R\text{-}\Delta^{++}$  (a  $\tilde{g}uuu$  state) lighter than 1.6 GeV.
- 397 The limits assume  $m_{\tilde{q}} = 100$  GeV. See their figure 3 for limits vs.  $m_{\tilde{q}}$ .
- 398 The gluino mass is defined by half the bound  $\tilde{g}\tilde{g}$  mass. If zero gluino mass gives a  $\tilde{g}\tilde{g}$  of mass about 1 GeV as suggested by various glueball mass estimates, then the low-mass bound can be replaced by zero. The high-mass bound is obtained by comparing the data with nonrelativistic potential-model estimates.
- 399 ALBRECHT 86C search for secondary decay vertices from  $\chi_{b1}(1P) \rightarrow \tilde{g}\tilde{g}g$  where  $\tilde{g}$ 's make long-lived hadrons. See their figure 4 for excluded region in the  $m_{\tilde{g}} - m_{\tilde{g}}$  and  $m_{\tilde{g}} - m_{\tilde{q}}$  plane. The lower  $m_{\tilde{g}}$  region below  $\sim 2$  GeV may be sensitive to fragmentation effects. Remark that the  $\tilde{g}$ -hadron mass is expected to be  $\sim 1$  GeV (glueball mass) in the zero  $\tilde{g}$  mass limit.
- 400 BADIER 86 looked for secondary decay vertices from long-lived  $\tilde{g}$ -hadrons produced at 300 GeV  $\pi^-$  beam dump. The quoted bound assumes  $\tilde{g}$ -hadron nucleon total cross section of  $10\mu\text{b}$ . See their figure 7 for excluded region in the  $m_{\tilde{g}} - m_{\tilde{q}}$  plane for several assumed total cross-section values.
- 401 BARNETT 86 rule out light gluinos ( $m = 3\text{--}5$  GeV) by calculating the monojet rate from gluino gluino gluon events (and from gluino gluino events) and by using UA1 data from  $p\bar{p}$  collisions at CERN.
- 402 VOLOSHIN 86 rules out stable gluino based on the cosmological argument that predicts too much hydrogen consisting of the charged stable hadron  $\tilde{g}uud$ . Quasi-stable ( $\tau >$

1.  $\times 10^{-7}$ s) light gluino of  $m_{\tilde{g}} < 3$  GeV is also ruled out by nonobservation of the stable charged particles,  $\tilde{g}uud$ , in high energy hadron collisions.
- 403 COOPER-SARKAR 85B is BEBC beam-dump. Gluinos decaying in dump would yield  $\tilde{\gamma}$ 's in the detector giving neutral-current-like interactions. For  $m_{\tilde{q}} > 330$  GeV, no limit is set.
- 404 DAWSON 85 first limit from neutral particle search. Second limit based on FNAL beam dump experiment.
- 405 FARRAR 85 points out that BALL 84 analysis applies only if the  $\tilde{g}$ 's decay before interacting, i.e.  $m_{\tilde{q}} < 80m_{\tilde{g}}^{1.5}$ . FARRAR 85 finds  $m_{\tilde{g}} < 0.5$  not excluded for  $m_{\tilde{q}} = 30$ –1000 GeV and  $m_{\tilde{g}} < 1.0$  not excluded for  $m_{\tilde{q}} = 100$ –500 GeV by BALL 84 experiment.
- 406 GOLDMAN 85 use nonobservation of a pseudoscalar  $\tilde{g}\text{-}\tilde{g}$  bound state in radiative  $\psi$  decay.
- 407 HABER 85 is based on survey of all previous searches sensitive to low mass  $\tilde{g}$ 's. Limit makes assumptions regarding the lifetime and electric charge of the lightest supersymmetric particle.
- 408 BALL 84 is FNAL beam dump experiment. Observed no interactions of  $\tilde{\gamma}$  in the calorimeter, where  $\tilde{\gamma}$ 's are expected to come from pair-produced  $\tilde{g}$ 's. Search for long-lived  $\tilde{\gamma}$  interacting in calorimeter 56m from target. Limit is for  $m_{\tilde{q}} = 40$  GeV and production cross section proportional to  $A^{0.72}$ . BALL 84 find no  $\tilde{g}$  allowed below 4.1 GeV at CL = 90%. Their figure 1 shows dependence on  $m_{\tilde{q}}$  and A. See also KANE 82.
- 409 BRICK 84 reanalyzed FNAL 147 GeV HBC data for  $R\text{-}\Delta(1232)^{++}$  with  $\tau > 10^{-9}$  s and  $p_{\text{lab}} > 2$  GeV. Set CL = 90% upper limits 6.1, 4.4, and 29 microbarns in  $pp$ ,  $\pi^+p$ ,  $K^+p$  collisions respectively.  $R\text{-}\Delta^{++}$  is defined as being  $\tilde{g}$  and 3 up quarks. If mass = 1.2–1.5 GeV, then limits may be lower than theory predictions.
- 410 FARRAR 84 argues that  $m_{\tilde{g}} < 100$  MeV is not ruled out if the lightest R-hadrons are long-lived. A long lifetime would occur if R-hadrons are lighter than  $\tilde{\gamma}$ 's or if  $m_{\tilde{q}} > 100$  GeV.
- 411 BERGSMA 83C is reanalysis of CERN-SPS beam-dump data. See their figure 1.
- 412 CHANOWITZ 83 find in bag-model that charged  $s$ -hadron exists which is stable against strong decay if  $m_{\tilde{g}} < 1$  GeV. This is important since tracks from decay of neutral  $s$ -hadron cannot be reconstructed to primary vertex because of missed  $\tilde{\gamma}$ . Charged  $s$ -hadron leaves track from vertex.
- 413 KANE 82 inferred above  $\tilde{g}$  mass limit from retroactive analysis of hadronic collision and beam dump experiments. Limits valid if  $\tilde{g}$  decays inside detector.

## LIGHT $\tilde{G}$ (Gravitino) MASS LIMITS FROM COLLIDER EXPERIMENTS

The following are bounds on light ( $\ll 1$  eV) gravitino indirectly inferred from its coupling to matter suppressed by the gravitino decay constant.

Unless otherwise stated, all limits assume that other supersymmetric particles besides the gravitino are too heavy to be produced. The gravitino is assumed to be undetected and to give rise to a missing energy ( $\cancel{E}$ ) signature.

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$> 1.09 \times 10^{-5}$	95	414 ABDALLAH	05B DLPH	$e^+e^- \rightarrow \tilde{G}\tilde{G}\gamma$
$> 1.35 \times 10^{-5}$	95	415 ACHARD	04E L3	$e^+e^- \rightarrow \tilde{G}\tilde{G}\gamma$
$> 1.3 \times 10^{-5}$	95	416 HEISTER	03C ALEP	$e^+e^- \rightarrow \tilde{G}\tilde{G}\gamma$
$> 11.7 \times 10^{-6}$	95	417 ACOSTA	02H CDF	
$> 8.7 \times 10^{-6}$	95	418 ABBIENDI,G	00D OPAL	$e^+e^- \rightarrow \tilde{G}\tilde{G}\gamma$

- >10.0 × 10<sup>-6</sup> 95 419 ABREU 00Z DLPH Superseded by ABDALLAH 05B  
 >11 × 10<sup>-6</sup> 95 420 AFFOLDER 00J CDF  $p\bar{p} \rightarrow \tilde{G}\tilde{G} + \text{jet}$   
 > 8.9 × 10<sup>-6</sup> 95 419 ACCIARRI 99R L3 Superseded by ACHARD 04E  
 > 7.9 × 10<sup>-6</sup> 95 421 ACCIARRI 98V L3  $e^+e^- \rightarrow \tilde{G}\tilde{G}\gamma$   
 > 8.3 × 10<sup>-6</sup> 95 421 BARATE 98J ALEP  $e^+e^- \rightarrow \tilde{G}\tilde{G}\gamma$
- 414 ABDALLAH 05B use data from  $\sqrt{s} = 180\text{--}208$  GeV. They look for events with a single photon +  $\cancel{E}$  final states from which a cross section limit of  $\sigma < 0.18$  pb at 208 GeV is obtained, allowing a limit on the mass to be set. Supersedes the results of ABREU 00Z.
- 415 ACHARD 04E use data from  $\sqrt{s} = 189\text{--}209$  GeV. They look for events with a single photon +  $\cancel{E}$  final states from which a limit on the Gravitino mass is set corresponding to  $\sqrt{F} > 238$  GeV. Supersedes the results of ACCIARRI 99R.
- 416 HEISTER 03C use the data from  $\sqrt{s} = 189\text{--}209$  GeV to search for  $\gamma\cancel{E}_T$  final states.
- 417 ACOSTA 02H looked in 87 pb<sup>-1</sup> of  $p\bar{p}$  collisions at  $\sqrt{s}=1.8$  TeV for events with a high- $E_t$  photon and  $\cancel{E}_T$ . They compared the data with a GMSB model where the final state could arise from  $q\bar{q} \rightarrow \tilde{G}\tilde{G}\gamma$ . Since the cross section for this process scales as  $1/|F|^4$ , a limit at 95% CL is derived on  $|F|^{1/2} > 221$  GeV. A model independent limit for the above topology is also given in the paper.
- 418 ABBIENDI,G 00D searches for  $\gamma\cancel{E}$  final states from  $\sqrt{s}=189$  GeV.
- 419 ABREU 00Z, ACCIARRI 99R search for  $\gamma\cancel{E}$  final states using data from  $\sqrt{s}=189$  GeV.
- 420 AFFOLDER 00J searches for final states with an energetic (from quark or gluon) and large  $\cancel{E}_T$  from undetected gravitinos.
- 421 Searches for  $\gamma\cancel{E}$  final states at  $\sqrt{s}=183$  GeV.

## Supersymmetry Miscellaneous Results

Results that do not appear under other headings or that make nonminimal assumptions.

VALUE	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
	422 ACOSTA	04E CDF	
	423 TCHIKILEV	04 ISTR	$K^- \rightarrow \pi^- \pi^0 P$
	424 AFFOLDER	02D CDF	$p\bar{p} \rightarrow \gamma b (\cancel{E}_T)$
	425 AFFOLDER	01H CDF	$p\bar{p} \rightarrow \gamma\gamma X$
	426 ABBOTT	00G D0	$p\bar{p} \rightarrow 3\ell + \cancel{E}_T, \cancel{E}, LL\bar{E}$
	427 ABREU,P	00C DLPH	$e^+e^- \rightarrow \gamma + S/P$
	428 ABACHI	97 D0	$\gamma\gamma X$
	429 BARBER	84B RVUE	
	430 HOFFMAN	83 CNTR	$\pi p \rightarrow n(e^+e^-)$
422	ACOSTA 04E looked in 107 pb <sup>-1</sup> of $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV for events with two same sign leptons without selection of other objects nor $\cancel{E}_T$ . No significant excess is observed compared to the Standard Model expectation and constraints are derived on the parameter space of MSUGRA models, see Figure 4.		
423	Looked for the scalar partner of a goldstino in decays $K^- \rightarrow \pi^- \pi^0 P$ from a 25 GeV $K^-$ beam produced at the IHEP 70 GeV proton synchrotron. The goldstino is assumed to be sufficiently long-lived to be invisible. A 90% CL upper limit on the decay branching ratio is set at $\sim 9.0 \times 10^{-6}$ for a goldstino mass range from 0 to 200 MeV, excluding the interval near $m(\pi^0)$ , where the limit is $\sim 3.5 \times 10^{-5}$ .		
424	AFFOLDER 02D looked in 85 pb <sup>-1</sup> of $p\bar{p}$ collisions at $\sqrt{s}=1.8$ TeV for events with a high- $E_T$ photon, and a $b$ -tagged jet with or without $\cancel{E}_T$ . They compared the data with models where the final state could arise from cascade decays of gluinos and/or squarks into $\tilde{\chi}^\pm$ and $\tilde{\chi}_2^0$ or direct associated production of $\tilde{\chi}_2^0 \tilde{\chi}_2^\pm$ , followed by $\tilde{\chi}_2^0 \rightarrow \gamma \tilde{\chi}_1^0$ or		



- a GMSB model where  $\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$ . It is concluded that the experimental sensitivity is insufficient to detect the associated production or the GMSB model, but some sensitivity may exist to the cascade decays. A model independent limit for the above topology is also given in the paper.
- 425 AFFOLDER 01H searches for  $p\bar{p} \rightarrow \gamma\gamma X$  events, where the di-photon system originates from sgoldstino production, in  $100 \text{ pb}^{-1}$  of data. Upper limits on the cross section times branching ratio are shown as function of the di-photon mass  $>70 \text{ GeV}$  in Fig. 5. Excluded regions are derived in the plane of the sgoldstino mass versus the supersymmetry breaking scale for two representative sets of parameter values, as shown in Figs. 6 and 7.
- 426 ABBOTT 00G searches for trilepton final states ( $\ell=e,\mu$ ) with  $\cancel{E}_T$  from the indirect decay of gauginos via  $LL\bar{E}$  couplings. Efficiencies are computed for all possible production and decay modes of SUSY particles in the framework of the Minimal Supergravity scenario. See Figs. 1–4 for excluded regions in the  $m_{1/2}$  versus  $m_0$  plane.
- 427 ABREU,P 00C look for the  $CP$ -even ( $S$ ) and  $CP$ -odd ( $P$ ) scalar partners of the goldstino, expected to be produced in association with a photon. The  $S/P$  decay into two photons or into two gluons and both the tri-photon and the photon + two jets topologies are investigated. Upper limits on the production cross section are shown in Fig. 5 and the excluded regions in Fig. 6. Data collected at  $\sqrt{s}=189\text{--}202 \text{ GeV}$ .
- 428 ABACHI 97 searched for  $p\bar{p} \rightarrow \gamma\gamma \cancel{E}_T + X$  as supersymmetry signature. It can be caused by selectron, sneutrino, or neutralino production with a radiative decay of their decay products. They placed limits on cross sections.
- 429 BARBER 84B consider that  $\tilde{\mu}$  and  $\tilde{e}$  may mix leading to  $\mu \rightarrow e\tilde{\gamma}\tilde{\gamma}$ . They discuss mass-mixing limits from decay dist. asym. in LBL-TRIUMF data and  $e^+$  polarization in SIN data.
- 430 HOFFMAN 83 set CL = 90% limit  $d\sigma/dt B(e^+e^-) < 3.5 \times 10^{-32} \text{ cm}^2/\text{GeV}^2$  for spin-1 partner of Goldstone fermions with  $140 < m < 160 \text{ MeV}$  decaying  $\rightarrow e^+e^-$  pair.

## REFERENCES FOR Supersymmetric Particle Searches

AKERIB	06	PR D73 011102R	D.S. Akerib <i>et al.</i>	(CDMS Collab.)
AKERIB	06A	PRL 96 011302	D.S. Akerib <i>et al.</i>	(CDMS Collab.)
ABAZOV	05A	PRL 94 041801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	05U	PRL 95 151805	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABDALLAH	05B	EPJ C38 395	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABULENCIA	05A	PRL 95 252001	A. Abulencia <i>et al.</i>	(CDF Collab.)
ACOSTA	05E	PR D71 031104R	D. Acosta <i>et al.</i>	(CDF Collab.)
ACOSTA	05R	PRL 95 131801	D. Acosta <i>et al.</i>	(CDF Collab.)
AKERIB	05	PR D72 052009	D.S. Akerib <i>et al.</i>	(CDMS Collab.)
AKTAS	05	PL B616 31	A. Aktas <i>et al.</i>	(H1 Collab.)
ALNER	05	PL B616 17	G.J. Alner <i>et al.</i>	(UK Dark Matter Collab.)
ALNER	05A	ASP 23 444	G.J. Alner <i>et al.</i>	(UK Dark Matter Collab.)
ANGLOHER	05	ASP 23 325	G. Angloher <i>et al.</i>	(CRESST-II Collab.)
BARNABE-HE...	05	PL B624 186	M. Barnabe-Heider <i>et al.</i>	(PICASSO Collab.)
BERGER	05	PR D71 014007	E.L. Berger <i>et al.</i>	
ELLIS	05	PR D71 095007	J. Ellis <i>et al.</i>	
GIRARD	05	PL B621 233	T.A. Girard <i>et al.</i>	(SIMPLE Collab.)
KLAPDOR-K...	05	PL B609 226	H.V. Klapdor-Kleingrothaus, I.V. Krivosheina, C. Tomei	
SANGLARD	05	PR D71 122002	V. Sanglard <i>et al.</i>	(EDELWEISS Collab.)
ABAZOV	04	PL B581 147	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	04B	PRL 93 011801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABBIENDI	04	EPJ C32 453	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	04F	EPJ C33 149	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	04H	EPJ C35 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	04N	PL B602 167	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABDALLAH	04H	EPJ C34 145	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABDALLAH	04M	EPJ C36 1	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
Also		EPJ C37 129 (erratum)	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ACHARD	04	PL B580 37	P. Achard <i>et al.</i>	(L3 Collab.)
ACHARD	04E	PL B587 16	P. Achard <i>et al.</i>	(L3)
ACOSTA	04B	PRL 92 051803	D. Acosta <i>et al.</i>	(CDF Collab.)
ACOSTA	04E	PRL 93 061802	D. Acosta <i>et al.</i>	(CDF Collab.)
AKERIB	04	PRL 93 211301	D. Akerib <i>et al.</i>	(CDMSII Collab.)

AKTAS	04B	PL B599 159	A. Aktas <i>et al.</i>	(H1 Collab.)
AKTAS	04D	EPJ C36 425	A. Aktas <i>et al.</i>	(H1 Collab.)
BELANGER	04	JHEP 0403 012	G. Belanger <i>et al.</i>	
BOTTINO	04	PR D69 037302	A. Bottino <i>et al.</i>	
DAS	04	PL B596 293	S.P. Das, A. Datta, M. Maity	
DESAI	04	PR D70 083523	S. Desai <i>et al.</i>	(Super-Kamiokande Collab.)
ELLIS	04	PR D69 015005	J. Ellis <i>et al.</i>	
ELLIS	04B	PR D70 055005	J. Ellis <i>et al.</i>	
GIULIANI	04	PL B588 151	F. Giuliani, T.A. Girard	
HEISTER	04	PL B583 247	A. Heister <i>et al.</i>	(ALEPH Collab.)
JANOT	04	PL B594 23	P. Janot	
PIERCE	04A	PR D70 075006	A. Pierce	
TCHIKILEV	04	PL B602 149	O.G. Tchikilev <i>et al.</i>	(ISTRA+ Coolab.)
ABBIENDI	03H	EPJ C29 479	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	03L	PL B572 8	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABDALLAH	03C	EPJ C26 505	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABDALLAH	03D	EPJ C27 153	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABDALLAH	03F	EPJ C28 15	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABDALLAH	03G	EPJ C29 285	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABDALLAH	03M	EPJ C31 421	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ACOSTA	03C	PRL 90 251801	D. Acosta <i>et al.</i>	(CDF Collab.)
ACOSTA	03E	PRL 91 171602	D. Acosta <i>et al.</i>	(CDF Collab.)
ADLOFF	03	PL B568 35	C. Adloff <i>et al.</i>	(H1 Collab.)
AHMED	03	ASP 19 691	B. Ahmed <i>et al.</i>	(UK Dark Matter Collab.)
AKERIB	03	PR D68 082002	D. Akerib <i>et al.</i>	(CDMS Collab.)
BAER	03	JCAP 0305 006	H. Baer, C. Balazs	
BAER	03A	JCAP 0309 007	H. Baer <i>et al.</i>	
BERGER	03	PL B552 223	E. Berger <i>et al.</i>	
BOTTINO	03	PR D68 043506	A. Bottino <i>et al.</i>	
BOTTINO	03A	PR D67 063519	A. Bottino, N. Fornengo, S. Scopel	
CHAKRAB...	03	PR D68 015005	S. Chakrabarti, M. Guchait, N.K. Mondal	
CHATTOPAD...	03	PR D68 035005	U. Chattopadhyay, A. Corsetti, P. Nath	
CHEKANOV	03B	PR D68 052004	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ELLIS	03	ASP 18 395	J. Ellis, K.A. Olive, Y. Santoso	
ELLIS	03B	NP B652 259	J. Ellis <i>et al.</i>	
ELLIS	03C	PL B565 176	J. Ellis <i>et al.</i>	
ELLIS	03D	PL B573 162	J. Ellis <i>et al.</i>	
ELLIS	03E	PR D67 123502	J. Ellis <i>et al.</i>	
HEISTER	03	EPJ C27 1	A. Heister <i>et al.</i>	(ALEPH)
HEISTER	03C	EPJ C28 1	A. Heister <i>et al.</i>	(ALEPH Collab.)
HEISTER	03G	EPJ C31 1	A. Heister <i>et al.</i>	(ALEPH Collab.)
HEISTER	03H	EPJ C31 327	A. Heister <i>et al.</i>	(ALEPH Collab.)
HOOPER	03	PL B562 18	D. Hooper, T. Plehn	
JANOT	03	PL B564 183	P. Janot	
KLAPDOR-K...	03	ASP 18 525	H.V. Klapdor-Kleingrothaus <i>et al.</i>	
LAHANAS	03	PL B568 55	A. Lahanas, D. Nanopoulos	
LEP	03	SLAC-R-701, LEPEWWG/2003-02		(LEP Collabs.)
ALEPH, DELPHI, L3, OPAL, the LEP EWWG, and the SLD HFEW				
TAKEDA	03	PL B572 145	A. Takeda <i>et al.</i>	
ABAZOV	02C	PRL 88 171802	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	02F	PRL 89 171801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	02G	PR D66 112001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	02H	PRL 89 261801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABBIENDI	02	EPJ C23 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	02B	PL B526 233	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	02H	PL B545 272	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
Also		PL B548 258 (erratum)	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABRAMS	02	PR D66 122003	D. Abrams <i>et al.</i>	(CDMS Collab.)
ACHARD	02	PL B524 65	P. Achard <i>et al.</i>	(L3 Collab.)
ACOSTA	02H	PRL 89 281801	D. Acosta <i>et al.</i>	(CDF Collab.)
AFFOLDER	02	PRL 88 041801	T. Affolder <i>et al.</i>	(CDF Collab.)
AFFOLDER	02D	PR D65 052006	T. Affolder <i>et al.</i>	(CDF Collab.)
ANGLOHER	02	ASP 18 43	G. Angloher <i>et al.</i>	(CRESST Collab.)
ARNOWITT	02	hep-ph/0211417	R. Arnowitt, B. Dutta	
BAEK	02	PL B541 161	S. Baek	
BAER	02	JHEP 0207 050	H. Baer <i>et al.</i>	
BECHER	02	PL B540 278	T. Becher <i>et al.</i>	
BENOIT	02	PL B545 43	A. Benoit <i>et al.</i>	(EDELWEISS Collab.)
CHEKANOV	02	PR D65 092004	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
CHEUNG	02B	PRL 89 221801	K. Cheung, W.-Y. Keung	
CHO	02	PRL 89 091801	G.-C. Cho	

ELLIS	02	PL B525 308	J. Ellis, D.V. Nanopoulos, K.A. Olive	
ELLIS	02B	PL B532 318	J. Ellis, A. Ferstl, K.A. Olive	
ELLIS	02C	PL B539 107	J. Ellis, K.A. Olive, Y. Santoso	
GHODBANE	02	NP B647 190	N. Ghodbane <i>et al.</i>	
HEISTER	02	PL B526 191	A. Heister <i>et al.</i>	(ALEPH Collab.)
HEISTER	02E	PL B526 206	A. Heister <i>et al.</i>	(ALEPH Collab.)
HEISTER	02F	EPJ C25 1	A. Heister <i>et al.</i>	(ALEPH Collab.)
HEISTER	02J	PL B533 223	A. Heister <i>et al.</i>	(ALEPH Collab.)
HEISTER	02K	PL B537 5	A. Heister <i>et al.</i>	(ALEPH Collab.)
HEISTER	02N	PL B544 73	A. Heister <i>et al.</i>	(ALEPH Collab.)
HEISTER	02R	EPJ C25 339	A. Heister <i>et al.</i>	(ALEPH Collab.)
KIM	02	PL B527 18	H.B. Kim <i>et al.</i>	
KIM	02B	JHEP 0212 034	Y.G. Kim <i>et al.</i>	
LAHANAS	02	EPJ C23 185	A. Lahanas, V.C. Spanos	
MORALES	02B	ASP 16 325	A. Morales <i>et al.</i>	(COSME Collab.)
MORALES	02C	PL B532 8	A. Morales <i>et al.</i>	(IGEX Collab.)
ABBIENDI	01	PL B501 12	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBOTT	01D	PR D63 091102	B. Abbott <i>et al.</i>	(D0 Collab.)
ABREU	01	EPJ C19 29	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	01B	EPJ C19 201	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	01C	PL B502 24	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	01D	PL B500 22	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	01G	PL B503 34	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	01	EPJ C19 397	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADAMS	01	PRL 87 041801	T. Adams <i>et al.</i>	(NuTeV Collab.)
ADLOFF	01B	EPJ C20 639	C. Adloff <i>et al.</i>	(H1 Collab.)
AFFOLDER	01B	PR D63 091101	T. Affolder <i>et al.</i>	(CDF Collab.)
AFFOLDER	01H	PR D64 092002	T. Affolder <i>et al.</i>	(CDF Collab.)
AFFOLDER	01J	PRL 87 251803	T. Affolder <i>et al.</i>	(CDF Collab.)
BALTZ	01	PRL 86 5004	E. Baltz, P. Gondolo	
BARATE	01	PL B499 67	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	01B	EPJ C19 415	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARGER	01C	PL B518 117	V. Barger, C. Kao	
BAUDIS	01	PR D63 022001	L. Baudis <i>et al.</i>	(Heidelberg-Moscow Collab.)
BENOIT	01	PL B513 15	A. Benoit <i>et al.</i>	(EDELWEISS Collab.)
BERGER	01	PRL 86 4231	E. Berger <i>et al.</i>	
BERNABEI	01	PL B509 197	R. Bernabei <i>et al.</i>	(DAMA Collab.)
BOTTINO	01	PR D63 125003	A. Bottino <i>et al.</i>	
BREITWEG	01	PR D63 052002	J. Breitweg <i>et al.</i>	(ZEUS Collab.)
CORSETTI	01	PR D64 125010	A. Corsetti, P. Nath	
DJOUADI	01	JHEP 0108 55	A. Djouadi, M. Drees, J.L. Kneur	
ELLIS	01B	PL B510 236	J. Ellis <i>et al.</i>	
ELLIS	01C	PR D63 065016	J. Ellis, A. Ferstl, K.A. Olive	
GOMEZ	01	PL B512 252	M.E. Gomez, J.D. Vergados	
LAHANAS	01	PL B518 94	A. Lahanas, D.V. Nanopoulos, V. Spanos	
ROSZKOWSKI	01	JHEP 0108 024	L. Roszkowski, R. Ruiz de Austri, T. Nihei	
SAVINOV	01	PR D63 051101	V. Savinov <i>et al.</i>	(CLEO Collab.)
ABBIENDI	00	EPJ C12 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	00G	EPJ C14 51	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	00H	EPJ C14 187	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
Also		EPJ C16 707 (erratum)	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	00J	EPJ C12 551	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	00R	EPJ C13 553	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI,G	00D	EPJ C18 253	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBOTT	00C	PRL 84 2088	B. Abbott <i>et al.</i>	(D0 Collab.)
ABBOTT	00G	PR D62 071701R	B. Abbott <i>et al.</i>	(D0 Collab.)
ABREU	00I	EPJ C13 591	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00J	PL B479 129	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00Q	PL B478 65	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00S	PL B485 45	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00T	PL B485 95	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00U	PL B487 36	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00V	EPJ C16 211	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00W	PL B489 38	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00Z	EPJ C17 53	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU,P	00C	PL B494 203	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU,P	00D	PL B496 59	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABUSAIDI	00	PRL 84 5699	R. Abusaidi <i>et al.</i>	(CDMS Collab.)
ACCIARRI	00C	EPJ C16 1	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	00D	PL B472 420	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	00K	PL B482 31	M. Acciarri <i>et al.</i>	(L3 Collab.)

ACCIARRI	00P	PL B489 81	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCOMANDO	00	NP B585 124	E. Accomando <i>et al.</i>	
AFFOLDER	00D	PRL 84 5704	T. Affolder <i>et al.</i>	(CDF Collab.)
AFFOLDER	00G	PRL 84 5273	T. Affolder <i>et al.</i>	(CDF Collab.)
AFFOLDER	00J	PRL 85 1378	T. Affolder <i>et al.</i>	(CDF Collab.)
AFFOLDER	00K	PRL 85 2056	T. Affolder <i>et al.</i>	(CDF Collab.)
BARATE	00G	EPJ C16 71	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	00H	EPJ C13 29	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	00I	EPJ C12 183	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	00P	PL B488 234	R. Barate <i>et al.</i>	(ALEPH Collab.)
BERNABEI	00	PL B480 23	R. Bernabei <i>et al.</i>	(DAMA Collab.)
BERNABEI	00C	EPJ C18 283	R. Bernabei <i>et al.</i>	(DAMA Collab.)
BERNABEI	00D	NJP 2 15	R. Bernabei <i>et al.</i>	(DAMA Collab.)
BOEHM	00B	PR D62 035012	C. Boehm, A. Djouadi, M. Drees	
BREITWEG	00E	EPJ C16 253	J. Breitweg <i>et al.</i>	(ZEUS Collab.)
CHO	00B	NP B574 623	G.-C. Cho, K. Hagiwara	
COLLAR	00	PRL 85 3083	J.I. Collar <i>et al.</i>	(SIMPLE Collab.)
ELLIS	00	PR D62 075010	J. Ellis <i>et al.</i>	
FENG	00	PL B482 388	J.L. Feng, K.T. Matchev, F. Wilczek	
LAHANAS	00	PR D62 023515	A. Lahanas, D.V. Nanopoulos, V.C. Spanos	
LEP	00	CERN-EP-2000-016	LEP Collabs.	(ALEPH, DELPHI, L3, OPAL, SLD+)
MAFI	00	PR D62 035003	A. Mafi, S. Raby	
MALTONI	00	PL B476 107	M. Maltoni <i>et al.</i>	
MORALES	00	PL B489 268	A. Morales <i>et al.</i>	(IGEX Collab.)
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i>	
SPOONER	00	PL B473 330	N.J.C. Spooner <i>et al.</i>	(UK Dark Matter Col.)
ABBIENDI	99	EPJ C6 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	99F	EPJ C8 23	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	99M	PL B456 95	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	99T	EPJ C11 619	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBOTT	99	PRL 82 29	B. Abbott <i>et al.</i>	(D0 Collab.)
ABBOTT	99F	PR D60 031101	B. Abbott <i>et al.</i>	(D0 Collab.)
ABBOTT	99J	PRL 83 2896	B. Abbott <i>et al.</i>	(D0 Collab.)
ABBOTT	99K	PRL 83 4476	B. Abbott <i>et al.</i>	(D0 Collab.)
ABBOTT	99L	PRL 83 4937	B. Abbott <i>et al.</i>	(D0 Collab.)
ABE	99I	PR D59 092002	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	99M	PRL 83 2133	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	99A	EPJ C11 383	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	99C	EPJ C6 385	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	99F	EPJ C7 595	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	99G	PL B446 62	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	99H	PL B456 283	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	99I	PL B459 354	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	99L	PL B462 354	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	99R	PL B470 268	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	99V	PL B471 308	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	99W	PL B471 280	M. Acciarri <i>et al.</i>	(L3 Collab.)
ALAVI-HARATI	99E	PRL 83 2128	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
AMBROSIO	99	PR D60 082002	M. Ambrosio <i>et al.</i>	(Macro Collab.)
BAER	99	PR D59 075002	H. Baer, K. Cheung, J.F. Gunion	
BARATE	99E	EPJ C7 383	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	99Q	PL B469 303	R. Barate <i>et al.</i>	(ALEPH Collab.)
BAUDIS	99	PR D59 022001	L. Baudis <i>et al.</i>	(Heidelberg-Moscow Collab.)
BELLI	99C	NP B563 97	P. Belli <i>et al.</i>	(DAMA Collab.)
BERNABEI	99	PL B450 448	R. Bernabei <i>et al.</i>	(DAMA Collab.)
FANTI	99	PL B446 117	V. Fanti <i>et al.</i>	(CERN NA48 Collab.)
MALTONI	99B	PL B463 230	M. Maltoni, M.I. Vysotsky	
OOTANI	99	PL B461 371	W. Ootani <i>et al.</i>	
ABBOTT	98	PRL 80 442	B. Abbott <i>et al.</i>	(D0 Collab.)
ABBOTT	98C	PRL 80 1591	B. Abbott <i>et al.</i>	(D0 Collab.)
ABBOTT	98E	PRL 80 2051	B. Abbott <i>et al.</i>	(D0 Collab.)
ABBOTT	98J	PRL 81 38	B. Abbott <i>et al.</i>	(D0 Collab.)
ABE	98J	PRL 80 5275	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	98S	PRL 81 4806	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	98	EPJ C1 1	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	98P	PL B444 491	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	98F	EPJ C4 207	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	98J	PL B433 163	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	98V	PL B444 503	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACKERSTAFF	98K	EPJ C4 47	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	98L	EPJ C2 213	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)

ACKERSTAFF	98P	PL B433 195	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	98V	EPJ C2 441	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	98H	PL B420 127	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98J	PL B429 201	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98K	PL B433 176	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98S	EPJ C4 433	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98X	EPJ C2 417	R. Barate <i>et al.</i>	(ALEPH Collab.)
BERNABEI	98	PL B424 195	R. Bernabei <i>et al.</i>	(DAMA Collab.)
BERNABEI	98C	PL B436 379	R. Bernabei <i>et al.</i>	(DAMA Collab.)
BREITWEG	98	PL B434 214	J. Breitweg <i>et al.</i>	(ZEUS Collab.)
ELLIS	98	PR D58 095002	J. Ellis <i>et al.</i>	
ELLIS	98B	PL B444 367	J. Ellis, T. Falk, K. Olive	
PDG	98	EPJ C3 1	C. Caso <i>et al.</i>	
ABACHI	97	PRL 78 2070	S. Abachi <i>et al.</i>	(D0 Collab.)
ABBOTT	97B	PRL 79 4321	B. Abbott <i>et al.</i>	(D0 Collab.)
ABE	97K	PR D56 R1357	F. Abe <i>et al.</i>	(CDF Collab.)
ACCIARRI	97U	PL B414 373	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACKERSTAFF	97H	PL B396 301	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ADAMS	97B	PRL 79 4083	J. Adams <i>et al.</i>	(FNAL KTeV Collab.)
ALBUQUERQUE...	97	PRL 78 3252	I.F. Albuquerque <i>et al.</i>	(FNAL E761 Collab.)
BAER	97	PR D57 567	H. Baer, M. Brhlik	
BARATE	97K	PL B405 379	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	97L	ZPHY C76 1	R. Barate <i>et al.</i>	(ALEPH Collab.)
BERNABEI	97	ASP 7 73	R. Bernabei <i>et al.</i>	(DAMA Collab.)
CARENA	97	PL B390 234	M. Carena, G.F. Giudice, C.E.M. Wagner	
CSIKOR	97	PRL 78 4335	F. Csikor, Z. Fodor	(EOTV, CERN)
DATTA	97	PL B395 54	A. Datta, M. Guchait, N. Parua	(ICTP, TATA)
DEGOUVEA	97	PL B400 117	A. de Gouvea, H. Murayama	
DERRICK	97	ZPHY C73 613	M. Derrick <i>et al.</i>	(ZEUS Collab.)
EDSJO	97	PR D56 1879	J. Edsjo, P. Gondolo	
ELLIS	97	PL B394 354	J. Ellis, J.L. Lopez, D.V. Nanopoulos	
HEWETT	97	PR D56 5703	J.L. Hewett, T.G. Rizzo, M.A. Doncheski	
KALINOWSKI	97	PL B400 112	J. Kalinowski, P. Zerwas	
TEREKHOV	97	PL B412 86	I. Terekhov	(ALAT)
ABACHI	96	PRL 76 2228	S. Abachi <i>et al.</i>	(D0 Collab.)
ABACHI	96B	PRL 76 2222	S. Abachi <i>et al.</i>	(D0 Collab.)
ABE	96	PRL 77 438	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96D	PRL 76 2006	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96K	PRL 76 4307	F. Abe <i>et al.</i>	(CDF Collab.)
AID	96	ZPHY C71 211	S. Aid <i>et al.</i>	(H1 Collab.)
AID	96C	PL B380 461	S. Aid <i>et al.</i>	(H1 Collab.)
ARNOWITT	96	PR D54 2374	R. Arnowitt, P. Nath	
BAER	96	PR D53 597	H. Baer, M. Brhlik	
BERGSTROM	96	ASP 5 263	L. Bergstrom, P. Gondolo	
CHO	96	PL B372 101	G.C. Cho, Y. Kizukuri, N. Oshimo	(TOKAH, OCH)
FARRAR	96	PRL 76 4111	G.R. Farrar	(RUTG)
LEWIN	96	ASP 6 87	J.D. Lewin, P.F. Smith	
TEREKHOV	96	PL B385 139	I. Terkhov, L. Clavelli	(ALAT)
ABACHI	95C	PRL 75 618	S. Abachi <i>et al.</i>	(D0 Collab.)
ABE	95N	PRL 74 3538	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	95T	PRL 75 613	F. Abe <i>et al.</i>	(CDF Collab.)
ACCIARRI	95E	PL B350 109	M. Acciarri <i>et al.</i>	(L3 Collab.)
AKERS	95A	ZPHY C65 367	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	95R	ZPHY C67 203	R. Akers <i>et al.</i>	(OPAL Collab.)
BEREZINSKY	95	ASP 5 1	V. Berezinsky <i>et al.</i>	
BUSKULIC	95E	PL B349 238	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
CLAVELLI	95	PR D51 1117	L. Clavelli, P.W. Coulter	(ALAT)
FALK	95	PL B354 99	T. Falk, K.A. Olive, M. Srednicki	(MINN, UCSB)
LOSECCO	95	PL B342 392	J.M. LoSecco	(NDAM)
AKERS	94K	PL B337 207	R. Akers <i>et al.</i>	(OPAL Collab.)
BECK	94	PL B336 141	M. Beck <i>et al.</i>	(MPIH, KIAE, SASSO)
CAKIR	94	PR D50 3268	M.B. Cakir, G.R. Farrar	(RUTG)
FALK	94	PL B339 248	T. Falk, K.A. Olive, M. Srednicki	(UCSB, MINN)
SHIRAI	94	PRL 72 3313	J. Shirai <i>et al.</i>	(VENUS Collab.)
ADRIANI	93M	PRPL 236 1	O. Adriani <i>et al.</i>	(L3 Collab.)
ALITTI	93	NP B400 3	J. Alitti <i>et al.</i>	(UA2 Collab.)
CLAVELLI	93	PR D47 1973	L. Clavelli, P.W. Coulter, K.J. Yuan	(ALAT)
DREES	93	PR D47 376	M. Drees, M.M. Nojiri	(DESY, SLAC)
DREES	93B	PR D48 3483	M. Drees, M.M. Nojiri	
FALK	93	PL B318 354	T. Falk <i>et al.</i>	(UCB, UCSB, MINN)
HEBBEKER	93	ZPHY C60 63	T. Hebbeker	(CERN)

KELLEY	93	PR D47 2461	S. Kelley <i>et al.</i>	(TAMU, ALAH)
LOPEZ	93C	PL B313 241	J.L. Lopez, D.V. Nanopoulos, X. Wang	(TAMU, HARC+)
MIZUTA	93	PL B298 120	S. Mizuta, M. Yamaguchi	(TOHO)
MORI	93	PR D48 5505	M. Mori <i>et al.</i>	(KEK, NIIG, TOKY, TOKA+)
ABE	92L	PRL 69 3439	F. Abe <i>et al.</i>	(CDF Collab.)
BOTTINO	92	MPL A7 733	A. Bottino <i>et al.</i>	(TORI, ZARA)
Also		PL B265 57	A. Bottino <i>et al.</i>	(TORI, INFN)
CLAVELLI	92	PR D46 2112	L. Clavelli	(ALAT)
DECAMP	92	PRPL 216 253	D. Decamp <i>et al.</i>	(ALEPH Collab.)
LOPEZ	92	NP B370 445	J.L. Lopez, D.V. Nanopoulos, K.J. Yuan	(TAMU)
MCDONALD	92	PL B283 80	J. McDonald, K.A. Olive, M. Srednicki	(LISB+)
ROY	92	PL B283 270	D.P. Roy	(CERN)
ABREU	91F	NP B367 511	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKESSON	91	ZPHY C52 219	T. Akesson <i>et al.</i>	(HELIOS Collab.)
ALEXANDER	91F	ZPHY C52 175	G. Alexander <i>et al.</i>	(OPAL Collab.)
ANTONIADIS	91	PL B262 109	I. Antoniadis, J. Ellis, D.V. Nanopoulos	(EPOL+)
BOTTINO	91	PL B265 57	A. Bottino <i>et al.</i>	(TORI, INFN)
GELMINI	91	NP B351 623	G.B. Gelmini, P. Gondolo, E. Roulet	(UCLA, TRST)
GRIEST	91	PR D43 3191	K. Griest, D. Seckel	
KAMIONKOW...	91	PR D44 3021	M. Kamionkowski	(CHIC, FNAL)
MORI	91B	PL B270 89	M. Mori <i>et al.</i>	(Kamiokande Collab.)
NOJIRI	91	PL B261 76	M.M. Nojiri	(KEK)
OLIVE	91	NP B355 208	K.A. Olive, M. Srednicki	(MINN, UCSB)
ROSZKOWSKI	91	PL B262 59	L. Roszkowski	(CERN)
SATO	91	PR D44 2220	N. Sato <i>et al.</i>	(Kamiokande Collab.)
ADACHI	90C	PL B244 352	I. Adachi <i>et al.</i>	(TOPAZ Collab.)
GRIEST	90	PR D41 3565	K. Griest, M. Kamionkowski, M.S. Turner	(UCB+)
BARBIERI	89C	NP B313 725	R. Barbieri, M. Frigeni, G. Giudice	
NAKAMURA	89	PR D39 1261	T.T. Nakamura <i>et al.</i>	(KYOT, TMTTC)
OLIVE	89	PL B230 78	K.A. Olive, M. Srednicki	(MINN, UCSB)
ELLIS	88D	NP B307 883	J. Ellis, R. Flores	
GRIEST	88B	PR D38 2357	K. Griest	
OLIVE	88	PL B205 553	K.A. Olive, M. Srednicki	(MINN, UCSB)
SREDNICKI	88	NP B310 693	M. Srednicki, R. Watkins, K.A. Olive	(MINN, UCSB)
ALBAJAR	87D	PL B198 261	C. Albajar <i>et al.</i>	(UA1 Collab.)
ANSARI	87D	PL B195 613	R. Ansari <i>et al.</i>	(UA2 Collab.)
ARNOLD	87	PL B186 435	R.G. Arnold <i>et al.</i>	(BRUX, DUUC, LOUC+)
NG	87	PL B188 138	K.W. Ng, K.A. Olive, M. Srednicki	(MINN, UCSB)
TUTS	87	PL B186 233	P.M. Tuts <i>et al.</i>	(CUSB Collab.)
ALBRECHT	86C	PL 167B 360	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BADIER	86	ZPHY C31 21	J. Badier <i>et al.</i>	(NA3 Collab.)
BARNETT	86	NP B267 625	R.M. Barnett, H.E. Haber, G.L. Kane	(LBL, UCSC+)
GAISSER	86	PR D34 2206	T.K. Gaisser, G. Steigman, S. Tilav	(BART, DELA)
VOLOSHIN	86	SJNP 43 495	M.B. Voloshin, L.B. Okun	(ITEP)
COOPER-...	85B	PL 160B 212	A.M. Cooper-Sarkar <i>et al.</i>	(WA66 Collab.)
DAWSON	85	PR D31 1581	S. Dawson, E. Eichten, C. Quigg	(LBL, FNAL)
FARRAR	85	PRL 55 895	G.R. Farrar	(RUTG)
GOLDMAN	85	Physica 15D 181	T. Goldman, H.E. Haber	(LANL, UCSC)
HABER	85	PRPL 117 75	H.E. Haber, G.L. Kane	(UCSC, MICH)
BALL	84	PRL 53 1314	R.C. Ball <i>et al.</i>	(MICH, FIRZ, OSU, FNAL+)
BARBER	84B	PL 139B 427	J.S. Barber, R.E. Shrock	(STON)
BRICK	84	PR D30 1134	D.H. Brick <i>et al.</i>	(BROW, CAVE, IIT+)
ELLIS	84	NP B238 453	J. Ellis <i>et al.</i>	(CERN)
FARRAR	84	PRL 53 1029	G.R. Farrar	(RUTG)
BERGSMA	83C	PL 121B 429	F. Bergsma <i>et al.</i>	(CHARM Collab.)
CHANOWITZ	83	PL 126B 225	M.S. Chanowitz, S. Sharpe	(UCB, LBL)
GOLDBERG	83	PRL 50 1419	H. Goldberg	(NEAS)
HOFFMAN	83	PR D28 660	C.M. Hoffman <i>et al.</i>	(LANL, ARZS)
KRAUSS	83	NP B227 556	L.M. Krauss	(HARV)
VYSOTSKII	83	SJNP 37 948	M.I. Vysotsky	(ITEP)
KANE	82	PL 112B 227	G.L. Kane, J.P. Leveille	(MICH)
CABIBBO	81	PL 105B 155	N. Cabibbo, G.R. Farrar, L. Maiani	(ROMA, RUTG)
FARRAR	78	PL 76B 575	G.R. Farrar, P. Fayet	(CIT)
Also		PL 79B 442	G.R. Farrar, P. Fayet	(CIT)