

B[±]/B⁰ ADMIXTURE

B DECAY MODES

The branching fraction measurements are for an admixture of B mesons at the $\Upsilon(4S)$. The values quoted assume that $B(\Upsilon(4S) \rightarrow B\bar{B}) = 100\%$.

For inclusive branching fractions, e.g., $B \rightarrow D^\pm$ anything, the values usually are multiplicities, not branching fractions. They can be greater than one.

\bar{B} modes are charge conjugates of the modes below. Reactions indicate the weak decay vertex and do not include mixing.

Mode	Fraction (Γ_j/Γ)	Scale factor/ Confidence level
Semileptonic and leptonic modes		
Γ_1 $B \rightarrow e^+ \nu_e$ anything	[a] (10.78 ± 0.18) %	
Γ_2 $B \rightarrow \bar{p} e^+ \nu_e$ anything	< 5.9	× 10 ⁻⁴ CL=90%
Γ_3 $B \rightarrow \mu^+ \nu_\mu$ anything	[a] (10.78 ± 0.18) %	
Γ_4 $B \rightarrow \ell^+ \nu_\ell$ anything	[a,b] (10.78 ± 0.18) %	
Γ_5 $B \rightarrow D^- \ell^+ \nu_\ell$ anything	[b] (2.8 ± 0.9) %	
Γ_6 $B \rightarrow \bar{D}^0 \ell^+ \nu_\ell$ anything	[b] (7.2 ± 1.5) %	
Γ_7 $B \rightarrow D^{*-} \ell^+ \nu_\ell$ anything	[c] (6.7 ± 1.3) × 10 ⁻³	
Γ_8 $B \rightarrow D^{*0} \ell^+ \nu_\ell$ anything		
Γ_9 $B \rightarrow \bar{D}^{**} \ell^+ \nu_\ell$	[b,d] (2.7 ± 0.7) %	
Γ_{10} $B \rightarrow \bar{D}_1(2420) \ell^+ \nu_\ell$ anything	(3.8 ± 1.3) × 10 ⁻³	S=2.4
Γ_{11} $B \rightarrow D\pi \ell^+ \nu_\ell$ anything + $D^* \pi \ell^+ \nu_\ell$ anything	(2.6 ± 0.5) %	S=1.5
Γ_{12} $B \rightarrow D\pi \ell^+ \nu_\ell$ anything	(1.5 ± 0.6) %	
Γ_{13} $B \rightarrow D^* \pi \ell^+ \nu_\ell$ anything	(1.9 ± 0.4) %	
Γ_{14} $B \rightarrow \bar{D}_2^*(2460) \ell^+ \nu_\ell$ anything	(4.4 ± 1.6) × 10 ⁻³	
Γ_{15} $B \rightarrow D^{*-} \pi^+ \ell^+ \nu_\ell$ anything	(1.00 ± 0.34) %	
Γ_{16} $B \rightarrow D_s^- \ell^+ \nu_\ell$ anything	[b] < 7	× 10 ⁻³ CL=90%
Γ_{17} $B \rightarrow D_s^- \ell^+ \nu_\ell K^+$ anything	[b] < 5	× 10 ⁻³ CL=90%
Γ_{18} $B \rightarrow D_s^- \ell^+ \nu_\ell K^0$ anything	[b] < 7	× 10 ⁻³ CL=90%
Γ_{19} $B \rightarrow \ell^+ \nu_\ell$ charm	(10.61 ± 0.17) %	
Γ_{20} $B \rightarrow X_u \ell^+ \nu_\ell$	(2.33 ± 0.22) × 10 ⁻³	
Γ_{21} $B \rightarrow K^+ \ell^+ \nu_\ell$ anything	[b] (6.2 ± 0.6) %	
Γ_{22} $B \rightarrow K^- \ell^+ \nu_\ell$ anything	[b] (10 ± 4) × 10 ⁻³	
Γ_{23} $B \rightarrow K^0 / \bar{K}^0 \ell^+ \nu_\ell$ anything	[b] (4.6 ± 0.5) %	

D, D^* , or D_s modes

Γ_{24}	$B \rightarrow D^\pm$ anything	(22.8 \pm 1.4) %	
Γ_{25}	$B \rightarrow D^0 / \bar{D}^0$ anything	(64.0 \pm 3.0) %	S=1.2
Γ_{26}	$B \rightarrow D^*(2010)^\pm$ anything	(22.5 \pm 1.5) %	
Γ_{27}	$B \rightarrow D^*(2007)^0$ anything	(26.0 \pm 2.7) %	
Γ_{28}	$B \rightarrow D_s^\pm$ anything	[e] (8.6 \pm 1.2) %	
Γ_{29}	$B \rightarrow D_s^{*\pm}$ anything	(6.5 \pm 1.2) %	
Γ_{30}	$B \rightarrow D_s^{*\pm} \bar{D}^*$	(3.4 \pm 0.7) %	
Γ_{31}	$B \rightarrow \bar{D} D_{s0}(2317)$		
Γ_{32}	$B \rightarrow \bar{D} D_{sJ}(2457)$		
Γ_{33}	$B \rightarrow D^{(*)} \bar{D}^{(*)} K^0 + D^{(*)} \bar{D}^{(*)} K^\pm$	[e,f] (7.1 \pm 2.7 / -1.7) %	
Γ_{34}	$b \rightarrow c \bar{c} s$	(22 \pm 4) %	
Γ_{35}	$B \rightarrow D_s^{(*)} \bar{D}^*$	[e,f] (4.0 \pm 0.6) %	
Γ_{36}	$B \rightarrow D^* D^*(2010)^\pm$	[e] < 5.9 $\times 10^{-3}$ CL=90%	
Γ_{37}	$B \rightarrow D D^*(2010)^\pm + D^* D^\pm$	[e] < 5.5 $\times 10^{-3}$ CL=90%	
Γ_{38}	$B \rightarrow D D^\pm$	[e] < 3.1 $\times 10^{-3}$ CL=90%	
Γ_{39}	$B \rightarrow D_s^{(*)\pm} \bar{D}^{(*)} X(n\pi^\pm)$	[e,f] (9 \pm 5 / -4) %	
Γ_{40}	$B \rightarrow D^*(2010)\gamma$	< 1.1 $\times 10^{-3}$ CL=90%	
Γ_{41}	$B \rightarrow D_s^+ \pi^-, D_s^{*+} \pi^-, D_s^+ \rho^-, D_s^{*+} \rho^-, D_s^+ \pi^0, D_s^{*+} \pi^0, D_s^+ \eta, D_s^{*+} \eta, D_s^+ \rho^0, D_s^{*+} \rho^0, D_s^+ \omega, D_s^{*+} \omega$	[e] < 4 $\times 10^{-4}$ CL=90%	
Γ_{42}	$B \rightarrow D_{s1}(2536)^+$ anything	< 9.5 $\times 10^{-3}$ CL=90%	

Charmonium modes

Γ_{43}	$B \rightarrow J/\psi(1S)$ anything	(1.094 \pm 0.032) %	S=1.1
Γ_{44}	$B \rightarrow J/\psi(1S)$ (direct) anything	(7.8 \pm 0.4) $\times 10^{-3}$	S=1.1
Γ_{45}	$B \rightarrow \psi(2S)$ anything	(3.07 \pm 0.21) $\times 10^{-3}$	
Γ_{46}	$B \rightarrow \chi_{c1}(1P)$ anything	(3.86 \pm 0.27) $\times 10^{-3}$	
Γ_{47}	$B \rightarrow \chi_{c1}(1P)$ (direct) anything	(3.18 \pm 0.25) $\times 10^{-3}$	
Γ_{48}	$B \rightarrow \chi_{c2}(1P)$ anything	(1.3 \pm 0.4) $\times 10^{-3}$	S=1.9
Γ_{49}	$B \rightarrow \chi_{c2}(1P)$ (direct) anything	(1.65 \pm 0.31) $\times 10^{-3}$	
Γ_{50}	$B \rightarrow \eta_c(1S)$ anything	< 9 $\times 10^{-3}$ CL=90%	
Γ_{51}	$B \rightarrow K Y(3940) \times B(Y(3940) \rightarrow \omega J/\psi)$	[g] (7.1 \pm 3.4) $\times 10^{-5}$	

K or K* modes

Γ_{52}	$B \rightarrow K^\pm$ anything	[e]	(78.9 \pm 2.5) %
Γ_{53}	$B \rightarrow K^+$ anything		(66 \pm 5) %
Γ_{54}	$B \rightarrow K^-$ anything		(13 \pm 4) %
Γ_{55}	$B \rightarrow K^0 / \bar{K}^0$ anything	[e]	(64 \pm 4) %
Γ_{56}	$B \rightarrow K^*(892)^\pm$ anything		(18 \pm 6) %
Γ_{57}	$B \rightarrow K^*(892)^0 / \bar{K}^*(892)^0$ anything	[e]	(14.6 \pm 2.6) %
Γ_{58}	$B \rightarrow K^*(892)\gamma$		(4.2 \pm 0.6) $\times 10^{-5}$
Γ_{59}	$B \rightarrow \eta K \gamma$		(8.5 \pm 1.8 / \pm 1.6) $\times 10^{-6}$
Γ_{60}	$B \rightarrow K_1(1400)\gamma$		< 1.27 $\times 10^{-4}$ CL=90%
Γ_{61}	$B \rightarrow K_2^*(1430)\gamma$		(1.7 \pm 0.6 / \pm 0.5) $\times 10^{-5}$
Γ_{62}	$B \rightarrow K_2(1770)\gamma$		< 1.2 $\times 10^{-3}$ CL=90%
Γ_{63}	$B \rightarrow K_3^*(1780)\gamma$		< 3.7 $\times 10^{-5}$ CL=90%
Γ_{64}	$B \rightarrow K_4^*(2045)\gamma$		< 1.0 $\times 10^{-3}$ CL=90%
Γ_{65}	$B \rightarrow K\eta'(958)$		(8.3 \pm 1.1) $\times 10^{-5}$
Γ_{66}	$B \rightarrow K^*(892)\eta'(958)$		< 2.2 $\times 10^{-5}$ CL=90%
Γ_{67}	$B \rightarrow K\eta$		< 5.2 $\times 10^{-6}$ CL=90%
Γ_{68}	$B \rightarrow K^*(892)\eta$		(1.8 \pm 0.5) $\times 10^{-5}$
Γ_{69}	$B \rightarrow K\phi\phi$		(2.3 \pm 0.9) $\times 10^{-6}$
Γ_{70}	$B \rightarrow \bar{b} \rightarrow \bar{s}\gamma$		(3.43 \pm 0.29) $\times 10^{-4}$
Γ_{71}	$B \rightarrow \bar{b} \rightarrow \bar{s}$ gluon		< 6.8 % CL=90%
Γ_{72}	$B \rightarrow \eta$ anything		< 4.4 $\times 10^{-4}$ CL=90%
Γ_{73}	$B \rightarrow \eta'$ anything		(4.2 \pm 0.9) $\times 10^{-4}$

Light unflavored meson modes

Γ_{74}	$B \rightarrow \rho\gamma$		< 1.9 $\times 10^{-6}$ CL=90%
Γ_{75}	$B \rightarrow \rho/\omega\gamma$		< 1.2 $\times 10^{-6}$ CL=90%
Γ_{76}	$B \rightarrow \pi^\pm$ anything	[e,h]	(358 \pm 7) %
Γ_{77}	$B \rightarrow \pi^0$ anything		(235 \pm 11) %
Γ_{78}	$B \rightarrow \eta$ anything		(17.6 \pm 1.6) %
Γ_{79}	$B \rightarrow \rho^0$ anything		(21 \pm 5) %
Γ_{80}	$B \rightarrow \omega$ anything		< 81 % CL=90%
Γ_{81}	$B \rightarrow \phi$ anything		(3.42 \pm 0.13) %
Γ_{82}	$B \rightarrow \phi K^*(892)$		< 2.2 $\times 10^{-5}$ CL=90%

Baryon modes

Γ_{83}	$B \rightarrow \Lambda_c^+ / \bar{\Lambda}_c^-$ anything		(6.4 \pm 1.1) %
Γ_{84}	$B \rightarrow \Lambda_c^+$ anything		
Γ_{85}	$B \rightarrow \bar{\Lambda}_c^-$ anything		
Γ_{86}	$B \rightarrow \bar{\Lambda}_c^- e^+$ anything		< 3.2 $\times 10^{-3}$ CL=90%
Γ_{87}	$B \rightarrow \bar{\Lambda}_c^- p$ anything		(3.6 \pm 0.7) %
Γ_{88}	$B \rightarrow \bar{\Lambda}_c^- p e^+ \nu_e$		< 1.5 $\times 10^{-3}$ CL=90%

Γ_{89}	$B \rightarrow \bar{\Sigma}_c^{--}$ anything		$(4.2 \pm 2.4) \times 10^{-3}$
Γ_{90}	$B \rightarrow \bar{\Sigma}_c^-$ anything		$< 9.6 \times 10^{-3}$ CL=90%
Γ_{91}	$B \rightarrow \bar{\Sigma}_c^0$ anything		$(4.6 \pm 2.4) \times 10^{-3}$
Γ_{92}	$B \rightarrow \bar{\Sigma}_c^0 N (N = p \text{ or } n)$		$< 1.5 \times 10^{-3}$ CL=90%
Γ_{93}	$B \rightarrow \Xi_c^0$ anything $\times B(\Xi_c^0 \rightarrow \Xi^- \pi^+)$		$(1.93 \pm 0.30) \times 10^{-4}$ S=1.1
Γ_{94}	$B \rightarrow \Xi_c^+$ anything $\times B(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+)$		$(4.5 \pm 1.3) \times 10^{-4}$
Γ_{95}	$B \rightarrow p/\bar{p}$ anything	[e]	$(8.0 \pm 0.4) \%$
Γ_{96}	$B \rightarrow p/\bar{p}$ (direct) anything	[e]	$(5.5 \pm 0.5) \%$
Γ_{97}	$B \rightarrow \Lambda/\bar{\Lambda}$ anything	[e]	$(4.0 \pm 0.5) \%$
Γ_{98}	$B \rightarrow \Lambda$ anything		
Γ_{99}	$B \rightarrow \bar{\Lambda}$ anything		
Γ_{100}	$B \rightarrow \Xi^-/\Xi^+$ anything	[e]	$(2.7 \pm 0.6) \times 10^{-3}$
Γ_{101}	$B \rightarrow$ baryons anything		$(6.8 \pm 0.6) \%$
Γ_{102}	$B \rightarrow p\bar{p}$ anything		$(2.47 \pm 0.23) \%$
Γ_{103}	$B \rightarrow \Lambda\bar{p}/\bar{\Lambda}p$ anything	[e]	$(2.5 \pm 0.4) \%$
Γ_{104}	$B \rightarrow \Lambda\bar{\Lambda}$ anything		$< 5 \times 10^{-3}$ CL=90%

**Lepton Family number (LF) violating modes or
 $\Delta B = 1$ weak neutral current (B1) modes**

Γ_{105}	$B \rightarrow s e^+ e^-$	B1	$(4.7 \pm 1.3) \times 10^{-6}$
Γ_{106}	$B \rightarrow s \mu^+ \mu^-$	B1	$(4.3 \pm 1.2) \times 10^{-6}$
Γ_{107}	$B \rightarrow s \ell^+ \ell^-$	B1 [b]	$(4.5 \pm 1.0) \times 10^{-6}$
Γ_{108}	$B \rightarrow K e^+ e^-$	B1	$(6.0 \pm 1.4) \times 10^{-7}$ S=1.1
Γ_{109}	$B \rightarrow K^*(892) e^+ e^-$	B1	$(1.24 \pm 0.37) \times 10^{-6}$
Γ_{110}	$B \rightarrow K \mu^+ \mu^-$	B1	$(4.7 \pm 1.1) \times 10^{-7}$
Γ_{111}	$B \rightarrow K^*(892) \mu^+ \mu^-$	B1	$(1.19 \pm 0.34) \times 10^{-6}$
Γ_{112}	$B \rightarrow K \ell^+ \ell^-$	B1	$(5.4 \pm 0.8) \times 10^{-7}$
Γ_{113}	$B \rightarrow K^*(892) \ell^+ \ell^-$	B1	$(1.05 \pm 0.20) \times 10^{-6}$
Γ_{114}	$B \rightarrow e^\pm \mu^\mp s$	LF [e]	$< 2.2 \times 10^{-5}$ CL=90%
Γ_{115}	$B \rightarrow \pi e^\pm \mu^\mp$	LF	$< 1.6 \times 10^{-6}$ CL=90%
Γ_{116}	$B \rightarrow \rho e^\pm \mu^\mp$	LF	$< 3.2 \times 10^{-6}$ CL=90%
Γ_{117}	$B \rightarrow K e^\pm \mu^\mp$	LF	$< 1.6 \times 10^{-6}$ CL=90%
Γ_{118}	$B \rightarrow K^*(892) e^\pm \mu^\mp$	LF	$< 6.2 \times 10^{-6}$ CL=90%

[a] These values are model dependent.

[b] An ℓ indicates an e or a μ mode, not a sum over these modes.

[c] Here "anything" means at least one particle observed.

- [d] D^{**} stands for the sum of the $D(1^1P_1)$, $D(1^3P_0)$, $D(1^3P_1)$, $D(1^3P_2)$, $D(2^1S_0)$, and $D(2^1S_1)$ resonances.
- [e] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [f] $D^{(*)}\bar{D}^{(*)}$ stands for the sum of $D^*\bar{D}^*$, $D^*\bar{D}$, $D\bar{D}^*$, and $D\bar{D}$.
- [g] $Y(3940)$ denotes a near-threshold enhancement in the $\omega J/\psi$ mass spectrum.
- [h] Inclusive branching fractions have a multiplicity definition and can be greater than 100%.

B^\pm/B^0 ADMIXTURE BRANCHING RATIOS

$\Gamma(\ell^+ \nu_\ell \text{ anything})/\Gamma_{\text{total}}$ Γ_4/Γ

These branching fraction values are model dependent.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account corrections between the measurements.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.1078 ± 0.0018 OUR EVALUATION

0.1081 ± 0.0014 OUR AVERAGE Includes data from the 2 datablocks that follow this one.

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.108 ± 0.002 ± 0.0056 ¹ HENDERSON 92 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

¹ HENDERSON 92 measurement employs e and μ . The systematic error contains 0.004 in quadrature from model dependence. The authors average a variation of the Isgur, Scora, Grinstein, and Wise model with that of the Altarelli-Cabibbo-Corbò-Maiani-Martinelli model for semileptonic decays to correct the acceptance.

$\Gamma(e^+ \nu_e \text{ anything})/\Gamma_{\text{total}}$ Γ_1/Γ

These branching fraction values are model dependent.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account corrections between the measurements.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

The data in this block is included in the average printed for a previous datablock.

0.1078 ± 0.0018 OUR EVALUATION

0.1081 ± 0.0014 OUR AVERAGE

0.1085 ± 0.0021 ± 0.0036 ² OKABE 05 BELL $e^+e^- \rightarrow \Upsilon(4S)$

0.1083 ± 0.0016 ± 0.0006 ³ AUBERT 04X BABR $e^+e^- \rightarrow \Upsilon(4S)$

0.1091 ± 0.0009 ± 0.0024 ⁴ MAHMOOD 04 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

0.097 ± 0.005 ± 0.004 ⁵ ALBRECHT 93H ARG $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.1036 ± 0.0006 ± 0.0023	⁶ AUBERT,B	04A BABR	$e^+e^- \rightarrow \Upsilon(4S)$
0.1087 ± 0.0018 ± 0.0030	⁷ AUBERT	03 BABR	Repl. by AUBERT 04X
0.109 ± 0.0012 ± 0.0049	⁸ ABE	02Y BELL	Repl. by OKABE 05
0.1049 ± 0.0017 ± 0.0043	⁹ BARISH	96B CLE2	Repl. by MAHMOOD 04
0.100 ± 0.004 ± 0.003	¹⁰ YANAGISAWA	91 CSB2	$e^+e^- \rightarrow \Upsilon(4S)$
0.103 ± 0.006 ± 0.002	¹¹ ALBRECHT	90H ARG	$e^+e^- \rightarrow \Upsilon(4S)$
0.117 ± 0.004 ± 0.010	¹² WACHS	89 CBAL	Direct e at $\Upsilon(4S)$
0.120 ± 0.007 ± 0.005	CHEN	84 CLEO	Direct e at $\Upsilon(4S)$
0.132 ± 0.008 ± 0.014	¹³ KLOPFEN...	83B CUSB	Direct e at $\Upsilon(4S)$

² The measurements are obtained for charged and neutral B mesons partial rates of semileptonic decay to electrons with momentum above 0.6 GeV/c in the B rest frame, and their ratio of $B(B^+ \rightarrow e^+ \nu_e X)/B(B^0 \rightarrow e^+ \nu_e X) = 1.08 \pm 0.05 \pm 0.02$.

³ The semileptonic branching ratio, $|V_{cb}|$ and other heavy-quark parameters are determined from a simultaneous fit to moments of the hadronic-mass and lepton-energy distribution.

⁴ Uses charge and angular correlations in $\Upsilon(4S)$ events with a high-momentum lepton and an additional electron.

⁵ ALBRECHT 93H analysis performed using tagged semileptonic decays of the B . This technique is almost model independent for the lepton branching ratio.

⁶ Uses the high-momentum lepton tag method and requires the electron energy above 0.6 GeV.

⁷ Uses the high-momentum lepton tag method. They also report $|V_{cb}| = 0.0423 \pm 0.0007(\text{exp}) \pm 0.0020(\text{theo.})$.

⁸ Uses the high-momentum lepton tag method. ABE 02Y also reports $|V_{cb}| = 0.0408 \pm 0.0010(\text{exp}) \pm 0.0025(\text{theo.})$. The second error is due to uncertainties of theoretical inputs.

⁹ BARISH 96B analysis performed using tagged semileptonic decays of the B . This technique is almost model independent for the lepton branching ratio.

¹⁰ YANAGISAWA 91 also measures an average semileptonic branching ratio at the $\Upsilon(5S)$ of 9.6–10.5% depending on assumptions about the relative production of different B meson species.

¹¹ ALBRECHT 90H uses the model of ALTARELLI 82 to correct over all lepton momenta. 0.099 ± 0.006 is obtained using ISGUR 89B.

¹² Using data above $p(e) = 2.4$ GeV, WACHS 89 determine $\sigma(B \rightarrow e \nu \text{up})/\sigma(B \rightarrow e \nu \text{charm}) < 0.065$ at 90% CL.

¹³ Ratio $\sigma(b \rightarrow e \nu \text{up})/\sigma(b \rightarrow e \nu \text{charm}) < 0.055$ at CL = 90%.

$\Gamma(\mu^+ \nu_\mu \text{ anything})/\Gamma_{\text{total}}$

Γ_3/Γ

These branching fraction values are model dependent.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account corrections between the measurements.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

The data in this block is included in the average printed for a previous datablock.

0.1078 ± 0.0018 OUR EVALUATION

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.100 ± 0.006 ± 0.002	¹⁴ ALBRECHT	90H ARG	$e^+e^- \rightarrow \Upsilon(4S)$
0.108 ± 0.006 ± 0.01	CHEN	84 CLEO	Direct μ at $\Upsilon(4S)$
0.112 ± 0.009 ± 0.01	LEVMAN	84 CUSB	Direct μ at $\Upsilon(4S)$

¹⁴ ALBRECHT 90H uses the model of ALTARELLI 82 to correct over all lepton momenta. 0.097 ± 0.006 is obtained using ISGUR 89B.

$\Gamma(\bar{p}e^+\nu_e \text{ anything})/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.9 $\times 10^{-4}$	90	¹⁵ ADAM	03B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.0016	90	ALBRECHT	90H ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹⁵ Based on $V-A$ model.

$\Gamma(D^-\ell^+\nu_\ell \text{ anything})/\Gamma(\ell^+\nu_\ell \text{ anything})$ Γ_5/Γ_4

$\ell = e \text{ or } \mu.$

VALUE	DOCUMENT ID	TECN	COMMENT
0.26 ± 0.07 ± 0.04	¹⁶ FULTON	91 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹⁶ FULTON 91 uses $B(D^+ \rightarrow K^-\pi^+\pi^+) = (9.1 \pm 1.3 \pm 0.4)\%$ as measured by MARK III.

$\Gamma(\bar{D}^0\ell^+\nu_\ell \text{ anything})/\Gamma(\ell^+\nu_\ell \text{ anything})$ Γ_6/Γ_4

$\ell = e \text{ or } \mu.$

VALUE	DOCUMENT ID	TECN	COMMENT
0.67 ± 0.09 ± 0.10	¹⁷ FULTON	91 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹⁷ FULTON 91 uses $B(D^0 \rightarrow K^-\pi^+) = (4.2 \pm 0.4 \pm 0.4)\%$ as measured by MARK III.

$\Gamma(D^{*-}\ell^+\nu_\ell \text{ anything})/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
0.67 ± 0.08 ± 0.10	ABDALLAH	04D DLPH	$e^+e^- \rightarrow Z^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.3 ± 0.1	¹⁸ BARISH	95 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
¹⁸ BARISH 95 use $B(D^0 \rightarrow K^-\pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$ and $B(D^{*+} \rightarrow D^0\pi^+) = (68.1 \pm 1.0 \pm 1.3)\%$.			

$\Gamma(D^{*0}\ell^+\nu_\ell \text{ anything})/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
0.6 ± 0.6 ± 0.1	¹⁹ BARISH	95 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹⁹ BARISH 95 use $B(D^0 \rightarrow K^-\pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$, $B(D^{*+} \rightarrow D^0\pi^+) = (68.1 \pm 1.0 \pm 1.3)\%$, $B(D^{*0} \rightarrow D^0\pi^0) = (63.6 \pm 2.3 \pm 3.3)\%$.

$\Gamma(\bar{D}^{**}\ell^+\nu_\ell)/\Gamma_{\text{total}}$ Γ_9/Γ

D^{**} stands for the sum of the $D(1^1P_1)$, $D(1^3P_0)$, $D(1^3P_1)$, $D(1^3P_2)$, $D(2^1S_0)$, and $D(2^1S_1)$ resonances. $\ell = e \text{ or } \mu$, not sum over e and μ modes.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.027 ± 0.005 ± 0.005	63	²⁰	ALBRECHT	93 ARG	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.028	95	²¹	BARISH	95 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
--------	----	---------------	--------	---------	-----------------------------------

- ²⁰ ALBRECHT 93 assumes the GISW model to correct for unseen modes. Using the BHKT model, the result becomes $0.023 \pm 0.006 \pm 0.004$. Assumes $B(D^{*+} \rightarrow D^0 \pi^+) = 68.1\%$, $B(D^0 \rightarrow K^- \pi^+) = 3.65\%$, $B(D^0 \rightarrow K^- \pi^+ \pi^- \pi^+) = 7.5\%$. We have taken their average e and μ value.
- ²¹ BARISH 95 use $B(D^0 \rightarrow K^- \pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$, assume all nonresonant channels are zero, and use GISW model for relative abundances of D^{**} states.

$\Gamma(\bar{D}_1(2420)\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0038 ± 0.0013 OUR AVERAGE	Error includes scale factor of 2.4.		
0.0033 ± 0.0006	²² ABAZOV	05O D0	$p\bar{p}$ at 1.96 TeV
0.0074 ± 0.0016	²³ BUSKULIC	97B ALEP	$e^+e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen ²⁴ BUSKULIC 95B ALEP Repl. by BUSKULIC 97B

²² Assumes $B(D_1 \rightarrow D^* \pi) = 1$, $B(D_1 \rightarrow D^* \pi^\pm) = 2/3$, and $B(b \rightarrow B) = 0.397$.

²³ BUSKULIC 97B assumes $B(D_1(2420) \rightarrow D^* \pi) = 1$, $B(D_1(2420) \rightarrow D^* \pi^\pm) = 2/3$, and $B(b \rightarrow B) = 0.378 \pm 0.022$.

²⁴ BUSKULIC 95B reports $f_B \times B(B \rightarrow \bar{D}_1(2420)^0 \ell^+ \nu_\ell \text{anything}) \times B(\bar{D}_1(2420)^0 \rightarrow \bar{D}^*(2010)^- \pi^+) = (2.04 \pm 0.58 \pm 0.34)10^{-3}$, where f_B is the production fraction for a single B charge state.

$[\Gamma(D\pi\ell^+\nu_\ell\text{anything}) + \Gamma(D^*\pi\ell^+\nu_\ell\text{anything})]/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.026 ± 0.005 OUR AVERAGE	Error includes scale factor of 1.5.		
0.0340 ± 0.0052 ± 0.0032	²⁵ ABREU	00R DLPH	$e^+e^- \rightarrow Z$
0.0226 ± 0.0029 ± 0.0033	²⁶ BUSKULIC	97B ALEP	$e^+e^- \rightarrow Z$

²⁵ Assumes no contribution from B_s and b baryons. Further assumes contributions from single pion ($D\pi$ and $D^*\pi$) states only, allowing isospin conservation to relate the relative π^0 and π^+ rates.

²⁶ BUSKULIC 97B assumes $B(b \rightarrow B) = 0.378 \pm 0.022$ and uses isospin invariance by assuming that all observed $D^0 \pi^+$, $D^{*0} \pi^+$, $D^+ \pi^-$, and $D^{*+} \pi^-$ are from D^{**} states. A correction has been applied to account for the production of B_s^0 and Λ_b^0 .

$\Gamma(D\pi\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0154 ± 0.0061	ABREU	00R DLPH	$e^+e^- \rightarrow Z$

$\Gamma(D^*\pi\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0186 ± 0.0038	ABREU	00R DLPH	$e^+e^- \rightarrow Z$

$\Gamma(\bar{D}_2^*(2460)\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.0044 ± 0.0016		²⁷ ABAZOV	05O D0	$p\bar{p}$ at 1.96 TeV
<0.0065	95	²⁸ BUSKULIC	97B ALEP	$e^+e^- \rightarrow Z$
not seen		²⁹ BUSKULIC	95B ALEP	$e^+e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

²⁷ Assumes $B(D_2^* \rightarrow D^* \pi^\pm) = 0.30 \pm 0.06$ and $B(b \rightarrow B) = 0.397$.

²⁸ A revised number based on BUSKULIC 97B which assumes $B(D_2^*(2460) \rightarrow D^* \pi^\pm) = 0.20$ and $B(b \rightarrow B) = 0.378 \pm 0.022$.

²⁹ BUSKULIC 95B reports $f_B \times B(B \rightarrow \bar{D}_2^*(2460)^0 \ell^+ \nu_\ell \text{ anything}) \times B(\bar{D}_2^*(2460)^0 \rightarrow \bar{D}^*(2010)^- \pi^+) \leq 0.81 \times 10^{-3}$ at CL=95%, where f_B is the production fraction for a single B charge state.

$$\frac{\Gamma(B \rightarrow \bar{D}_2^*(2460) \ell^+ \nu_\ell \text{ anything}) \times B(D_2^*(2460) \rightarrow D^{*-} \pi^+)}{\Gamma(B \rightarrow \bar{D}_1(2420) \ell^+ \nu_\ell \text{ anything}) \times B(\bar{D}_1(2420) \rightarrow D^{*-} \pi^+)}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.39±0.09±0.12	ABAZOV	050 D0	$p\bar{p}$ at 1.96 TeV

$$\Gamma(D^{*-} \pi^+ \ell^+ \nu_\ell \text{ anything}) / \Gamma_{\text{total}} \quad \Gamma_{15} / \Gamma$$

Includes resonant and nonresonant contributions.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
10.0±2.7±2.1	³⁰ BUSKULIC	95B ALEP	$e^+ e^- \rightarrow Z$

³⁰ BUSKULIC 95B reports $f_B \times B(B \rightarrow \bar{D}^*(2010)^- \pi^+ \ell^+ \nu_\ell \text{ anything}) = (3.7 \pm 1.0 \pm 0.7) 10^{-3}$. Above value assumes $f_B = 0.37 \pm 0.03$.

$$\Gamma(D_s^- \ell^+ \nu_\ell \text{ anything}) / \Gamma_{\text{total}} \quad \Gamma_{16} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.007	90	³¹ ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

³¹ ALBRECHT 93E reports < 0.012 for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.044$.

$$\Gamma(D_s^- \ell^+ \nu_\ell K^+ \text{ anything}) / \Gamma_{\text{total}} \quad \Gamma_{17} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.005	90	³² ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

³² ALBRECHT 93E reports < 0.008 for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.044$.

$$\Gamma(D_s^- \ell^+ \nu_\ell K^0 \text{ anything}) / \Gamma_{\text{total}} \quad \Gamma_{18} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.007	90	³³ ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

³³ ALBRECHT 93E reports < 0.012 for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.044$.

$$\Gamma(\ell^+ \nu_\ell \text{ charm}) / \Gamma_{\text{total}} \quad \Gamma_{19} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.1061±0.0016±0.0006	³⁴ AUBERT	04X BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

³⁴ The semileptonic branching ratio, $|V_{cb}|$ and other heavy-quark parameters are determined from a simultaneous fit to moments of the hadronic-mass and lepton-energy distribution.

$\Gamma(X_u \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_{20} / Γ

VALUE (units 10^{-3}) DOCUMENT ID TECN COMMENT

2.33 ± 0.22 OUR AVERAGE

2.27 ± 0.26 ^{+0.37} _{-0.33}	35	AUBERT	06H	BABR	$e^+ e^- \rightarrow \gamma(4S)$
2.53 ± 0.24 ± 0.24	36	AUBERT,B	05X	BABR	$e^+ e^- \rightarrow \gamma(4S)$
2.80 ± 0.52 ± 0.41	37	LIMOSANI	05	BELL	$e^+ e^- \rightarrow \gamma(4S)$
1.77 ± 0.29 ± 0.38	38	BORNHEIM	02	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.24 ± 0.27 ± 0.47	39,40	AUBERT	04I	BABR	Repl. by AUBERT,B 05X
--------------------	-------	--------	-----	------	-----------------------

³⁵ Obtained from the partial rate $\Delta B = (0.572 \pm 0.041 \pm 0.065) \times 10^{-3}$ for the electron momentum interval of 2.0–2.6 GeV/c based on BLNP method.

³⁶ Determined from the partial rate $\Delta B = (3.54 \pm 0.33 \pm 0.34) \times 10^{-4}$ measured for electron energy > 2 GeV and hadronic mass squared < 3.5 GeV², and calculated acceptance 0.14 in that region. The V_{ub} is measured as $(3.95 \pm 0.26^{+0.63}_{-0.49}) \times 10^{-3}$.

³⁷ Uses electrons in the momentum interval 1.9–2.6 GeV/c in the center-of-mass frame. The V_{ub} is found to be $(5.08 \pm 0.47^{+0.49}_{-0.48}) \times 10^{-3}$.

³⁸ BORNHEIM 02 uses the observed yield of leptons from semileptonic B decays in the end-point momentum interval 2.2–2.6 GeV/c with recent CLEO-2 data on $B \rightarrow X_S \gamma$. The V_{ub} is found to be $(4.08 \pm 0.34 \pm 0.53) \times 10^{-3}$.

³⁹ Used BaBar measurement of Semileptonic branching fraction $B(B \rightarrow X \ell \nu_\ell) = (10.87 \pm 0.18 \pm 0.30)\%$ to convert the ratio of rates to branching fraction.

⁴⁰ The third error includes the systematics and theoretical errors summed in quadrature.

$\Gamma(X_u \ell^+ \nu_\ell) / \Gamma(\ell^+ \nu_\ell \text{ anything})$ Γ_{20} / Γ_4

ℓ denotes e or μ , not the sum. These experiments measure this ratio in very limited momentum intervals.

VALUE (units 10^{-2}) CL% EVTS DOCUMENT ID TECN COMMENT

2.06 ± 0.25 ± 0.42

			41	AUBERT	04I	BABR	$e^+ e^- \rightarrow \gamma(4S)$	
				42	ALBRECHT	94C	ARG	$e^+ e^- \rightarrow \gamma(4S)$
		107		43	BARTELT	93B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
		77		44	ALBRECHT	91C	ARG	$e^+ e^- \rightarrow \gamma(4S)$
		41		45	ALBRECHT	90	ARG	$e^+ e^- \rightarrow \gamma(4S)$
		76		46	FULTON	90	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
<4.0	90			47	BEHRENDIS	87	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
<4.0	90				CHEN	84	CLEO	Direct e at $\gamma(4S)$
<5.5	90				KLOPFEN...	83B	CUSB	Direct e at $\gamma(4S)$

⁴¹ The third error includes the systematics and theoretical errors summed in quadrature.

⁴² ALBRECHT 94C find $\Gamma(b \rightarrow c) / \Gamma(b \rightarrow \text{all}) = 0.99 \pm 0.02 \pm 0.04$.

⁴³ BARTELT 93B (CLEO II) measures an excess of $107 \pm 15 \pm 11$ leptons in the lepton momentum interval 2.3–2.6 GeV/c which is attributed to $b \rightarrow u \ell \nu_\ell$. This corresponds to a model-dependent partial branching ratio ΔB_{ub} between $(1.15 \pm 0.16 \pm 0.15) \times 10^{-4}$, as evaluated using the KS model (KOERNER 88), and $(1.54 \pm 0.22 \pm 0.20) \times 10^{-4}$ using the ACCMM model (ARTUSO 93). The corresponding values of $|V_{ub}|/|V_{cb}|$ are 0.056 ± 0.006 and 0.076 ± 0.008 , respectively.

⁴⁴ ALBRECHT 91C result supersedes ALBRECHT 90. Two events are fully reconstructed providing evidence for the $b \rightarrow u$ transition. Using the model of ALTARELLI 82, they obtain $|V_{ub}/V_{cb}| = 0.11 \pm 0.012$ from 77 leptons in the 2.3–2.6 GeV momentum range.

⁴⁵ ALBRECHT 90 observes 41 ± 10 excess e and μ (lepton) events in the momentum interval $p = 2.3\text{--}2.6$ GeV signaling the presence of the $b \rightarrow u$ transition. The events correspond to a model-dependent measurement of $|V_{ub}/V_{cb}| = 0.10 \pm 0.01$.

⁴⁶ FULTON 90 observe 76 ± 20 excess e and μ (lepton) events in the momentum interval $p = 2.4\text{--}2.6$ GeV signaling the presence of the $b \rightarrow u$ transition. The average branching ratio, $(1.8 \pm 0.4 \pm 0.3) \times 10^{-4}$, corresponds to a model-dependent measurement of approximately $|V_{ub}/V_{cb}| = 0.1$ using $B(b \rightarrow c\ell\nu) = 10.2 \pm 0.2 \pm 0.7\%$.

⁴⁷ The quoted possible limits range from 0.018 to 0.04 for the ratio, depending on which model or momentum range is chosen. We select the most conservative limit they have calculated. This corresponds to a limit on $|V_{ub}|/|V_{cb}| < 0.20$. While the endpoint technique employed is more robust than their previous results in CHEN 84, these results do not provide a numerical improvement in the limit.

$\Gamma(K^+ \ell^+ \nu_\ell \text{ anything}) / \Gamma(\ell^+ \nu_\ell \text{ anything})$ Γ_{21}/Γ_4

ℓ denotes e or μ , not the sum.

VALUE	DOCUMENT ID	TECN	COMMENT
0.58 ± 0.05 OUR AVERAGE			
0.594 ± 0.021 ± 0.056	ALBRECHT	94C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.54 ± 0.07 ± 0.06	⁴⁸ ALAM	87B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

⁴⁸ ALAM 87B measurement relies on lepton-kaon correlations.

$\Gamma(K^- \ell^+ \nu_\ell \text{ anything}) / \Gamma(\ell^+ \nu_\ell \text{ anything})$ Γ_{22}/Γ_4

ℓ denotes e or μ , not the sum.

VALUE	DOCUMENT ID	TECN	COMMENT
0.092 ± 0.035 OUR AVERAGE			
0.086 ± 0.011 ± 0.044	ALBRECHT	94C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.10 ± 0.05 ± 0.02	⁴⁹ ALAM	87B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

⁴⁹ ALAM 87B measurement relies on lepton-kaon correlations.

$\Gamma(K^0 / \bar{K}^0 \ell^+ \nu_\ell \text{ anything}) / \Gamma(\ell^+ \nu_\ell \text{ anything})$ Γ_{23}/Γ_4

ℓ denotes e or μ , not the sum. Sum over K^0 and \bar{K}^0 states.

VALUE	DOCUMENT ID	TECN	COMMENT
0.42 ± 0.05 OUR AVERAGE			
0.452 ± 0.038 ± 0.056	⁵⁰ ALBRECHT	94C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.39 ± 0.06 ± 0.04	⁵¹ ALAM	87B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

⁵⁰ ALBRECHT 94C assume a K^0/\bar{K}^0 multiplicity twice that of K_S^0 .

⁵¹ ALAM 87B measurement relies on lepton-kaon correlations.

$\langle n_c \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
1.10 ± 0.05	⁵² GIBBONS	97B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.98 ± 0.16 ± 0.12	⁵³ ALAM	87B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
--------------------	--------------------	----------	------------------------------------

⁵² GIBBONS 97B from charm counting using $B(D_s^+ \rightarrow \phi\pi) = 0.036 \pm 0.009$ and $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = 0.044 \pm 0.006$.

⁵³ From the difference between K^- and K^+ widths. ALAM 87B measurement relies on lepton-kaon correlations. It does not consider the possibility of $B\bar{B}$ mixing. We have thus removed it from the average.

$\Gamma(D^\pm \text{ anything})/\Gamma_{\text{total}}$ Γ_{24}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.228±0.014 OUR AVERAGE				
0.227±0.012±0.008		54 GIBBONS	97B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.24 ±0.04 ±0.01		55 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.22 ±0.05 ±0.01		56 ALBRECHT	91H ARG	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.20 ±0.05 ±0.01	20k	57 BORTOLETTO87	CLEO	Sup. by BORTOLETTO 92
<p>54 GIBBONS 97B reports $[B(B \rightarrow D^\pm \text{ anything}) \times B(D^+ \rightarrow K^- \pi^+ \pi^+)] = 0.0216 \pm 0.0008 \pm 0.00082$. We divide by our best value $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.51 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p> <p>55 BORTOLETTO 92 reports $[B(B \rightarrow D^\pm \text{ anything}) \times B(D^+ \rightarrow K^- \pi^+ \pi^+)] = 0.0226 \pm 0.0030 \pm 0.0018$. We divide by our best value $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.51 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p> <p>56 ALBRECHT 91H reports $[B(B \rightarrow D^\pm \text{ anything}) \times B(D^+ \rightarrow K^- \pi^+ \pi^+)] = 0.0209 \pm 0.0027 \pm 0.0040$. We divide by our best value $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.51 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p> <p>57 BORTOLETTO 87 reports $[B(B \rightarrow D^\pm \text{ anything}) \times B(D^+ \rightarrow K^- \pi^+ \pi^+)] = 0.019 \pm 0.004 \pm 0.002$. We divide by our best value $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.51 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

$\Gamma(D^0/\bar{D}^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{25}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.640±0.030 OUR AVERAGE				
Error includes scale factor of 1.2.				
0.661±0.025±0.012		58 GIBBONS	97B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.61 ±0.05 ±0.01		59 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.51 ±0.08 ±0.01		60 ALBRECHT	91H ARG	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.55 ±0.07 ±0.01	21k	61 BORTOLETTO87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.63 ±0.19 ±0.01		62 GREEN	83 CLEO	Repl. by BORTOLETTO 87
<p>58 GIBBONS 97B reports $[B(B \rightarrow D^0/\bar{D}^0 \text{ anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.0251 \pm 0.0006 \pm 0.00075$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.80 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p> <p>59 BORTOLETTO 92 reports $[B(B \rightarrow D^0/\bar{D}^0 \text{ anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.0233 \pm 0.0012 \pm 0.0014$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.80 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p> <p>60 ALBRECHT 91H reports $[B(B \rightarrow D^0/\bar{D}^0 \text{ anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.0194 \pm 0.0015 \pm 0.0025$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.80 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p> <p>61 BORTOLETTO 87 reports $[B(B \rightarrow D^0/\bar{D}^0 \text{ anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.0210 \pm 0.0015 \pm 0.0021$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.80 \pm 0.07) \times 10^{-2}$.</p>				

Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶² GREEN 83 reports $[B(B \rightarrow D^0/\bar{D}^0 \text{ anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.024 \pm 0.006 \pm 0.004$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.80 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^*(2010)^\pm \text{ anything})/\Gamma_{\text{total}}$ **Γ_{26}/Γ**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.225 ± 0.015 OUR AVERAGE				
0.247 ± 0.019 ± 0.01		63 GIBBONS	97B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.205 ± 0.019 ± 0.007		64 ALBRECHT	96D ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.230 ± 0.028 ± 0.009		65 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.283 ± 0.053 ± 0.002		66 ALBRECHT	91H ARG	Sup. by AL-BRECHT 96D
0.22 ± 0.04 ^{+0.07} / _{-0.04}	5200	67 BORTOLETTO87	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
0.27 ± 0.06 ^{+0.08} / _{-0.06}	510	68 CSORNA	85 CLEO	Repl. by BORTOLETTO 87

⁶³ GIBBONS 97B reports $B(B \rightarrow D^*(2010)^+ \text{ anything}) = 0.239 \pm 0.015 \pm 0.014 \pm 0.009$ using CLEO measured D and D^* branching fractions. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶⁴ ALBRECHT 96D reports $B(B \rightarrow D^*(2010)^+ \text{ anything}) = 0.196 \pm 0.019$ using CLEO measured $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.681 \pm 0.01 \pm 0.013$, $B(D^0 \rightarrow K^- \pi^+) = 0.0401 \pm 0.0014$, $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-) = 0.081 \pm 0.005$. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶⁵ BORTOLETTO 92 reports $B(B \rightarrow D^*(2010)^+ \text{ anything}) = 0.25 \pm 0.03 \pm 0.04$ using MARK II $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$ and $B(D^0 \rightarrow K^- \pi^+) = 0.042 \pm 0.008$. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶⁶ ALBRECHT 91H reports $0.348 \pm 0.060 \pm 0.035$ for $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.55 \pm 0.04$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Uses the PDG 90 $B(D^0 \rightarrow K^- \pi^+) = 0.0371 \pm 0.0025$.

⁶⁷ BORTOLETTO 87 uses old MARK III (BALTRUSAITIS 86E) branching ratios $B(D^0 \rightarrow K^- \pi^+) = 0.056 \pm 0.004 \pm 0.003$ and also assumes $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.60^{+0.08}_{-0.15}$. The product branching ratio for $B(B \rightarrow D^*(2010)^+) B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ is $0.13 \pm 0.02 \pm 0.012$. Superseded by BORTOLETTO 92.

⁶⁸ $V-A$ momentum spectrum used to extrapolate below $p = 1$ GeV. We correct the value assuming $B(D^0 \rightarrow K^- \pi^+) = 0.042 \pm 0.006$ and $B(D^{*+} \rightarrow D^0 \pi^+) = 0.6^{+0.08}_{-0.15}$. The product branching fraction is $B(B \rightarrow D^{*+} X) \cdot B(D^{*+} \rightarrow \pi^+ D^0) \cdot B(D^0 \rightarrow K^- \pi^+) = (68 \pm 15 \pm 9) \times 10^{-4}$.

$$\Gamma(D^*(2007)^0 \text{ anything})/\Gamma_{\text{total}} \qquad \Gamma_{27}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.260±0.023±0.015	⁶⁹ GIBBONS	97B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

⁶⁹ GIBBONS 97B reports $B(B \rightarrow D^*(2007)^0 \text{ anything}) = 0.247 \pm 0.012 \pm 0.018 \pm 0.018$ using CLEO measured D and D^* branching fractions. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(D_s^\pm \text{ anything})/\Gamma_{\text{total}} \qquad \Gamma_{28}/\Gamma$$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.086±0.012 OUR AVERAGE				

0.089±0.005±0.012 ⁷⁰ AUBERT 02G BABR $e^+e^- \rightarrow \Upsilon(4S)$

0.096±0.008±0.013 ⁷¹ GIBAUT 96 CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

0.066±0.011±0.009 ⁷² ALBRECHT 92G ARG $e^+e^- \rightarrow \Upsilon(4S)$

0.070±0.011±0.009 257 ⁷³ BORTOLETTO90 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

0.086±0.023±0.011 ⁷⁴ HAAS 86 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.095±0.025±0.012 ⁷⁵ ALBRECHT 87H ARG $e^+e^- \rightarrow \Upsilon(4S)$

⁷⁰ AUBERT 02G reports $[B(B \rightarrow D_s^\pm \text{ anything}) \times B(D_s^+ \rightarrow \phi\pi^+)] = 0.00393 \pm 0.00007 \pm 0.00021$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁷¹ GIBAUT 96 reports $0.1211 \pm 0.0039 \pm 0.0088$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁷² ALBRECHT 92G reports $[B(B \rightarrow D_s^\pm \text{ anything}) \times B(D_s^+ \rightarrow \phi\pi^+)] = 0.00292 \pm 0.00039 \pm 0.00031$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁷³ BORTOLETTO 90 reports $[B(B \rightarrow D_s^\pm \text{ anything}) \times B(D_s^+ \rightarrow \phi\pi^+)] = 0.00306 \pm 0.00047$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁷⁴ HAAS 86 reports $[B(B \rightarrow D_s^\pm \text{ anything}) \times B(D_s^+ \rightarrow \phi\pi^+)] = 0.0038 \pm 0.0010$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. 64 ± 22% decays are 2-body.

⁷⁵ ALBRECHT 87H reports $[B(B \rightarrow D_s^\pm \text{ anything}) \times B(D_s^+ \rightarrow \phi\pi^+)] = 0.0042 \pm 0.0009 \pm 0.0006$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. 46 ± 16% of $B \rightarrow D_s X$ decays are 2-body. Superseded by ALBRECHT 92G.

$$\Gamma(D_s^{*\pm} \text{ anything})/\Gamma_{\text{total}} \qquad \Gamma_{29}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.065±0.009±0.008	⁷⁶ AUBERT	02G BABR	$e^+e^- \rightarrow \Upsilon(4S)$

⁷⁶ AUBERT 02G reports $[B(B \rightarrow D_s^{*\pm} \text{anything}) \times B(D_s^+ \rightarrow \phi\pi^+)] = 0.00284 \pm 0.00029 \pm 0.00025$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^{*\pm} \bar{D}^{(*)}) / \Gamma(D_s^{*\pm} \text{anything})$ $\Gamma_{30} / \Gamma_{29}$
 Sum over modes

VALUE	DOCUMENT ID	TECN	COMMENT
0.533 ± 0.037 ± 0.037	AUBERT	02G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(\bar{D} D_{s0}(2317)) / \Gamma_{\text{total}}$ Γ_{31} / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	⁷⁷ KROKOVNY	03B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

⁷⁷ The product branching ratio for $B(B \rightarrow \bar{D} D_{s0}(2317)^+) \times B(D_{s0}(2317)^+ \rightarrow D_s \pi^0)$ is measured to be $(8.5_{-1.9}^{+2.1} \pm 2.6) \times 10^{-4}$.

$\Gamma(\bar{D} D_{sJ}(2457)) / \Gamma_{\text{total}}$ Γ_{32} / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	⁷⁸ KROKOVNY	03B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

⁷⁸ The product branching ratio for $B(B \rightarrow \bar{D} D_{sJ}(2457)^+) \times B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \pi^0, D_s^+ \gamma)$ are measured to be $(17.8_{-3.9}^{+4.5} \pm 5.3) \times 10^{-4}$ and $(6.7_{-1.2}^{+1.3} \pm 2.0) \times 10^{-4}$, respectively.

$[\Gamma(D^{(*)} \bar{D}^{(*)} K^0) + \Gamma(D^{(*)} \bar{D}^{(*)} K^\pm)] / \Gamma_{\text{total}}$ Γ_{33} / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.071 +0.025 +0.010 -0.015 -0.009	⁷⁹ BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

⁷⁹ The systematic error includes the uncertainties due to the charm branching ratios.

$\Gamma(c\bar{c}s) / \Gamma_{\text{total}}$ Γ_{34} / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.219 ± 0.037	⁸⁰ COAN	98 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

⁸⁰ COAN 98 uses D - ℓ correlation.

$\Gamma(D_s^{(*)} \bar{D}^{(*)}) / \Gamma(D_s^\pm \text{anything})$ $\Gamma_{35} / \Gamma_{28}$
 Sum over modes.

VALUE	DOCUMENT ID	TECN	COMMENT
0.469 ± 0.017 OUR AVERAGE			
0.464 ± 0.013 ± 0.015	AUBERT	02G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.56 +0.21 +0.09 -0.15 -0.08	⁸¹ BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$
0.457 ± 0.019 ± 0.037	GIBAUT	96 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.58 ± 0.07 ± 0.09	ALBRECHT	92G ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.56 ± 0.10	BORTOLETTO90	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

⁸¹ BARATE 98Q measures $B(B \rightarrow D_s^{(*)} \bar{D}^{(*)}) = 0.056_{-0.015}^{+0.021} + 0.009_{-0.008} + 0.019_{-0.011}$, where the third error results from the uncertainty on the different D branching ratios and is dominated by the uncertainty on $B(D_s^+ \rightarrow \phi\pi^+)$. We divide $B(B \rightarrow D_s^{(*)} \bar{D}^{(*)})$ by our best value of $B(B \rightarrow D_s \text{anything}) = 0.1 \pm 0.025$.

$\Gamma(D^* D^*(2010)^\pm)/\Gamma_{\text{total}}$					Γ_{36}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<5.9 \times 10^{-3}$	90	BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$	

$[\Gamma(D D^*(2010)^\pm) + \Gamma(D^* D^\pm)]/\Gamma_{\text{total}}$					Γ_{37}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<5.5 \times 10^{-3}$	90	BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$	

$\Gamma(D D^\pm)/\Gamma_{\text{total}}$					Γ_{38}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<3.1 \times 10^{-3}$	90	BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$	

$\Gamma(D_s^{(*)\pm} \bar{D}^{(*)} \chi(n\pi^\pm))/\Gamma_{\text{total}}$					Γ_{39}/Γ
VALUE		DOCUMENT ID	TECN	COMMENT	
$0.094^{+0.040+0.034}_{-0.031-0.024}$		⁸² BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$	

⁸² The systematic error includes the uncertainties due to the charm branching ratios.

$\Gamma(D^*(2010)\gamma)/\Gamma_{\text{total}}$					Γ_{40}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.1 \times 10^{-3}$	90	⁸³ LESIAK	92 CBAL	$e^+ e^- \rightarrow \Upsilon(4S)$	

⁸³ LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s-quark hadronization.

$\Gamma(D_s^+ \pi^-, D_s^{*+} \pi^-, D_s^+ \rho^-, D_s^{*+} \rho^-, D_s^+ \pi^0, D_s^{*+} \pi^0, D_s^+ \eta, D_s^{*+} \eta, D_s^+ \rho^0, D_s^{*+} \rho^0, D_s^+ \omega, D_s^{*+} \omega)/\Gamma_{\text{total}}$					Γ_{41}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0004	90	⁸⁴ ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$	

Sum over modes.

⁸⁴ ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$. This branching ratio limit provides a model-dependent upper limit $|V_{ub}|/|V_{cb}| < 0.16$ at CL=90%.

$\Gamma(D_{s1}(2536)^+ \text{ anything})/\Gamma_{\text{total}}$					Γ_{42}/Γ
---	--	--	--	--	----------------------

$D_{s1}(2536)^+$ is the narrow *P*-wave D_s^+ meson with $J^P = 1^+$.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0095	90	⁸⁵ BISHAI	98 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$	

⁸⁵ Assuming factorization, the decay constant $f_{D_{s1}^+}$ is at least a factor of 2.5 times smaller than $f_{D_s^+}$.

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.094 ± 0.032 OUR AVERAGE		Error includes scale factor of 1.1.		
1.057 ± 0.012 ± 0.040		⁸⁶ AUBERT	03F BABR	$e^+e^- \rightarrow \Upsilon(4S)$
1.121 ± 0.013 ± 0.042		ANDERSON	02 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
1.30 ± 0.45 ± 0.01	27	⁸⁷ MASCHMANN	90 CBAL	$e^+e^- \rightarrow \Upsilon(4S)$
1.24 ± 0.27 ± 0.01	120	⁸⁸ ALBRECHT	87D ARG	$e^+e^- \rightarrow \Upsilon(4S)$
1.36 ± 0.24 ± 0.01	52	⁸⁹ ALAM	86 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.13 ± 0.06 ± 0.01	1489	⁹⁰ BALEST	95B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
1.4 ^{+0.6} _{-0.5}	7	⁹¹ ALBRECHT	85H ARG	$e^+e^- \rightarrow \Upsilon(4S)$
1.1 ± 0.21 ± 0.23	46	⁹² HAAS	85 CLEO	Repl. by ALAM 86

⁸⁶ AUBERT 03F also reports the momentum distribution and helicity of $J/\psi \rightarrow \ell^+ \ell^-$ in the $\Upsilon(4S)$ center-of-mass frame.

⁸⁷ MASCHMANN 90 reports $(1.12 \pm 0.33 \pm 0.25) \times 10^{-2}$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁸⁸ ALBRECHT 87D reports $(1.07 \pm 0.16 \pm 0.22) \times 10^{-2}$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. ALBRECHT 87D find the branching ratio for J/ψ not from $\psi(2S)$ to be 0.0081 ± 0.0023 .

⁸⁹ ALAM 86 reports $(1.09 \pm 0.16 \pm 0.21) \times 10^{-2}$ for $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.074 \pm 0.012$. We rescale to our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.93 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁹⁰ BALEST 95B reports $(1.12 \pm 0.04 \pm 0.06) \times 10^{-2}$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0599 \pm 0.0025$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.. They measure $J/\psi(1S) \rightarrow e^+e^-$ and $\mu^+\mu^-$ and use PDG 1994 values for the branching fractions. The rescaling is the same for either mode so we use e^+e^- .

⁹¹ Statistical and systematic errors were added in quadrature. ALBRECHT 85H also report a CL = 90% limit of 0.007 for $B \rightarrow J/\psi(1S) + X$ where $m_X < 1$ GeV.

⁹² Dimuon and dielectron events used.

 $\Gamma(J/\psi(1S)\text{(direct) anything})/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0078 ± 0.0004 OUR AVERAGE	Error includes scale factor of 1.1.		
0.00740 ± 0.00023 ± 0.00043	⁹³ AUBERT	03F BABR	$e^+e^- \rightarrow \Upsilon(4S)$
0.00813 ± 0.00017 ± 0.00037	⁹⁴ ANDERSON	02 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.0080 ± 0.0008	⁹⁵ BALEST	95B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

- ⁹³ AUBERT 03F also reports the helicity of $J/\psi \rightarrow \ell^+ \ell^-$ produced directly in B decay.
- ⁹⁴ Also reports the measurement of $J/\psi \rightarrow \ell^+ \ell^-$ polarization produced directly from B decay.
- ⁹⁵ BALEST 95B assume PDG 1994 values for sub mode branching ratios. $J/\psi(1S)$ mesons are reconstructed in $J/\psi(1S) \rightarrow e^+ e^-$ and $J/\psi(1S) \rightarrow \mu^+ \mu^-$. The $B \rightarrow J/\psi(1S) X$ branching ratio contains $J/\psi(1S)$ mesons directly from B decays and also from feeddown through $\psi(2S) \rightarrow J/\psi(1S)$, $\chi_{c1}(1P) \rightarrow J/\psi(1S)$, or $\chi_{c2}(1P) \rightarrow J/\psi(1S)$. Using the measured inclusive rates, BALEST 95B corrects for the feeddown and finds the $B \rightarrow J/\psi(1S)$ (direct) X branching ratio.

$\Gamma(\psi(2S)\text{anything})/\Gamma_{\text{total}}$ Γ_{45}/Γ

<u>VALUE</u>	<u>EVTs</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.00307 ± 0.00021 OUR AVERAGE				
0.00297 ± 0.00020 ± 0.00020		AUBERT	03F BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.00316 ± 0.00014 ± 0.00028		⁹⁶ ANDERSON	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.0046 ± 0.0017 ± 0.0011	8	ALBRECHT	87D ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.0034 ± 0.0004 ± 0.0003	240	⁹⁷ BALEST	95B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

⁹⁶ Also reports the measurement of $\psi(2S) \rightarrow \ell^+ \ell^-$ polarization produced directly from B decay.

⁹⁷ BALEST 95B assume PDG 1994 values for sub mode branching ratios. They find $B(B \rightarrow \psi(2S)X, \psi(2S) \rightarrow \ell^+ \ell^-) = 0.30 \pm 0.05 \pm 0.04$ and $B(B \rightarrow \psi(2S)X, \psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = 0.37 \pm 0.05 \pm 0.05$. Weighted average is quoted for $B(B \rightarrow \psi(2S)X)$.

$\Gamma(\chi_{c1}(1P)\text{anything})/\Gamma_{\text{total}}$ Γ_{46}/Γ

<u>VALUE</u>	<u>EVTs</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.00386 ± 0.00027 OUR AVERAGE				
0.00367 ± 0.00035 ± 0.00044		AUBERT	03F BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.00363 ± 0.00022 ± 0.00034		⁹⁸ ABE	02L BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
0.00435 ± 0.00029 ± 0.00040		ANDERSON	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.00317 ± 0.00034 ± 0.00017		⁹⁹ CHEN	01 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.0040 ± 0.0006 ± 0.0004	112	¹⁰⁰ BALEST	95B CLE2	Repl. by CHEN 01
0.0105 ± 0.0035 ± 0.0025		¹⁰¹ ALBRECHT	92E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

⁹⁸ ABE 02L uses PDG 01 values for $B(J/\psi(1S) \rightarrow \ell^+ \ell^-)$ and $B(\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma)$.

⁹⁹ CHEN 01 reports $0.00414 \pm 0.00031 \pm 0.00040$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (35.6 \pm 1.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

¹⁰⁰ BALEST 95B assume $B(\chi_{c1}(1P) \rightarrow J/\psi(1S)\gamma) = (27.3 \pm 1.6) \times 10^{-2}$, the PDG 1994 value. Fit to ψ -photon invariant mass distribution allows for a $\chi_{c1}(1P)$ and a $\chi_{c2}(1P)$ component.

¹⁰¹ ALBRECHT 92E assumes no $\chi_{c2}(1P)$ production.

$\Gamma(\chi_{c1}(1P)(\text{direct}) \text{ anything})/\Gamma_{\text{total}}$ Γ_{47}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.00318 ± 0.00025 OUR AVERAGE

0.00341 ± 0.00035 ± 0.00042	AUBERT	03F BABR	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------------	--------	----------	-----------------------------------

0.00332 ± 0.00022 ± 0.00034	102 ABE	02L BELL	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------------	---------	----------	-----------------------------------

0.00294 ± 0.00035 ± 0.00016	103 CHEN	01 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------------	----------	---------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0037 ± 0.0007	104 BALEST	95B CLE2	Repl. by CHEN 01
-----------------	------------	----------	------------------

102 ABE 02L uses PDG 01 values for $B(J/\psi(1S) \rightarrow \ell^+\ell^-)$ and $B(\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma)$.

103 CHEN 01 reports $0.00383 \pm 0.00031 \pm 0.00040$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (35.6 \pm 1.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

104 BALEST 95B assume PDG 1994 values. $J/\psi(1S)$ mesons are reconstructed in the e^+e^- and $\mu^+\mu^-$ modes. The $B \rightarrow \chi_{c1}(1P)X$ branching ratio contains $\chi_{c1}(1P)$ mesons directly from B decays and also from feeddown through $\psi(2S) \rightarrow \chi_{c1}(1P)\gamma$. Using the measured inclusive rates, BALEST 95B corrects for the feeddown and finds the $B \rightarrow \chi_{c1}(1P)(\text{direct}) X$ branching ratio.

$\Gamma(\chi_{c2}(1P)\text{anything})/\Gamma_{\text{total}}$ Γ_{48}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	-------------	--------------------	-------------	----------------

13 ± 4 OUR AVERAGE			Error includes scale factor of 1.9. See the ideogram below.
---------------------------	--	--	---

21.0 ± 4.5 ± 3.1	AUBERT	03F BABR	$e^+e^- \rightarrow \Upsilon(4S)$
------------------	--------	----------	-----------------------------------

18.0 ^{+2.3} _{-2.8} ± 2.6	105 ABE	02L BELL	$e^+e^- \rightarrow \Upsilon(4S)$
--	---------	----------	-----------------------------------

6.5 ± 3.3 ± 0.3	106 CHEN	01 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------	----------	---------	-----------------------------------


• • • We do not use the following data for averages, fits, limits, etc. • • •

<38	90	35	107 BALEST 95B CLE2 Repl. by CHEN 01
-------	----	----	--------------------------------------

105 ABE 02L uses PDG 01 values for $B(J/\psi(1S) \rightarrow \ell^+\ell^-)$ and $B(\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma)$.

106 CHEN 01 reports $(9.8 \pm 4.8 \pm 1.5) \times 10^{-4}$ for $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$. We rescale to our best value $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (20.2 \pm 1.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

107 BALEST 95B assume $B(\chi_{c2}(1P) \rightarrow J/\psi(1S)\gamma) = (13.5 \pm 1.1) \times 10^{-2}$, the PDG 1994 value. $J/\psi(1S)$ mesons are reconstructed in the e^+e^- and $\mu^+\mu^-$ modes, and PDG 1994 branching fractions are used. If interpreted as signal, the 35 ± 13 events correspond to $B(B \rightarrow \chi_{c2}(1P)X) = (0.25 \pm 0.10 \pm 0.03) \times 10^{-2}$.



$\Gamma(\chi_{c2}(1P) \text{ (direct) anything}) / \Gamma_{\text{total}} \quad \Gamma_{49} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
0.00165 ± 0.00031 OUR AVERAGE			
$0.00190 \pm 0.00045 \pm 0.00029$	AUBERT	03F BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.00153^{+0.00023}_{-0.00028} \pm 0.00027$	108 ABE	02L BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹⁰⁸ ABE 02L uses PDG 01 values for $B(J/\psi(1S) \rightarrow \ell^+ \ell^-)$ and $B(\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma)$.

$\Gamma(\eta_c(1S) \text{ anything}) / \Gamma_{\text{total}} \quad \Gamma_{50} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.009	90	109 BALEST	95B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹⁰⁹ BALEST 95B assume PDG 1994 values for sub mode branching ratios. $J/\psi(1S)$ mesons are reconstructed in $J/\psi(1S) \rightarrow e^+ e^-$ and $J/\psi(1S) \rightarrow \mu^+ \mu^-$. Search region $2960 < m_{\eta_c(1S)} < 3010 \text{ MeV}/c^2$.

$\Gamma(K Y(3940) \times B(Y(3940) \rightarrow \omega J/\psi)) / \Gamma_{\text{total}} \quad \Gamma_{51} / \Gamma$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$7.1 \pm 1.3 \pm 3.1$	110 CHOI	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹¹⁰ CHOI 05 reports the observation of a near-threshold enhancement in the $\omega J/\psi$ mass spectrum in exclusive $B \rightarrow K \omega J/\psi$. The new state, denoted as $Y(3940)$, has a mass of $3943 \pm 11 \pm 13 \text{ GeV}/c^2$ and a width $\Gamma = 87 \pm 22 \pm 26 \text{ MeV}$.

$\Gamma(K^\pm \text{ anything})/\Gamma_{\text{total}}$ Γ_{52}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.789 ± 0.025 OUR AVERAGE			
0.82 ± 0.01 ± 0.05	ALBRECHT	94C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.775 ± 0.015 ± 0.025	¹¹¹ ALBRECHT	93I ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.85 ± 0.07 ± 0.09	ALAM	87B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
seen	¹¹² BRODY	82 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
seen	¹¹³ GIANNINI	82 CUSB	$e^+ e^- \rightarrow \Upsilon(4S)$
¹¹¹ ALBRECHT 93I value is not independent of the sum of $B \rightarrow K^+$ anything and $B \rightarrow K^-$ anything ALBRECHT 94C values.			
¹¹² Assuming $\Upsilon(4S) \rightarrow B\bar{B}$, a total of $3.38 \pm 0.34 \pm 0.68$ kaons per $\Upsilon(4S)$ decay is found (the second error is systematic). In the context of the standard B -decay model, this leads to a value for $(b\text{-quark} \rightarrow c\text{-quark})/(b\text{-quark} \rightarrow \text{all})$ of $1.09 \pm 0.33 \pm 0.13$.			
¹¹³ GIANNINI 82 at CESR-CUSB observed $1.58 \pm 0.35 K^0$ per hadronic event much higher than 0.82 ± 0.10 below threshold. Consistent with predominant $b \rightarrow cX$ decay.			

 $\Gamma(K^+ \text{ anything})/\Gamma_{\text{total}}$ Γ_{53}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.66 ± 0.05	¹¹⁴ ALBRECHT	94C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.620 ± 0.013 ± 0.038	¹¹⁵ ALBRECHT	94C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.66 ± 0.05 ± 0.07	¹¹⁵ ALAM	87B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
¹¹⁴ Measurement relies on lepton-kaon correlations. It is for the weak decay vertex and does not include mixing of the neutral B meson. Mixing effects were corrected for by assuming a mixing parameter r of $(18.1 \pm 4.3)\%$.			
¹¹⁵ Measurement relies on lepton-kaon correlations. It includes production through mixing of the neutral B meson.			

 $\Gamma(K^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{54}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.13 ± 0.04	¹¹⁶ ALBRECHT	94C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.165 ± 0.011 ± 0.036	¹¹⁷ ALBRECHT	94C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.19 ± 0.05 ± 0.02	¹¹⁷ ALAM	87B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
¹¹⁶ Measurement relies on lepton-kaon correlations. It is for the weak decay vertex and does not include mixing of the neutral B meson. Mixing effects were corrected for by assuming a mixing parameter r of $(18.1 \pm 4.3)\%$.			
¹¹⁷ Measurement relies on lepton-kaon correlations. It includes production through mixing of the neutral B meson.			

 $\Gamma(K^0/\bar{K}^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{55}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.64 ± 0.04 OUR AVERAGE			
0.642 ± 0.010 ± 0.042	¹¹⁸ ALBRECHT	94C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.63 ± 0.06 ± 0.06	ALAM	87B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
¹¹⁸ ALBRECHT 94C assume a K^0/\bar{K}^0 multiplicity twice that of K_S^0 .			

$\Gamma(K^*(892)^\pm \text{ anything})/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.182 \pm 0.054 \pm 0.024$	ALBRECHT 94J	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^*(892)^0 / \bar{K}^*(892)^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.146 \pm 0.016 \pm 0.020$	ALBRECHT 94J	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^*(892)\gamma)/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$4.24 \pm 0.54 \pm 0.32$		119 COAN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 150	90	120 LESIAK	92 CBAL	$e^+ e^- \rightarrow \Upsilon(4S)$
< 24	90	ALBRECHT	88H ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

119 An average of $B(B^+ \rightarrow K^*(892)^+ \gamma)$ and $B(B^0 \rightarrow K^*(892)^0 \gamma)$ measurements reported in COAN 00 by assuming full correlated systematic errors.

120 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s \gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s-quark hadronization.

$\Gamma(\eta K \gamma)/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$8.5 \pm 1.3^{+1.2}_{-0.9}$	121 NISHIDA	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

121 $m_{\eta K} < 2.4 \text{ GeV}/c^2$

$\Delta_{0+}(B \rightarrow K^*(892)\gamma)$

Δ_{0+} describes the isospin asymmetry between $\Gamma(B^0 \rightarrow K^*(892)^0 \gamma)$ and $\Gamma(B^+ \rightarrow K^*(892)^+ \gamma)$.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.050 \pm 0.045 \pm 0.037$	122 AUBERT,BE	04A BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

122 Uses the production ratio of charged and neutral B from $\Upsilon(4S)$ decays $R^{+/0} = 1.006 \pm 0.048$ and the lifetime ratio of $\tau_{B^+} / \tau_{B^0} = 1.083 \pm 0.017$. The 90% CL interval is $-0.046 < \Delta_{0+} < 0.146$.

$\Gamma(K_1(1400)\gamma)/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 12.7×10^{-5}	90	123 COAN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.6×10^{-3}	90	124 LESIAK	92 CBAL	$e^+ e^- \rightarrow \Upsilon(4S)$
< 4.1×10^{-4}	90	ALBRECHT	88H ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

123 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

124 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s \gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s-quark hadronization.

$\Gamma(K_2^*(1430)\gamma)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$1.66^{+0.59}_{-0.53} \pm 0.13$		125 COAN	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<83 90 ALBRECHT 88H ARG $e^+e^- \rightarrow \Upsilon(4S)$

125 COAN 00 obtains a fitted signal yield of $15.9^{+5.7}_{-5.2}$ events. A search for contamination by $K^*(1410)$ yielded a rate consistent with 0; the central value assumes no contamination.

$\Gamma(K_2(1770)\gamma)/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-3}$	90	126 LESIAK	92 CBAL	$e^+e^- \rightarrow \Upsilon(4S)$

126 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s-quark hadronization.

$\Gamma(K_3^*(1780)\gamma)/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.7 \times 10^{-5}$	90	127 NISHIDA	05 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.0×10^{-3} 90 ALBRECHT 88H ARG $e^+e^- \rightarrow \Upsilon(4S)$

127 Uses $B(K_3^*(1780) \rightarrow \eta K) = 0.11^{+0.05}_{-0.04}$.

$\Gamma(K_4^*(2045)\gamma)/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-3}$	90	128 LESIAK	92 CBAL	$e^+e^- \rightarrow \Upsilon(4S)$

128 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s-quark hadronization.

$\Gamma(K\eta'(958))/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$(8.3^{+0.9}_{-0.8} \pm 0.7) \times 10^{-5}$		129 RICHICHI	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

129 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)\eta'(958))/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-5}$	90	130 RICHICHI	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

130 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K\eta)/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-6}$	90	131 RICHICHI	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

131 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)\eta)/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$(1.80^{+0.49}_{-0.43} \pm 0.18) \times 10^{-5}$	132 RICHICHI	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹³² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K\phi\phi)/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$2.3^{+0.9}_{-0.8} \pm 0.3$	133 HUANG	03 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹³³ Assumes equal production of charged and neutral B meson pairs and isospin symmetry.

$\Gamma(\bar{b} \rightarrow \bar{s}\gamma)/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.43 ± 0.29 OUR AVERAGE			

3.49 ± 0.20 $^{+0.59}_{-0.46}$ 134,135 AUBERT,B 05R BABR $e^+e^- \rightarrow \Upsilon(4S)$

3.50 ± 0.32 ± 0.31 135,136 KOPPENBURG04 BELL $e^+e^- \rightarrow \Upsilon(4S)$

3.29 ± 0.44 ± 0.29 135,137 CHEN 01C CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.36 ± 0.53 $^{+0.65}_{-0.68}$ 138 ABE 01F BELL Repl. by KOPPENBURG 04

2.32 ± 0.57 ± 0.35 ALAM 95 CLE2 Repl. by CHEN 01C

¹³⁴ The measurement reported is $3.27 \pm 0.18^{+0.55}_{-0.42}$ for $E_\gamma > 1.9$ GeV.

¹³⁵ We correct it to $E_\gamma > 1.6$ GeV using the method of hep-ph/0507253 (average of three theoretical models).

¹³⁶ The measurement reported is $3.21 \pm 0.43^{+0.32}_{-0.29}$ for $E_\gamma > 2.0$ GeV.

¹³⁷ The measurement reported is $3.55 \pm 0.32 \pm 0.32$ for $E_\gamma > 1.8$ GeV.

¹³⁸ ABE 01F reports their systematic errors $\pm 0.42^{+0.50}_{-0.54}$, where the second error is due to the theoretical uncertainty. We combine them in quadrature.

$\Gamma(\bar{b} \rightarrow \bar{s}g\text{luon})/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.068	90		139 COAN	98 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.08 2 140 ALBRECHT 95D ARG $e^+e^- \rightarrow \Upsilon(4S)$

¹³⁹ COAN 98 uses D - ℓ correlation.

¹⁴⁰ ALBRECHT 95D use full reconstruction of one B decay as tag. Two candidate events for charmless B decay can be interpreted as either $b \rightarrow sg\text{luon}$ or $b \rightarrow u$ transition. If interpreted as $b \rightarrow sg\text{luon}$ they find a branching ratio of ~ 0.026 or the upper limit quoted above. Result is highly model dependent.

$\Gamma(\eta\text{anything})/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.4 × 10⁻⁴	90	141 BROWDER	98 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹⁴¹ BROWDER 98 search for high momentum $B \rightarrow \eta X_s$ between 2.1 and 2.7 GeV/ c .

$\Gamma(\eta' \text{ anything})/\Gamma_{\text{total}}$ **Γ_{73}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.2 ± 0.9 OUR AVERAGE			
$3.9 \pm 0.8 \pm 0.9$	142 AUBERT, B	04F BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$4.6 \pm 1.1 \pm 0.6$	143 BONVICINI	03 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$6.2 \pm 1.6^{+1.3}_{-2.0}$	144 BROWDER	98 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹⁴² The reported branching ratio is for high momentum η between 2.0 and 2.7 GeV in the $\Upsilon(4S)$ center-of-mass frame. Xs represents a recoil system consisting of a kaon and zero to four pions.

¹⁴³ BONVICINI 03 observed a signal of 61.2 ± 13.9 events in $B \rightarrow \eta' X_{nc}$ production for high momentum η' between 2.0 and 2.7 GeV/c in the $\Upsilon(4S)$ center-of-mass frame. The X_{nc} denotes "charmless" hadronic states recoiling against η' . The second error combines systematic and background subtraction uncertainties in quadrature.

¹⁴⁴ BROWDER 98 observed a signal of 39.0 ± 11.6 events in high momentum $B \rightarrow \eta' X_s$ production between 2.0 and 2.7 GeV/c. The branching fraction is based on the interpretation of $b \rightarrow sg$, where the last error includes additional uncertainties due to the color-suppressed $b \rightarrow$ backgrounds.

 $\Gamma(\rho\gamma)/\Gamma_{\text{total}}$ **Γ_{74}/Γ**

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.9 \times 10^{-6}$	90	145 AUBERT	04C BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 1.4 \times 10^{-5}$	90	146 COAN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹⁴⁵ Assumes $\Gamma(B \rightarrow \rho\gamma) = \Gamma(B^+ \rightarrow \rho^+\gamma) = 2 \Gamma(B^0 \rightarrow \rho^0\gamma)$ and uses lifetime ratio of $\tau_{B^+}/\tau_{B^0} = 1.083 \pm 0.017$.

¹⁴⁶ COAN 00 reports $B(B \rightarrow \rho\gamma)/B(B \rightarrow K^*(892)\gamma) < 0.32$ at 90%CL and scaled by the central value of $B(B \rightarrow K^*(892)\gamma) = (4.24 \pm 0.54 \pm 0.32) \times 10^{-5}$.

 $\Gamma(\rho/\omega\gamma)/\Gamma_{\text{total}}$ **Γ_{75}/Γ**

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.2 \times 10^{-6}$	90	AUBERT	05 BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 1.4 \times 10^{-6}$	90	MOHAPATRA	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

 $\Gamma(\rho/\omega\gamma)/\Gamma(K^*(892)\gamma)$ **Γ_{75}/Γ_{58}**

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 0.035	90	147 MOHAPATRA	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹⁴⁷ A limit of $|V_{td}/V_{ts}| < 0.22$ at 90% CL is also obtained from the measurement.

 $\Gamma(\pi^\pm \text{ anything})/\Gamma_{\text{total}}$ **Γ_{76}/Γ**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.585 \pm 0.025 \pm 0.070$	148 ALBRECHT	93I ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹⁴⁸ ALBRECHT 93 excludes π^\pm from K_S^0 and Λ decays. If included, they find $4.105 \pm 0.025 \pm 0.080$.

$\Gamma(\pi^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE		DOCUMENT ID	TECN	COMMENT
2.35±0.02±0.11	149	ABE	01J BELL	$e^+e^- \rightarrow \Upsilon(4S)$

149 From fully inclusive π^0 yield with no corrections from decays of K_S^0 or other particles.

$\Gamma(\eta \text{ anything})/\Gamma_{\text{total}}$ Γ_{78}/Γ

VALUE		DOCUMENT ID	TECN	COMMENT
0.176±0.011±0.012		KUBOTA	96 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\rho^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{79}/Γ

VALUE		DOCUMENT ID	TECN	COMMENT
0.208±0.042±0.032		ALBRECHT	94J ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\omega \text{ anything})/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.81	90	ALBRECHT	94J ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE		DOCUMENT ID	TECN	COMMENT
0.0342±0.0013 OUR AVERAGE				
0.0341±0.0006±0.0012		AUBERT	04S BABR	$e^+e^- \rightarrow \Upsilon(4S)$
0.0390±0.0030±0.0035		ALBRECHT	94J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
0.023 ±0.006 ±0.005		BORTOLETTO	86 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\phi K^*(892))/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.2 × 10⁻⁵	90	150 BERGFELD	98 CLE2	

150 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\Lambda_c^+ / \bar{\Lambda}_c^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.064±0.008±0.008		151 CRAWFORD	92 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.14 ±0.09		152 ALBRECHT	88E ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<0.112	90	153 ALAM	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

151 CRAWFORD 92 result derived from lepton baryon correlations. Assumes all charmed baryons in B^0 and B^\pm decay are Λ_c .

152 ALBRECHT 88E measured $B(B \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (0.30 \pm 0.12 \pm 0.06)\%$ and used $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (2.2 \pm 1.0)\%$ from ABRAMS 80 to obtain above number.

153 Assuming all baryons result from charmed baryons, ALAM 86 conclude the branching fraction is $7.4 \pm 2.9\%$. The limit given above is model independent.

$\Gamma(\Lambda_c^+ \text{ anything})/\Gamma(\bar{\Lambda}_c^- \text{ anything})$ Γ_{84}/Γ_{85}

VALUE		DOCUMENT ID	TECN	COMMENT
0.19±0.13±0.04	154	AMMAR	97 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

154 AMMAR 97 uses a high-momentum lepton tag ($P_\ell > 1.4 \text{ GeV}/c^2$).

$\Gamma(\bar{\Lambda}_c^- e^+ \text{ anything})/\Gamma(\Lambda_c^+ / \bar{\Lambda}_c^- \text{ anything})$ Γ_{86}/Γ_{83}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.05	90	155 BONVICINI	98 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

155 BONVICINI 98 uses the electron with momentum above 0.6 GeV/c.

$\Gamma(\bar{\Lambda}_c^- p \text{ anything})/\Gamma(\Lambda_c^+ / \bar{\Lambda}_c^- \text{ anything})$ Γ_{87}/Γ_{83}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.57 \pm 0.05 \pm 0.05$	BONVICINI	98 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(\bar{\Lambda}_c^- p e^+ \nu_e)/\Gamma(\bar{\Lambda}_c^- p \text{ anything})$ Γ_{88}/Γ_{87}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	90	156 BONVICINI	98 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

156 BONVICINI 98 uses the electron with momentum above 0.6 GeV/c.

$\Gamma(\bar{\Sigma}_c^{--} \text{ anything})/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.0042 \pm 0.0021 \pm 0.0011$	77	157 PROCARIO	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

157 PROCARIO 94 reports $[B(B \rightarrow \bar{\Sigma}_c^{--} \text{ anything}) \times B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = 0.00021 \pm 0.00008 \pm 0.00007$. We divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{\Sigma}_c^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{90}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.010	90	158 PROCARIO	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

158 PROCARIO 94 reports $[B(B \rightarrow \bar{\Sigma}_c^- \text{ anything}) \times B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = < 0.00048$. We divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.050$.

$\Gamma(\bar{\Sigma}_c^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.0046 \pm 0.0021 \pm 0.0012$	76	159 PROCARIO	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

159 PROCARIO 94 reports $[B(B \rightarrow \bar{\Sigma}_c^0 \text{ anything}) \times B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = 0.00023 \pm 0.00008 \pm 0.00007$. We divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{\Sigma}_c^0 N (N = p \text{ or } n))/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0015	90	160 PROCARIO	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

160 PROCARIO 94 reports < 0.0017 for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.043$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.050$.

$\Gamma(\Xi_c^0 \text{ anything} \times B(\Xi_c^0 \rightarrow \Xi^- \pi^+))/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.193 ± 0.030 OUR AVERAGE	Error includes scale factor of 1.1.		
0.211 ± 0.019 ± 0.025	161 AUBERT,B	05M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.144 ± 0.048 ± 0.021	162 BARISH	97 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹⁶¹ The yield is obtained by requiring the momentum $P < 2.15$ GeV/c.

¹⁶² BARISH 97 find $79 \pm 27 \Xi_c^0$ events.

$\Gamma(\Xi_c^+ \text{ anything} \times B(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+))/\Gamma_{\text{total}}$ Γ_{94}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.453 ± 0.096^{+0.085}_{-0.065}	163 BARISH	97 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹⁶³ BARISH 97 find $125 \pm 28 \Xi_c^+$ events.

$\Gamma(p/\bar{p} \text{ anything})/\Gamma_{\text{total}}$ Γ_{95}/Γ

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.080 ± 0.004 OUR AVERAGE				
0.080 ± 0.005 ± 0.005		ALBRECHT	93I ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.080 ± 0.005 ± 0.003		CRAWFORD	92 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
0.082 ± 0.005 ^{+0.013} _{-0.010}	2163	¹⁶⁴ ALBRECHT	89K ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.021 ¹⁶⁵ ALAM 83B CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

¹⁶⁴ ALBRECHT 89K include direct and nondirect protons.

¹⁶⁵ ALAM 83B reported their result as $> 0.036 \pm 0.006 \pm 0.009$. Data are consistent with equal yields of p and \bar{p} . Using assumed yields below cut, $B(B \rightarrow p + X) = 0.03$ not including protons from Λ decays.

$\Gamma(p/\bar{p}(\text{direct}) \text{ anything})/\Gamma_{\text{total}}$ Γ_{96}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.055 ± 0.005 OUR AVERAGE				
0.055 ± 0.005 ± 0.0035		ALBRECHT	93I ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.056 ± 0.006 ± 0.005		CRAWFORD	92 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
0.055 ± 0.016	1220	¹⁶⁶ ALBRECHT	89K ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹⁶⁶ ALBRECHT 89K subtract contribution of Λ decay from the inclusive proton yield.

$\Gamma(\Lambda/\bar{\Lambda} \text{ anything})/\Gamma_{\text{total}}$ Γ_{97}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.040 ± 0.005 OUR AVERAGE				
0.038 ± 0.004 ± 0.006	2998	CRAWFORD	92 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
0.042 ± 0.005 ± 0.006	943	ALBRECHT	89K ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.022 ± 0.003 ± 0.0022 ¹⁶⁷ ACKERSTAFF 97N OPAL $e^+ e^- \rightarrow Z$

>0.011 ¹⁶⁸ ALAM 83B CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

¹⁶⁷ ACKERSTAFF 97N assumes $B(b \rightarrow B) = 0.868 \pm 0.041$, *i.e.*, an admixture of B^0 , B^\pm , and B_s .

¹⁶⁸ ALAM 83B reported their result as $> 0.022 \pm 0.007 \pm 0.004$. Values are for $(B(\Lambda X) + B(\bar{\Lambda} X))/2$. Data are consistent with equal yields of p and \bar{p} . Using assumed yields below cut, $B(B \rightarrow \Lambda X) = 0.03$.

$\Gamma(\Lambda \text{ anything})/\Gamma(\bar{\Lambda} \text{ anything})$

Γ_{98}/Γ_{99}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.43 ± 0.09 ± 0.07	169 AMMAR	97 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
169 AMMAR 97 uses a high-momentum lepton tag ($P_\ell > 1.4 \text{ GeV}/c^2$).			

$\Gamma(\Xi^- / \Xi^+ \text{ anything})/\Gamma_{\text{total}}$

Γ_{100}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0027 ± 0.0006 OUR AVERAGE				
0.0027 ± 0.0005 ± 0.0004	147	CRAWFORD	92 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.0028 ± 0.0014	54	ALBRECHT	89K ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\text{baryons anything})/\Gamma_{\text{total}}$

Γ_{101}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.068 ± 0.005 ± 0.003	170 ALBRECHT	920 ARG	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.076 ± 0.014	171 ALBRECHT	89K ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹⁷⁰ ALBRECHT 920 result is from simultaneous analysis of p and Λ yields, $p\bar{p}$ and $\Lambda\bar{\Lambda}$ correlations, and various lepton-baryon and lepton-baryon-antibaryon correlations. Supersedes ALBRECHT 89K.

¹⁷¹ ALBRECHT 89K obtain this result by adding their their measurements (5.5 ± 1.6)% for direct protons and ($4.2 \pm 0.5 \pm 0.6$)% for inclusive Λ production. They then assume (5.5 ± 1.6)% for neutron production and add it in also. Since each B decay has two baryons, they divide by 2 to obtain (7.6 ± 1.4)%.

$\Gamma(p\bar{p} \text{ anything})/\Gamma_{\text{total}}$

Γ_{102}/Γ

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0247 ± 0.0023 OUR AVERAGE				
0.024 ± 0.001 ± 0.004		CRAWFORD	92 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.025 ± 0.002 ± 0.002	918	ALBRECHT	89K ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(p\bar{p} \text{ anything})/\Gamma(p/\bar{p} \text{ anything})$

Γ_{102}/Γ_{95}

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.30 ± 0.02 ± 0.05	172 CRAWFORD	92 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹⁷² CRAWFORD 92 value is not independent of their $\Gamma(p\bar{p} \text{ anything})/\Gamma_{\text{total}}$ value.

$\Gamma(\Lambda\bar{p}/\bar{\Lambda}p \text{ anything})/\Gamma_{\text{total}}$

Γ_{103}/Γ

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.025 ± 0.004 OUR AVERAGE				
0.029 ± 0.005 ± 0.005		CRAWFORD	92 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.023 ± 0.004 ± 0.003	165	ALBRECHT	89K ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\Lambda\bar{p}/\bar{\Lambda}p\text{anything})/\Gamma(\Lambda/\bar{\Lambda}\text{anything})$ Γ_{103}/Γ_{97}

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.76 \pm 0.11 \pm 0.08$	¹⁷³ CRAWFORD	92	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
¹⁷³ CRAWFORD ⁹² value is not independent of their $[\Gamma(\Lambda\bar{p}\text{anything})+\Gamma(\bar{\Lambda}p\text{anything})]/\Gamma_{\text{total}}$ value.			

$\Gamma(\Lambda\bar{\Lambda}\text{anything})/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.005	90		CRAWFORD	92	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<0.0088	90	12	ALBRECHT	89K	ARG $e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\Lambda\bar{\Lambda}\text{anything})/\Gamma(\Lambda/\bar{\Lambda}\text{anything})$ Γ_{104}/Γ_{97}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.13	90	¹⁷⁴ CRAWFORD	92	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
¹⁷⁴ CRAWFORD ⁹² value is not independent of their $\Gamma(\Lambda\bar{\Lambda}\text{anything})/\Gamma_{\text{total}}$ value.				

$\Gamma(se^+e^-)/\Gamma_{\text{total}}$ Γ_{105}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
4.7 ± 1.3 OUR AVERAGE				
$4.04 \pm 1.30^{+0.87}_{-0.83}$		¹⁷⁵ IWASAKI	05	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$6.0 \pm 1.7 \pm 1.3$		¹⁷⁶ AUBERT,B	04I	BABR $e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$5.0 \pm 2.3^{+1.3}_{-1.1}$		¹⁷⁶ KANEKO	03	BELL Repl. by IWASAKI 05
< 57	90	GLENN	98	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
<50000	90	BEBEK	81	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
¹⁷⁵ Requires $M_{\ell^+\ell^-} > 0.2 \text{ GeV}/c^2$.				
¹⁷⁶ Requires $M_{e^+e^-} > 0.2 \text{ GeV}/c^2$.				

$\Gamma(s\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{106}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
4.3 ± 1.2 OUR AVERAGE				
$4.13 \pm 1.05^{+0.85}_{-0.81}$		¹⁷⁷ IWASAKI	05	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$5.0 \pm 2.8 \pm 1.2$		AUBERT,B	04I	BABR $e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$7.9 \pm 2.1^{+2.1}_{-1.5}$		KANEKO	03	BELL Repl. by IWASAKI 05
< 58	90	GLENN	98	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
<17000	90	CHADWICK	81	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
¹⁷⁷ Requires $M_{\ell^+\ell^-} > 0.2 \text{ GeV}/c^2$.				

$[\Gamma(se^+e^-) + \Gamma(s\mu^+\mu^-)]/\Gamma_{\text{total}}$ $(\Gamma_{105} + \Gamma_{106})/\Gamma$

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.2 × 10⁻⁵	90	GLENN	98 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.0024	90	¹⁷⁸ BEAN	87 CLEO	Repl. by GLENN 98
<0.0062	90	¹⁷⁹ AVERY	84 CLEO	Repl. by BEAN 87

¹⁷⁸BEAN 87 reports $[(\mu^+\mu^-) + (e^+e^-)]/2$ and we converted it.

¹⁷⁹Determine ratio of B^+ to B^0 semileptonic decays to be in the range 0.25–2.9.

$\Gamma(sl^+\ell^-)/\Gamma_{\text{total}}$ Γ_{107}/Γ

Test for $\Delta B = 1$ weak neutral current.

VALUE (units 10 ⁻⁶)	DOCUMENT ID	TECN	COMMENT
4.5 ± 1.0 OUR AVERAGE			
4.11 ± 0.83 ^{+0.85} _{-0.81}	¹⁸⁰ IWASAKI	05 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
5.6 ± 1.5 ± 1.3	¹⁸¹ AUBERT,B	04i BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
6.1 ± 1.4 ^{+1.4} _{-1.1}	¹⁸¹ KANEKO	03 BELL	Repl. by IWASAKI 05

¹⁸⁰Requires $M_{\ell^+\ell^-} > 0.2 \text{ GeV}/c^2$.

¹⁸¹Requires $M_{e^+e^-} > 0.2 \text{ GeV}/c^2$.

$\Gamma(Ke^+e^-)/\Gamma_{\text{total}}$ Γ_{108}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10 ⁻⁷)	CL%	DOCUMENT ID	TECN	COMMENT
6.0^{+1.4}_{-1.2} OUR AVERAGE				Error includes scale factor of 1.1.
7.4 ^{+1.8} _{-1.6} ± 0.5		¹⁸² AUBERT	03U BABR	$e^+e^- \rightarrow \Upsilon(4S)$
4.8 ^{+1.5} _{-1.3} ± 0.3		^{182,183} ISHIKAWA	03 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13 90 ABE 02 BELL Repl. by ISHIKAWA 03

¹⁸²Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

¹⁸³The second error is a total of systematic uncertainties including model dependence.

$\Gamma(K^*(892)e^+e^-)/\Gamma_{\text{total}}$ Γ_{109}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10 ⁻⁶)	CL%	DOCUMENT ID	TECN	COMMENT
1.24^{+0.37}_{-0.32} OUR AVERAGE				
0.98 ^{+0.50} _{-0.42} ± 0.11		¹⁸⁴ AUBERT	03U BABR	$e^+e^- \rightarrow \Upsilon(4S)$
1.49 ^{+0.52} _{-0.46} ± 0.12 ± 0.13		¹⁸⁵ ISHIKAWA	03 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5.6 90 ABE 02 BELL Repl. by ISHIKAWA 03

184 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

185 Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$. The second error is a total of systematic uncertainties including model dependence.

$\Gamma(K\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{110}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

(4.7 $^{+1.1}_{-1.0}$) $\times 10^{-7}$ OUR AVERAGE

(4.5 $^{+2.3}_{-1.9}$ ± 0.4) $\times 10^{-7}$	186 AUBERT	03U BABR	$e^+e^- \rightarrow \Upsilon(4S)$
--	------------	----------	-----------------------------------

(4.8 $^{+1.2}_{-1.1}$ ± 0.4) $\times 10^{-7}$	187 ISHIKAWA	03 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
--	--------------	---------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

(0.99 $^{+0.40}_{-0.32}$ $^{+0.13}_{-0.14}$) $\times 10^{-6}$	ABE	02 BELL	Repl. by ISHIKAWA 03
--	-----	---------	----------------------

186 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

187 Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$. The second error is a total of systematic uncertainties including model dependence.

$\Gamma(K^*(892)\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{111}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

1.19 $^{+0.34}_{-0.29}$ OUR AVERAGE

1.27 $^{+0.76}_{-0.61}$ ± 0.16	188 AUBERT	03U BABR	$e^+e^- \rightarrow \Upsilon(4S)$
------------------------------------	------------	----------	-----------------------------------

1.17 $^{+0.36}_{-0.31}$ ± 0.10	189 ISHIKAWA	03 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
------------------------------------	--------------	---------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.1	90	ABE	02 BELL Repl. by ISHIKAWA 03
---------	----	-----	------------------------------

188 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

189 Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$. The second error is a total of systematic uncertainties including model dependence.

$\Gamma(K\ell^+\ell^-)/\Gamma_{\text{total}}$ Γ_{112}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

0.54 ± 0.08 OUR AVERAGE

0.65 $^{+0.14}_{-0.13}$ ± 0.04	190 AUBERT	03U BABR	$e^+e^- \rightarrow \Upsilon(4S)$
------------------------------------	------------	----------	-----------------------------------

0.48 $^{+0.10}_{-0.09}$ ± 0.03	191 ISHIKAWA	03 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
------------------------------------	--------------	---------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.75 $^{+0.25}_{-0.21}$ ± 0.06	192 ABE	02 BELL	Repl. by ISHIKAWA 03
------------------------------------	---------	---------	----------------------

< 0.51	90	193 AUBERT	02L BABR $e^+e^- \rightarrow \Upsilon(4S)$
----------	----	------------	--

< 1.7	90	194 ANDERSON	01B CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
---------	----	--------------	--

- 190 Assumes all four $B \rightarrow K\ell^+\ell^-$ modes having equal partial widths in the fit.
 191 Assumes equal production rate for charge and neutral B meson pairs, isospin invariance, lepton universality for $B \rightarrow K\ell^+\ell^-$, and $B(B \rightarrow K^*(892)\mu^+\mu^-) = 1.33$. The second error is total systematic uncertainties including model dependence.
 192 Assumes lepton universality.
 193 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
 194 The result is for di-lepton masses above 0.5 GeV.

$\Gamma(K^*(892)\ell^+\ell^-)/\Gamma_{\text{total}}$ Γ_{113}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
1.05 ± 0.20 OUR AVERAGE				
$0.88^{+0.33}_{-0.29} \pm 0.10$		195 AUBERT	03U BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$1.15^{+0.26}_{-0.24} \pm 0.08$		196 ISHIKAWA	03 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

- | | | | | |
|------|----|----------------|----------|-----------------------------------|
| <3.1 | 90 | 197,198 AUBERT | 02L BABR | Repl. by AUBERT 03U |
| <3.3 | 90 | 199 ANDERSON | 01B CLE2 | $e^+e^- \rightarrow \Upsilon(4S)$ |
- 195 Assumes the partial width ratio of electron and muon modes to be $\Gamma(B \rightarrow K^*(892)e^+e^-)/\Gamma(B \rightarrow K^*(892)\mu^+\mu^-) = 1.33$.
 196 Assumes equal production rate for charge and neutral B meson pairs, isospin invariance, lepton universality for $B \rightarrow K\ell^+\ell^-$, and $B(B \rightarrow K^*(892)\mu^+\mu^-) = 1.33$. The second error is total systematic uncertainties including model dependence.
 197 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
 198 For averaging $K^*(892)\mu^+\mu^-$ and $K^*(892)e^+e^-$ modes, AUBERT 02L assumed $B(B \rightarrow K^*(892)e^+e^-)/B(B \rightarrow K^*(892)\mu^+\mu^-) = 1.2$.
 199 The result is for di-lepton masses above 0.5 GeV.

$\Gamma(e^\pm\mu^\mp s)/\Gamma_{\text{total}}$ Γ_{114}/Γ

Test for lepton family number conservation. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.2 × 10⁻⁵	90	GLENN	98 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\pi e^\pm\mu^\mp)/\Gamma_{\text{total}}$ Γ_{115}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.6 × 10⁻⁶	90	200 EDWARDS	02B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

200 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\rho e^\pm\mu^\mp)/\Gamma_{\text{total}}$ Γ_{116}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.2 × 10⁻⁶	90	201 EDWARDS	02B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

201 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{117}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.6 \times 10^{-6}$	90	202 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

²⁰² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892) e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{118}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.2 \times 10^{-6}$	90	203 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

²⁰³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

CP VIOLATION

A_{CP} is defined as

$$\frac{B(B \rightarrow \bar{f}) - B(\bar{B} \rightarrow f)}{B(B \rightarrow f) + B(\bar{B} \rightarrow f)}$$

the CP -violation charge asymmetry of inclusive B^\pm and B^0 decay.

$A_{CP}(B \rightarrow K^*(892)\gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.010 ± 0.028 OUR AVERAGE			

$-0.013 \pm 0.036 \pm 0.010$ ²⁰⁴ AUBERT, BE 04A BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$-0.015 \pm 0.044 \pm 0.012$ ²⁰⁵ NAKAO 04 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

$+0.08 \pm 0.13 \pm 0.03$ ²⁰⁵ COAN 00 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.044 \pm 0.076 \pm 0.012$ ²⁰⁶ AUBERT 02C BABR Repl. by AUBERT, BE 04A

²⁰⁴ Corresponds to a 90% CL allowed region, $-0.074 < A_{CP} < 0.049$.

²⁰⁵ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

²⁰⁶ A 90% CL range is $-0.170 < A_{CP} < 0.082$.

$A_{CP}(B \rightarrow s\gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.00 ± 0.04 OUR AVERAGE			

$0.025 \pm 0.050 \pm 0.015$ ²⁰⁷ AUBERT, B 04E BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$0.002 \pm 0.050 \pm 0.030$ ²⁰⁸ NISHIDA 04 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

$-0.079 \pm 0.108 \pm 0.022$ ²⁰⁹ COAN 01 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

²⁰⁷ Corresponds to $-0.06 < A_{CP} < +0.11$ at 90% CL.

²⁰⁸ This measurement is performed inclusively for recoil mass X_s less than 2.1 GeV, which corresponds to $-0.093 < A_{CP} < 0.096$ at 90% CL.

²⁰⁹ Corresponds to $-0.27 < A_{CP} < 0.10$ at 90% CL.

$A_{CP}(b \rightarrow X_s \ell^+ \ell^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.22 \pm 0.26 \pm 0.02$	²¹⁰ AUBERT, B 04I	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

²¹⁰ The final state flavor is determined by the kaon and pion charges where modes with $X_s = K_S^0, K_S^0 \pi^0$ or $K_S^0 \pi^+ \pi^-$ are not used.

ISOSPIN ASYMMETRY Δ_{0-} is defined as

$$\frac{\Gamma(B^0 \rightarrow f_d) - \Gamma(B^+ \rightarrow f_u)}{\Gamma(B^0 \rightarrow f) + \Gamma(B^+ \rightarrow f)},$$

the isospin asymmetry of inclusive neutral and charged B decay.

 $\Delta_{0-}(B(B \rightarrow X_s \gamma))$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.006 \pm 0.058 \pm 0.026$	AUBERT,B	05R BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

 $B \rightarrow X_c \ell \nu$ HADRONIC MASS MOMENTS **$\langle M_X^2 - \overline{M}_D^2 \rangle$ (First Moments)**

VALUE (GeV ²)	DOCUMENT ID	TECN	COMMENT
0.36 ± 0.08 OUR AVERAGE	Error includes scale factor of 1.8.		
0.467 ± 0.038 ± 0.068	²¹¹ ACOSTA	05F CDF	$p\bar{p}$ at 1.96 TeV
0.293 ± 0.012 ± 0.058	²¹² CSORNA	04 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.251 ± 0.023 ± 0.062 ²¹³ CRONIN-HEN..01B CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$ ²¹¹ Moments are measured with a minimum lepton momentum of 0.7 GeV/c in the B rest frame;²¹² Uses minimum lepton energy of 1.5 GeV and also reports moments with $E_\ell > 1.0$ GeV.²¹³ The leptons are required to have $P_1 > 1.5$ GeV/c. **$\langle (M_X^2 - \overline{M}_X^2)^2 \rangle$ (Second Moments)**

VALUE (GeV ⁴)	DOCUMENT ID	TECN	COMMENT
0.71 ± 0.17 OUR AVERAGE	Error includes scale factor of 1.3.		
1.05 ± 0.26 ± 0.13	²¹⁴ ACOSTA	05F CDF	$p\bar{p}$ at 1.96 TeV
0.629 ± 0.031 ± 0.143	²¹⁵ CSORNA	04 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.576 ± 0.048 ± 0.168 ²¹⁶ CRONIN-HEN..01B CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$ ²¹⁴ Moments are measured with a minimum lepton momentum of 0.7 GeV/c in the B rest frame;²¹⁵ Uses minimum lepton energy of 1.5 GeV and also reports moments with $E_\ell > 1.0$ GeV.²¹⁶ The leptons are required to have $P_1 > 1.5$ GeV/c. **$\langle (M_X^2 - \overline{M}_D^2)^2 \rangle$ (Second Moments)**

VALUE (GeV ⁴)	DOCUMENT ID	TECN	COMMENT
0.639 ± 0.056 ± 0.178	²¹⁷ CRONIN-HEN..01B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

²¹⁷ The leptons are required to have $P_1 > 1.5$ GeV/c. **$B \rightarrow X_c \ell \nu$ LEPTON MOMENTUM MOMENTS** **$R_0(\Gamma_{E_\ell > 1.7 \text{ GeV}} / \Gamma_{E_\ell > 1.5 \text{ GeV}})$**

VALUE	DOCUMENT ID	TECN	COMMENT
0.6187 ± 0.0014 ± 0.0016	²¹⁸ MAHMOOD	03 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

²¹⁸ The leptons are required to have $E_\ell > 1.5$ GeV in the B rest frame.

$R_1 (\langle E_l \rangle_{E_l > 1.5 \text{ GeV}})$


VALUE	DOCUMENT ID	TECN	COMMENT
1.7797 ± 0.0018 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.		

$1.7743 \pm 0.0019 \pm 0.0014$	219 AUBERT,B	04A BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$1.7792 \pm 0.0021 \pm 0.0027$	220 MAHMOOD	04 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
$1.7810 \pm 0.0007 \pm 0.0009$	221 MAHMOOD	03 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

219 The leptons are required to have $E_l > 1.5$ GeV in the B rest frame. The result with $E_l > 0.6$ GeV is also given.

220 Uses $E_e > 1.5$ GeV and also reports moments with other minimum minimum E_e conditions, as low as $E_e > 0.6$ GeV.

221 The leptons are required to have $E_l > 1.5$ GeV in the B rest frame.



$R_2 (\langle E_l^2 - \bar{E}_l^2 \rangle_{E_l > 1.5 \text{ GeV}})$

VALUE (10^{-3} GeV^2)	DOCUMENT ID	TECN	COMMENT
30.8 ± 0.8 OUR AVERAGE			

$30.3 \pm 0.9 \pm 0.5$	222 AUBERT,B	04A BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$31.6 \pm 0.8 \pm 1.0$	223 MAHMOOD	04 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

222 The leptons are required to have $E_l > 1.5$ GeV in the B rest frame. The result with $E_l > 0.6$ GeV is also given.

223 Uses $E_e > 1.5$ GeV and also reports moments with other minimum minimum E_e conditions, as low as $E_e > 0.6$ GeV.

$R_3 (\langle E_l^3 - \bar{E}_l^3 \rangle_{E_l > 1.5 \text{ GeV}})$

VALUE (10^{-3} GeV^3)	DOCUMENT ID	TECN	COMMENT
2.12 ± 0.47 ± 0.20	224 AUBERT,B	04A BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

224 The leptons are required to have $E_l > 1.5 \text{ GeV}$ in the B rest frame. The result with $E_l > 0.6 \text{ GeV}$ is also given.

 B^\pm/B^0 ADMIXTURE REFERENCES

AUBERT	06H	PR D73 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABAZOV	05O	PRL 95 171803	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ACOSTA	05F	PR D71 051103R	D. Acosta <i>et al.</i>	(CDF Collab.)
AUBERT	05	PRL 94 011801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05M	PRL 95 142003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05R	PR D72 052004	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05X	PRL 95 111801	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHOI	05	PRL 94 182002	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
IWASAKI	05	PR D72 092005	M. Iwasaki <i>et al.</i>	(BELLE Collab.)
LIMOSANI	05	PL B621 28	A. Limosani <i>et al.</i>	(BELLE Collab.)
MOHAPATRA	05	PR D72 011101R	D. Mohapatra <i>et al.</i>	(BELLE Collab.)
NISHIDA	05	PL B610 23	S. Nishida <i>et al.</i>	(BELLE Collab.)
OKABE	05	PL B614 27	T. Okabe <i>et al.</i>	(BELLE Collab.)
ABDALLAH	04D	EPJ C33 213	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
AUBERT	04C	PRL 92 111801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	04I	PRL 92 071802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04S	PR D69 052005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04X	PRL 93 011803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04A	PR D69 111104R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04E	PRL 93 021804	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04F	PRL 93 061801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04I	PRL 93 081802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	04A	PR D70 112006	B. Aubert <i>et al.</i>	(BABAR Collab.)
CSORNA	04	PR D70 032002	S.E. Csorna <i>et al.</i>	(CLEO Collab.)
KOPPENBURG	04	PRL 93 061803	P. Koppenburg <i>et al.</i>	(BELLE Collab.)
MAHMOOD	04	PR D70 032003	A.H. Mahmodd <i>et al.</i>	(CLEO Collab.)
NAKAO	04	PR D69 112001	M. Nakao <i>et al.</i>	(BELLE Collab.)
NISHIDA	04	PRL 93 031803	S. Nishida <i>et al.</i>	(BELLE Collab.)
ADAM	03B	PR D68 012004	N.E. Adam <i>et al.</i>	(CLEO Collab.)
AUBERT	03	PR D67 031101R	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03F	PR D67 032002	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03U	PRL 91 221802	B. Aubert <i>et al.</i>	(BaBar Collab.)
BONVICINI	03	PR D68 011101R	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
ISHIKAWA	03	PRL 91 261601	A. Ishikawa <i>et al.</i>	(BELLE Collab.)
KANEKO	03	PRL 90 021801	J. Kaneko <i>et al.</i>	(BELLE Collab.)
KROKOVNY	03B	PRL 91 262002	P. Krokovny <i>et al.</i>	(BELLE Collab.)
MAHMOOD	03	PR D67 072001	A.H. Mahmodd <i>et al.</i>	(CLEO Collab.)
ABE	02	PRL 88 021801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02L	PRL 89 011803	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02Y	PL B547 181	K. Abe <i>et al.</i>	(BELLE Collab.)
ANDERSON	02	PRL 89 282001	S. Anderson <i>et al.</i>	(CLEO Collab.)
AUBERT	02C	PRL 88 101805	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02G	PR D65 091104R	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02L	PRL 88 241801	B. Aubert <i>et al.</i>	(BaBar Collab.)
BORNHEIM	02	PRL 88 231803	A. Bornheim <i>et al.</i>	(CLEO Collab.)
EDWARDS	02B	PR D65 111102R	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ABE	01F	PL B511 151	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01J	PR D64 072001	K. Abe <i>et al.</i>	(BELLE Collab.)
ANDERSON	01B	PRL 87 181803	S. Anderson <i>et al.</i>	(CLEO Collab.)
CHEN	01	PR D63 031102	S. Chen <i>et al.</i>	(CLEO Collab.)
CHEN	01C	PRL 87 251807	S. Chen <i>et al.</i>	(CLEO Collab.)
COAN	01	PRL 86 5661	T.E. Coan <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN...	01B	PRL 87 251808	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PDG	01	Unofficial 2001 WWW edition		
ABREU	00R	PL B475 407	P. Abreu <i>et al.</i>	(DELPHI Collab.)
COAN	00	PRL 84 5283	T.E. Coan <i>et al.</i>	(CLEO Collab.)
RICHICHI	00	PRL 85 520	S.J. Richichi <i>et al.</i>	(CLEO Collab.)

BARATE	98Q	EPJ C4 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
BERGFELD	98	PRL 81 272	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BISHAI	98	PR D57 3847	M. Bishai <i>et al.</i>	(CLEO Collab.)
BONVICINI	98	PR D57 6604	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BROWDER	98	PRL 81 1786	T.E. Browder <i>et al.</i>	(CLEO Collab.)
COAN	98	PRL 80 1150	T.E. Coan <i>et al.</i>	(CLEO Collab.)
GLENN	98	PRL 80 2289	S. Glenn <i>et al.</i>	(CLEO Collab.)
ACKERSTAFF	97N	ZPHY C74 423	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
AMMAR	97	PR D55 13	R. Ammar <i>et al.</i>	(CLEO Collab.)
BARISH	97	PRL 79 3599	B. Barish <i>et al.</i>	(CLEO Collab.)
BUSKULIC	97B	ZPHY C73 601	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
GIBBONS	97B	PR D56 3783	L. Gibbons <i>et al.</i>	(CLEO Collab.)
ALBRECHT	96D	PL B374 256	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARISH	96B	PRL 76 1570	B.C. Barish <i>et al.</i>	(CLEO Collab.)
GIBAUT	96	PR D53 4734	D. Gibaut <i>et al.</i>	(CLEO Collab.)
KUBOTA	96	PR D53 6033	Y. Kubota <i>et al.</i>	(CLEO Collab.)
PDG	96	PR D54 1	R. M. Barnett <i>et al.</i>	
ALAM	95	PRL 74 2885	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	95D	PL B353 554	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BALEST	95B	PR D52 2661	R. Balest <i>et al.</i>	(CLEO Collab.)
BARISH	95	PR D51 1014	B.C. Barish <i>et al.</i>	(CLEO Collab.)
BUSKULIC	95B	PL B345 103	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ALBRECHT	94C	ZPHY C62 371	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	94J	ZPHY C61 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
PROCARIO	94	PRL 73 1472	M. Procaro <i>et al.</i>	(CLEO Collab.)
ALBRECHT	93	ZPHY C57 533	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	93E	ZPHY C60 11	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	93H	PL B318 397	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	93I	ZPHY C58 191	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	93B	PL B319 365	J. Alexander <i>et al.</i>	(CLEO Collab.)
ARTUSO	93	PL B311 307	M. Artuso	(SYRA)
BARTELT	93B	PRL 71 4111	J.E. Bartelt <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92E	PL B277 209	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92G	ZPHY C54 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92O	ZPHY C56 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO	92	PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
CRAWFORD	92	PR D45 752	G. Crawford <i>et al.</i>	(CLEO Collab.)
HENDERSON	92	PR D45 2212	S. Henderson <i>et al.</i>	(CLEO Collab.)
LESIAK	92	ZPHY C55 33	T. Lesiak <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	91C	PL B255 297	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	91H	ZPHY C52 353	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
FULTON	91	PR D43 651	R. Fulton <i>et al.</i>	(CLEO Collab.)
YANAGISAWA	91	PRL 66 2436	C. Yanagisawa <i>et al.</i>	(CUSB II Collab.)
ALBRECHT	90	PL B234 409	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	90H	PL B249 359	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO	90	PRL 64 2117	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
Also				
FULTON	90	PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
FULTON	90	PRL 64 16	R. Fulton <i>et al.</i>	(CLEO Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
PDG	90	PL B239	J.J. Hernandez <i>et al.</i>	(IFIC, BOST, CIT+)
ALBRECHT	89K	ZPHY C42 519	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ISGUR	89B	PR D39 799	N. Isgur <i>et al.</i>	(TNT0, CIT)
WACHS	89	ZPHY C42 33	K. Wachs <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	88E	PL B210 263	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	88H	PL B210 258	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KOERNER	88	ZPHY C38 511	J.G. Korner, G.A. Schuler	(MANZ, DESY)
ALAM	87	PRL 59 22	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALAM	87B	PRL 58 1814	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	87D	PL B199 451	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87H	PL B187 425	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BEAN	87	PR D35 3533	A. Bean <i>et al.</i>	(CLEO Collab.)
BEHRENDTS	87	PRL 59 407	S. Behrendts <i>et al.</i>	(CLEO Collab.)
BORTOLETTO	87	PR D35 19	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
ALAM	86	PR D34 3279	M.S. Alam <i>et al.</i>	(CLEO Collab.)
BALTRUSAIT...	86E	PRL 56 2140	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BORTOLETTO	86	PRL 56 800	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
HAAS	86	PRL 56 2781	J. Haas <i>et al.</i>	(CLEO Collab.)
ALBRECHT	85H	PL 162B 395	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
CSORNA	85	PRL 54 1894	S.E. Csorna <i>et al.</i>	(CLEO Collab.)
HAAS	85	PRL 55 1248	J. Haas <i>et al.</i>	(CLEO Collab.)
AVERY	84	PRL 53 1309	P. Avery <i>et al.</i>	(CLEO Collab.)

CHEN	84	PRL 52 1084	A. Chen <i>et al.</i>	(CLEO Collab.)
LEVMAN	84	PL 141B 271	G.M. Levman <i>et al.</i>	(CUSB Collab.)
ALAM	83B	PRL 51 1143	M.S. Alam <i>et al.</i>	(CLEO Collab.)
GREEN	83	PRL 51 347	J. Green <i>et al.</i>	(CLEO Collab.)
KLOPFEN...	83B	PL 130B 444	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
ALTARELLI	82	NP B208 365	G. Altarelli <i>et al.</i>	(ROMA, INFN, FRAS)
BRODY	82	PRL 48 1070	A.D. Brody <i>et al.</i>	(CLEO Collab.)
GIANNINI	82	NP B206 1	G. Giannini <i>et al.</i>	(CUSB Collab.)
BEBEK	81	PRL 46 84	C. Bebek <i>et al.</i>	(CLEO Collab.)
CHADWICK	81	PRL 46 88	K. Chadwick <i>et al.</i>	(CLEO Collab.)
ABRAMS	80	PRL 44 10	G.S. Abrams <i>et al.</i>	(SLAC, LBL)
