

# $\Upsilon(1S)$

$$J^{PC} = 0^{-}(1^{- -})$$

## $\Upsilon(1S)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>9460.30 ± 0.26 OUR AVERAGE</b>	Error includes scale factor of 3.3.		
9460.51 ± 0.09 ± 0.05	<sup>1</sup> ARTAMONOV 00	MD1	$e^+ e^- \rightarrow$ hadrons
9459.97 ± 0.11 ± 0.07	MACKAY 84	REDE	$e^+ e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
9460.60 ± 0.09 ± 0.05	<sup>2,3</sup> BARU 92B	REDE	$e^+ e^- \rightarrow$ hadrons
9460.59 ± 0.12	BARU 86	REDE	$e^+ e^- \rightarrow$ hadrons
9460.6 ± 0.4	<sup>3,4</sup> ARTAMONOV 84	REDE	$e^+ e^- \rightarrow$ hadrons
<sup>1</sup> Reanalysis of BARU 92B and ARTAMONOV 84 using new electron mass (COHEN 87).			
<sup>2</sup> Superseding BARU 86.			
<sup>3</sup> Superseded by ARTAMONOV 00.			
<sup>4</sup> Value includes data of ARTAMONOV 82.			

## $\Upsilon(1S)$ WIDTH

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>
<b>54.02 ± 1.25 OUR EVALUATION</b>	See the Note on "Width Determinations of the $\Upsilon$ States"

## $\Upsilon(1S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $\tau^+ \tau^-$	(2.60 ± 0.10) %	
$\Gamma_2$ $e^+ e^-$	(2.38 ± 0.11) %	
$\Gamma_3$ $\mu^+ \mu^-$	(2.48 ± 0.05) %	
<b>Hadronic decays</b>		
$\Gamma_4$ $\eta'(958)$ anything	(2.94 ± 0.24) %	
$\Gamma_5$ $J/\psi(1S)$ anything	(6.5 ± 0.7) × 10 <sup>-4</sup>	
$\Gamma_6$ $\chi_{c0}$ anything	< 5 × 10 <sup>-3</sup>	90%
$\Gamma_7$ $\chi_{c1}$ anything	(2.3 ± 0.7) × 10 <sup>-4</sup>	
$\Gamma_8$ $\chi_{c2}$ anything	(3.4 ± 1.0) × 10 <sup>-4</sup>	
$\Gamma_9$ $\psi(2S)$ anything	(2.7 ± 0.9) × 10 <sup>-4</sup>	
$\Gamma_{10}$ $\rho\pi$	< 2 × 10 <sup>-4</sup>	90%
$\Gamma_{11}$ $\pi^+ \pi^-$	< 5 × 10 <sup>-4</sup>	90%
$\Gamma_{12}$ $K^+ K^-$	< 5 × 10 <sup>-4</sup>	90%
$\Gamma_{13}$ $p\bar{p}$	< 5 × 10 <sup>-4</sup>	90%
$\Gamma_{14}$ $\pi^0 \pi^+ \pi^-$	< 1.84 × 10 <sup>-5</sup>	90%
$\Gamma_{15}$ $D^*(2010)^\pm$ anything		
$\Gamma_{16}$ $\bar{d}$ anything	(2.86 ± 0.28) × 10 <sup>-5</sup>	

### Radiative decays

$\Gamma_{17}$	$\gamma\pi^+\pi^-$	$(6.3 \pm 1.8) \times 10^{-5}$	
$\Gamma_{18}$	$\gamma\pi^0\pi^0$	$(1.7 \pm 0.7) \times 10^{-5}$	
$\Gamma_{19}$	$\gamma\pi^0\eta$	$< 2.4 \times 10^{-6}$	90%
$\Gamma_{20}$	$K^+K^-$ with $2 < m_{K^+K^-} < 3$ GeV	$(1.14 \pm 0.13) \times 10^{-5}$	
$\Gamma_{21}$	$\gamma p\bar{p}$ with $2 < m_{p\bar{p}} < 3$ GeV	$< 6 \times 10^{-6}$	90%
$\Gamma_{22}$	$\gamma 2h^+2h^-$	$(7.0 \pm 1.5) \times 10^{-4}$	
$\Gamma_{23}$	$\gamma 3h^+3h^-$	$(5.4 \pm 2.0) \times 10^{-4}$	
$\Gamma_{24}$	$\gamma 4h^+4h^-$	$(7.4 \pm 3.5) \times 10^{-4}$	
$\Gamma_{25}$	$\gamma\pi^+\pi^-K^+K^-$	$(2.9 \pm 0.9) \times 10^{-4}$	
$\Gamma_{26}$	$\gamma 2\pi^+2\pi^-$	$(2.5 \pm 0.9) \times 10^{-4}$	
$\Gamma_{27}$	$\gamma 3\pi^+3\pi^-$	$(2.5 \pm 1.2) \times 10^{-4}$	
$\Gamma_{28}$	$\gamma 2\pi^+2\pi^-K^+K^-$	$(2.4 \pm 1.2) \times 10^{-4}$	
$\Gamma_{29}$	$\gamma\pi^+\pi^-p\bar{p}$	$(1.5 \pm 0.6) \times 10^{-4}$	
$\Gamma_{30}$	$\gamma 2\pi^+2\pi^-p\bar{p}$	$(4 \pm 6) \times 10^{-5}$	
$\Gamma_{31}$	$\gamma 2K^+2K^-$	$(2.0 \pm 2.0) \times 10^{-5}$	
$\Gamma_{32}$	$\gamma\eta'(958)$	$< 1.6 \times 10^{-5}$	90%
$\Gamma_{33}$	$\gamma\eta$	$< 2.1 \times 10^{-5}$	90%
$\Gamma_{34}$	$\gamma f_0(980)$	$< 3 \times 10^{-5}$	90%
$\Gamma_{35}$	$\gamma f_2'(1525)$	$(3.7^{+1.2}_{-1.1}) \times 10^{-5}$	
$\Gamma_{36}$	$\gamma f_2(1270)$	$(1.01 \pm 0.09) \times 10^{-4}$	
$\Gamma_{37}$	$\gamma\eta(1405)$	$< 8.2 \times 10^{-5}$	90%
$\Gamma_{38}$	$\gamma f_0(1500)$	$< 1.5 \times 10^{-5}$	90%
$\Gamma_{39}$	$\gamma f_0(1710)$	$< 2.6 \times 10^{-4}$	90%
$\Gamma_{40}$	$\gamma f_0(1710) \rightarrow \gamma K^+K^-$	$< 7 \times 10^{-6}$	90%
$\Gamma_{41}$	$\gamma f_0(1710) \rightarrow \gamma\pi^0\pi^0$	$< 1.4 \times 10^{-6}$	90%
$\Gamma_{42}$	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$	$< 1.8 \times 10^{-6}$	90%
$\Gamma_{43}$	$\gamma f_4(2050)$	$< 5.3 \times 10^{-5}$	90%
$\Gamma_{44}$	$\gamma f_0(2200) \rightarrow \gamma K^+K^-$	$< 2 \times 10^{-4}$	90%
$\Gamma_{45}$	$\gamma f_J(2220) \rightarrow \gamma K^+K^-$	$< 8 \times 10^{-7}$	90%
$\Gamma_{46}$	$\gamma f_J(2220) \rightarrow \gamma\pi^+\pi^-$	$< 6 \times 10^{-7}$	90%
$\Gamma_{47}$	$\gamma f_J(2220) \rightarrow \gamma p\bar{p}$	$< 1.1 \times 10^{-6}$	90%
$\Gamma_{48}$	$\gamma\eta(2225) \rightarrow \gamma\phi\phi$	$< 3 \times 10^{-3}$	90%
$\Gamma_{49}$	$\gamma X$ ( $X$ = pseudoscalar with $m < 7.2$ GeV)	$< 3 \times 10^{-5}$	90%
$\Gamma_{50}$	$\gamma X\bar{X}$ ( $X\bar{X}$ = vectors with $m < 3.1$ GeV)	$< 1 \times 10^{-3}$	90%

### Other decays

$\Gamma_{51}$	invisible	$< 2.5 \times 10^{-3}$	90%
---------------	-----------	------------------------	-----

### $\Upsilon(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(e^+e^-) \times \Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$				$\Gamma_2\Gamma_3/\Gamma$
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>31.2±1.6±1.7</b>	KOBEL	92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_0\Gamma_2/\Gamma$
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>1.240±0.016 OUR AVERAGE</b>				
1.252±0.004±0.019	<sup>5</sup> ROSNER	06	CLEO	9.5 $e^+e^- \rightarrow \text{hadrons}$
1.187±0.023±0.031	<sup>5</sup> BARU	92B	MD1	$e^+e^- \rightarrow \text{hadrons}$
1.23 ±0.02 ±0.05	<sup>5</sup> JAKUBOWSKI	88	CBAL	$e^+e^- \rightarrow \text{hadrons}$
1.37 ±0.06 ±0.09	<sup>6</sup> GILES	84B	CLEO	$e^+e^- \rightarrow \text{hadrons}$
1.23 ±0.08 ±0.04	<sup>6</sup> ALBRECHT	82	DASP	$e^+e^- \rightarrow \text{hadrons}$
1.13 ±0.07 ±0.11	<sup>6</sup> NICZYPORUK	82	LENA	$e^+e^- \rightarrow \text{hadrons}$
1.09 ±0.25	<sup>6</sup> BOCK	80	CNTR	$e^+e^- \rightarrow \text{hadrons}$
1.35 ±0.14	<sup>7</sup> BERGER	79	PLUT	$e^+e^- \rightarrow \text{hadrons}$

<sup>5</sup> Radiative corrections evaluated following KURAEV 85.

<sup>6</sup> Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

<sup>7</sup> Radiative corrections reevaluated by ALEXANDER 89 using  $B(\mu\mu) = 0.026$ .

### $\Upsilon(1S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$				$\Gamma_2$
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>			
<b>1.340±0.018 OUR EVALUATION</b>				

### $\Upsilon(1S)$ BRANCHING RATIOS

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$				$\Gamma_1/\Gamma$
<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.60±0.10 OUR AVERAGE</b>				
2.53±0.13±0.05	60k	<sup>8</sup> BESSON	07	CLEO $e^+e^- \rightarrow \Upsilon(1S) \rightarrow \tau^+\tau^-$
2.61±0.12 <sup>+0.09</sup> <sub>-0.13</sub>	25k	CINABRO	94B	CLE2 $e^+e^- \rightarrow \tau^+\tau^-$
2.7 ±0.4 ±0.2		<sup>9</sup> ALBRECHT	85C	ARG $\Upsilon(2S) \rightarrow \pi^+\pi^-\tau^+\tau^-$
3.4 ±0.4 ±0.4		GILES	83	CLEO $e^+e^- \rightarrow \tau^+\tau^-$

<sup>8</sup> BESSON 07 reports  $[B(\Upsilon(1S) \rightarrow \tau^+\tau^-) / B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = 1.02 \pm 0.02 \pm 0.05$ .

We multiply by our best value  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>9</sup> Using  $B(\Upsilon(1S) \rightarrow ee) = B(\Upsilon(1S) \rightarrow \mu\mu) = 0.0256$ ; not used for width evaluations.

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_2/\Gamma$
<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.38±0.11 OUR AVERAGE</b>				
2.29±0.08±0.11		ALEXANDER	98	CLE2 $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
2.42±0.14±0.14	307	ALBRECHT	87	ARG $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
2.8 ±0.3 ±0.2	826	BESSON	84	CLEO $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
5.1 ±3.0		BERGER	80C	PLUT $e^+e^- \rightarrow e^+e^-$

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$			$\Gamma_3/\Gamma$		
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>0.0248 ± 0.0005 OUR AVERAGE</b>					
0.0249 ± 0.0002 ± 0.0007	345k	ADAMS	05	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0249 ± 0.0008 ± 0.0013		ALEXANDER	98	CLE2	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
0.0212 ± 0.0020 ± 0.0010		<sup>10</sup> BARU	92	MD1	$e^+e^- \rightarrow \mu^+\mu^-$
0.0231 ± 0.0012 ± 0.0010		<sup>10</sup> KOBEL	92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$
0.0252 ± 0.0007 ± 0.0007		CHEN	89B	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0261 ± 0.0009 ± 0.0011		KAARSBERG	89	CSB2	$e^+e^- \rightarrow \mu^+\mu^-$
0.0230 ± 0.0025 ± 0.0013	86	ALBRECHT	87	ARG	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
0.029 ± 0.003 ± 0.002	864	BESSION	84	CLEO	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
0.027 ± 0.003 ± 0.003		ANDREWS	83	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.032 ± 0.013 ± 0.003		ALBRECHT	82	DASP	$e^+e^- \rightarrow \mu^+\mu^-$
0.038 ± 0.015 ± 0.002		NICZYPORUK	82	LENA	$e^+e^- \rightarrow \mu^+\mu^-$
0.014 +0.034 -0.014		BOCK	80	CNTR	$e^+e^- \rightarrow \mu^+\mu^-$
0.022 ± 0.020		BERGER	79	PLUT	$e^+e^- \rightarrow \mu^+\mu^-$

<sup>10</sup> Taking into account interference between the resonance and continuum.

$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$			$\Gamma_1/\Gamma_3$		
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>1.02 ± 0.02 ± 0.05</b>					
	60k	BESSION	07	CLEO	$e^+e^- \rightarrow \Upsilon(1S)$

$\Gamma(\eta'(958) \text{ anything})/\Gamma_{\text{total}}$			$\Gamma_4/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<b>0.0294 ± 0.0024 OUR AVERAGE</b>					
0.030 ± 0.002 ± 0.002	AQUINES	06A	CLE3	$\Upsilon(1S) \rightarrow \eta' \text{ anything}$	
0.028 ± 0.004 ± 0.002	ARTUSO	03	CLE2	$\Upsilon(1S) \rightarrow \eta' \text{ anything}$	

$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$			$\Gamma_5/\Gamma$		
<u>VALUE (units 10<sup>-3</sup>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.65 ± 0.07 OUR AVERAGE</b>					
0.64 ± 0.04 ± 0.06		730 ± 40	BRIERE	04	CLEO $e^+e^- \rightarrow J/\psi X$
1.1 ± 0.4 ± 0.2			<sup>11</sup> FULTON	89	CLEO $e^+e^- \rightarrow \mu^+\mu^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.68	90	ALBRECHT	92J	ARG	$e^+e^- \rightarrow e^+e^- X, \mu^+\mu^- X$
<1.7	90	MASCHMANN	90	CBAL	$e^+e^- \rightarrow \text{hadrons}$
<20	90	NICZYPORUK	83	LENA	

<sup>11</sup> Using  $B((J/\psi) \rightarrow \mu^+\mu^-) = (6.9 \pm 0.9)\%$ .

$\Gamma(\chi_{c0} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$			$\Gamma_6/\Gamma_5$		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;7.4</b>					
	90	BRIERE	04	CLEO	$e^+e^- \rightarrow J/\psi X$

**$\Gamma(\chi_{c1} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$   $\Gamma_7/\Gamma_5$**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.35±0.08±0.06</b>	52 ± 12	BRIERE	04 CLEO	$e^+ e^- \rightarrow J/\psi X$

**$\Gamma(\chi_{c2} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$   $\Gamma_8/\Gamma_5$**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.52±0.12±0.09</b>	47 ± 11	BRIERE	04 CLEO	$e^+ e^- \rightarrow J/\psi X$

**$\Gamma(\psi(2S) \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$   $\Gamma_9/\Gamma_5$**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.41±0.11±0.08</b>	42 ± 11	BRIERE	04 CLEO	$e^+ e^- \rightarrow J/\psi \pi^+ \pi^- X$

**$\Gamma(\rho\pi)/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$**

<u>VALUE (units 10<sup>-4</sup>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; 2</b>	90	FULTON	90B	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<10	90	BLINOV	90 MD1	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$
<21	90	NICZYPORUK	83 LENA	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$

**$\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$**

<u>VALUE (units 10<sup>-4</sup>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;5</b>	90	BARU	92 MD1	$\Upsilon(1S) \rightarrow \pi^+ \pi^-$

**$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$**

<u>VALUE (units 10<sup>-4</sup>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;5</b>	90	BARU	92 MD1	$\Upsilon(1S) \rightarrow K^+ K^-$

**$\Gamma(p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$**

<u>VALUE (units 10<sup>-4</sup>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;5</b>	90	<sup>12</sup> BARU	96 MD1	$\Upsilon(1S) \rightarrow p\bar{p}$

<sup>12</sup>Supersedes BARU 92 in this node.

**$\Gamma(\pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$**

<u>VALUE (units 10<sup>-5</sup>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;1.84</b>	90	ANASTASSOV	99 CLE2	$e^+ e^- \rightarrow \text{hadrons}$

**$\Gamma(D^*(2010)^\pm \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$**

<u>VALUE (units 10<sup>-3</sup>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<19	90	<sup>13</sup> ALBRECHT	92J ARG	$e^+ e^- \rightarrow D^0 \pi^\pm X$

<sup>13</sup>For  $x_p > 0.2$ .

**$\Gamma(\bar{d} \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$**

<u>VALUE (units 10<sup>-5</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.86±0.19±0.21</b>	455	ASNER	07 CLEO	$e^+ e^- \rightarrow \bar{d} X$

$\Gamma(ggg, gg\gamma \rightarrow \bar{d} \text{ anything})/\Gamma(ggg, gg\gamma \rightarrow \text{anything})$

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>3.36 \pm 0.23 \pm 0.25</math></b>	455	ASNER 07	CLEO	$e^+ e^- \rightarrow \bar{d} X$

$\Gamma(\gamma\pi^+\pi^-)/\Gamma_{\text{total}}$

$\Gamma_{17}/\Gamma$

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>6.3 \pm 1.2 \pm 1.3</math></b>	<sup>14</sup> ANASTASSOV 99	CLE2	$e^+ e^- \rightarrow \text{hadrons}$

<sup>14</sup> For  $m_{\pi\pi} > 1$  GeV.

$\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$

$\Gamma_{18}/\Gamma$

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.7 \pm 0.6 \pm 0.3</math></b>	<sup>15</sup> ANASTASSOV 99	CLE2	$e^+ e^- \rightarrow \text{hadrons}$

<sup>15</sup> For  $m_{\pi\pi} > 1$  GeV.

$\Gamma(\gamma\pi^0\eta)/\Gamma_{\text{total}}$

$\Gamma_{19}/\Gamma$

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 2.4</math></b>	90	<sup>16</sup> BESSON 07A	CLEO	$e^+ e^- \rightarrow \gamma(1S)$

<sup>16</sup> BESSON 07A obtained this limit for  $0.7 < m_{\pi^0\eta} < 3$  GeV.

$\Gamma(K^+K^- \text{ with } 2 < m_{K^+K^-} < 3 \text{ GeV})/\Gamma_{\text{total}}$

$\Gamma_{20}/\Gamma$

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.14 \pm 0.08 \pm 0.10</math></b>	90	ATHAR 06	CLE3	$\gamma(1S) \rightarrow \gamma K^+ K^-$

$\Gamma(\gamma p\bar{p} \text{ with } 2 < m_{p\bar{p}} < 3 \text{ GeV})/\Gamma_{\text{total}}$

$\Gamma_{21}/\Gamma$

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 0.6</math></b>	90	ATHAR 06	CLE3	$\gamma(1S) \rightarrow \gamma p\bar{p}$

$\Gamma(\gamma 2h^+ 2h^-)/\Gamma_{\text{total}}$

$\Gamma_{22}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>7.0 \pm 1.1 \pm 1.0</math></b>	$80 \pm 12$	FULTON 90B	CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 3h^+ 3h^-)/\Gamma_{\text{total}}$

$\Gamma_{23}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>5.4 \pm 1.5 \pm 1.3</math></b>	$39 \pm 11$	FULTON 90B	CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 4h^+ 4h^-)/\Gamma_{\text{total}}$

$\Gamma_{24}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>7.4 \pm 2.5 \pm 2.5</math></b>	$36 \pm 12$	FULTON 90B	CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$

$\Gamma_{25}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.9 \pm 0.7 \pm 0.6</math></b>	$29 \pm 8$	FULTON 90B	CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{26}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.5 \pm 0.7 \pm 0.5</math></b>	$26 \pm 7$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{27}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.5 \pm 0.9 \pm 0.8</math></b>	$17 \pm 5$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{28}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.4 \pm 0.9 \pm 0.8</math></b>	$18 \pm 7$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma \pi^+ \pi^- \rho \bar{\rho})/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.5 \pm 0.5 \pm 0.3</math></b>	$22 \pm 6$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2\pi^+ 2\pi^- \rho \bar{\rho})/\Gamma_{\text{total}}$   $\Gamma_{30}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.4 \pm 0.4 \pm 0.4</math></b>	$7 \pm 6$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2K^+ 2K^-)/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.2 \pm 0.2</math></b>	$2 \pm 2$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma \eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 1.6</math></b>	90	RICHICHI 01B	CLE2	$\Upsilon(1S) \rightarrow \gamma \eta' \rightarrow \gamma \eta \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 130$	90	SCHMITT 88	CBAL	$\Upsilon(1S) \rightarrow \gamma X$

$\Gamma(\gamma \eta)/\Gamma_{\text{total}}$   $\Gamma_{33}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 2.1</math></b>	90	MASEK 02	CLEO	$\Upsilon(1S) \rightarrow \gamma \eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 28.2$	90	MASEK 02	CLEO	$\eta \rightarrow \gamma \gamma$
$< 6.7$	90	MASEK 02	CLEO	$\eta \rightarrow \pi^0 \pi^0 \pi^0$
$< 2.6$	90	MASEK 02	CLEO	$\eta \rightarrow \pi^+ \pi^- \pi^0$
$< 35$	90	SCHMITT 88	CBAL	$\Upsilon(1S) \rightarrow \gamma X$

$\Gamma(\gamma f_0(980))/\Gamma_{\text{total}}$   $\Gamma_{34}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 3</math></b>	90	<sup>17</sup> ATHAR 06	CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

<sup>17</sup> Assuming  $B(f_0(980) \rightarrow \pi \pi) = 1$ .

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$   $\Gamma_{35}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$3.7^{+0.9}_{-0.7} \pm 0.8$		ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<14	90	<sup>18</sup> FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
<19.4	90	<sup>18</sup> ALBRECHT	89	ARG $\Upsilon(1S) \rightarrow \gamma K^+ K^-$

<sup>18</sup> Assuming  $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$ .

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$   $\Gamma_{36}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>10.1 ± 0.9 OUR AVERAGE</b>				

$10.5 \pm 1.6^{+1.9}_{-1.8}$		<sup>19</sup> BESSON	07A	CLE3 $\Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$
$10.2 \pm 0.8 \pm 0.7$		ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
$8.1 \pm 2.3^{+2.9}_{-2.7}$		<sup>20</sup> ANASTASSOV	99	CLE2 $e^+ e^- \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90	<sup>20</sup> FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<13	90	<sup>20</sup> ALBRECHT	89	ARG $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<81	90	SCHMITT	88	CBAL $\Upsilon(1S) \rightarrow \gamma X$

<sup>19</sup> Using  $B(f_2(1270) \rightarrow \pi^0 \pi^0) = B(f_2(1270) \rightarrow \pi \pi)/3$  and  $B(f_2(1270) \rightarrow \pi \pi) = (0.845^{+0.025}_{-0.012})\%$ .

<sup>20</sup> Using  $B(f_2(1270) \rightarrow \pi \pi) = 0.84$ .

$\Gamma(\gamma \eta(1405))/\Gamma_{\text{total}}$   $\Gamma_{37}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;8.2</b>	90	<sup>21</sup> FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K^\pm \pi^\mp K_S^0$

<sup>21</sup> Includes unknown branching ratio of  $\eta(1405) \rightarrow K^\pm \pi^\mp K_S^0$ .

$\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$   $\Gamma_{38}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.5</b>	90	<sup>22</sup> BESSON	07A	CLEO $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.1	90	<sup>23</sup> BESSON	07A	CLEO $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \eta \eta$
------	----	----------------------	-----	--

<sup>22</sup> Using  $B(f_0(1500) \rightarrow \pi^0 \pi^0) = B(f_0(1500) \rightarrow \pi \pi)/3$  and  $B(f_0(1500) \rightarrow \pi \pi) = (0.349 \pm 0.023)\%$ .

<sup>23</sup> Calculated by us using  $B(f_0(1500) \rightarrow \eta \eta) = (5.1 \pm 0.9)\%$ .

$\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 2.6</b>	90	<sup>24</sup> ALBRECHT	89	ARG $\Upsilon(1S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.3	90	<sup>24</sup> FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
<19	90	<sup>24</sup> FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K_S^0 K_S^0$
< 8	90	<sup>25</sup> ALBRECHT	89	ARG $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<24	90	<sup>26</sup> SCHMITT	88	CBAL $\Upsilon(1S) \rightarrow \gamma X$



<sup>24</sup> Assuming  $B(f_0(1710) \rightarrow K\bar{K}) = 0.38$ .

<sup>25</sup> Assuming  $B(f_0(1710) \rightarrow \pi\pi) = 0.04$ .

<sup>26</sup> Assuming  $B(f_0(1710) \rightarrow \eta\eta) = 0.18$ .

**$\Gamma(\gamma f_0(1710) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{40}/\Gamma$**

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	ATHAR 06	CLEO	$e^+e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$

**$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^0 \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{41}/\Gamma$**

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	BESSON 07A	CLEO	$e^+e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$

**$\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$   $\Gamma_{42}/\Gamma$**

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.8	90	BESSON 07A	CLEO	$e^+e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \eta \eta$

**$\Gamma(\gamma f_4(2050))/\Gamma_{\text{total}}$   $\Gamma_{43}/\Gamma$**

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<5.3	90	<sup>27</sup> ATHAR 06	CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

<sup>27</sup> Assuming  $B(f_4(2050) \rightarrow \pi\pi) = 0.17$ .

**$\Gamma(\gamma f_0(2200) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{44}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0002	90	BARU 89	MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

**$\Gamma(\gamma f_J(2220) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{45}/\Gamma$**

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 8	90	ATHAR 06	CLE3	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 160	90	MASEK 02	CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 150	90	FULTON 90B	CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 290	90	ALBRECHT 89	ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
<2000	90	BARU 89	MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

**$\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{46}/\Gamma$**

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 6	90	ATHAR 06	CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<120	90	MASEK 02	CLEO	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
------	----	----------	------	---

**$\Gamma(\gamma f_J(2220) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$   $\Gamma_{47}/\Gamma$**

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 11	90	ATHAR 06	CLE3	$\Upsilon(1S) \rightarrow \gamma p \bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<160	90	MASEK 02	CLEO	$\Upsilon(1S) \rightarrow \gamma p \bar{p}$
------	----	----------	------	---

$\Gamma(\gamma\eta(2225) \rightarrow \gamma\phi\phi)/\Gamma_{\text{total}}$   $\Gamma_{48}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.003	90	BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^- K^+ K^-$

$\Gamma(\gamma X)/\Gamma_{\text{total}}$   $\Gamma_{49}/\Gamma$

( $X$  = pseudoscalar with  $m < 7.2$  GeV)

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	<sup>28</sup> BALEST	95 CLEO	$e^+ e^- \rightarrow \gamma + X$

<sup>28</sup> For a noninteracting pseudoscalar  $X$  with mass  $< 7.2$  GeV.

$\Gamma(\gamma X \bar{X})/\Gamma_{\text{total}}$   $\Gamma_{50}/\Gamma$

( $X \bar{X}$  = vectors with  $m < 3.1$  GeV)

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1	90	<sup>29</sup> BALEST	95 CLEO	$e^+ e^- \rightarrow \gamma + X \bar{X}$

<sup>29</sup> For a noninteracting vector  $X$  with mass  $< 3.1$  GeV.

$\Gamma(\text{invisible})/\Gamma_{\text{total}}$   $\Gamma_{51}/\Gamma$

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.25	90	TAJIMA	07 BELL	$\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.39	90	RUBIN	07 CLEO	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$
-------	----	-------	---------	---

## $\Upsilon(1S)$ REFERENCES

ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
BESSON	07A	PR D75 072001	D. Besson <i>et al.</i>	(CLEO Collab.)
RUBIN	07	PR D75 031104	P. Rubin <i>et al.</i>	(CLEO Collab.)
TAJIMA	07	PRL 98 132001	O. Tajima <i>et al.</i>	(BELLE Collab.)
AQUINES	06A	PR D74 092006	O. Aquines <i>et al.</i>	(CLEO Collab.)
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
BRIERE	04	PR D70 072001	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	03	PR D67 052003	M. Artuso <i>et al.</i>	(CLEO Collab.)
MASEK	02	PR D65 072002	G. Masek <i>et al.</i>	(CLEO Collab.)
RICHICHI	01B	PRL 87 141801	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
ANASTASSOV	99	PRL 82 286	A. Anastassov <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)
BALEST	95	PR D51 2053	R. Balest <i>et al.</i>	(CLEO Collab.)
CINABRO	94B	PL B340 129	D. Cinabro <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92J	ZPHY C55 25	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARU	92	ZPHY C54 229	S.E. Baru <i>et al.</i>	(NOVO)
BARU	92B	ZPHY C56 547	S.E. Baru <i>et al.</i>	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
BLINOV	90	PL B245 311	A.E. Blinov <i>et al.</i>	(NOVO)
FULTON	90B	PR D41 1401	R. Fulton <i>et al.</i>	(CLEO Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
BARU	89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
FULTON	89	PL B224 445	R. Fulton <i>et al.</i>	(CLEO Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)

BUCHMUEL...	88	HE $e^+e^-$ Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC
SCHMITT	88	ZPHY C40 199	P. Schmitt <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
BARU	86	ZPHY C30 551	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85C	PL 154B 452	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
MACKAY	84	PR D29 2483	W.W. MacKay <i>et al.</i>	(CUSB Collab.)
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)
GILES	83	PRL 50 877	R. Giles <i>et al.</i>	(HARV, OSU, ROCH, RUTG+)
NICZYPORUK	83	ZPHY C17 197	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
ARTAMONOV	82	PL 118B 225	A.S. Artamonov <i>et al.</i>	(NOVO)
NICZYPORUK	82	ZPHY C15 299	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BERGER	80C	PL 93B 497	C. Berger <i>et al.</i>	(PLUTO Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)
BERGER	79	ZPHY C1 343	C. Berger <i>et al.</i>	(PLUTO Collab.)

### OTHER RELATED PAPERS

BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
KOENIGS...	86	DESY 86/136	K. Koenigsmann	(DESY)
ALBRECHT	84	PL 134B 137	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
ARTAMONOV	82	PL 118B 225	A.S. Artamonov <i>et al.</i>	(NOVO)
BERGER	78	PL 76B 243	C. Berger <i>et al.</i>	(PLUTO Collab.)
BIELEIN	78	PL 78B 360	J.K. Bienlein <i>et al.</i>	(DESY, HAMB, HEIDP+)
DARDEN	78	PL 76B 246	C.W. Darden <i>et al.</i>	(DESY, DORT, HEIDH+)
GARELICK	78	PR D18 945	D.A. Garelick <i>et al.</i>	(NEAS, WASH, TUFTS)
KAPLAN	78	PRL 40 435	D.M. Kaplan <i>et al.</i>	(STON, FNAL, COLU)
YOH	78	PRL 41 684	J.K. Yoh <i>et al.</i>	(COLU, FNAL, STON)
COBB	77	PL 72B 273	J.H. Cobb <i>et al.</i>	(BNL, CERN, SYRA, YALE)
HERB	77	PRL 39 252	S.W. Herb <i>et al.</i>	(COLU, FNAL, STON)
INNES	77	PRL 39 1240	W.R. Innes <i>et al.</i>	(COLU, FNAL, STON)