

BOTTOM, CHARMED MESONS ($B = C = \pm 1$)

$$B_c^+ = c\bar{b}, B_c^- = \bar{c}b, \quad \text{similarly for } B_c^{*'}\text{'s}$$

B_c^\pm

$$I(J^P) = 0(0^-)$$

I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

$$\text{Mass } m = 6.277 \pm 0.006 \text{ GeV} \quad (S = 1.6)$$

$$\text{Mean life } \tau = (0.453 \pm 0.041) \times 10^{-12} \text{ s}$$

B_c^- modes are charge conjugates of the modes below.

B_c^+ DECAY MODES $\times B(\bar{b} \rightarrow B_c)$	Fraction (Γ_i/Γ)	Confidence level	P (MeV/c)
--	--------------------------------	------------------	----------------

The following quantities are not pure branching ratios; rather the fraction $\Gamma_i/\Gamma \times B(\bar{b} \rightarrow B_c)$.

$J/\psi(1S)\ell^+\nu_\ell$ anything	$(5.2_{-2.1}^{+2.4}) \times 10^{-5}$		—
$J/\psi(1S)\pi^+$	$< 8.2 \times 10^{-5}$	90%	2372
$J/\psi(1S)\pi^+\pi^+\pi^-$	$< 5.7 \times 10^{-4}$	90%	2352
$J/\psi(1S)a_1(1260)$	$< 1.2 \times 10^{-3}$	90%	2171
$D^*(2010)^+\bar{D}^0$	$< 6.2 \times 10^{-3}$	90%	2468