

$$I(J^P) = \frac{1}{2}(0^-)$$

A REVIEW GOES HERE – Check our WWW List of Reviews

K^\pm MASS


<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
493.677±0.016 OUR FIT	Error includes scale factor of 2.8.			
493.677±0.013 OUR AVERAGE	Error includes scale factor of 2.4. See the ideogram below.			
493.696±0.007	¹ DENISOV	91	CNTR –	Kaonic atoms
493.636±0.011	² GALL	88	CNTR –	Kaonic atoms
493.640±0.054	LUM	81	CNTR –	Kaonic atoms
493.670±0.029	BARKOV	79	EMUL ±	$e^+e^- \rightarrow K^+K^-$
493.657±0.020	² CHENG	75	CNTR –	Kaonic atoms
493.691±0.040	BACKENSTO...73	CNTR –		Kaonic atoms
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
493.631±0.007	GALL	88	CNTR –	K^- Pb (9→ 8)
493.675±0.026	GALL	88	CNTR –	K^- Pb (11→ 10)
493.709±0.073	GALL	88	CNTR –	K^- W (9→ 8)
493.806±0.095	GALL	88	CNTR –	K^- W (11→ 10)
493.640±0.022±0.008	³ CHENG	75	CNTR –	K^- Pb (9→ 8)
493.658±0.019±0.012	³ CHENG	75	CNTR –	K^- Pb (10→ 9)
493.638±0.035±0.016	³ CHENG	75	CNTR –	K^- Pb (11→ 10)
493.753±0.042±0.021	³ CHENG	75	CNTR –	K^- Pb (12→ 11)
493.742±0.081±0.027	³ CHENG	75	CNTR –	K^- Pb (13→ 12)

¹ Error increased from 0.0059 based on the error analysis in IVANOV 92.

² This value is the authors' combination of all of the separate transitions listed for this paper.

³ The CHENG 75 values for separate transitions were calculated from their Table 7 transition energies. The first error includes a 20% systematic error in the noncircular contaminant shift. The second error is due to a ±5 eV uncertainty in the theoretical transition energies.

WEIGHTED AVERAGE
 493.677 ± 0.013 (Error scaled by 2.4)



$m_{K^+} - m_{K^-}$

Test of *CPT*.


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG
-0.032 ± 0.090	1.5M	⁴ FORD	72	ASPK ±

⁴ FORD 72 uses $m_{\pi^+} - m_{\pi^-} = +28 \pm 70$ keV.

K^\pm MEAN LIFE

VALUE (10^{-8} s)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1.2380 ± 0.0021 OUR FIT		Error includes scale factor of 1.9.			
1.2379 ± 0.0021 OUR AVERAGE		Error includes scale factor of 1.9. See the ideogram below.			
1.2347 ± 0.0030	15M	⁵ AMBROSINO	08	KLOE ±	$\phi \rightarrow K^+ K^-$
1.2451 ± 0.0030	250k	KOPTEV	95	CNTR	<i>K</i> at rest, U target
1.2368 ± 0.0041	150k	KOPTEV	95	CNTR	<i>K</i> at rest, Cu target
1.2380 ± 0.0016	3M	OTT	71	CNTR +	<i>K</i> at rest
1.2272 ± 0.0036		LOBKOWICZ	69	CNTR +	<i>K</i> in flight
1.2443 ± 0.0038		FITCH	65B	CNTR +	<i>K</i> at rest
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1.2415 ± 0.0024	400k	⁶ KOPTEV	95	CNTR	<i>K</i> at rest
1.221 ± 0.011		FORD	67	CNTR ±	
1.231 ± 0.011		BOYARSKI	62	CNTR +	

- ⁵ Result obtained by averaging the decay length and decay time analyses taking correlations into account.
⁶ KOPTEV 95 report this weighted average of their U-target and Cu-target results, where they have weighted by $1/\sigma$ rather than $1/\sigma^2$.



$$(\tau_{K^+} - \tau_{K^-}) / \tau_{\text{average}}$$

This quantity is a measure of *CPT* invariance in weak interactions.

VALUE (%)	DOCUMENT ID	TECN
0.10 ± 0.09 OUR AVERAGE	Error includes scale factor of 1.2.	
-0.4 ± 0.4	AMBROSINO 08	KLOE
0.090 ± 0.078	LOBKOWICZ 69	CNTR
0.47 ± 0.30	FORD 67	CNTR

A REVIEW GOES HERE – Check our WWW List of Reviews

K^+ DECAY MODES

K^- modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
------	--------------------------------	-----------------------------------

Leptonic and semileptonic modes

Γ_1	$K^+ \rightarrow e^+ \nu_e$	$(1.584 \pm 0.020) \times 10^{-5}$	
Γ_2	$K^+ \rightarrow \mu^+ \nu_\mu$	$(63.55 \pm 0.11) \%$	S=1.2
Γ_3	$K^+ \rightarrow \pi^0 e^+ \nu_e$	$(5.07 \pm 0.04) \%$	S=2.1
	Called K_{e3}^+ .		
Γ_4	$K^+ \rightarrow \pi^0 \mu^+ \nu_\mu$	$(3.353 \pm 0.034) \%$	S=1.8
	Called $K_{\mu 3}^+$.		
Γ_5	$K^+ \rightarrow \pi^0 \pi^0 e^+ \nu_e$	$(2.2 \pm 0.4) \times 10^{-5}$	
Γ_6	$K^+ \rightarrow \pi^+ \pi^- e^+ \nu_e$	$(4.09 \pm 0.10) \times 10^{-5}$	
Γ_7	$K^+ \rightarrow \pi^+ \pi^- \mu^+ \nu_\mu$	$(1.4 \pm 0.9) \times 10^{-5}$	
Γ_8	$K^+ \rightarrow \pi^0 \pi^0 \pi^0 e^+ \nu_e$	$< 3.5 \times 10^{-6}$	CL=90%

Hadronic modes

Γ_9	$K^+ \rightarrow \pi^+ \pi^0$	$(20.66 \pm 0.08) \%$	S=1.2
Γ_{10}	$K^+ \rightarrow \pi^+ \pi^0 \pi^0$	$(1.761 \pm 0.022) \%$	S=1.1
Γ_{11}	$K^+ \rightarrow \pi^+ \pi^+ \pi^-$	$(5.59 \pm 0.04) \%$	S=1.3

Leptonic and semileptonic modes with photons

Γ_{12}	$K^+ \rightarrow \mu^+ \nu_\mu \gamma$	[a,b] $(6.2 \pm 0.8) \times 10^{-3}$	
Γ_{13}	$K^+ \rightarrow \mu^+ \nu_\mu \gamma (SD^+)$	[c,d] $(1.33 \pm 0.22) \times 10^{-5}$	
Γ_{14}	$K^+ \rightarrow \mu^+ \nu_\mu \gamma (SD^+ INT)$	[c,d] $< 2.7 \times 10^{-5}$	CL=90%
Γ_{15}	$K^+ \rightarrow \mu^+ \nu_\mu \gamma (SD^- + SD^- INT)$	[c,d] $< 2.6 \times 10^{-4}$	CL=90%
Γ_{16}	$K^+ \rightarrow e^+ \nu_e \gamma$	$(9.4 \pm 0.4) \times 10^{-6}$	
Γ_{17}	$K^+ \rightarrow \pi^0 e^+ \nu_e \gamma$	[a,b] $(2.56 \pm 0.16) \times 10^{-4}$	
Γ_{18}	$K^+ \rightarrow \pi^0 e^+ \nu_e \gamma (SD)$	[c,d] $< 5.3 \times 10^{-5}$	CL=90%
Γ_{19}	$K^+ \rightarrow \pi^0 \mu^+ \nu_\mu \gamma$	[a,b] $(1.5 \pm 0.4) \times 10^{-5}$	
Γ_{20}	$K^+ \rightarrow \pi^0 \pi^0 e^+ \nu_e \gamma$	$< 5 \times 10^{-6}$	CL=90%

Hadronic modes with photons or $\ell\bar{\ell}$ pairs

Γ_{21}	$K^+ \rightarrow \pi^+ \pi^0 \gamma$	[a,b] $(2.75 \pm 0.15) \times 10^{-4}$	
Γ_{22}	$K^+ \rightarrow \pi^+ \pi^0 \gamma (DE)$	[a,e] $(4.3 \pm 0.7) \times 10^{-6}$	
Γ_{23}	$K^+ \rightarrow \pi^+ \pi^0 \pi^0 \gamma$	[a,b] $(7.6 \pm_{-3.0}^{6.0}) \times 10^{-6}$	
Γ_{24}	$K^+ \rightarrow \pi^+ \pi^+ \pi^- \gamma$	[a,b] $(1.04 \pm 0.31) \times 10^{-4}$	
Γ_{25}	$K^+ \rightarrow \pi^+ \gamma \gamma$	[a] $(1.10 \pm 0.32) \times 10^{-6}$	
Γ_{26}	$K^+ \rightarrow \pi^+ 3\gamma$	[a] $< 1.0 \times 10^{-4}$	CL=90%
Γ_{27}	$K^\pm \rightarrow \pi^+ e^+ e^- \gamma$	$(1.19 \pm 0.13) \times 10^{-8}$	

Leptonic modes with $\ell\bar{\ell}$ pairs

Γ_{28}	$K^+ \rightarrow e^+ \nu_e \nu \bar{\nu}$	$< 6 \times 10^{-5}$	CL=90%
Γ_{29}	$K^+ \rightarrow \mu^+ \nu_\mu \nu \bar{\nu}$	$< 6.0 \times 10^{-6}$	CL=90%
Γ_{30}	$K^+ \rightarrow e^+ \nu_e e^+ e^-$	$(2.48 \pm 0.20) \times 10^{-8}$	
Γ_{31}	$K^+ \rightarrow \mu^+ \nu_\mu e^+ e^-$	$(7.06 \pm 0.31) \times 10^{-8}$	
Γ_{32}	$K^+ \rightarrow e^+ \nu_e \mu^+ \mu^-$	$(1.7 \pm 0.5) \times 10^{-8}$	
Γ_{33}	$K^+ \rightarrow \mu^+ \nu_\mu \mu^+ \mu^-$	$< 4.1 \times 10^{-7}$	CL=90%

**Lepton Family number (LF), Lepton number (L), $\Delta S = \Delta Q$ (SQ)
violating modes, or $\Delta S = 1$ weak neutral current (SI) modes**

Γ_{34}	$K^+ \rightarrow \pi^+ \pi^+ e^- \bar{\nu}_e$	SQ	< 1.2	$\times 10^{-8}$	CL=90%
Γ_{35}	$K^+ \rightarrow \pi^+ \pi^+ \mu^- \bar{\nu}_\mu$	SQ	< 3.0	$\times 10^{-6}$	CL=95%
Γ_{36}	$K^+ \rightarrow \pi^+ e^+ e^-$	SI	(3.00 \pm 0.09)	$\times 10^{-7}$	
Γ_{37}	$K^+ \rightarrow \pi^+ \mu^+ \mu^-$	SI	(8.1 \pm 1.4)	$\times 10^{-8}$	S=2.7
Γ_{38}	$K^+ \rightarrow \pi^+ \nu \bar{\nu}$	SI	(1.7 \pm 1.1)	$\times 10^{-10}$	
Γ_{39}	$K^+ \rightarrow \pi^+ \pi^0 \nu \bar{\nu}$	SI	< 4.3	$\times 10^{-5}$	CL=90%
Γ_{40}	$K^+ \rightarrow \mu^- \nu e^+ e^+$	LF	< 2.0	$\times 10^{-8}$	CL=90%
Γ_{41}	$K^+ \rightarrow \mu^+ \nu_e$	LF	[f] < 4	$\times 10^{-3}$	CL=90%
Γ_{42}	$K^+ \rightarrow \pi^+ \mu^+ e^-$	LF	< 1.3	$\times 10^{-11}$	CL=90%
Γ_{43}	$K^+ \rightarrow \pi^+ \mu^- e^+$	LF	< 5.2	$\times 10^{-10}$	CL=90%
Γ_{44}	$K^+ \rightarrow \pi^- \mu^+ e^+$	L	< 5.0	$\times 10^{-10}$	CL=90%
Γ_{45}	$K^+ \rightarrow \pi^- e^+ e^+$	L	< 6.4	$\times 10^{-10}$	CL=90%
Γ_{46}	$K^+ \rightarrow \pi^- \mu^+ \mu^+$	L	[f] < 3.0	$\times 10^{-9}$	CL=90%
Γ_{47}	$K^+ \rightarrow \mu^+ \bar{\nu}_e$	L	[f] < 3.3	$\times 10^{-3}$	CL=90%
Γ_{48}	$K^+ \rightarrow \pi^0 e^+ \bar{\nu}_e$	L	< 3	$\times 10^{-3}$	CL=90%
Γ_{49}	$K^+ \rightarrow \pi^+ \gamma$		[g] < 2.3	$\times 10^{-9}$	CL=90%

[a] See the Particle Listings below for the energy limits used in this measurement.

[b] Most of this radiative mode, the low-momentum γ part, is also included in the parent mode listed without γ 's.

[c] Structure-dependent part.

[d] See the "Note on $\pi^\pm \rightarrow l^\pm \nu \gamma$ and $K^\pm \rightarrow l^\pm \nu \gamma$ Form Factors" in the π^\pm Particle Listings for definitions and details.

[e] Direct-emission branching fraction.

[f] Derived from an analysis of neutrino-oscillation experiments.

[g] Violates angular-momentum conservation.

CONSTRAINED FIT INFORMATION

An overall fit to the mean life, a decay rate, and 13 branching ratios uses 32 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 51.8$ for 25 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_3	-64						
x_4	-62	90					
x_5	-3	4	3				
x_9	-65	1	-1	0			
x_{10}	-13	-6	-6	0	-6		
x_{11}	-21	-9	-9	0	-10	3	
Γ	5	2	2	0	2	-1	-24
	x_2	x_3	x_4	x_5	x_9	x_{10}	x_{11}

	Mode	Rate (10^8 s^{-1})	Scale factor
Γ_2	$K^+ \rightarrow \mu^+ \nu_\mu$	0.5133 ± 0.0013	1.5
Γ_3	$K^+ \rightarrow \pi^0 e^+ \nu_e$	0.0410 ± 0.0004	2.1
	Called K_{e3}^+ .		
Γ_4	$K^+ \rightarrow \pi^0 \mu^+ \nu_\mu$	0.02708 ± 0.00028	1.9
	Called $K_{\mu 3}^+$.		
Γ_5	$K^+ \rightarrow \pi^0 \pi^0 e^+ \nu_e$	$(1.77 \begin{smallmatrix} +0.35 \\ -0.30 \end{smallmatrix}) \times 10^{-5}$	
Γ_9	$K^+ \rightarrow \pi^+ \pi^0$	0.1669 ± 0.0007	1.3
Γ_{10}	$K^+ \rightarrow \pi^+ \pi^0 \pi^0$	0.01423 ± 0.00018	1.1
Γ_{11}	$K^+ \rightarrow \pi^+ \pi^+ \pi^-$	0.04518 ± 0.00029	1.2

K^\pm DECAY RATES

$\Gamma(\mu^+ \nu_\mu)$

Γ_2

VALUE (10^6 s^{-1}) DOCUMENT ID TECN CHG

51.33 ± 0.13 OUR FIT Error includes scale factor of 1.5.

• • • We do not use the following data for averages, fits, limits, etc. • • •

51.2 ± 0.8 FORD 67 CNTR ±

$\Gamma(\pi^+ \pi^+ \pi^-)$

Γ_{11}

<u>VALUE</u> (10^6 s^{-1})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
--	-------------	--------------------	-------------	------------

4.518 ± 0.029 OUR FIT Error includes scale factor of 1.2.

4.511 ± 0.024 ⁷FORD 70 ASPK

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.529 ± 0.032 3.2M ⁷FORD 70 ASPK

4.496 ± 0.030 ⁷FORD 67 CNTR ±

⁷ First FORD 70 value is second FORD 70 combined with FORD 67.

$(\Gamma(K^+) - \Gamma(K^-)) / \Gamma(K)$

$K^\pm \rightarrow \mu^\pm \nu_\mu$ RATE DIFFERENCE/AVERAGE

Test of *CPT* conservation.

<u>VALUE</u> (%)	<u>DOCUMENT ID</u>	<u>TECN</u>
------------------	--------------------	-------------

-0.54 ± 0.41 FORD 67 CNTR

$K^\pm \rightarrow \pi^\pm \pi^+ \pi^-$ RATE DIFFERENCE/AVERAGE

Test of *CP* conservation.

<u>VALUE</u> (%)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
------------------	-------------	--------------------	-------------	------------

0.08 ± 0.12 ⁸FORD 70 ASPK

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.02 ± 0.16 ⁹SMITH 73 ASPK ±

0.10 ± 0.14 3.2M ⁸FORD 70 ASPK

-0.50 ± 0.90 FLETCHER 67 OSPK

-0.04 ± 0.21 ⁸FORD 67 CNTR

⁸ First FORD 70 value is second FORD 70 combined with FORD 67.

⁹ SMITH 73 value of $K^\pm \rightarrow \pi^\pm \pi^+ \pi^-$ rate difference is derived from SMITH 73 value of $K^\pm \rightarrow \pi^\pm 2\pi^0$ rate difference.

$K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$ RATE DIFFERENCE/AVERAGE

Test of *CP* conservation.

<u>VALUE</u> (%)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
------------------	-------------	--------------------	-------------	------------

0.0 ± 0.6 OUR AVERAGE

0.08 ± 0.58 SMITH 73 ASPK ±

-1.1 ± 1.8 1802 HERZO 69 OSPK

$K^\pm \rightarrow \pi^\pm \pi^0$ RATE DIFFERENCE/AVERAGE

Test of *CPT* conservation.

<u>VALUE</u> (%)	<u>DOCUMENT ID</u>	<u>TECN</u>
------------------	--------------------	-------------

0.8 ± 1.2 HERZO 69 OSPK

$K^\pm \rightarrow \pi^\pm \pi^0 \gamma$ RATE DIFFERENCE/AVERAGE

Test of *CP* conservation.

<u>VALUE</u> (%)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
------------------	-------------	--------------------	-------------	------------	----------------

0.9 ± 3.3 OUR AVERAGE

0.8 ± 5.8 2461 SMITH 76 WIRE ± E_π 55–90 MeV

1.0 ± 4.0 4000 ABRAMS 73B ASPK ± E_π 51–100 MeV

K^+ BRANCHING RATIOS**Leptonic and semileptonic modes**

$$\Gamma(e^+\nu_e)/\Gamma(\mu^+\nu_\mu)$$

$$\Gamma_1/\Gamma_2$$

See the note on "Decay Constants of Charged Pseudoscalar Mesons" in the D_s^+ Listings.

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
2.493±0.025±0.019	13.8K	¹⁰ AMBROSINO 09E	KLOE	±
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.51 ±0.15	404	HEINTZE	76	SPEC +
2.37 ±0.17	534	HEARD	75B	SPEC +
2.42 ±0.42	112	CLARK	72	OSPK +

¹⁰The ratio is defined to include internal-bremsstrahlung, ignoring direct-emission contributions. AMBROSINO 09E determined the ratio from the measurement of $\Gamma(K \rightarrow e\nu(\gamma))$, $E_\gamma < 10$ MeV / $\Gamma(K \rightarrow \mu\nu(\gamma))$. 89.8% of $K \rightarrow e\nu(\gamma)$ events had $E_\gamma < 10$ MeV.

$$\Gamma(\mu^+\nu_\mu)/\Gamma_{\text{total}}$$

$$\Gamma_2/\Gamma$$

See the note on "Decay Constants of Charged Pseudoscalar Mesons" in the D_s^+ Listings.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
63.55±0.11 OUR FIT					Error includes scale factor of 1.2.
63.60±0.16 OUR AVERAGE					
63.66±0.09±0.15	865k	¹¹ AMBROSINO 06A	KLOE	+	
63.24±0.44	62k	CHIANG 72	OSPK	+	1.84 GeV/c K^+

¹¹Fully inclusive. Used tagged kaons from ϕ decays.

$$\Gamma(\pi^0 e^+\nu_e)/\Gamma_{\text{total}}$$

$$\Gamma_3/\Gamma$$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
5.07 ±0.04 OUR FIT					Error includes scale factor of 2.1.
4.94 ±0.05 OUR AVERAGE					
4.965±0.038±0.037		¹² AMBROSINO 08A	KLOE	±	
4.86 ±0.10	3516	CHIANG 72	OSPK	+	1.84 GeV/c K^+
• • • We do not use the following data for averages, fits, limits, etc. • • •					
4.7 ±0.3	429	SHAKLEE	64	HLBC +	
5.0 ±0.5		ROE	61	HLBC +	

¹²Depends on K^+ lifetime τ . AMBROSINO 08A uses PDG 06 value of $\tau = (1.2385 \pm 0.0024) \times 10^{-8}$ sec. The correlation between K_{e3}^+ and $K_{\mu3}^+$ branching fraction measurements is 62.7%.

$$\Gamma(\pi^0 e^+\nu_e)/\Gamma(\mu^+\nu_\mu)$$

$$\Gamma_3/\Gamma_2$$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
0.0798±0.0008 OUR FIT				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.069 ±0.006	350	ZELLER	69	ASPK +
0.0775±0.0033	960	BOTTERILL	68C	ASPK +
0.069 ±0.006	561	GARLAND	68	OSPK +
0.0791±0.0054	295	¹³ AUERBACH	67	OSPK +

¹³ AUERBACH 67 changed from 0.0797 ± 0.0054 . See comment with ratio $\Gamma(\pi^0 \mu^+ \nu_\mu) / \Gamma(\mu^+ \nu_\mu)$. The value 0.0785 ± 0.0025 given in AUERBACH 67 is an average of AUERBACH 67 $\Gamma(\pi^0 e^+ \nu_e) / \Gamma(\mu^+ \nu_\mu)$ and CESTER 66 $\Gamma(\pi^0 e^+ \nu_e) / [\Gamma(\mu^+ \nu_\mu) + \Gamma(\pi^+ \pi^0)]$.

$$\Gamma(\pi^0 e^+ \nu_e) / [\Gamma(\mu^+ \nu_\mu) + \Gamma(\pi^+ \pi^0)] \quad \Gamma_3 / (\Gamma_2 + \Gamma_9)$$

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN CHG

6.02 ± 0.06 OUR FIT Error includes scale factor of 2.1.

6.02 ± 0.15 OUR AVERAGE

6.16 ± 0.22 5110 ESCHSTRUTH 68 OSPK +

5.89 ± 0.21 1679 CESTER 66 OSPK +

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.92 ± 0.65 ¹⁴ WEISSENBE... 76 SPEC +

¹⁴ Value calculated from WEISSENBERG 76 ($\pi^0 e \nu$), ($\mu \nu$), and ($\pi \pi^0$) values to eliminate dependence on our 1974 ($\pi 2\pi^0$) and ($\pi \pi^+ \pi^-$) fractions.

$$\Gamma(\pi^0 e^+ \nu_e) / [\Gamma(\pi^0 \mu^+ \nu_\mu) + \Gamma(\pi^+ \pi^0) + \Gamma(\pi^+ \pi^0 \pi^0)] \quad \Gamma_3 / (\Gamma_4 + \Gamma_9 + \Gamma_{10})$$

VALUE EVTS DOCUMENT ID TECN CHG

0.1968 ± 0.0016 OUR FIT Error includes scale factor of 2.4.

0.1962 ± 0.0008 ± 0.0035 71k SHER 03 B865 +

$$\Gamma(\pi^0 e^+ \nu_e) / \Gamma(\pi^+ \pi^0) \quad \Gamma_3 / \Gamma_9$$

VALUE EVTS DOCUMENT ID TECN CHG COMMENT

0.2455 ± 0.0023 OUR FIT Error includes scale factor of 2.6.

0.2470 ± 0.0009 ± 0.0004 87k BATLEY 07A NA48 ±

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.221 ± 0.012 786 ¹⁵ LUCAS 73B HBC - Dalitz pairs only

¹⁵ LUCAS 73B gives $N(K_{e3}) = 786 \pm 3.1\%$, $N(2\pi) = 3564 \pm 3.1\%$. We use these values to obtain quoted result.

$$\Gamma(\pi^0 e^+ \nu_e) / \Gamma(\pi^+ \pi^+ \pi^-) \quad \Gamma_3 / \Gamma_{11}$$

VALUE EVTS DOCUMENT ID TECN CHG

0.907 ± 0.010 OUR FIT Error includes scale factor of 1.6.

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.867 ± 0.027 2768 BARMIN 87 XEBC +

0.856 ± 0.040 2827 BRAUN 75 HLBC +

0.850 ± 0.019 4385 ¹⁶ HAIDT 71 HLBC +

0.846 ± 0.021 4385 ¹⁶ EICHTEN 68 HLBC +

0.94 ± 0.09 854 BELLOTTI 67B HLBC

0.90 ± 0.06 230 BORREANI 64 HBC +

¹⁶ HAIDT 71 is a reanalysis of EICHTEN 68. Not included in average because of large discrepancy in $\Gamma(\pi^0 \mu^+ \nu) / \Gamma(\pi^0 e^+ \nu)$ with more precise results.

$\Gamma(\pi^0 \mu^+ \nu_\mu) / \Gamma_{\text{total}}$ Γ_4 / Γ

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN CHG COMMENT

3.353 ± 0.034 OUR FIT Error includes scale factor of 1.8.

3.24 ± 0.04 OUR AVERAGE

3.233 ± 0.029 ± 0.026 ¹⁷ AMBROSINO 08A KLOE ±
 3.33 ± 0.16 2345 CHIANG 72 OSPK + 1.84 GeV/c K^+

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.8 ± 0.4 ¹⁸ TAYLOR 59 EMUL +

¹⁷ Depends on K^+ lifetime τ . AMBROSINO 08A uses PDG 06 value of $\tau = (1.2385 \pm 0.0024) \times 10^{-8}$ sec. The correlation between K_{e3}^+ and $K_{\mu 3}^+$ branching fraction measurements is 62.7%.

¹⁸ Earlier experiments not averaged.

$\Gamma(\pi^0 \mu^+ \nu_\mu) / \Gamma(\mu^+ \nu_\mu)$ Γ_4 / Γ_2

VALUE EVTS DOCUMENT ID TECN CHG

0.0528 ± 0.0006 OUR FIT Error includes scale factor of 1.8.

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.054 ± 0.009 240 ZELLER 69 ASPK +

0.0480 ± 0.0037 424 ¹⁹ GARLAND 68 OSPK +

0.0486 ± 0.0040 307 ²⁰ AUERBACH 67 OSPK +

¹⁹ GARLAND 68 changed from 0.055 ± 0.004 in agreement with μ -spectrum calculation of GAILLARD 70 appendix B. L.G.Pondrom, (private communication 73).

²⁰ AUERBACH 67 changed from 0.0602 ± 0.0046 by erratum which brings the μ -spectrum calculation into agreement with GAILLARD 70 appendix B.

$\Gamma(\pi^0 \mu^+ \nu_\mu) / \Gamma(\pi^0 e^+ \nu_e)$ Γ_4 / Γ_3

VALUE EVTS DOCUMENT ID TECN CHG COMMENT

0.6608 ± 0.0030 OUR FIT Error includes scale factor of 1.1.

0.6618 ± 0.0027 OUR AVERAGE

0.6511 ± 0.0064 ²¹ AMBROSINO 08A KLOE ±

0.663 ± 0.003 ± 0.001 77k BATLEY 07A NA48 ±

0.671 ± 0.007 ± 0.008 24k HORIE 01 SPEC

0.670 ± 0.014 ²² HEINTZE 77 SPEC +

0.667 ± 0.017 5601 BOTTERILL 68B ASPK +

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.608 ± 0.014 1585 ²³ BRAUN 75 HLBC +

0.705 ± 0.063 554 ²⁴ LUCAS 73B HBC - Dalitz pairs only

0.698 ± 0.025 3480 ²⁵ CHIANG 72 OSPK + 1.84 GeV/c K^+

0.596 ± 0.025 ²⁶ HAIDT 71 HLBC +

0.604 ± 0.022 1398 ²⁶ EICHTEN 68 HLBC

0.703 ± 0.056 1509 CALLAHAN 66B HLBC

²¹ Not used in the fit. This result enters the fit via correlation of K_{e3}^+ and $K_{\mu 3}^+$ branching fraction measurements of AMBROSINO 08A.

²² HEINTZE 77 value from fit to λ_0 . Assumes μ -e universality.

²³ BRAUN 75 value is from form factor fit. Assumes μ -e universality.

²⁴ LUCAS 73B gives $N(K_{\mu 3}) = 554 \pm 7.6\%$, $N(K_{e3}) = 786 \pm 3.1\%$. We divide.

²⁵ CHIANG 72 $\Gamma(\pi^0 \mu^+ \nu_\mu) / \Gamma(\pi^0 e^+ \nu_e)$ is statistically independent of CHIANG 72 $\Gamma(\pi^0 \mu^+ \nu_\mu) / \Gamma_{\text{total}}$ and $\Gamma(\pi^0 e^+ \nu_e) / \Gamma_{\text{total}}$.

²⁶ HAIDT 71 is a reanalysis of EICHTEN 68. Not included in average because of large discrepancy with more precise results.

$$\frac{\Gamma(\pi^0 \mu^+ \nu_\mu) + \Gamma(\pi^+ \pi^0)}{\Gamma_{\text{total}}} \quad (\Gamma_4 + \Gamma_9) / \Gamma$$

We combine these two modes for experiments measuring them in xenon bubble chamber because of difficulties of separating them there.

<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
--------------------------------------	-------------	--------------------	-------------	------------

24.02 ± 0.08 OUR FIT Error includes scale factor of 1.2.

• • • We do not use the following data for averages, fits, limits, etc. • • •

25.4 ± 0.9	886	SHAKLEE	64	HLBC	+
------------	-----	---------	----	------	---

23.4 ± 1.1		ROE	61	HLBC	+
------------	--	-----	----	------	---

$$\frac{\Gamma(\pi^0 \mu^+ \nu_\mu)}{\Gamma(\pi^+ \pi^0)} \quad \Gamma_4 / \Gamma_9$$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
--------------	-------------	--------------------	-------------	------------

0.1637 ± 0.0006 ± 0.0003 77k BATLEY 07A NA48 ±

$$\frac{\Gamma(\pi^0 \mu^+ \nu_\mu)}{\Gamma(\pi^+ \pi^+ \pi^-)} \quad \Gamma_4 / \Gamma_{11}$$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	------------	----------------

0.599 ± 0.007 OUR FIT Error includes scale factor of 1.6.

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.503 ± 0.019	1505	²⁷ HAIDT	71	HLBC	+
---------------	------	---------------------	----	------	---

0.510 ± 0.017	1505	²⁷ EICHTEN	68	HLBC	+
---------------	------	-----------------------	----	------	---

0.63 ± 0.07	2845	²⁸ BISI	65B	BC	+	HBC+HLBC
-------------	------	--------------------	-----	----	---	----------

²⁷ HAIDT 71 is a reanalysis of EICHTEN 68. Not included in average because of large discrepancy in $\Gamma(\pi^0 \mu^+ \nu) / \Gamma(\pi^0 e^+ \nu)$ with more precise results.

²⁸ Error enlarged for background problems. See GAILLARD 70.

$$\frac{\Gamma(\pi^0 \pi^0 e^+ \nu_e)}{\Gamma_{\text{total}}} \quad \Gamma_5 / \Gamma$$

<u>VALUE (units 10⁻⁵)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
--------------------------------------	-------------	--------------------	-------------	------------

2.2 ± 0.4 OUR FIT

2.54 ± 0.89	10	BARMIN	88B	HLBC	+
--------------------	----	--------	-----	------	---

$$\frac{\Gamma(\pi^0 \pi^0 e^+ \nu_e)}{\Gamma(\pi^0 e^+ \nu_e)} \quad \Gamma_5 / \Gamma_3$$

<u>VALUE (units 10⁻⁴)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
--------------------------------------	-------------	--------------------	-------------	------------

4.3^{+0.9}_{-0.7} OUR FIT

4.1^{+1.0}_{-0.7} OUR AVERAGE

4.2 ^{+1.0} _{-0.9}	25	BOLOTOV	86B	CALO	-
-------------------------------------	----	---------	-----	------	---

3.8 ^{+5.0} _{-1.2}	2	LJUNG	73	HLBC	+
-------------------------------------	---	-------	----	------	---

$$\frac{\Gamma(\pi^+ \pi^- e^+ \nu_e)}{\Gamma(\pi^+ \pi^+ \pi^-)} \quad \Gamma_6 / \Gamma_{11}$$

<u>VALUE (units 10⁻⁴)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
--------------------------------------	-------------	--------------------	-------------	------------

7.31 ± 0.16 OUR AVERAGE

7.35 ± 0.01 ± 0.19	388k	²⁹ PISLAK	01	B865	
--------------------	------	----------------------	----	------	--

7.21 ± 0.32	30k	ROSSELET	77	SPEC	+
-------------	-----	----------	----	------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.36 ± 0.68	500	BOURQUIN	71	ASPK	
-------------	-----	----------	----	------	--

7.0 ± 0.9	106	SCHWEINB...	71	HLBC	+
-----------	-----	-------------	----	------	---

5.83 ± 0.63	269	ELY	69	HLBC	+
-------------	-----	-----	----	------	---

²⁹ PISLAK 01 reports $\Gamma(\pi^+\pi^-\mu^+\nu_\mu)/\Gamma_{\text{total}} = (4.109 \pm 0.008 \pm 0.110) \times 10^{-5}$ using the PDG 00 value $\Gamma(\pi^+\pi^+\pi^-)/\Gamma_{\text{total}} = (5.59 \pm 0.05) \times 10^{-2}$. We divide by the PDG value and unfold its error from the systematic error. PISLAK 03 gives additional details on the branching ratio measurement and gives improved errors on the S -wave π - π scattering length: $a_0^0 = 0.216 \pm 0.013(\text{stat.}) \pm 0.002(\text{syst.}) \pm 0.002(\text{theor.})$.

$\Gamma(\pi^+\pi^-\mu^+\nu_\mu)/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.77^{+0.54}_{-0.50}$	1	CLINE	65	FBC +

$\Gamma(\pi^+\pi^-\mu^+\nu_\mu)/\Gamma(\pi^+\pi^+\pi^-)$ Γ_7/Γ_{11}

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
2.57 ± 1.55	7	BISI	67	DBC +
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 2.5	1	GREINER	64	EMUL +

$\Gamma(\pi^0\pi^0\pi^0e^+\nu_e)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
<3.5	90	0	BOLOTOV	88	SPEC -
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<9	90	0	BARMIN	92	XEBC +

————— Hadronic modes —————

$\Gamma(\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
20.66 ± 0.08 OUR FIT	Error includes scale factor of 1.2.				
20.70 ± 0.16 OUR AVERAGE	Error includes scale factor of 1.8.				
$20.65 \pm 0.05 \pm 0.08$	1.4M	³⁰ AMBROSINO	08E	KLOE +	$\phi \rightarrow K^+K^-$
21.18 ± 0.28	16k	CHIANG	72	OSPK +	1.84 GeV/c K^+
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
21.0 ± 0.6		CALLAHAN	65	HLBC	See Γ_9/Γ_{11}
³⁰ Fully inclusive of final-state radiation. The branching ratio is evaluated using K^+ lifetime, $\tau = 12.385$ ns.					

$\Gamma(\pi^+\pi^0)/\Gamma(\pi^+\pi^+\pi^-)$ Γ_9/Γ_{11}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
3.694 ± 0.029 OUR FIT	Error includes scale factor of 1.2.			
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3.96 ± 0.15	1045	CALLAHAN	66	FBC +

$\Gamma(\pi^+\pi^0)/\Gamma(\mu^+\nu_\mu)$ Γ_9/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	------------	----------------

0.3252±0.0016 OUR FIT Error includes scale factor of 1.2.

0.3325±0.0032 OUR AVERAGE

0.3329±0.0047±0.0010	45k	USHER	92	SPEC	+	$p\bar{p}$ at rest
----------------------	-----	-------	----	------	---	--------------------

0.3355±0.0057		³¹ WEISSENBE...	76	SPEC	+	
---------------	--	----------------------------	----	------	---	--

0.3277±0.0065	4517	³² AUERBACH	67	OSPK	+	
---------------	------	------------------------	----	------	---	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.328 ±0.005	25k	³¹ WEISSENBE...	74	STRC	+	
--------------	-----	----------------------------	----	------	---	--

0.305 ±0.018	1600	ZELLER	69	ASPK	+	
--------------	------	--------	----	------	---	--

³¹ WEISSENBERG 76 revises WEISSENBERG 74.

³² AUERBACH 67 changed from 0.3253 ± 0.0065. See comment with ratio $\Gamma(\pi^0\mu^+\nu_\mu)/\Gamma(\mu^+\nu_\mu)$.

$\Gamma(\pi^+\pi^0\pi^0)/\Gamma_{total}$ Γ_{10}/Γ

<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------------------------------	-------------	--------------------	-------------	------------	----------------

1.761±0.022 OUR FIT Error includes scale factor of 1.1.

1.775±0.028 OUR AVERAGE Error includes scale factor of 1.2.

1.763±0.013±0.022		ALOISIO	04A	KLOE	±	
-------------------	--	---------	-----	------	---	--

1.84 ±0.06	1307	CHIANG	72	OSPK	+	1.84 GeV/c K^+
------------	------	--------	----	------	---	------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.53 ±0.11	198	³³ PANDOULAS	70	EMUL	+	
------------	-----	-------------------------	----	------	---	--

1.8 ±0.2	108	SHAKLEE	64	HLBC	+	
----------	-----	---------	----	------	---	--

1.7 ±0.2		ROE	61	HLBC	+	
----------	--	-----	----	------	---	--

1.5 ±0.2		³⁴ TAYLOR	59	EMUL	+	
----------	--	----------------------	----	------	---	--

³³ Includes events of TAYLOR 59.

³⁴ Earlier experiments not averaged.

$\Gamma(\pi^+\pi^0\pi^0)/\Gamma(\pi^+\pi^0)$ Γ_{10}/Γ_9

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	------------	----------------

0.0852±0.0011 OUR FIT Error includes scale factor of 1.1.

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.081 ±0.005	574	³⁵ LUCAS	73B	HBC	-	Dalitz pairs only
--------------	-----	---------------------	-----	-----	---	-------------------

³⁵ LUCAS 73B gives $N(\pi 2\pi^0) = 574 \pm 5.9\%$, $N(2\pi) = 3564 \pm 3.1\%$. We quote $0.5N(\pi 2\pi^0)/N(2\pi)$ where 0.5 is because only Dalitz pair π^0 's were used.

$\Gamma(\pi^+\pi^0\pi^0)/\Gamma(\pi^+\pi^+\pi^-)$ Γ_{10}/Γ_{11}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	------------	----------------

0.315±0.004 OUR FIT Error includes scale factor of 1.1.

0.303±0.009	2027	BISI	65	BC	+	HBC+HLBC
--------------------	------	------	----	----	---	----------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.393±0.099	17	YOUNG	65	EMUL	+	
-------------	----	-------	----	------	---	--

$\Gamma(\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{11}/Γ**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	------------	----------------

5.59±0.04 OUR FIT Error includes scale factor of 1.3.

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.56±0.20	2330	³⁶ CHIANG	72	OSPK +	1.84 GeV/c K^+
5.34±0.21	693	³⁷ PANDOULAS	70	EMUL +	
5.71±0.15		DEMARCO	65	HBC	
6.0 ±0.4	44	YOUNG	65	EMUL +	
5.54±0.12	2332	CALLAHAN	64	HLBC +	
5.1 ±0.2	540	SHAKLEE	64	HLBC +	
5.7 ±0.3		ROE	61	HLBC +	

³⁶ Value is not independent of CHIANG 72 $\Gamma(\mu^+\nu_\mu)/\Gamma_{\text{total}}$, $\Gamma(\pi^+\pi^0)/\Gamma_{\text{total}}$, $\Gamma(\pi^+\pi^0\pi^0)/\Gamma_{\text{total}}$, $\Gamma(\pi^0\mu^+\nu_\mu)/\Gamma_{\text{total}}$, and $\Gamma(\pi^0e^+\nu_e)/\Gamma_{\text{total}}$.

³⁷ Includes events of TAYLOR 59.

————— **Leptonic and semileptonic modes with photons** —————

$\Gamma(\mu^+\nu_\mu\gamma)/\Gamma_{\text{total}}$ **Γ_{12}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	------------	----------------

6.2±0.8 OUR AVERAGE

6.6±1.5	38,39	DEMIDOV	90	XEBC	$P(\mu) < 231.5 \text{ MeV}/c$
6.0±0.9		BARMIN	88	HLBC +	$P(\mu) < 231.5 \text{ MeV}/c$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.5±0.8	39,40	DEMIDOV	90	XEBC	$E(\gamma) > 20 \text{ MeV}$
3.2±0.5	57	⁴¹ BARMIN	88	HLBC +	$E(\gamma) > 20 \text{ MeV}$
5.4±0.3		⁴² AKIBA	85	SPEC	$P(\mu) < 231.5 \text{ MeV}/c$

³⁸ $P(\mu)$ cut given in DEMIDOV 90 paper, 235.1 MeV/c, is a misprint according to authors (private communication).

³⁹ DEMIDOV 90 quotes only inner bremsstrahlung (IB) part.

⁴⁰ Not independent of above DEMIDOV 90 value. Cuts differ.

⁴¹ Not independent of above BARMIN 88 value. Cuts differ.

⁴² Assumes μ - e universality and uses constraints from $K \rightarrow e\nu\gamma$.

$\Gamma(\mu^+\nu_\mu\gamma(\text{SD}^+))/\Gamma_{\text{total}}$ **Γ_{13}/Γ**

Structure-dependent part with $+\gamma$ helicity (SD^+ term). See the “Note on $\pi^\pm \rightarrow \ell^\pm\nu\gamma$ and $K^\pm \rightarrow \ell^\pm\nu\gamma$ Form Factors” in the π^\pm section of the Particle Data Listings above.

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
---	------------	-------------	--------------------	-------------

1.33±0.12±0.18 2588 ⁴³ ADLER 00B B787

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.0	90	AKIBA	85	SPEC
------	----	-------	----	------

⁴³ ADLER 00B obtains the branching ratio by extrapolating the measurement in the kinematic region $E_\mu > 137 \text{ MeV}$, $E_\gamma > 90 \text{ MeV}$ to the full SD^+ phase-space. Also reports $|F_V + F_A| = 0.165 \pm 0.007 \pm 0.011$ and $-0.04 < F_V - F_A < 0.24$ at 90% CL.

$\Gamma(\mu^+ \nu_\mu \gamma(\text{SD}^+ \text{INT}))/\Gamma_{\text{total}}$ Γ_{14}/Γ

Interference term between internal Bremsstrahlung and SD^+ term. See the “Note on $\pi^\pm \rightarrow \ell^\pm \nu_\ell \gamma$ and $K^\pm \rightarrow \ell^\pm \nu_\ell \gamma$ Form Factors” in the π^\pm section of the Particle Data Listings above.

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
<2.7	90	AKIBA	85 SPEC

$\Gamma(\mu^+ \nu_\mu \gamma(\text{SD}^- + \text{SD}^- \text{INT}))/\Gamma_{\text{total}}$ Γ_{15}/Γ

Sum of structure-dependent part with $-\gamma$ helicity (SD^- term) and interference term between internal Bremsstrahlung and SD^- term. See the “Note on $\pi^\pm \rightarrow \ell^\pm \nu_\ell \gamma$ and $K^\pm \rightarrow \ell^\pm \nu_\ell \gamma$ Form Factors” in the π^\pm section of the Particle Data Listings above.

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
<2.6	90	44 AKIBA	85 SPEC

⁴⁴ Assumes μ -e universality and uses constraints from $K \rightarrow e \nu \gamma$.

$\Gamma(e^+ \nu_e \gamma)/\Gamma(\mu^+ \nu_\mu)$ Γ_{16}/Γ_2


<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
$1.483 \pm 0.066 \pm 0.013$	1.4K	⁴⁵ AMBROSINO	09E	KLOE	\pm E_γ in 10–250 MeV, $p_e > 200$ MeV/c

⁴⁵ AMBROSINO 09E measured the differential width $dR_\gamma/dE_\gamma = (1/\Gamma(K \rightarrow \mu\nu)) (d\Gamma(K \rightarrow e\nu\gamma)/dE_\gamma)$. Result obtained by integrating the differential width over E_γ from 10 to 250 MeV.

$\Gamma(\pi^0 e^+ \nu_e \gamma)/\Gamma(\pi^0 e^+ \nu_e)$ Γ_{17}/Γ_3

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
0.505 ± 0.032 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.			
$0.47 \pm 0.02 \pm 0.03$	4476	⁴⁶ AKIMENKO	07	ISTR	$-$ $E_\gamma > 10$ MeV, $0.6 < \cos(\theta_{e\gamma}) < 0.9$
0.46 ± 0.08	82	⁴⁷ BARMIN	91	XEBC	$E_\gamma > 10$ MeV, $0.6 < \cos(\theta_{e\gamma}) < 0.9$
0.56 ± 0.04	192	⁴⁸ BOLOTOV	86B	CALO	$-$ $E_\gamma > 10$ MeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$1.81 \pm 0.03 \pm 0.07$	4476	⁴⁶ AKIMENKO	07	ISTR	$-$ $E_\gamma > 10$ MeV, $\theta_{e\gamma} > 10^\circ$
$0.63 \pm 0.02 \pm 0.03$	4476	⁴⁶ AKIMENKO	07	ISTR	$-$ $E_\gamma > 30$ MeV, $\theta_{e\gamma} > 20^\circ$
1.51 ± 0.25	82	⁴⁷ BARMIN	91	XEBC	$E_\gamma > 10$ MeV, $\cos(\theta_{e\gamma}) < 0.98$
0.48 ± 0.20	16	⁴⁹ LJUNG	73	HLBC	$+$ $E_\gamma > 30$ MeV
$0.22^{+0.15}_{-0.10}$		⁴⁹ LJUNG	73	HLBC	$+$ $E_\gamma > 30$ MeV
0.76 ± 0.28	13	⁵⁰ ROMANO	71	HLBC	$E_\gamma > 10$ MeV
0.53 ± 0.22		⁵⁰ ROMANO	71	HLBC	$+$ $E_\gamma > 30$ MeV
1.2 ± 0.8		BELLOTTI	67	HLBC	$E_\gamma > 30$ MeV

- 46 AKIMENKO 07 provides values for three kinematic regions. For averaging, we use value with $E_\gamma > 10$ MeV and $0.6 < \cos(\theta_{e\gamma}) < 0.9$.
- 47 BARMIN 91 quotes branching ratio $\Gamma(K \rightarrow e\pi^0\nu\gamma)/\Gamma_{\text{all}}$. The measured normalization is $[\Gamma(K \rightarrow e\pi^0\nu) + \Gamma(K \rightarrow \pi^+\pi^+\pi^-)]$. For comparison with other experiments we used $\Gamma(K \rightarrow e\pi^0\nu)/\Gamma_{\text{all}} = 0.0482$ to calculate the values quoted here.
- 48 $\cos(\theta_{e\gamma})$ between 0.6 and 0.9.
- 49 First LJUNG 73 value is for $\cos(\theta_{e\gamma}) < 0.9$, second value is for $\cos(\theta_{e\gamma})$ between 0.6 and 0.9 for comparison with ROMANO 71.
- 50 Both ROMANO 71 values are for $\cos(\theta_{e\gamma})$ between 0.6 and 0.9. Second value is for comparison with second LJUNG 73 value. We use lowest E_γ cut for Summary Table value. See ROMANO 71 for E_γ dependence.



$\Gamma(\pi^0 e^+ \nu_e \gamma(\text{SD}))/\Gamma_{\text{total}}$
Structure-dependent part.

Γ_{18}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	CHG
<5.3	90	BOLOTOV 86B	CALO	-

$\Gamma(\pi^0 \mu^+ \nu_\mu \gamma)/\Gamma_{\text{total}}$

Γ_{19}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1.46±0.22±0.32		153	51 TCHIKILEV 07	ISTR	-	30 < E_γ < 60 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.4 ±0.5 ±0.6	125	SHIMIZU	06	K470	+	$E_\gamma > 30$ MeV; $\Theta_{\mu\gamma} > 20^\circ$	
<6.1	90	0	LJUNG	73	HLBC	+	$E(\gamma) > 30$ MeV

⁵¹ Obtained from measuring $B(K_{\mu 3\gamma}) / B(K_{\mu 3})$ and using PDG 02 value $B(K_{\mu 3}) = 3.27\%$.
 $B(K_{\mu 3\gamma}) = (8.82 \pm 0.94 \pm 0.86) \times 10^{-5}$ is obtained for $5 \text{ MeV} < E_\gamma < 30 \text{ MeV}$.

$\Gamma(\pi^0 \pi^0 e^+ \nu_e \gamma) / \Gamma_{\text{total}}$			Γ_{20} / Γ			
VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<5	90	0	BARMIN	92	XEBC	+ $E_\gamma > 10 \text{ MeV}$

————— Hadronic modes with photons —————

$\Gamma(\pi^+ \pi^0 \gamma) / \Gamma_{\text{total}}$			Γ_{21} / Γ			
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT

2.75 ± 0.15 OUR AVERAGE

2.71 ± 0.45		140	BOLOTOV	87	WIRE	− T_{π^-} 55–90 MeV
2.87 ± 0.32		2461	SMITH	76	WIRE	± T_{π^\pm} 55–90 MeV
2.71 ± 0.19		2100	ABRAMS	72	ASPK	± T_{π^+} 55–90 MeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
1.5 $^{+1.1}_{-0.6}$			⁵² LJUNG	73	HLBC	+ T_{π^+} 55–80 MeV
2.6 $^{+1.5}_{-1.1}$			⁵² LJUNG	73	HLBC	+ T_{π^+} 55–90 MeV
6.8 $^{+3.7}_{-2.1}$		17	⁵² LJUNG	73	HLBC	+ T_{π^+} 55–102 MeV
2.4 ± 0.8		24	EDWARDS	72	OSPK	T_{π^+} 58–90 MeV
<1.0		0	⁵³ MALTSEV	70	HLBC	+ T_{π^+} <55 MeV
<1.9	90	0	EMMERSON	69	OSPK	T_{π^+} 55–80 MeV
2.2 ± 0.7		18	CLINE	64	FBC	+ T_{π^+} 55–80 MeV

⁵² The LJUNG 73 values are not independent.

⁵³ MALTSEV 70 selects low π^+ energy to enhance direct emission contribution.

$\Gamma(\pi^+ \pi^0 \gamma(\text{DE})) / \Gamma_{\text{total}}$			Γ_{22} / Γ			
---	--	--	------------------------	--	--	--

Direct emission (DE) part of $\Gamma(\pi^+ \pi^0 \gamma) / \Gamma_{\text{total}}$, assuming that interference (INT) component is zero.

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
--------------------------	------	-------------	------	-----	---------

4.3 ± 0.7 OUR AVERAGE

3.8 ± 0.8 ± 0.7	10k	ALIEV	06	K470	+ T_{π^+} 55–90 MeV
3.7 ± 3.9 ± 1.0	930	UVAROV	06	ISTR	− T_{π^-} 55–90 MeV
4.7 ± 0.8 ± 0.3	20k	⁵⁴ ADLER	00C	B787	+ T_{π^+} 55–90 MeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
3.2 ± 1.3 ± 1.0	4k	ALIEV	03	K470	+ T_{π^+} 55–90 MeV
6.1 ± 2.5 ± 1.9	4k	ALIEV	03	K470	+ T_{π^+} full range
20.5 ± 4.6 $^{+3.9}_{-2.3}$		BOLOTOV	87	WIRE	− T_{π^-} 55–90 MeV
15.6 ± 3.5 ± 5.0		ABRAMS	72	ASPK	± T_{π^\pm} 55–90 MeV

⁵⁴ ADLER 00C measures the INT component to be $(-0.4 \pm 1.6)\%$ of the inner bremsstrahlung (IB) component.

$\Gamma(\pi^+ \pi^0 \pi^0 \gamma) / \Gamma(\pi^+ \pi^0 \pi^0)$			$\Gamma_{23} / \Gamma_{10}$			
VALUE (units 10^{-4})	DOCUMENT ID	TECN	CHG	COMMENT		

4.3 $^{+3.2}_{-1.7}$	BOLOTOV	85	SPEC	− $E(\gamma) > 10 \text{ MeV}$
--	---------	----	------	--------------------------------

$\Gamma(\pi^+\pi^+\pi^-\gamma)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1.04±0.31 OUR AVERAGE					
1.10±0.48	7	BARMIN	89	XEBC	$E(\gamma) > 5$ MeV
1.0 ±0.4		STAMER	65	EMUL +	$E(\gamma) > 11$ MeV

$\Gamma(\pi^+\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-7})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
11±3 ±1		31	⁵⁵ KITCHING	97	B787	
●●● We do not use the following data for averages, fits, limits, etc. ●●●						
< 0.083	90		⁵⁶ ARTAMONOV	05	B949 +	$P_\pi > 213$ MeV/c
< 10	90	0	ATIYA	90B	B787	T_π 117–127 MeV
< 84	90	0	ASANO	82	CNTR +	T_π 117–127 MeV
−420 ± 520		0	ABRAMS	77	SPEC +	$T_\pi < 92$ MeV
< 350	90	0	LJUNG	73	HLBC +	6–102, 114–127 MeV
< 500	90	0	KLEMS	71	OSPK +	$T_\pi < 117$ MeV
−100 ± 600			CHEN	68	OSPK +	T_π 60–90 MeV

⁵⁵KITCHING 97 is extrapolated from their model-independent branching fraction ($6.0 \pm 1.5 \pm 0.7$) $\times 10^{-7}$ for $100 \text{ MeV}/c < P_{\pi^+} < 180 \text{ MeV}/c$ using Chiral Perturbation Theory.

⁵⁶ARTAMONOV 05 limit assumes ChPT with $\hat{c} = 1.8$ with unitarity corrections. With $\hat{c} = 1.6$ and no unitarity corrections they obtain $< 2.3 \times 10^{-8}$ at 90% CL. This partial branching ratio is predicted to be 6.10×10^{-9} and 0.49×10^{-9} for the cases with and without unitarity correction.

$\Gamma(\pi^+3\gamma)/\Gamma_{\text{total}}$ Γ_{26}/Γ

Values given here assume a phase space pion energy spectrum.

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<1.0	90	ASANO	82	CNTR +	$T(\pi)$ 117–127 MeV

●●● We do not use the following data for averages, fits, limits, etc. ●●●

<3.0	90	KLEMS	71	OSPK +	$T(\pi) > 117$ MeV
------	----	-------	----	--------	--------------------

$\Gamma(\pi^+e^+e^-\gamma)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
1.19±0.12±0.04	113	⁵⁷ BATLEY	08	NA48 $m_{ee\gamma} > 260$ MeV

⁵⁷BATLEY 08 also reports the Chiral Perturbation Theory parameter $\hat{c} = 0.9 \pm 0.45$ obtained using the shape of the $e^+e^-\gamma$ invariant mass spectrum. By extrapolating the theoretical amplitude to $m_{ee\gamma} < 260$ MeV, it obtains the inclusive $B(K^+ \rightarrow \pi^+e^+e^-\gamma) = (1.29 \pm 0.13 \pm 0.03) \times 10^{-8}$, where the first error is the combined statistical and systematic errors and the second error is from the uncertainty in \hat{c} .

———— Leptonic modes with $l\bar{l}$ pairs ————

$\Gamma(e^+\nu_e\nu\bar{\nu})/\Gamma(e^+\nu_e)$ Γ_{28}/Γ_1

VALUE	CL%	EVTS	DOCUMENT ID	TECN	CHG
<3.8	90	0	HEINTZE	79	SPEC +

$\Gamma(\mu^+ \nu_\mu \nu \bar{\nu})/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	CHG
<6.0	90	0	⁵⁸ PANG	73	CNTR +

⁵⁸ PANG 73 assumes μ spectrum from ν - ν interaction of BARDIN 70.

$\Gamma(e^+ \nu_e e^+ e^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2.48 ± 0.14 ± 0.14	410	POBLAGUEV 02	B865	+	$m_{ee} > 150$ MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

20 ± 20	4	DIAMANT-...	76	SPEC	+	$m_{e^+e^-} > 140$ MeV
---------	---	-------------	----	------	---	------------------------

$\Gamma(\mu^+ \nu_\mu e^+ e^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
7.06 ± 0.16 ± 0.26	2.7k	POBLAGUEV 02	B865	+	$m_{ee} > 145$ MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

100 ± 30	14	DIAMANT-...	76	SPEC	+	$m_{e^+e^-} > 140$ MeV
----------	----	-------------	----	------	---	------------------------

$\Gamma(e^+ \nu_e \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-8})	CL%	DOCUMENT ID	TECN
1.72 ± 0.45		MA 06	B865

• • • We do not use the following data for averages, fits, limits, etc. • • •

<50	90	ADLER 98	B787
-----	----	----------	------

$\Gamma(\mu^+ \nu_\mu \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	CHG
<4.1	90	ATIYA 89	B787	+

———— Lepton Family number (*LF*), Lepton number (*L*), $\Delta S = \Delta Q$ (*SQ*)
 ————— violating modes, or $\Delta S = 1$ weak neutral current (*S1*) modes —————

$\Gamma(\pi^+ \pi^+ e^- \bar{\nu}_e)/\Gamma_{\text{total}}$ Γ_{34}/Γ

Test of $\Delta S = \Delta Q$ rule.

VALUE (units 10^{-7})	CL%	EVTS	DOCUMENT ID	TECN	CHG
--------------------------	-----	------	-------------	------	-----

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9.0	95	0	SCHWEINB...	71	HLBC +
< 6.9	95	0	ELY	69	HLBC +
< 20.	95		BIRGE	65	FBC +

$\Gamma(\pi^+ \pi^+ e^- \bar{\nu}_e)/\Gamma(\pi^+ \pi^- e^+ \nu_e)$ Γ_{34}/Γ_6

Test of $\Delta S = \Delta Q$ rule.

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN
< 3	90	3	⁵⁹ BLOCH	76 SPEC

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 130.	95	0	BOURQUIN 71	ASPK
--------	----	---	-------------	------

⁵⁹ BLOCH 76 quotes 3.6×10^{-4} at CL = 95%, we convert.

$\Gamma(\pi^+ \pi^+ \mu^- \bar{\nu}_\mu) / \Gamma_{\text{total}}$ Γ_{35} / Γ
 Test of $\Delta S = \Delta Q$ rule.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	CHG
<3.0	95	0	BIRGE	65	FBC +

$\Gamma(\pi^+ e^+ e^-) / \Gamma_{\text{total}}$ Γ_{36} / Γ
 Test for $\Delta S = 1$ weak neutral current. Allowed by combined first-order weak and electromagnetic interactions.

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	CHG
3.00 ± 0.09 OUR AVERAGE				
3.11 ± 0.04 ± 0.12	7253	⁶⁰ BATLEY	09 NA48	±
2.94 ± 0.05 ± 0.14	10300	⁶¹ APPEL	99 SPEC	+
2.75 ± 0.23 ± 0.13	500	⁶² ALLIEGRO	92 SPEC	+
2.7 ± 0.5	41	⁶³ BLOCH	75 SPEC	+

⁶⁰ Value extrapolated from a measurement in the region $z = (m_{ee}/m_K)^2 > 0.08$. BATLEY 09 also evaluated the shape of the form factor using four different theoretical models.

⁶¹ APPEL 99 establishes vector nature of this decay and determines form factor $f(Z) = f_0(1 + \delta Z)$, $Z = M_{ee}^2/m_K^2$, $\delta = 2.14 \pm 0.13 \pm 0.15$.

⁶² ALLIEGRO 92 assumes a vector interaction with a form factor given by $\lambda = 0.105 \pm 0.035 \pm 0.015$ and a correlation coefficient of -0.82 .

⁶³ BLOCH 75 assumes a vector interaction.

$\Gamma(\pi^+ \mu^+ \mu^-) / \Gamma_{\text{total}}$ Γ_{37} / Γ
 Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-8})	CL%	EVTS	DOCUMENT ID	TECN	CHG
8.1 ± 1.4 OUR AVERAGE Error includes scale factor of 2.7. See the ideogram below.					

9.8 ± 1.0 ± 0.5	110	⁶⁴ PARK	02 HYCP	±
9.22 ± 0.60 ± 0.49	402	⁶⁵ MA	00 B865	+
5.0 ± 0.4 ± 0.9	207	⁶⁶ ADLER	97C B787	+

• • • We do not use the following data for averages, fits, limits, etc. • • •


9.7 ± 1.2 ± 0.4	65	PARK	02 HYCP	+
10.0 ± 1.9 ± 0.7	35	PARK	02 HYCP	-
<23	90	ATIYA	89 B787	+

⁶⁴ PARK 02 "±" result comes from combining $K^+ \rightarrow \pi^+ \mu^+ \mu^-$ and $K^- \rightarrow \pi^- \mu^+ \mu^-$, assuming CP is conserved.

⁶⁵ MA 00 establishes vector nature of this decay and determines form factor $f(Z) = f_0(1 + \delta Z)$, $Z = M_{\mu\mu}^2/m_K^2$, $\delta = 2.45^{+1.30}_{-0.95}$.

⁶⁶ ADLER 97C gives systematic error 0.7×10^{-8} and theoretical uncertainty 0.6×10^{-8} , which we combine in quadrature to obtain our second error.

WEIGHTED AVERAGE
 8.1 ± 1.4 (Error scaled by 2.7)



$\Gamma(\pi^+ \mu^+ \mu^-) / \Gamma_{\text{total}}$

Γ_{37} / Γ

$\Gamma(\pi^+ \nu \bar{\nu}) / \Gamma_{\text{total}}$

Γ_{38} / Γ

Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interactions. Branching ratio values are extrapolated from the momentum or energy regions shown in the comments assuming Standard Model phase space except for those labeled "Scalar" or "Tensor" to indicate the assumed non-Standard-Model interaction.

VALUE (units 10^{-9})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$0.173^{+0.115}_{-0.105}$		7	67 ARTAMONOV 08	B949	+	$140 < P_\pi < 199$ MeV, $211 < P_\pi < 229$ MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.789^{+0.926}_{-0.510}$		3	68 ARTAMONOV 08	B949	+	$140 < P_\pi < 199$ MeV
< 2.2	90	1	69 ADLER	04 B787	+	$211 < P_\pi < 229$ MeV
< 2.7	90		ADLER	04 B787	+	Scalar
< 1.8	90		ADLER	04 B787	+	Tensor
$0.147^{+0.130}_{-0.089}$		3	70 ANISIMOVSK...04	B949	+	$211 < P_\pi < 229$ MeV
$0.157^{+0.175}_{-0.082}$		2	ADLER	02 B787	+	$P_\pi > 211$ MeV/c
< 4.2	90	1	ADLER	02C B787	+	$140 < P_\pi < 195$ MeV
< 4.7	90		71 ADLER	02C B787	+	Scalar
< 2.5	90		71 ADLER	02C B787	+	Tensor
$0.15^{+0.34}_{-0.12}$		1	ADLER	00 B787		In ADLER 02
$0.42^{+0.97}_{-0.35}$		1	ADLER	97 B787		

< 2.4	90	ADLER	96	B787	
< 7.5	90	ATIYA	93	B787	+ T(π) 115–127 MeV
< 5.2	90	⁷² ATIYA	93	B787	+
< 17	90	0	ATIYA	93B B787	+ T(π) 60–100 MeV
< 34	90	ATIYA	90	B787	+
<140	90	ASANO	81B	CNTR	+ T(π) 116–127 MeV

⁶⁷ Value obtained combining ANISIMOVSKY 04, ADLER 04, and the present ARTAMONOV 08 results.

⁶⁸ Observed 3 events with an estimated background of $0.93 \pm 0.17^{+0.32}_{-0.24}$. Signal-to-background ratio for each of these 3 events is 0.20, 0.42, and 0.47.

⁶⁹ Value obtained combining the previous result ADLER 02C with 1 event and the present result with 0 events to obtain an expected background 1.22 ± 0.24 events and 1 event observed.

⁷⁰ Value obtained combining the previous E787 result ADLER 02 with 2 events and the present E949 with 1 event. The additional event has a signal-to-background ratio 0.9. Superseded by ARTAMONOV 08.

⁷¹ Superseded by ADLER 04.

⁷² Combining ATIYA 93 and ATIYA 93B results. Superseded by ADLER 96.

$\Gamma(\pi^+ \pi^0 \nu \bar{\nu})/\Gamma_{\text{total}}$ Γ_{39}/Γ

Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN
<4.3	90	⁷³ ADLER	01 SPEC

⁷³ Search region defined by $90 \text{ MeV}/c < P_{\pi^+} < 188 \text{ MeV}/c$ and $135 \text{ MeV} < E_{\pi^0} < 180 \text{ MeV}$.

$\Gamma(\mu^- \nu e^+ e^+)/\Gamma(\pi^+ \pi^- e^+ \nu_e)$ Γ_{40}/Γ_6

Test of lepton family number conservation.

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	CHG
<0.5	90	0	⁷⁴ DIAMANT-...	76	SPEC +

⁷⁴ DIAMANT-BERGER 76 quotes this result times our 1975 $\pi^+ \pi^- e \nu$ BR ratio.

$\Gamma(\mu^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{41}/Γ

Forbidden by lepton family number conservation.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.004	90	0	⁷⁵ LYONS	81	HLBC 200 GeV K^+ narrow band ν beam

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.012	90	⁷⁵ COOPER	82	HLBC	Wideband ν beam
--------	----	----------------------	----	------	---------------------

⁷⁵ COOPER 82 and LYONS 81 limits on ν_e observation are here interpreted as limits on lepton family number violation in the absence of mixing.

$\Gamma(\pi^+ \mu^+ e^-)/\Gamma_{\text{total}}$ Γ_{42}/Γ

Test of lepton family number conservation.

VALUE (units 10^{-10})	CL%	DOCUMENT ID	TECN	CHG
<0.13	90	⁷⁶ SHER	05	RVUE +

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.21	90	SHER	05	B865	+
<0.39	90	APPEL	00	B865	+
<2.1	90	LEE	90	SPEC	+

⁷⁶ This result combines SHER 05 1998 data, APPEL 00 1996 data, and data from BERGMAN 97 and PISLAK 97 theses, all from BNL-E865, with LEE 90 BNL-E777 data.

$\Gamma(\pi^+ \mu^- e^+)/\Gamma_{\text{total}}$ Γ_{43}/Γ

Test of lepton family number conservation.

<u>VALUE (units 10^{-10})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
< 5.2	90	0	APPEL	00B B865	+

• • • We do not use the following data for averages, fits, limits, etc. • • •

<70	90	0	⁷⁷ DIAMANT-...	76	SPEC	+
-----	----	---	---------------------------	----	------	---

⁷⁷ Measurement actually applies to the sum of the $\pi^+ \mu^- e^+$ and $\pi^- \mu^+ e^+$ modes.

$\Gamma(\pi^- \mu^+ e^+)/\Gamma_{\text{total}}$ Γ_{44}/Γ

Test of total lepton number conservation.

<u>VALUE (units 10^{-10})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
< 5.0	90	0	APPEL	00B B865	+

• • • We do not use the following data for averages, fits, limits, etc. • • •

<70	90	0	⁷⁸ DIAMANT-...	76	SPEC	+
-----	----	---	---------------------------	----	------	---

⁷⁸ Measurement actually applies to the sum of the $\pi^+ \mu^- e^+$ and $\pi^- \mu^+ e^+$ modes.

$\Gamma(\pi^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{45}/Γ

Test of total lepton number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
< 6.4×10^{-10}	90	0	APPEL	00B B865	+

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9.2×10^{-9}	90	0	DIAMANT-...	76	SPEC	+
< 1.5×10^{-5}			CHANG	68	HBC	-

$\Gamma(\pi^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{46}/Γ

Forbidden by total lepton number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
< 3.0×10^{-9}	90	0	APPEL	00B B865	+

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.5×10^{-4}	90		⁷⁹ LITTENBERG	92	HBC
------------------------	----	--	--------------------------	----	-----

⁷⁹ LITTENBERG 92 is from retroactive data analysis of CHANG 68 bubble chamber data.

$\Gamma(\mu^+ \bar{\nu}_e)/\Gamma_{\text{total}}$ Γ_{47}/Γ

Forbidden by total lepton number conservation.

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.3	90	⁸⁰ COOPER	82	HLBC Wideband ν beam

⁸⁰ COOPER 82 limit on $\bar{\nu}_e$ observation is here interpreted as a limit on lepton number violation in the absence of mixing.

$\Gamma(\pi^0 e^+ \bar{\nu}_e)/\Gamma_{\text{total}}$ Γ_{48}/Γ

Forbidden by total lepton number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.003	90	⁸¹ COOPER	82	HLBC Wideband ν beam

⁸¹ COOPER 82 limit on $\bar{\nu}_e$ observation is here interpreted as a limit on lepton number violation in the absence of mixing.

$\Gamma(\pi^+\gamma)/\Gamma_{\text{total}}$

Γ_{49}/Γ

Violates angular momentum conservation and gauge invariance. Current interest in this decay is as a search for non-commutative space-time effects as discussed in ARTAMONOV 05 and for exotic physics such as a vacuum expectation value of a new vector field, non-local Superstring effects, or departures from Lorentz invariance, as discussed in ADLER 02B.

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	CHG
< 2.3	90	ARTAMONOV 05	B949	+
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 360	90	ADLER	02B B787	+
<1400	90	ASANO	82 CNTR	+
<4000	90	⁸² KLEMS	71 OSPK	+

⁸² Test of model of Selleri, Nuovo Cimento **60A** 291 (1969).

K^+ LONGITUDINAL POLARIZATION OF EMITTED μ^+

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
< -0.990	90	⁸³ AOKI	94 SPEC	+	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< -0.990	90	IMAZATO	92 SPEC	+	Repl. by AOKI 94
-0.970 ± 0.047		⁸⁴ YAMANAKA	86 SPEC	+	
-1.0 ± 0.1		⁸⁴ CUTTS	69 SPRK	+	
-0.96 ± 0.12		⁸⁴ COOMBES	57 CNTR	+	

⁸³ AOKI 94 measures $\xi P_\mu = -0.9996 \pm 0.0030 \pm 0.0048$. The above limit is obtained by summing the statistical and systematic errors in quadrature, normalizing to the physically significant region ($|\xi P_\mu| < 1$) and assuming that $\xi=1$, its maximum value.

⁸⁴ Assumes $\xi=1$.

A REVIEW GOES HERE – Check our WWW List of Reviews

ENERGY DEPENDENCE OF K^\pm DALITZ PLOT

$$|\text{matrix element}|^2 = 1 + gu + hu^2 + kv^2$$

where $u = (s_3 - s_0) / m_\pi^2$ and $v = (s_2 - s_1) / m_\pi^2$

LINEAR COEFFICIENT g FOR $K^\pm \rightarrow \pi^\pm \pi^+ \pi^-$

Some experiments use Dalitz variables x and y . In the comments we give $a_y =$ coefficient of y term. See note above on “Dalitz Plot Parameters for $K \rightarrow 3\pi$ Decays.” For discussion of the conversion of a_y to g , see the earlier version of the same note in the Review published in Physics Letters **111B** 70 (1982).

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-0.21134 ± 0.00017	471M	⁸⁵ BATLEY	07B NA48	±	

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.2221 ±0.0065	225k	DEVAUX	77	SPEC	+	$a_y = .2814 \pm .0082$
-0.199 ±0.008	81k	⁸⁶ LUCAS	73	HBC	-	$a_y = 0.252 \pm 0.011$
-0.2157 ±0.0028	750k	FORD	72	ASPK	+	$a_y = .2734 \pm .0035$
-0.2186 ±0.0028	750k	FORD	72	ASPK	-	$a_y = .2770 \pm .0035$
-0.200 ±0.009	39819	⁸⁷ HOFFMASTER	72	HLBC	+	
-0.196 ±0.012	17898	⁸⁸ GRAUMAN	70	HLBC	+	$a_y = 0.228 \pm 0.030$
-0.193 ±0.010	50919	MAST	69	HBC	-	$a_y = 0.244 \pm 0.013$
-0.218 ±0.016	9994	⁸⁹ BUTLER	68	HBC	+	$a_y = 0.277 \pm 0.020$
-0.190 ±0.023	5778	^{89,90} MOSCOSO	68	HBC	-	$a_y = 0.242 \pm 0.029$
-0.22 ±0.024	5428	^{89,90} ZINCHENKO	67	HBC	+	$a_y = 0.28 \pm 0.03$
-0.220 ±0.035	1347	⁹¹ FERRO-LUZZI	61	HBC	-	$a_y = 0.28 \pm 0.045$

⁸⁵ Final state strong interaction and radiative corrections not included in the fit.

⁸⁶ Quadratic dependence is required by K_L^0 experiments.

⁸⁷ HOFFMASTER 72 includes GRAUMAN 70 data.

⁸⁸ Emulsion data added — all events included by HOFFMASTER 72.

⁸⁹ Experiments with large errors not included in average.

⁹⁰ Also includes DBC events.

⁹¹ No radiative corrections included.

QUADRATIC COEFFICIENT h FOR $K^\pm \rightarrow \pi^\pm \pi^+ \pi^-$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
1.848 ± 0.040	471M	⁹² BATLEY	07B NA48	±

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.06 ±1.43	225k	DEVAUX	77	SPEC	+
1.87 ±0.62	750k	FORD	72	ASPK	+
1.25 ±0.62	750k	FORD	72	ASPK	-
-0.9 ±1.4	39819	HOFFMASTER	72	HLBC	+
-0.1 ±1.2	50919	MAST	69	HBC	-

⁹² Final state strong interaction and radiative corrections not included in the fit.

QUADRATIC COEFFICIENT k FOR $K^\pm \rightarrow \pi^\pm \pi^+ \pi^-$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
- 4.63 ± 0.14	471M	⁹³ BATLEY	07B NA48	±

• • • We do not use the following data for averages, fits, limits, etc. • • •

-20.5 ± 3.9	225k	DEVAUX	77	SPEC	+
- 7.5 ± 1.9	750k	FORD	72	ASPK	+
- 8.3 ± 1.9	750k	FORD	72	ASPK	-
-10.5 ± 4.5	39819	HOFFMASTER	72	HLBC	+
-14 ±12	50919	MAST	69	HBC	-

⁹³ Final state strong interaction and radiative corrections not included in the fit.

$(g_+ - g_-) / (g_+ + g_-)$ FOR $K^\pm \rightarrow \pi^\pm \pi^+ \pi^-$

This is a *CP* violating asymmetry between linear coefficients g_+ for $K^+ \rightarrow \pi^+ \pi^+ \pi^-$ decay and g_- for $K^- \rightarrow \pi^- \pi^+ \pi^-$ decay.

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$-1.5 \pm 1.5 \pm 1.6$	3.1G	⁹⁴ BATLEY	07E NA48

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.7 \pm 2.1 \pm 2.0$	1.7G	⁹⁵ BATLEY	06 NA48
-70.0 ± 53	3.2M	FORD	70 ASPK

⁹⁴ BATLEY 07E includes data from BATLEY 06. Uses quadratic parametrization and value $g_+ + g_- = 2g$ from BATLEY 07B. This measurement neglects any possible charge asymmetries in higher order slope parameters h or k .

⁹⁵ This measurement neglects any possible charge asymmetries in higher order slope parameters h or k .

LINEAR COEFFICIENT g FOR $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$

Unless otherwise stated, all experiments include terms quadratic in $(s_3 - s_0) / m_{\pi^+}^2$. See note above on "Dalitz Plot Parameters for $K \rightarrow 3\pi$ Decays."

See BATUSOV 98 for a discussion of the discrepancy between their result and others, especially BOLOTOV 86. At this time we have no way to resolve the discrepancy so we depend on the large scale factor as a warning.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
0.626 ± 0.007 OUR AVERAGE					

$0.6259 \pm 0.0043 \pm 0.0093$ 493k AKOPDZHAN..05B TNF \pm

$0.627 \pm 0.004 \pm 0.010$ 252k ^{96,97} AJINENKO 03B ISTR $-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.736 \pm 0.014 \pm 0.012$ 33k BATUSOV 98 SPEC $+$

0.582 ± 0.021 43k BOLOTOV 86 CALO $-$

0.670 ± 0.054 3263 BRAUN 76B HLBC $+$

0.630 ± 0.038 5635 SHEAFF 75 HLBC $+$

0.510 ± 0.060 27k SMITH 75 WIRE $+$

0.67 ± 0.06 1365 AUBERT 72 HLBC $+$

0.544 ± 0.048 4048 DAVISON 69 HLBC $+$ Also emulsion

⁹⁶ Measured using in-flight decays of the 25 GeV negative secondary beam.

⁹⁷ They form new world averages $g_- = (0.617 \pm 0.018)$ and $g_+ = (0.684 \pm 0.033)$ which give $\Delta g_{\pi^0} = 0.051 \pm 0.028$.

QUADRATIC COEFFICIENT h FOR $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
0.052 ± 0.008 OUR AVERAGE					

$0.0551 \pm 0.0044 \pm 0.0086$ 493k AKOPDZHAN..05B TNF \pm

$0.046 \pm 0.004 \pm 0.012$ 252k ⁹⁸ AJINENKO 03B ISTR $-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.128 \pm 0.015 \pm 0.024$ 33k BATUSOV 98 SPEC $+$

0.037 ± 0.024 43k BOLOTOV 86 CALO $-$

0.152 ± 0.082 3263 BRAUN 76B HLBC $+$

0.041 ± 0.030 5635 SHEAFF 75 HLBC $+$

0.009 ± 0.040 27k SMITH 75 WIRE $+$

-0.01 ± 0.08 1365 AUBERT 72 HLBC $+$

0.026 ± 0.050 4048 DAVISON 69 HLBC $+$ Also emulsion

⁹⁸ Measured using in-flight decays of the 25 GeV negative secondary beam.

QUADRATIC COEFFICIENT k FOR $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$

VALUE	EVTS	DOCUMENT ID	TECN	CHG
0.0054 ± 0.0035 OUR AVERAGE				
Error includes scale factor of 2.5.				
0.0082 ± 0.0011 ± 0.0014	493k	AKOPDZHAN..05B	TNF	±
0.001 ± 0.001 ± 0.002	252k	⁹⁹ AJINENKO	03B ISTR	–

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0197 ± 0.0045 ± 0.0029	33k	BATUSOV	98 SPEC	+
--------------------------	-----	---------	---------	---

⁹⁹ Measured using in-flight decays of the 25 GeV negative secondary beam.

$(g_+ - g_-) / (g_+ + g_-)$ FOR $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$

A nonzero value for this quantity indicates CP violation.

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN
1.8 ± 1.8 OUR AVERAGE			
1.8 ± 1.7 ± 0.6	91.3M	¹⁰⁰ BATLEY	07E NA48
2 ± 18 ± 5	619k	¹⁰¹ AKOPDZHAN..05	TNF

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 ± 2.2 ± 1.3	47M	¹⁰² BATLEY	06A NA48
-----------------	-----	-----------------------	----------

¹⁰⁰ BATLEY 07E includes data from BATLEY 06A. Uses quadratic parametrization and PDG 06 value $g = 0.626 \pm 0.007$ to obtain $g_+ - g_- = (2.2 \pm 2.1 \pm 0.7) \times 10^{-4}$. Neglects any possible charge asymmetries in higher order slope parameters h or k .

¹⁰¹ Asymmetry obtained assuming that $g_+ + g_- = 2 \times 0.652$ (PDG 02) and that asymmetries in h and k are zero.

¹⁰² Linear and quadratic slopes from PDG 04 are used. Any possible charge asymmetries in higher order slope parameters h or k are neglected.

ALTERNATIVE PARAMETRIZATIONS OF $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$ DALITZ PLOT

The following functional form for the matrix element suggested by $\pi\pi$ rescattering in $K^+ \rightarrow \pi^+ \pi^+ \pi^- \rightarrow \pi^+ \pi^0 \pi^0$ is used for this fit (CABIBBO 04A, CABIBBO 05): Matrix element = $M_0 + M_1$ where $M_0 = 1 + (1/2)g_0 u + (1/2)h' u^2 + (1/2)k_0 v^2$ with $u = (s_3 - s_0)/(m_{\pi^+})^2$, $v = (s_2 - s_1)/(m_{\pi^+})^2$ and where M_1 takes into account the non-analytic piece due to $\pi\pi$ rescattering amplitudes a_0 and a_2 ; The parameters g_0 and h' are related to the parameters g and h of the matrix element squared given in the previous section by the approximations $g_0 \sim g^{PDG}$ and $h' \sim h^{PDG} - (g/2)^2$ and $k_0 \sim k^{PDG}$.

In addition, we also consider the effective field theory framework of COLANGELO 06A and BISSEGGER 09 to extract g_{BB} and h_{BB}^l .

LINEAR COEFFICIENT g_0 FOR $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$

VALUE	EVTS	DOCUMENT ID	TECN	CHG
0.6525 ± 0.0009 ± 0.0033	60M	¹⁰³ BATLEY	09A NA48	±
0.645 ± 0.004 ± 0.009	23M	¹⁰⁴ BATLEY	06B NA48	±

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹⁰³ This fit is obtained with the CABIBBO 05 matrix element in the $2\pi^0$ invariant mass squared range $0.074094 < m_{2\pi^0}^2 < 0.104244 \text{ GeV}^2$. Electromagnetic corrections and CHPT constraints for $\pi\pi$ phase shifts (a_0 and a_2) have been used. Also measured $(a_0 - a_2) m_{\pi^+} = 0.2646 \pm 0.0021 \pm 0.0023$, where k_0 was kept fixed in the fit at -0.0099 .

¹⁰⁴ Superseded by BATLEY 09A. This fit is obtained with the CABIBBO 05 matrix element in the $2\pi^0$ invariant mass squared range $0.074 \text{ GeV}^2 < m_{2\pi^0}^2 < 0.097 \text{ GeV}^2$, assuming $k = 0$ (no term proportional to $(s_2 - s_1)^2$) and excluding the kinematic region around the cusp ($m_{2\pi^0}^2 = (2m_{\pi^+})^2 \pm 0.000525 \text{ GeV}^2$). Also π - π phase shifts a_0 and a_2 are measured: $(a_0 - a_2) m_{\pi^+} = 0.268 \pm 0.010 \pm 0.004 \pm 0.013(\text{external})$ and $a_2 m_{\pi^+} = -0.041 \pm 0.022 \pm 0.014$.

QUADRATIC COEFFICIENT h' FOR $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$

VALUE	EVTS	DOCUMENT ID	TECN	CHG
$-0.0433 \pm 0.0008 \pm 0.0026$	60M	¹⁰⁵ BATLEY	09A NA48	\pm

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.047 \pm 0.012 \pm 0.011$	23M	¹⁰⁶ BATLEY	06B NA48	\pm
------------------------------	-----	-----------------------	----------	-------

¹⁰⁵ This fit is obtained with the CABIBBO 05 matrix element in the $2\pi^0$ invariant mass squared range $0.074094 < m_{2\pi^0}^2 < 0.104244 \text{ GeV}^2$. Electromagnetic corrections and CHPT constraints for $\pi\pi$ phase shifts (a_0 and a_2) have been used. Also measured $(a_0 - a_2) m_{\pi^+} = 0.2646 \pm 0.0021 \pm 0.0023$, where k_0 was kept fixed in the fit at -0.0099 .

¹⁰⁶ Superseded by BATLEY 09A. This fit is obtained with the CABIBBO 05 matrix element in the $2\pi^0$ invariant mass squared range $0.074 \text{ GeV}^2 < m_{2\pi^0}^2 < 0.097 \text{ GeV}^2$, assuming $k = 0$ (no term proportional to $(s_2 - s_1)^2$) and excluding the kinematic region around the cusp ($m_{2\pi^0}^2 = (2m_{\pi^+})^2 \pm 0.000525 \text{ GeV}^2$). Also π - π phase shifts a_0 and a_2 are measured: $(a_0 - a_2) m_{\pi^+} = 0.268 \pm 0.010 \pm 0.004 \pm 0.013(\text{external})$ and $a_2 m_{\pi^+} = -0.041 \pm 0.022 \pm 0.014$.

QUADRATIC COEFFICIENT k_0 FOR $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$

VALUE	EVTS	DOCUMENT ID	TECN	CHG
$0.0095 \pm 0.00017 \pm 0.00048$	60M	¹⁰⁷ BATLEY	09A NA48	\pm

¹⁰⁷ Assumed $a_2 m_{\pi^+} = -0.0044$ in the fit.

LINEAR COEFFICIENT g_{BB} FOR $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$

VALUE	EVTS	DOCUMENT ID	TECN	CHG
$0.6219 \pm 0.0009 \pm 0.0033$	60M	¹⁰⁸ BATLEY	09A NA48	\pm

¹⁰⁸ This fit is obtained using parametrizations of COLANGELO 06A and BISSEGGGER 09 in the $2\pi^0$ invariant mass squared range $0.074094 < m_{2\pi^0}^2 < 0.104244 \text{ GeV}^2$. Electromagnetic corrections and CHPT constraints for $\pi\pi$ phase shifts (a_0 and a_2) have been used. Also measured $(a_0 - a_2) m_{\pi^+} = 0.2633 \pm 0.0024 \pm 0.0024$, where k_0 was kept fixed in the fit at 0.0085.

QUADRATIC COEFFICIENT h'_{BB} FOR $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$

VALUE	EVTS	DOCUMENT ID	TECN	CHG
$-0.0520 \pm 0.0009 \pm 0.0026$	60M	¹⁰⁹ BATLEY	09A NA48	\pm

¹⁰⁹This fit is obtained using parametrizations of COLANGELO 06A and BISSEGGER 09 in the $2\pi^0$ invariant mass squared range $0.074094 < m_{2\pi^0}^2 < 0.104244 \text{ GeV}^2$. Electromagnetic corrections and CHPT constraints for $\pi\pi$ phase shifts (a_0 and a_2) have been used. Also measured $(a_0 - a_2) m_{\pi^+} = 0.2633 \pm 0.0024 \pm 0.0024$, where k_0 was kept fixed in the fit at 0.0085.

A REVIEW GOES HERE – Check our WWW List of Reviews

K_{e3}^{\pm} FORM FACTORS

In the form factor comments, the following symbols are used.

f_+ and f_- are form factors for the vector matrix element.

f_S and f_T refer to the scalar and tensor term.

$$f_0 = f_+ + f_- t / (m_{K^+}^2 - m_{\pi^0}^2).$$

t = momentum transfer to the π .

λ_+ and λ_0 are the linear expansion coefficients of f_+ and f_0 :

$$f_+(t) = f_+(0) (1 + \lambda_+ t / m_{\pi^+}^2)$$

For quadratic expansion

$$f_+(t) = f_+(0) (1 + \lambda'_+ t / m_{\pi^+}^2 + \frac{\lambda''_+}{2} t^2 / m_{\pi^+}^4)$$

as used by KTeV. If there is a non-vanishing quadratic term, then λ_+ represents an average slope, which is then different from λ'_+ .

NA48 and ISTRA quadratic expansion coefficients are converted with $\lambda'_+{}^{PDG} = \lambda_+{}^{NA48}$ and $\lambda''_+{}^{PDG} = 2 \lambda'_+{}^{NA48}$

$$\lambda'_+{}^{PDG} = \left(\frac{m_{\pi^+}}{m_{\pi^0}}\right)^2 \lambda_+{}^{ISTRA} \text{ and}$$

$$\lambda''_+{}^{PDG} = 2 \left(\frac{m_{\pi^+}}{m_{\pi^0}}\right)^4 \lambda'_+{}^{ISTRA}$$

ISTRA linear expansion coefficients are converted with

$$\lambda_+{}^{PDG} = \left(\frac{m_{\pi^+}}{m_{\pi^0}}\right)^2 \lambda_+{}^{ISTRA} \text{ and } \lambda_0{}^{PDG} = \left(\frac{m_{\pi^+}}{m_{\pi^0}}\right)^2 \lambda_0{}^{ISTRA}$$

The pole parametrization is

$$f_+(t) = f_+(0) \left(\frac{M_V^2}{M_V^2 - t}\right)$$

$$f_0(t) = f_0(0) \left(\frac{M_S^2}{M_S^2 - t}\right)$$

where M_V and M_S are the vector and scalar pole masses.

The following abbreviations are used:

DP = Dalitz plot analysis.

PI = π spectrum analysis.

MU = μ spectrum analysis.

POL = μ polarization analysis.

BR = $K_{\mu 3}^{\pm} / K_{e 3}^{\pm}$ branching ratio analysis.

E = positron or electron spectrum analysis.

RC = radiative corrections.

λ_+ (LINEAR ENERGY DEPENDENCE OF f_+ IN K_{e3}^\pm DECAY)

These results are for a linear expansion only. See the next section for fits including a quadratic term. For radiative correction of the K_{e3}^\pm Dalitz plot, see GINSBERG 67, BECHERRAWY 70, CIRIGLIANO 02, CIRIGLIANO 04, and ANDRE 07. Results labeled OUR FIT are discussed in the review “ $K_{\ell 3}^\pm$ and $K_{\ell 3}^0$ Form Factors” above. For earlier, lower statistics results, see the 2004 edition of this review, Physics Letters **B592** 1 (2004).

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2.97 ± 0.05 OUR FIT	Assuming μ -e universality				
2.98 ± 0.05 OUR AVERAGE					
3.044 ± 0.083 ± 0.074	1.1M	AKOPDZANOV 09	TNF	±	
2.966 ± 0.050 ± 0.034	919k	¹¹⁰ YUSHCHENKO 04B	ISTR	–	DP
2.78 ± 0.26 ± 0.30	41k	SHIMIZU 00	SPEC	+	DP
2.84 ± 0.27 ± 0.20	32k	¹¹¹ AKIMENKO 91	SPEC		PI, no RC
2.9 ± 0.4	62k	¹¹² BOLOTOV 88	SPEC		PI, no RC
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.06 ± 0.09 ± 0.06	550k	^{110,113} AJINENKO 03C	ISTR	–	DP
2.93 ± 0.15 ± 0.2	130k	¹¹³ AJINENKO 02	SPEC		DP

¹¹⁰ Rescaled to agree with our conventions as noted above.

¹¹¹ AKIMENKO 91 state that radiative corrections would raise λ_+ by 0.0013.

¹¹² BOLOTOV 88 state radiative corrections of GINSBERG 67 would raise λ_+ by 0.002.

¹¹³ Superseded by YUSHCHENKO 04B.

λ_+ (LINEAR ENERGY DEPENDENCE OF f_+ IN $K_{\mu 3}^\pm$ DECAY)

Results labeled OUR FIT are discussed in the review “ $K_{\ell 3}^\pm$ and $K_{\ell 3}^0$ Form Factors” above. For earlier, lower statistics results, see the 2004 edition of this review, Physics Letters **B592** 1 (2004).

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2.97 ± 0.05 OUR FIT	Assuming μ -e universality				
2.96 ± 0.17 OUR FIT	Not assuming μ -e universality				
2.96 ± 0.14 ± 0.10	540k	¹¹⁴ YUSHCHENKO04	ISTR	–	DP
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.21 ± 0.45	112k	¹¹⁵ AJINENKO 03	ISTR	–	DP

¹¹⁴ Rescaled to agree with our conventions as noted above.

¹¹⁵ Superseded by YUSHCHENKO 04.

λ_0 (LINEAR ENERGY DEPENDENCE OF f_0 IN $K_{\mu 3}^\pm$ DECAY)

Results labeled OUR FIT are discussed in the review “ $K_{\ell 3}^\pm$ and $K_{\ell 3}^0$ Form Factors” above. For earlier, lower statistics results, see the 2004 edition of this review, Physics Letters **B592** 1 (2004).

VALUE (units 10^{-2})	$d\lambda_0/d\lambda_+$	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1.95 ± 0.12 OUR FIT	Assuming μ -e universality					
1.96 ± 0.13 OUR FIT	Not assuming μ -e universality					
+1.96 ± 0.12 ± 0.06	–0.348	540k	¹¹⁶ YUSHCHENKO04	ISTR	–	DP

• • • We do not use the following data for averages, fits, limits, etc. • • •

+2.09±0.45	-0.46	112k	117	AJINENKO	03	ISTR	-	DP
+1.9 ±0.64		24k	118	HORIE	01	SPEC	+	BR
+1.9 ±1.0	+0.03	55k	119	HEINTZE	77	SPEC	+	BR

116 Rescaled to agree with our conventions as noted above.

117 Superseded by YUSHCHENKO 04.

118 HORIE 01 assumes μ - e universality in $K_{\ell 3}^+$ decay and uses SHIMIZU 00 value $\lambda=0.0278 \pm 0.0040$ from K_{e3}^{\pm} decay.

119 HEINTZE 77 uses $\lambda_+ = 0.029 \pm 0.003$. $d\lambda_0/d\lambda_+$ estimated by us.

λ'_+ (LINEAR K_{e3}^{\pm} FORM FACTOR FROM QUADRATIC FIT)

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2.485±0.163±0.034	919k	120,121	YUSHCHENKO04B	ISTR	- DP

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.07 ±0.21	550k	120,122	AJINENKO	03C	ISTR	-	DP
------------	------	---------	----------	-----	------	---	----

120 Rescaled to agree with our conventions as noted above.

121 YUSHCHENKO 04B λ'_+ and λ''_+ are strongly correlated with coefficient $\rho(\lambda'_+, \lambda''_+) = -0.95$.

122 Superseded by YUSHCHENKO 04B.

λ''_+ (QUADRATIC K_{e3}^{\pm} FORM FACTOR)

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.192±0.062±0.071	919k	123,124	YUSHCHENKO04B	ISTR	- DP

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.5 ±0.7 ±1.5	550k	123,125	AJINENKO	03C	ISTR	-	DP
----------------	------	---------	----------	-----	------	---	----

123 Rescaled to agree with our conventions as noted above.

124 YUSHCHENKO 04B λ'_+ and λ''_+ are strongly correlated with coefficient $\rho(\lambda'_+, \lambda''_+) = -0.95$.

125 Superseded by YUSHCHENKO 04B.

$|f_S/f_+|$ FOR K_{e3}^{\pm} DECAY

Ratio of scalar to f_+ couplings.

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
--------------------------	-----	------	-------------	------	-----	---------

-0.3 ^{+0.8}/_{-0.7} OUR AVERAGE

-0.37 ^{+0.66} / _{-0.56} ±0.41	919k	YUSHCHENKO04B	ISTR	-	$\lambda'_+, \lambda''_+, f_S$ fit
---	------	---------------	------	---	------------------------------------

0.2 ±2.6 ±1.4	41k	SHIMIZU	00	SPEC	+ λ_+, f_S, f_T fit
---------------	-----	---------	----	------	-----------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.2 ^{+2.0} / _{-2.2} ±0.3	550k	126	AJINENKO	03C	ISTR	- λ_+, f_S, f_T fit
--	------	-----	----------	-----	------	-----------------------------

-1.9 ^{+2.5} / _{-1.6}	130k	126	AJINENKO	02	SPEC	λ_+, f_S fit
--	------	-----	----------	----	------	----------------------

7.0 ±1.6 ±1.6	32k	AKIMENKO	91	SPEC	$\lambda_+, f_S, f_T, \phi$ fit
---------------	-----	----------	----	------	---------------------------------

0 ± 10	2827	127	BRAUN	75	HLBC	+
--------	------	-----	-------	----	------	---

< 13	90	4017	CHIANG	72	OSPK	+
------	----	------	--------	----	------	---

14 ⁺³ / ₋₄	2707	127	STEINER	71	HLBC	+ $\lambda_+, f_S, f_T, \phi$ fit
----------------------------------	------	-----	---------	----	------	-----------------------------------

< 23	90	BOTTERILL	68C	ASPK		
------	----	-----------	-----	------	--	--

< 18	90	BELLOTTI	67B	HLBC		
------	----	----------	-----	------	--	--

< 30	95	KALMUS	67	HLBC	+	
------	----	--------	----	------	---	--

¹²⁶ Superseded by YUSHCHENKO 04B.

¹²⁷ Statistical errors only.

$|f_T/f_+|$ FOR K_{e3}^\pm DECAY

Ratio of tensor to f_+ couplings.

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
– 1.2± 2.3 OUR AVERAGE						
– 1.2± 2.1± 1.1		919k	YUSHCHENKO04B	ISTR	–	$\lambda'_+, \lambda''_+, f_T$ fit
1 ±14 ± 9		41k	SHIMIZU	00	SPEC	+ λ_+, f_S, f_T fit
• • • We do not use the following data for averages, fits, limits, etc. • • •						
2.1 ⁺ _– 6.4 [±] _{7.5} ± 2.6		550k	¹²⁸ AJINENKO	03C	ISTR	– λ_+, f_S, f_T fit
– 4.5 ⁺ _– 6.0 _{5.7}		130k	¹²⁸ AJINENKO	02	SPEC	λ_+, f_T fit
53 ⁺ _{–10} ±10		32k	AKIMENKO	91	SPEC	$\lambda_+, f_S, f_T, \phi$ fit
7 ±37		2827	¹²⁹ BRAUN	75	HLBC	+
< 75	90	4017	CHIANG	72	OSPK	+
24 ⁺ _{–14} ±16		2707	¹²⁹ STEINER	71	HLBC	+ $\lambda_+, f_S, f_T, \phi$ fit
< 58	90		BOTTERILL	68C	ASPK	
< 58	90		BELLOTTI	67B	HLBC	
< 110	95		KALMUS	67	HLBC	+

¹²⁸ Superseded by YUSHCHENKO 04B.

¹²⁹ Statistical errors only.

f_S/f_+ FOR $K_{\mu 3}^\pm$ DECAY

Ratio of scalar to f_+ couplings.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.17±0.14±0.54	540k	¹³⁰ YUSHCHENKO04	ISTR	–	DP
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.4 ±0.5 ±0.5	112k	¹³¹ AJINENKO	03	ISTR	– DP

¹³⁰ The second error is the theoretical error from the uncertainty in the chiral perturbation theory prediction for λ_0 , ± 0.0053 , combined in quadrature with the systematic error ± 0.0009 .

¹³¹ The second error is the theoretical error from the uncertainty in the chiral perturbation theory prediction for λ_0 . Superseded by YUSHCHENKO 04.

f_T/f_+ FOR $K_{\mu 3}^\pm$ DECAY

Ratio of tensor to f_+ couplings.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
–0.07± 0.71±0.20	540k	YUSHCHENKO04	ISTR	–	DP
• • • We do not use the following data for averages, fits, limits, etc. • • •					
–2.1 ± 2.8 ±1.4	112k	¹³² AJINENKO	03	ISTR	– DP
2 ±12	1585	BRAUN	75	HLBC	

¹³² The second error is the theoretical error from the uncertainty in the chiral perturbation theory prediction for λ_0 . Superseded by YUSHCHENKO 04.

$K_{\ell 4}^{\pm}$ FORM FACTORS

Based on the parametrizations of AMOROS 99, the $K_{\ell 4}^{\pm}$ form factors can be expressed as

$$F_s = f_s + f'_s q^2 + f''_s q^4 + f'_e S_e / 4m_\pi^2$$

$$F_p = f_p + f'_p q^2$$

$$G_p = g_p + g'_p q^2$$

$$H_p = h_p + h'_p q^2$$

where $q^2 = (S_\pi / 4m_\pi^2)$, S_π is the invariant mass squared of the dipion, and S_e is the invariant mass squared of the dilepton.

f_s FOR $K^{\pm} \rightarrow \pi^+ \pi^- e^{\pm} \nu$ DECAY

VALUE	EVTS	DOCUMENT ID	TECN	CHG
$5.75 \pm 0.02 \pm 0.08$	400k	¹³³ PISLAK	03 B865	+

¹³³ Radiative corrections included. Using Roy equations and not including isospin breaking, PISLAK 03 obtains the following $\pi\pi$ scattering lengths $a_0^0 = 0.228 \pm 0.012 \pm 0.004^{+0.012}_{-0.016}$ (theor.) and $a_0^2 = -0.037 \pm 0.023 \pm 0.008 \pm 0.003$ (theor.).

f'_s FOR $K^{\pm} \rightarrow \pi^+ \pi^- e^{\pm} \nu$ DECAY

VALUE	EVTS	DOCUMENT ID	TECN	CHG
0.99 ± 0.06 OUR AVERAGE				
$0.99 \pm 0.06 \pm 0.01$	670k	^{134,135} BATLEY	08A NA48	\pm
$1.06 \pm 0.10 \pm 0.40$	400k	¹³⁶ PISLAK	03 B865	+

¹³⁴ BATLEY 08A reports $[f'_s] / [f_s] = (17.2 \pm 0.9 \pm 0.6) \times 10^{-2}$ which we multiply by our best value $f_s = 5.75 \pm 0.08$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹³⁵ Radiative corrections included. Using Roy equations and not including isospin breaking, BATLEY 08A obtains the following $\pi\pi$ scattering length $a_0^0 = 0.233 \pm 0.016 \pm 0.007$ $a_0^2 = -0.0471 \pm 0.011 \pm 0.004$.

¹³⁶ Radiative corrections included. Using Roy equations and not including isospin breaking, PISLAK 03 obtains the following $\pi\pi$ scattering lengths $a_0^0 = 0.228 \pm 0.012 \pm 0.004^{+0.012}_{-0.016}$ (theor.) and $a_0^2 = -0.037 \pm 0.023 \pm 0.008 \pm 0.003$ (theor.).

f''_s FOR $K^{\pm} \rightarrow \pi^+ \pi^- e^{\pm} \nu$ DECAY

VALUE	EVTS	DOCUMENT ID	TECN	CHG
-0.52 ± 0.07 OUR AVERAGE				
$-0.52 \pm 0.07 \pm 0.01$	670k	^{137,138} BATLEY	08A NA48	\pm
$-0.59 \pm 0.12 \pm 0.40$	400k	¹³⁹ PISLAK	03 B865	+

¹³⁷ BATLEY 08A reports $[f''_s] / [f_s] = (-9.0 \pm 0.9 \pm 0.7) \times 10^{-2}$ which we multiply by our best value $f_s = 5.75 \pm 0.08$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹³⁸ Radiative corrections included. Using Roy equations and not including isospin breaking, BATLEY 08A obtains the following $\pi\pi$ scattering length $a_0^0 = 0.233 \pm 0.016 \pm 0.007$ $a_0^2 = -0.0471 \pm 0.011 \pm 0.004$.

¹³⁹ Radiative corrections included. Using Roy equations and not including isospin breaking, PISLAK 03 obtains the following $\pi\pi$ scattering lengths $a_0^0 = 0.228 \pm 0.012 \pm 0.004^{+0.012}_{-0.016}$ (theor.) and $a_0^2 = -0.037 \pm 0.023 \pm 0.008 \pm 0.003$ (theor.).

f'_e FOR $K^\pm \rightarrow \pi^+ \pi^- e^\pm \nu$ DECAY

VALUE	EVTS	DOCUMENT ID	TECN	CHG
0.47±0.07±0.01	670k	140,141 BATLEY	08A NA48	±

¹⁴⁰ BATLEY 08A reports $[f'_e] / [f_s] = (8.1 \pm 0.8 \pm 0.9) \times 10^{-2}$ which we multiply by our best value $f_s = 5.75 \pm 0.08$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁴¹ Radiative corrections included. Using Roy equations and not including isospin breaking, BATLEY 08A obtains the following $\pi\pi$ scattering length $a_0^0 = 0.233 \pm 0.016 \pm 0.007$
 $a_0^2 = -0.0471 \pm 0.011 \pm 0.004$.

f_p FOR $K^\pm \rightarrow \pi^+ \pi^- e^\pm \nu$ DECAY

VALUE	EVTS	DOCUMENT ID	TECN	CHG
-0.276±0.033±0.004	670k	142,143 BATLEY	08A NA48	±

¹⁴² BATLEY 08A reports $[f_p] / [f_s] = (-4.8 \pm 0.4 \pm 0.4) \times 10^{-2}$ which we multiply by our best value $f_s = 5.75 \pm 0.08$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁴³ Radiative corrections included. Using Roy equations and not including isospin breaking, BATLEY 08A obtains the following $\pi\pi$ scattering length $a_0^0 = 0.233 \pm 0.016 \pm 0.007$
 $a_0^2 = -0.0471 \pm 0.011 \pm 0.004$.

g_p FOR $K^\pm \rightarrow \pi^+ \pi^- e^\pm \nu$ DECAY

VALUE	EVTS	DOCUMENT ID	TECN	CHG
4.78±0.17 OUR AVERAGE				Error includes scale factor of 2.4.

5.02±0.10±0.07	670k	144,145 BATLEY	08A NA48	±
----------------	------	----------------	----------	---

4.66±0.05±0.07	400k	146 PISLAK	03 B865	+
----------------	------	------------	---------	---

¹⁴⁴ BATLEY 08A reports $[g_p] / [f_s] = (87.3 \pm 1.3 \pm 1.2) \times 10^{-2}$ which we multiply by our best value $f_s = 5.75 \pm 0.08$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁴⁵ Radiative corrections included. Using Roy equations and not including isospin breaking, BATLEY 08A obtains the following $\pi\pi$ scattering length $a_0^0 = 0.233 \pm 0.016 \pm 0.007$
 $a_0^2 = -0.0471 \pm 0.011 \pm 0.004$.

¹⁴⁶ Radiative corrections included. Using Roy equations and not including isospin breaking, PISLAK 03 obtains the following $\pi\pi$ scattering lengths $a_0^0 = 0.228 \pm 0.012 \pm 0.004$ $^{+0.012}_{-0.016}$ (theor.) and $a_0^2 = -0.037 \pm 0.023 \pm 0.008 \pm 0.003$ (theor.).

g'_p FOR $K^\pm \rightarrow \pi^+ \pi^- e^\pm \nu$ DECAY

VALUE	EVTS	DOCUMENT ID	TECN	CHG
0.60±0.10 OUR AVERAGE				Error includes scale factor of 1.1.

0.47±0.15±0.01	670k	147,148 BATLEY	08A NA48	±
----------------	------	----------------	----------	---

0.67±0.10±0.04	400k	149 PISLAK	03 B865	+
----------------	------	------------	---------	---

¹⁴⁷ BATLEY 08A reports $[g'_p] / [f_s] = (8.1 \pm 2.2 \pm 1.5) \times 10^{-2}$ which we multiply by our best value $f_s = 5.75 \pm 0.08$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁴⁸ Radiative corrections included. Using Roy equations and not including isospin breaking, BATLEY 08A obtains the following $\pi\pi$ scattering length $a_0^0 = 0.233 \pm 0.016 \pm 0.007$
 $a_0^2 = -0.0471 \pm 0.011 \pm 0.004$.

149 Radiative corrections included. Using Roy equations and not including isospin breaking, PISLAK 03 obtains the following $\pi\pi$ scattering lengths $a_0^0 = 0.228 \pm 0.012 \pm 0.004^{+0.012}_{-0.016}$ (theor.) and $a_0^2 = -0.037 \pm 0.023 \pm 0.008 \pm 0.003$ (theor.).

h_p FOR $K^\pm \rightarrow \pi^+\pi^-e^\pm\nu$ DECAY

VALUE	EVTS	DOCUMENT ID	TECN	CHG
-2.46 ± 0.22 OUR AVERAGE		Error includes scale factor of 1.9.		
$-2.36 \pm 0.12 \pm 0.03$	670k	150,151 BATLEY	08A NA48	\pm
$-2.95 \pm 0.19 \pm 0.20$	400k	152 PISLAK	03 B865	$+$

150 BATLEY 08A reports $[h_p] / [f_s] = (-41.1 \pm 1.9 \pm 0.8) \times 10^{-2}$ which we multiply by our best value $f_s = 5.75 \pm 0.08$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

151 Radiative corrections included. Using Roy equations and not including isospin breaking, BATLEY 08A obtains the following $\pi\pi$ scattering length $a_0^0 = 0.233 \pm 0.016 \pm 0.007$ and $a_0^2 = -0.0471 \pm 0.011 \pm 0.004$.

152 Radiative corrections included. Using Roy equations and not including isospin breaking, PISLAK 03 obtains the following $\pi\pi$ scattering lengths $a_0^0 = 0.228 \pm 0.012 \pm 0.004^{+0.012}_{-0.016}$ (theor.) and $a_0^2 = -0.037 \pm 0.023 \pm 0.008 \pm 0.003$ (theor.).

DECAY FORM FACTOR FOR $K^\pm \rightarrow \pi^0\pi^0e^\pm\nu$

Given in BOLOTOV 86B, BARMIN 88B, and SHIMIZU 04.

$K^\pm \rightarrow \ell^\pm\nu\gamma$ FORM FACTORS

For definitions of the axial-vector F_A and vector F_V form factor, see the "Note on $\pi^\pm \rightarrow \ell^\pm\nu\gamma$ and $K^\pm \rightarrow \ell^\pm\nu\gamma$ Form Factors" in the π^\pm section. In the kaon literature, often different definitions $a_K = F_A/m_K$ and $v_K = F_V/m_K$ are used.


$F_A + F_V$, SUM OF AXIAL-VECTOR AND VECTOR FORM FACTOR FOR $K \rightarrow e\nu_e\gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.133 ± 0.008 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
$0.125 \pm 0.007 \pm 0.001$	1.4K	153 AMBROSINO	09E KLOE	E_γ in 10–250 MeV, $p_e > 200$ MeV/c
0.147 ± 0.011	51	154 HEINTZE	79 SPEC	
$0.150^{+0.018}_{-0.023}$	56	155 HEARD	75 SPEC	

153 Vector form factor fitted with a linear function, $V(x) = F_V (1 + \lambda(1-x))$, $x = 2E_\gamma/m_K$. The fitted value of $\lambda = 0.38 \pm 0.20 \pm 0.02$ with a correlation of -0.93 between $(F_V + F_A)$ and λ .

154 HEINTZE 79 quotes absolute value of $|F_A + F_V| \sin\theta_c$. We use $\sin\theta_c = V_{us} = 0.2205$.

155 HEARD 75 quotes absolute value of $|F_A + F_V| \sin\theta_c$. We use $\sin\theta_c = V_{us} = 0.2205$.



$F_A + F_V$, SUM OF AXIAL-VECTOR AND VECTOR FORM FACTOR
 FOR $K \rightarrow e\nu_e\gamma$

**$F_A + F_V$, SUM OF AXIAL-VECTOR AND VECTOR FORM FACTOR FOR
 $K \rightarrow \mu\nu_\mu\gamma$**

VALUE	CL%	EVTS	DOCUMENT ID	TECN	CHG
$0.165 \pm 0.007 \pm 0.011$		2588	¹⁵⁶ ADLER	00B B787	+

• • • We do not use the following data for averages, fits, limits, etc. • • •

-1.2 to 1.1	90		DEMIDOV	90	XEBC
< 0.23	90	¹⁵⁶	AKIBA	85	SPEC

¹⁵⁶ Quotes absolute value. Sign not determined.

**$F_A - F_V$, DIFFERENCE OF AXIAL-VECTOR AND VECTOR FORM FAC-
 TOR FOR $K \rightarrow e\nu_e\gamma$**

VALUE	CL%	EVTS	DOCUMENT ID	TECN	CHG
< 0.49		90	¹⁵⁷ HEINTZE	79	SPEC

¹⁵⁷ HEINTZE 79 quotes $|F_A - F_V| < \sqrt{11} |F_A + F_V|$.

**$F_A - F_V$, DIFFERENCE OF AXIAL-VECTOR AND VECTOR FORM FAC-
 TOR FOR $K \rightarrow \mu\nu_\mu\gamma$**

VALUE	CL%	EVTS	DOCUMENT ID	TECN	CHG
-0.24 to 0.04	90	2588	ADLER	00B B787	+

• • • We do not use the following data for averages, fits, limits, etc. • • •

-2.2 to 0.6	90		DEMIDOV	90	XEBC
-2.5 to 0.3	90		AKIBA	85	SPEC

K^\pm CHARGE RADIUS

<u>VALUE (fm)</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
0.560±0.031 OUR AVERAGE		
0.580±0.040	AMENDOLIA 86B	$K e \rightarrow K e$
0.530±0.050	DALLY 80	$K e \rightarrow K e$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●		
0.620±0.037	BLATNIK 79	VMD + dispersion relations

CP VIOLATION TESTS IN K^+ AND K^- DECAYS

$$\Delta(K_{\pi ee}^\pm) = \frac{\Gamma(K_{\pi ee}^+) - \Gamma(K_{\pi ee}^-)}{\Gamma(K_{\pi ee}^+) + \Gamma(K_{\pi ee}^-)}$$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
-2.2±1.5±0.6	158 BATLEY 09	NA48

¹⁵⁸ This implies an upper limit of 2.1×10^{-2} at 90% CL.

$$\Delta(K_{\pi\mu\mu}^\pm) = \frac{\Gamma(K_{\pi\mu\mu}^+) - \Gamma(K_{\pi\mu\mu}^-)}{\Gamma(K_{\pi\mu\mu}^+) + \Gamma(K_{\pi\mu\mu}^-)}$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
-0.02±0.11±0.04	PARK 02	HYCP

T VIOLATION TESTS IN K^+ AND K^- DECAYS

P_T in $K^+ \rightarrow \pi^0 \mu^+ \nu_\mu$

T-violating muon polarization. Sensitive to new sources of CP violation beyond the Standard Model.

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
-1.7±2.3±1.1	159	ABE 04F	K246	+

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

-4.2±4.9±0.9	3.9M	ABE 99S	K246	+
--------------	------	---------	------	---

¹⁵⁹ Includes three sets of data: 96-97 (ABE 99S), 98, and 99-00 totaling about three times the ABE 99S data sample. Corresponds to $P_T < 5.0 \times 10^{-3}$ at 90% CL.

P_T in $K^+ \rightarrow \mu^+ \nu_\mu \gamma$

T-violating muon polarization. Sensitive to new sources of CP violation beyond the Standard Model.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>
-0.64±1.85±0.10	114k	¹⁶⁰ ANISIMOVSK..03	K246	+

¹⁶⁰ Muons stopped and polarization measured from decay to positrons.

$\text{Im}(\xi)$ in $K^+ \rightarrow \pi^0 \mu^+ \nu_\mu$ DECAY (from transverse μ pol.)

Test of T reversal invariance.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
-0.006 ±0.008 OUR AVERAGE					

-0.0053±0.0071±0.0036	¹⁶¹	ABE 04F	K246	+	
-----------------------	----------------	---------	------	---	--

-0.016 ±0.025	20M	CAMPBELL 81	CNTR	+	Pol.
---------------	-----	-------------	------	---	------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

-0.013 ±0.016 ±0.003	3.9M	ABE 99S	CNTR	+	$p_T K^+$ at rest
----------------------	------	---------	------	---	-------------------

161 Includes three sets of data: 96-97 (ABE 99S), 98, and 99-00 totaling about three times the ABE 99S data sample. Corresponds to $\text{Im}(\xi) < 0.016$ at 90% CL.

K^\pm REFERENCES

AKOPDZANOV	09	PAN 71 2074	G.A. Akopdzanov <i>et al.</i>	(IHEP)
		Translated from YAF 71	2108.	
AMBROSINO	09E	EPJ C64 627	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
Also		EPJ C65 703 (erratum)	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
BATLEY	09	PL B677 246	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
BATLEY	09A	EPJ C64 589	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
BISSEGGER	09	NP B806 178	M. Bissegger <i>et al.</i>	
AMBROSINO	08	JHEP 0801 073	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AMBROSINO	08A	JHEP 0802 098	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AMBROSINO	08E	PL B666 305	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
ARTAMONOV	08	PRL 101 191802	A.V. Artamonov <i>et al.</i>	(BNL E949 Collab.)
Also		PR D79 092004	A.V. Artamonov <i>et al.</i>	(BNL E949 Collab.)
BATLEY	08	PL B659 493	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
BATLEY	08A	EPJ C54 411	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
AKIMENKO	07	PAN 70 702	S.A. Akimenko <i>et al.</i>	(ISTRA+ Collab.)
		Translated from YAF 70	734.	
ANDRE	07	ANP 322 2518	T. Andre	(EFI)
BATLEY	07A	EPJ C50 329	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
Also		EPJ C52 1021 (erratum)	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
BATLEY	07B	PL B649 349	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
BATLEY	07E	EPJ C52 875	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
TCHIKILEV	07	PAN 70 29	O.G. Tchikilev <i>et al.</i>	(ISTRA+ Collab.)
ALIEV	06	EPJ C46 61	M.A. Aliev <i>et al.</i>	(KEK E470 Collab.)
AMBROSINO	06A	PL B632 76	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
BATLEY	06	PL B634 474	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
BATLEY	06A	PL B638 22	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
Also		PL B640 297 (erratum)	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
BATLEY	06B	PL B633 173	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
COLANGELO	06A	PL B638 187	G. Colangelo <i>et al.</i>	
MA	06	PR D73 037101	H. Ma <i>et al.</i>	(BNL E865 Collab.)
PDG	06	JPG 33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
SHIMIZU	06	PL B633 190	S. Shimizu <i>et al.</i>	(KEK E470 Collab.)
UVAROV	06	PAN 69 26	V.A. Uvarov <i>et al.</i>	(ISTRA+ Collab.)
AKOPDZHAN...	05	EPJ C40 343	G.A. Akopdzhanov <i>et al.</i>	(IHEP)
Also		PAN 68 948	G.A. Akopdzhanov <i>et al.</i>	(IHEP)
		Translated from YAF 68	986.	
AKOPDZHAN...	05B	JETPL 82 675	G.A. Akopdzhanov <i>et al.</i>	(IHEP)
		Translated from ZETFP	82 771.	
ARTAMONOV	05	PL B623 192	A.V. Artamonov <i>et al.</i>	(BNL E949 Collab.)
CABIBBO	05	JHEP 0503 021	N. Cabibbo, G. Isidori	(CERN, ROMAI, FRAS)
SHER	05	PR D72 012005	A. Sher <i>et al.</i>	(BNL E865 Collab.)
ABE	04F	PRL 93 131601	M. Abe <i>et al.</i>	(KEK E246 Collab.)
Also		PR D73 072005	M. Abe <i>et al.</i>	(KEK E246 Collab.)
ADLER	04	PR D70 037102	S. Adler <i>et al.</i>	(BNL E787 Collab.)
ALOISIO	04A	PL B597 139	A. Aloisio <i>et al.</i>	(KLOE Collab.)
ANISIMOVSK...	04	PRL 93 031801	V.V. Anisimovsky <i>et al.</i>	(BNL E949 Collab.)
Also		PR D77 052003	S. Adler <i>et al.</i>	(BNL E949 Collab.)
CABIBBO	04A	PRL 93 121801	N. Cabibbo	(CERN, ROMAI)
CIRIGLIANO	04	EPJ C35 53	V. Cirigliano, H. Neufeld, H. Pichl	(CIT, VALE+)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
SHIMIZU	04	PR D70 037101	S. Shimizu <i>et al.</i>	(KEK E470 Collab.)
YUSHCHENKO	04	PL B581 31	O.P. Yushchenko <i>et al.</i>	(INRM, INRM)
YUSHCHENKO	04B	PL B589 111	O.P. Yushchenko <i>et al.</i>	(INRM)
AJINENKO	03	PAN 66 105	I.V. Ajinenko <i>et al.</i>	(IHEP, INRM)
		Translated from YAF 66	107.	
AJINENKO	03B	PL B567 159	I.V. Ajinenko <i>et al.</i>	(IHEP, INRM)
AJINENKO	03C	PL B574 14	I.V. Ajinenko <i>et al.</i>	(IHEP, INRM)
ALIEV	03	PL B554 7	M.A. Aliev <i>et al.</i>	(KEK E470 Collab.)
ANISIMOVSK...	03	PL B562 166	V.V. Anisimovsky <i>et al.</i>	
PISLAK	03	PR D67 072004	S. Pislak <i>et al.</i>	(BNL E865 Collab.)
SHER	03	PRL 91 261802	A. Sher <i>et al.</i>	(BNL E865 Collab.)
ADLER	02	PRL 88 041803	S. Adler <i>et al.</i>	(BNL E787 Collab.)
ADLER	02B	PR D65 052009	S. Adler <i>et al.</i>	(BNL E787 Collab.)
ADLER	02C	PL B537 211	S. Adler <i>et al.</i>	(BNL E787 Collab.)
AJINENKO	02	PAN 65 2064	I.V. Ajinenko <i>et al.</i>	(IHEP, INRM)
		Translated from YAF 65	2125.	

CIRIGLIANO	02	EPJ C23 121	V. Cirigliano <i>et al.</i>	(VIEN, VALE, MARS)
PARK	02	PRL 88 111801	H.K. Park <i>et al.</i>	(FNAL HyperCP Collab.)
PDG	02	PR D66 010001	K. Hagiwara <i>et al.</i>	
POBLAGUEV	02	PRL 89 061803	A.A. Poblaguev <i>et al.</i>	(BNL 865 Collab.)
ADLER	01	PR D63 032004	S. Adler <i>et al.</i>	(BNL E787 Collab.)
HORIE	01	PL B513 311	K. Horie <i>et al.</i>	(KEK E426 Collab.)
PISLAK	01	PRL 87 221801	S. Pislak <i>et al.</i>	(BNL E865 Collab.)
Also		PR D67 072004	S. Pislak <i>et al.</i>	(BNL E865 Collab.)
ADLER	00	PRL 84 3768	S. Adler <i>et al.</i>	(BNL E787 Collab.)
ADLER	00B	PRL 85 2256	S. Adler <i>et al.</i>	(BNL E787 Collab.)
ADLER	00C	PRL 85 4856	S. Adler <i>et al.</i>	(BNL E787 Collab.)
APPEL	00	PRL 85 2450	R. Appel <i>et al.</i>	(BNL 865 Collab.)
Also		Thesis, Yale Univ.	D.R. Bergman	
Also		Thesis, Univ. Zurich	S. Pislak	
APPEL	00B	PRL 85 2877	R. Appel <i>et al.</i>	(BNL 865 Collab.)
MA	00	PRL 84 2580	H. Ma <i>et al.</i>	(BNL 865 Collab.)
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i>	
SHIMIZU	00	PL B495 33	S. Shimizu <i>et al.</i>	(KEK E246 Collab.)
ABE	99S	PRL 83 4253	M. Abe <i>et al.</i>	(KEK E246 Collab.)
AMOROS	99	JPG 25 1607	G. Amoros, J. Bijnens	(LUND, HELS)
APPEL	99	PRL 83 4482	R. Appel <i>et al.</i>	(BNL 865 Collab.)
ADLER	98	PR D58 012003	S. Adler <i>et al.</i>	(BNL E787 Collab.)
BATUSOV	98	NP B516 3	V.Y. Batusov <i>et al.</i>	
ADLER	97	PRL 79 2204	S. Adler <i>et al.</i>	(BNL E787 Collab.)
ADLER	97C	PRL 79 4756	S. Adler <i>et al.</i>	(BNL E787 Collab.)
BERGMAN	97	Thesis, Yale Univ.	D.R. Bergman	
KITCHING	97	PRL 79 4079	P. Kitching <i>et al.</i>	(BNL E787 Collab.)
PISLAK	97	Thesis, Univ. Zurich	S. Pislak	
ADLER	96	PRL 76 1421	S. Adler <i>et al.</i>	(BNL E787 Collab.)
KOPTEV	95	JETPL 61 877	V.P. Koptev <i>et al.</i>	(PNPI)
		Translated from ZETFP 61 865.		
AOKI	94	PR D50 69	M. Aoki <i>et al.</i>	(INUS, KEK, TOKMS)
ATIYA	93	PRL 70 2521	M.S. Atiya <i>et al.</i>	(BNL E787 Collab.)
Also		PRL 71 305 (erratum)	M.S. Atiya <i>et al.</i>	(BNL E787 Collab.)
ATIYA	93B	PR D48 R1	M.S. Atiya <i>et al.</i>	(BNL E787 Collab.)
ALLIEGRO	92	PRL 68 278	C. Alliegro <i>et al.</i>	(BNL, FNAL, PSI+)
BARMIN	92	SJNP 55 547	V.V. Barmin <i>et al.</i>	(ITEP)
		Translated from YAF 55 976.		
IMAZATO	92	PRL 69 877	J. Imazato <i>et al.</i>	(KEK, INUS, TOKY+)
IVANOV	92	THESIS	Yu.M. Ivanov	(PNPI)
LITTENBERG	92	PRL 68 443	L.S. Littenberg, R.E. Shrock	(BNL, STON)
USHER	92	PR D45 3961	T. Usher <i>et al.</i>	(UCI)
AKIMENKO	91	PL B259 225	S.A. Akimenko <i>et al.</i>	(SERP, JINR, TBIL+)
BARMIN	91	SJNP 53 606	V.V. Barmin <i>et al.</i>	(ITEP)
		Translated from YAF 53 981.		
DENISOV	91	JETPL 54 558	A.S. Denisov <i>et al.</i>	(PNPI)
		Translated from ZETFP 54 557.		
Also		THESIS	Yu.M. Ivanov	(PNPI)
ATIYA	90	PRL 64 21	M.S. Atiya <i>et al.</i>	(BNL E787 Collab.)
ATIYA	90B	PRL 65 1188	M.S. Atiya <i>et al.</i>	(BNL E787 Collab.)
DEMIDOV	90	SJNP 52 1006	V.S. Demidov <i>et al.</i>	(ITEP)
		Translated from YAF 52 1595.		
LEE	90	PRL 64 165	A.M. Lee <i>et al.</i>	(BNL, FNAL, VILL, WASH+)
ATIYA	89	PRL 63 2177	M.S. Atiya <i>et al.</i>	(BNL E787 Collab.)
BARMIN	89	SJNP 50 421	V.V. Barmin <i>et al.</i>	(ITEP)
		Translated from YAF 50 679.		
BARMIN	88	SJNP 47 643	V.V. Barmin <i>et al.</i>	(ITEP)
		Translated from YAF 47 1011.		
BARMIN	88B	SJNP 48 1032	V.V. Barmin <i>et al.</i>	(ITEP)
		Translated from YAF 48 1719.		
BOLOTOV	88	JETPL 47 7	V.N. Bolotov <i>et al.</i>	(ASCI)
		Translated from ZETFP 47 8.		
GALL	88	PRL 60 186	K.P. Gall <i>et al.</i>	(BOST, MIT, WILL, CIT+)
BARMIN	87	SJNP 45 62	V.V. Barmin <i>et al.</i>	(ITEP)
		Translated from YAF 45 97.		
BOLOTOV	87	SJNP 45 1023	V.N. Bolotov <i>et al.</i>	(INRM)
		Translated from YAF 45 1652.		
AMENDOLIA	86B	PL B178 435	S.R. Amendolia <i>et al.</i>	(CERN NA7 Collab.)
BOLOTOV	86	SJNP 44 73	V.N. Bolotov <i>et al.</i>	(INRM)
		Translated from YAF 44 117.		
BOLOTOV	86B	SJNP 44 68	V.N. Bolotov <i>et al.</i>	(INRM)
		Translated from YAF 44 108.		

YAMANAKA	86	PR D34 85	T. Yamanaka <i>et al.</i>	(KEK, TOKY)
Also		PRL 52 329	R.S. Hayano <i>et al.</i>	(TOKY, KEK)
AKIBA	85	PR D32 2911	Y. Akiba <i>et al.</i>	(TOKY, TINT, TSUK, KEK)
BOLOTOV	85	JETPL 42 481	V.N. Bolotov <i>et al.</i>	(INRM)
		Translated from ZETFP 42 390.		
ASANO	82	PL 113B 195	Y. Asano <i>et al.</i>	(KEK, TOKY, INUS, OSAK)
COOPER	82	PL 112B 97	A.M. Cooper <i>et al.</i>	(RL)
PDG	82B	PL 111B 70	M. Roos <i>et al.</i>	(HELS, CIT, CERN)
ASANO	81B	PL 107B 159	Y. Asano <i>et al.</i>	(KEK, TOKY, INUS, OSAK)
CAMPBELL	81	PRL 47 1032	M.K. Campbell <i>et al.</i>	(YALE, BNL)
Also		PR D27 1056	S.R. Blatt <i>et al.</i>	(YALE, BNL)
LUM	81	PR D23 2522	G.K. Lum <i>et al.</i>	(LBL, NBS+)
LYONS	81	ZPHY C10 215	L. Lyons, C. Albajar, G. Myatt	(OXF)
DALLY	80	PRL 45 232	E.B. Dally <i>et al.</i>	(UCLA+)
BARKOV	79	NP B148 53	L.M. Barkov <i>et al.</i>	(NOVO, KIAE)
BLATNIK	79	LNC 24 39	S. Blatnik, J. Stahov, C.B. Lang	(TUZL, GRAZ)
HEINTZE	79	NP B149 365	J. Heintze <i>et al.</i>	(HEIDP, CERN)
ABRAMS	77	PR D15 22	R.J. Abrams <i>et al.</i>	(BNL)
DEVAUX	77	NP B126 11	B. Devaux <i>et al.</i>	(SACL, GEVA)
HEINTZE	77	PL 70B 482	J. Heintze <i>et al.</i>	(HEIDP, CERN)
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)
BLOCH	76	PL 60B 393	P. Bloch <i>et al.</i>	(GEVA, SACL)
BRAUN	76B	LNC 17 521	H.M. Braun <i>et al.</i>	(AACH3, BARI, BELG+)
DIAMANT-...	76	PL 62B 485	A.M. Diamant-Berger <i>et al.</i>	(SACL, GEVA)
HEINTZE	76	PL 60B 302	J. Heintze <i>et al.</i>	(HEIDP)
SMITH	76	NP B109 173	K.M. Smith <i>et al.</i>	(GLAS, LIVP, OXF+)
WEISSENBE...	76	NP B115 55	A.O. Weissenberg <i>et al.</i>	(ITEP, LEBD)
BLOCH	75	PL 56B 201	P. Bloch <i>et al.</i>	(SACL, GEVA)
BRAUN	75	NP B89 210	H.M. Braun <i>et al.</i>	(AACH3, BARI, BRUX+)
CHENG	75	NP A254 381	S.C. Cheng <i>et al.</i>	(COLU, YALE)
HEARD	75	PL 55B 324	K.S. Heard <i>et al.</i>	(CERN, HEIDH)
HEARD	75B	PL 55B 327	K.S. Heard <i>et al.</i>	(CERN, HEIDH)
SHEAFF	75	PR D12 2570	M. Sheaff	(WISC)
SMITH	75	NP B91 45	K.M. Smith <i>et al.</i>	(GLAS, LIVP, OXF+)
WEISSENBE...	74	PL 48B 474	A.O. Weissenberg <i>et al.</i>	(ITEP, LEBD)
ABRAMS	73B	PRL 30 500	R.J. Abrams <i>et al.</i>	(BNL)
BACKENSTO...	73	PL 43B 431	G. Backenstoss <i>et al.</i>	(CERN, KARLK, KARLE+)
LJUNG	73	PR D8 1307	D. Ljung, D. Cline	(WISC)
Also		PRL 28 523	D. Ljung	(WISC)
Also		PRL 28 1287	D. Cline, D. Ljung	(WISC)
Also		PRL 23 326	U. Camerini <i>et al.</i>	(WISC)
LUCAS	73	PR D8 719	P.W. Lucas, H.D. Taft, W.J. Willis	(YALE)
LUCAS	73B	PR D8 727	P.W. Lucas, H.D. Taft, W.J. Willis	(YALE)
PANG	73	PR D8 1989	C.Y. Pang <i>et al.</i>	(EFI, ARIZ, LBL)
Also		PL 40B 699	G.D. Cable <i>et al.</i>	(EFI, LBL)
SMITH	73	NP B60 411	K.M. Smith <i>et al.</i>	(GLAS, LIVP, OXF+)
ABRAMS	72	PRL 29 1118	R.J. Abrams <i>et al.</i>	(BNL)
AUBERT	72	NC 12A 509	B. Aubert <i>et al.</i>	(ORSAY, BRUX, EPOL)
CHIANG	72	PR D6 1254	I.H. Chiang <i>et al.</i>	(ROCH, WISC)
CLARK	72	PRL 29 1274	A.R. Clark <i>et al.</i>	(LBL)
EDWARDS	72	PR D5 2720	R.T. Edwards <i>et al.</i>	(ILL)
FORD	72	PL 38B 335	W.T. Ford <i>et al.</i>	(PRIN)
HOFFMASTER	72	NP B36 1	S. Hoffmaster <i>et al.</i>	(STEV, SETO, LEHI)
BOURQUIN	71	PL 36B 615	M.H. Bourquin <i>et al.</i>	(GEVA, SACL)
HAIDT	71	PR D3 10	D. Haidt	(AACH, BARI, CERN, EPOL, NIJM+)
Also		PL 29B 691	D. Haidt <i>et al.</i>	(AACH, BARI, CERN, EPOL+)
KLEMS	71	PR D4 66	J.H. Klems, R.H. Hildebrand, R. Stiening	(CHIC+)
Also		PRL 24 1086	J.H. Klems, R.H. Hildebrand, R. Stiening	(LRL+)
Also		PRL 25 473	J.H. Klems, R.H. Hildebrand, R. Stiening	(LRL+)
OTT	71	PR D3 52	R.J. Ott, T.W. Pritchard	(LOQM)
ROMANO	71	PL 36B 525	F. Romano <i>et al.</i>	(BARI, CERN, ORSAY)
SCHWEINB...	71	PL 36B 246	W. Schweinberger	(AACH, BELG, CERN, NIJM+)
STEINER	71	PL 36B 521	H.J. Steiner	(AACH, BARI, CERN, EPOL, ORSAY+)
BARDIN	70	PL 32B 121	D.Y. Bardin, S.N. Bilenky, B.M. Pontecorvo	(JINR)
BECHERRAWY	70	PR D1 1452	T. Becherrawy	(ROCH)
FORD	70	PRL 25 1370	W.T. Ford <i>et al.</i>	(PRIN)
GAILLARD	70	CERN 70-14	J.M. Gaillard, L.M. Chounet	(CERN, ORSAY)
GRAUMAN	70	PR D1 1277	J. Grauman <i>et al.</i>	(STEV, SETO, LEHI)
Also		PRL 23 737	J.U. Grauman <i>et al.</i>	(STEV, SETO, LEHI)
MALTSEV	70	SJNP 10 678	E.I. Maltsev <i>et al.</i>	(JINR)
		Translated from YAF 10 1195.		

PANDOULAS	70	PR D2 1205	D. Pandoulas <i>et al.</i>	(STEV, SETO)
CUTTS	69	PR 184 1380	D. Cutts <i>et al.</i>	(LRL, MIT)
Also		PRL 20 955	D. Cutts <i>et al.</i>	(LRL, MIT)
DAVISON	69	PR 180 1333	D.C. Davison <i>et al.</i>	(UCR)
ELY	69	PR 180 1319	R.P.J. Ely <i>et al.</i>	(LOUC, WISC, LRL)
EMMERSON	69	PRL 23 393	J.M.L. Emmerson, T.W. Quirk	(OXF)
HERZO	69	PR 186 1403	D. Herzo <i>et al.</i>	(ILL)
LOBKOWICZ	69	PR 185 1676	F. Lobkowicz <i>et al.</i>	(ROCH, BNL)
Also		PRL 17 548	F. Lobkowicz <i>et al.</i>	(ROCH, BNL)
MAST	69	PR 183 1200	T.S. Mast <i>et al.</i>	(LRL)
SELLERI	69	NC 60A 291	F. Selleri	
ZELLER	69	PR 182 1420	M.E. Zeller <i>et al.</i>	(UCLA, LRL)
BOTTERILL	68B	PRL 21 766	D.R. Botterill <i>et al.</i>	(OXF)
BOTTERILL	68C	PR 174 1661	D.R. Botterill <i>et al.</i>	(OXF)
BUTLER	68	UCRL 18420	W.D. Butler <i>et al.</i>	(LRL)
CHANG	68	PRL 20 510	C.Y. Chang <i>et al.</i>	(UMD, RUTG)
CHEN	68	PRL 20 73	M. Chen <i>et al.</i>	(LRL, MIT)
EICHTEN	68	PL 27B 586	T. Eichten	(AACH, BARI, CERN, EPOL, ORSAY+)
ESCHSTRUTH	68	PR 165 1487	P.T. Eschstruth <i>et al.</i>	(PRIN, PENN)
GARLAND	68	PR 167 1225	R. Garland <i>et al.</i>	(COLU, RUTG, WISC)
MOSCOSO	68	Thesis	L. Moscoso	(ORSAY)
AUERBACH	67	PR 155 1505	L.B. Auerbach <i>et al.</i>	(PENN, PRIN)
Also		PR D9 3216	L.B. Auerbach	
Erratum.				
BELLOTTI	67	Heidelberg Conf.	E. Bellotti, A. Pullia	(MILA)
BELLOTTI	67B	NC 52A 1287	E. Bellotti, E. Fiorini, A. Pullia	(MILA)
Also		PL 20 690	E. Bellotti <i>et al.</i>	(MILA)
BISI	67	PL 25B 572	V. Bisi <i>et al.</i>	(TORI)
FLETCHER	67	PRL 19 98	C.R. Fletcher <i>et al.</i>	(ILL)
FORD	67	PRL 18 1214	W.T. Ford <i>et al.</i>	(PRIN)
GINSBERG	67	PR 162 1570	E.S. Ginsberg	(MASB)
KALMUS	67	PR 159 1187	G.E. Kalmus, A. Kernan	(LRL)
ZINCHENKO	67	Thesis Rutgers	A.I. Zinchenko	(RUTG)
CALLAHAN	66	NC 44A 90	A.C. Callahan	(WISC)
CALLAHAN	66B	PR 150 1153	A.C. Callahan <i>et al.</i>	(WISC, LRL, UCR+)
CESTER	66	PL 21 343	R. Cester <i>et al.</i>	(PPA)
See footnote 1 in		AUERBACH 67.		
Also		PR 155 1505	L.B. Auerbach <i>et al.</i>	(PENN, PRIN)
BIRGE	65	PR 139 B1600	R.W. Birge <i>et al.</i>	(LRL, WISC)
BISI	65	NC 35 768	V. Bisi <i>et al.</i>	(TORI)
BISI	65B	PR 139 B1068	V. Bisi <i>et al.</i>	(TORI)
CALLAHAN	65	PRL 15 129	A. Callahan, D. Cline	(WISC)
CLINE	65	PL 15 293	D. Cline, W.F. Fry	(WISC)
DEMARCO	65	PR 140B 1430	A. de Marco, C. Grosso, G. Rinaudo	(TORI, CERN)
FITCH	65B	PR 140B 1088	V.L. Fitch, C.A. Quarles, H.C. Wilkins	(PRIN+)
STAMER	65	PR 138 B440	P. Stamer <i>et al.</i>	(STEV)
YOUNG	65	Thesis UCRL 16362	P.S. Young	(LRL)
Also		PR 156 1464	P.S. Young, W.Z. Osborne, W.H. Barkas	(LRL)
BORREANI	64	PL 12 123	G. Borreani, G. Rinaudo, A.E. Werbrouck	(TORI)
CALLAHAN	64	PR 136 B1463	A. Callahan, R. March, R. Stark	(WISC)
CLINE	64	PRL 13 101	D. Cline, W.F. Fry	(WISC)
GREINER	64	PRL 13 284	D.E. Greiner, W.Z. Osborne, W.H. Barkas	(LRL)
SHAKLEE	64	PR 136 B1423	F.S. Shaklee <i>et al.</i>	(MICH)
BOYARSKI	62	PR 128 2398	A.M. Boyarski <i>et al.</i>	(MIT)
FERRO-LUZZI	61	NC 22 1087	M. Ferro-Luzzi <i>et al.</i>	(LRL)
ROE	61	PRL 7 346	B.P. Roe <i>et al.</i>	(MICH, LRL)
TAYLOR	59	PR 114 359	S. Taylor <i>et al.</i>	(COLU)
COOMBES	57	PR 108 1348	C.A. Coombes <i>et al.</i>	(LBL)

OTHER RELATED PAPERS

LITTENBERG	93	ARNPS 43 729	L.S. Littenberg, G. Valencia	(BNL, FNAL)
		Rare and Radiative Kaon Decays		
RITCHIE	93	RMP 65 1149	J.L. Ritchie, S.G. Wojcicki	
		"Rare K Decays"		
BATTISTON	92	PRPL 214 293	R. Battiston <i>et al.</i>	(PGIA, CERN, TRSTT)
		Status and Perspectives of K Decay Physics		
BRYMAN	89	IJMP A4 79	D.A. Bryman	(TRIU)
		"Rare Kaon Decays"		
CHOUNET	72	PRPL 4C 199	L.M. Chounet, J.M. Gaillard, M.K. Gaillard	(ORSAY+)
FEARING	70	PR D2 542	H.W. Fearing, E. Fischbach, J. Smith	(STON, BOHR)

HAIDT	69B	PL 29B 696	D. Haidt <i>et al.</i>	(AACH, BARI, CERN, EPOL+)
CRONIN	68B	Vienna Conf. 241	J.W. Cronin	(PRIN)
Rapporteur talk.				
WILLIS	67	Heidelberg Conf. 273	W.J. Willis	(YALE)
Rapporteur talk.				
CABIBBO	66	Berkeley Conf. 33	N. Cabibbo	(CERN)
ADAIR	64	PL 12 67	R.K. Adair, L.B. Leipuner	(YALE, BNL)
CABIBBO	64	PL 9 352	N. Cabibbo, A. Maksymowicz	(CERN)
Also		PL 11 360	N. Cabibbo, A. Maksymowicz	(CERN)
Also		PL 14 72	N. Cabibbo, A. Maksymowicz	(CERN)
BIRGE	63	PRL 11 35	R.W. Birge <i>et al.</i>	(LRL, WISC, BARI)
BLOCK	62B	CERN Conf. 371	M.M. Block, L. Lendinara, L. Monari	(NWES, BGNA)
BRENE	61	NP 22 553	N. Brene, L. Egardt, B. Qvist	(NORD)
