

$f_2(1910)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE


We list here three different peaks with close masses and widths seen in the mass distributions of $\omega\omega$, $\eta\eta'$, and K^+K^- final states. ALDE 91B argues that they are of different nature.

$f_2(1910)$ MASS

$f_2(1910)$ $\omega\omega$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1903 ± 9 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.		
1890 ± 10	¹ AMELIN	06	VES 36 $\pi^- p \rightarrow \omega\omega n$
1934 ± 20	ANISOVICH	00J	SPEC
1897 ± 11	BARBERIS	00F	450 $pp \rightarrow p_f\omega\omega p_s$
1924 ± 14	ALDE	90	GAM2 38 $\pi^- p \rightarrow \omega\omega n$

¹Supersedes BELADIDZE 92B.



$f_2(1910) \eta\eta'$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1934±16	² BARBERIS	00A	450 $p p \rightarrow p_f \eta \eta' p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1911±10	ALDE	91B	GAM2 38 $\pi^- p \rightarrow \eta \eta' n$
² Also compatible with $J^{PC}=1^-+$.			


$f_2(1910) K^+ K^-$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1941±18	AMSLER	06	CBAR 1.64 $\bar{p} p \rightarrow K^+ K^- \pi^0$

$f_2(1910)$ WIDTH

$f_2(1910) \omega\omega$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
196±31 OUR AVERAGE	Error includes scale factor of 2.3. See the ideogram below.		
165±19	³ AMELIN	06	VES 36 $\pi^- p \rightarrow \omega\omega n$
271±25	ANISOVICH	00J	SPEC
202±32	BARBERIS	00F	450 $p p \rightarrow p_f \omega\omega p_s$
91±50	ALDE	90	GAM2 38 $\pi^- p \rightarrow \omega\omega n$
³ Supersedes BELADIDZE 92B.			



$f_2(1910) \eta\eta'$ MODE

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
141±41	⁴ BARBERIS	00A	450 $p p \rightarrow p_f \eta \eta' p_S$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
90±35	ALDE	91B	GAM2 38 $\pi^- p \rightarrow \eta \eta' n$
⁴ Also compatible with $J^{PC}=1^-+$.			

$f_2(1910) K^+ K^-$ MODE

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
120±40	AMSLER	06	CBAR 1.64 $\bar{p} p \rightarrow K^+ K^- \pi^0$

$f_2(1910)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi^0 \pi^0$	
$\Gamma_2 K^+ K^-$	seen
$\Gamma_3 K_S^0 K_S^0$	
$\Gamma_4 \eta \eta$	seen
$\Gamma_5 \omega \omega$	seen
$\Gamma_6 \eta \eta'$	seen
$\Gamma_7 \eta' \eta'$	
$\Gamma_8 \rho \rho$	seen

$f_2(1910)$ BRANCHING RATIOS

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$	Γ_2/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	AMSLER	06	CBAR 1.64 $\bar{p} p \rightarrow K^+ K^- \pi^0$

$\Gamma(\pi^0 \pi^0)/\Gamma(\eta \eta')$	Γ_1/Γ_6		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.1	ALDE	89	GAM2 38 $\pi^- p \rightarrow \eta \eta' n$

$\Gamma(K_S^0 K_S^0)/\Gamma(\eta \eta')$	Γ_3/Γ_6			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.066	90	BALOSHIN	86	SPEC 40 $\pi p \rightarrow K_S^0 K_S^0 n$

$\Gamma(\eta \eta)/\Gamma(\eta \eta')$	Γ_4/Γ_6			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.05	90	ALDE	91B	GAM2 38 $\pi^- p \rightarrow \eta \eta' n$

$\Gamma(\omega\omega)/\Gamma(\eta\eta')$

Γ_5/Γ_6

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •		
2.6 ± 0.6	BARBERIS 00F	450 $pp \rightarrow p_f \omega \omega p_S$

$\Gamma(\eta'\eta')/\Gamma_{\text{total}}$

Γ_7/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
probably not seen	BARBERIS 00A		450 $pp \rightarrow p_f \eta' \eta' p_S$
possibly seen	BELADIDZE 92D	VES	37 $\pi^- p \rightarrow \eta' \eta' n$

$\Gamma(\rho\rho)/\Gamma(\omega\omega)$

Γ_8/Γ_5

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •		
2.6 ± 0.4	BARBERIS 00F	450 $pp \rightarrow p_f \omega \omega p_S$

$f_2(1910)$ REFERENCES

AMELIN	06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.)
		Translated from YAF 69 715.		
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(CBAR Collab.)
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	
BARBERIS	00A	PL B471 429	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00F	PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BELADIDZE	92B	ZPHY C54 367	G.M. Beladidze <i>et al.</i>	(VES Collab.)
BELADIDZE	92D	ZPHY C57 13	G.M. Beladidze <i>et al.</i>	(VES Collab.)
ALDE	91B	SJNP 54 455	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
		Translated from YAF 54 751.		
Also		PL B276 375	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)
ALDE	90	PL B241 600	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
ALDE	89	PL B216 447	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)
Also		SJNP 48 1035	D.M. Alde <i>et al.</i>	(BELG, SERP, LANL, LAPP)
		Translated from YAF 48 1724.		
BALOSHIN	86	SJNP 43 959	O.N. Baloshin <i>et al.</i>	(ITEP)
		Translated from YAF 43 1487.		