

# $\Sigma(1560)$ Bumps

$$I(J^P) = 1(?^?) \quad \text{Status: } **$$

## OMITTED FROM SUMMARY TABLE

This entry lists peaks reported in mass spectra around 1560 MeV without implying that they are necessarily related.

DIONISI 78B observes a 6 standard-deviation enhancement at 1553 MeV in the charged  $\Lambda/\Sigma\pi$  mass spectra from  $K^- p \rightarrow (\Lambda/\Sigma)\pi K\bar{K}$  at 4.2 GeV/c. In a CERN ISR experiment, LOCKMAN 78 reports a narrow 6 standard-deviation enhancement at 1572 MeV in  $\Lambda\pi^\pm$  from the reaction  $pp \rightarrow \Lambda\pi^+\pi^- X$ . These enhancements are unlikely to be associated with the  $\Sigma(1580)$  (which has not been confirmed by several recent experiments – see the next entry in the Listings).

CARROLL 76 observes a bump at 1550 MeV (as well as one at 1580 MeV) in the isospin-1  $\bar{K}N$  total cross section, but uncertainties in cross section measurements outside the mass range of the experiment preclude estimating its significance.

See also MEADOWS 80 for a review of this state.

### $\Sigma(1560)$ MASS (PRODUCTION EXPERIMENTS)

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<b><math>\approx 1560</math> OUR ESTIMATE</b>					
1553 $\pm$ 7	121	DIONISI	78B	HBC	$\pm$ $K^- p \rightarrow (Y\pi)K\bar{K}$
1572 $\pm$ 4	40	LOCKMAN	78	SPEC	$\pm$ $pp \rightarrow \Lambda\pi^+\pi^- X$

### $\Sigma(1560)$ WIDTH (PRODUCTION EXPERIMENTS)

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
79 $\pm$ 30	121	DIONISI	78B	HBC	$\pm$ $K^- p \rightarrow (Y\pi)K\bar{K}$
15 $\pm$ 6	40	<sup>1</sup> LOCKMAN	78	SPEC	$\pm$ $pp \rightarrow \Lambda\pi^+\pi^- X$

### $\Sigma(1560)$ DECAY MODES (PRODUCTION EXPERIMENTS)

Mode	Fraction ( $\Gamma_j/\Gamma$ )
$\Gamma_1$ $\Lambda\pi$	seen
$\Gamma_2$ $\Sigma\pi$	

## $\Sigma(1560)$ BRANCHING RATIOS (PRODUCTION EXPERIMENTS)

$\Gamma(\Sigma\pi)/[\Gamma(\Lambda\pi) + \Gamma(\Sigma\pi)]$	$\Gamma_2/(\Gamma_1+\Gamma_2)$				
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	
0.35 ± 0.12	DIONISI	78B	HBC	±	$K^- p \rightarrow (Y\pi) K \bar{K}$

$\Gamma(\Lambda\pi)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$				
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	
<b>seen</b>	LOCKMAN	78	SPEC	±	$pp \rightarrow \Lambda\pi^+\pi^- X$

## $\Sigma(1560)$ FOOTNOTES (PRODUCTION EXPERIMENTS)

<sup>1</sup> The width observed by LOCKMAN 78 is consistent with experimental resolution.

## $\Sigma(1560)$ REFERENCES (PRODUCTION EXPERIMENTS)

MEADOWS	80	Toronto Conf. 283	B.T. Meadows	
DIONISI	78B	PL 78B 154	C. Dionisi, R. Armenteros, J. Diaz	(CINC)
LOCKMAN	78	Saclay DPHPE 78-01	W. Lockman <i>et al.</i>	(CERN, AMST+) I
CARROLL	76	PRL 37 806	A.S. Carroll <i>et al.</i>	(UCLA, SACL) I
				(BNL) I