



$$I(J^P) = 0(0^-)$$


The angular distributions of the decays of the ϕ and $\bar{K}^*(892)^0$ in the $\phi\pi^+$ and $K^+\bar{K}^*(892)^0$ modes strongly indicate that the spin is zero. The parity given is that expected of a $c\bar{s}$ ground state.

D_s^\pm MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements. Measurements of the D_s^\pm mass with an error greater than 10 MeV are omitted from the fit and average. A number of early measurements have been omitted altogether.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1968.45 ± 0.33 OUR FIT		Error includes scale factor of 1.3.		
1969.0 ± 1.4 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
1967.0 ± 1.0 ± 1.0	54	BARLAG	90C ACCM	π^- Cu 230 GeV
1969.3 ± 1.4 ± 1.4		ALBRECHT	88 ARG	e^+e^- 9.4–10.6 GeV
1972.7 ± 1.5 ± 1.0	21	BECKER	87B SILI	200 GeV π, K, p
1972.4 ± 3.7 ± 3.7	27	BLAYLOCK	87 MRK3	e^+e^- 4.14 GeV
1963 ± 3 ± 3	30	DERRICK	85B HRS	e^+e^- 29 GeV
1970 ± 5 ± 5	104	CHEN	83C CLEO	e^+e^- 10.5 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1968.3 ± 0.7 ± 0.7	290	¹ ANJOS	88 E691	Photoproduction
1980 ± 15	6	USHIDA	86 EMUL	ν wideband
1973.6 ± 2.6 ± 3.0	163	ALBRECHT	85D ARG	e^+e^- 10 GeV
1948 ± 28 ± 10	65	AIHARA	84D TPC	e^+e^- 29 GeV
1975 ± 9 ± 10	49	ALTHOFF	84 TASS	e^+e^- 14–25 GeV
1975 ± 4	3	BAILEY	84 ACCM	hadron ⁺ Be → $\phi\pi^+X$

¹ ANJOS 88 enters the fit via $m_{D_s^\pm} - m_{D^\pm}$ (see below).



$m_{D_s^\pm} - m_{D^\pm}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
98.88±0.30 OUR FIT	Error includes scale factor of 1.4.			
98.85±0.25 OUR AVERAGE	Error includes scale factor of 1.1.			
99.41±0.38±0.21		ACOSTA	03D CDF2	$\bar{p}p, \sqrt{s}=1.96$ TeV
98.4 ±0.1 ±0.3	48k	AUBERT	02G BABR	$e^+e^- \approx \Upsilon(4S)$
99.5 ±0.6 ±0.3		BROWN	94 CLE2	$e^+e^- \approx \Upsilon(4S)$
98.5 ±1.5	555	CHEN	89 CLEO	e^+e^- 10.5 GeV
99.0 ±0.8	290	ANJOS	88 E691	Photoproduction


D_s^\pm MEAN LIFE

Measurements with an error greater than 100×10^{-15} s or with fewer than 100 events have been omitted from the Listings.

VALUE (10^{-15} s)	EVTS	DOCUMENT ID	TECN	COMMENT
500 ± 7 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
507.4± 5.5± 5.1	13.6k	LINK	05J FOCS	$\phi\pi^+$ and $\bar{K}^{*0}K^+$
472.5±17.2± 6.6	760	IORI	01 SELX	600 GeV Σ^-, π^-, p
518 ±14 ± 7	1662	AITALA	99 E791	π^- nucleus, 500 GeV

$486.3 \pm 15.0^{+4.9}_{-5.1}$	2167	² BONVICINI	99	CLE2	$e^+e^- \approx \Upsilon(4S)$
$475 \pm 20 \pm 7$	900	FRABETTI	93F	E687	$\gamma \text{Be}, \phi\pi^+$
$500 \pm 60 \pm 30$	104	FRABETTI	90	E687	$\gamma \text{Be}, \phi\pi^+$
$470 \pm 40 \pm 20$	228	RAAB	88	E691	Photoproduction

² BONVICINI 99 obtains 1.19 ± 0.04 for the ratio of D_s^+ to D^0 lifetimes.



D_s^+ DECAY MODES

Unless otherwise noted, the branching fractions for modes with a resonance in the final state include all the decay modes of the resonance. D_s^- modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Inclusive modes		
Γ_1 e^+ semileptonic	[a] (6.5 ± 0.4) %	
Γ_2 π^+ anything	(119.3 ± 1.4) %	
Γ_3 π^- anything	(43.2 ± 0.9) %	
Γ_4 π^0 anything	(123 ± 7) %	
Γ_5 K^- anything	(18.7 ± 0.5) %	
Γ_6 K^+ anything	(28.9 ± 0.7) %	
Γ_7 K_S^0 anything	(19.0 ± 1.1) %	
Γ_8 η anything	[b] (29.9 ± 2.8) %	

Γ_9	ω anything	(6.1 \pm 1.4) %	
Γ_{10}	η' anything	[c] (11.7 \pm 1.8) %	
Γ_{11}	$f_0(980)$ anything, $f_0 \rightarrow \pi^+ \pi^-$	< 1.3 %	CL=90%
Γ_{12}	ϕ anything	(15.7 \pm 1.0) %	
Γ_{13}	$K^+ K^-$ anything	(15.8 \pm 0.7) %	
Γ_{14}	$K_S^0 K^+$ anything	(5.8 \pm 0.5) %	
Γ_{15}	$K_S^0 K^-$ anything	(1.9 \pm 0.4) %	
Γ_{16}	$2K_S^0$ anything	(1.70 \pm 0.32) %	
Γ_{17}	$2K^+$ anything	< 2.6 $\times 10^{-3}$	CL=90%
Γ_{18}	$2K^-$ anything	< 6 $\times 10^{-4}$	CL=90%

Leptonic and semileptonic modes

Γ_{19}	$e^+ \nu_e$	< 1.2 $\times 10^{-4}$	CL=90%
Γ_{20}	$\mu^+ \nu_\mu$	(5.90 \pm 0.33) $\times 10^{-3}$	
Γ_{21}	$\tau^+ \nu_\tau$	(5.43 \pm 0.31) %	
Γ_{22}	$K^+ K^- e^+ \nu_e$	—	
Γ_{23}	$\phi e^+ \nu_e$	[d] (2.49 \pm 0.14) %	
Γ_{24}	$\eta e^+ \nu_e + \eta'(958) e^+ \nu_e$	[d] (3.66 \pm 0.37) %	
Γ_{25}	$\eta e^+ \nu_e$	[d] (2.67 \pm 0.29) %	S=1.1
Γ_{26}	$\eta'(958) e^+ \nu_e$	[d] (9.9 \pm 2.3) $\times 10^{-3}$	
Γ_{27}	$K^0 e^+ \nu_e$	(3.7 \pm 1.0) $\times 10^{-3}$	
Γ_{28}	$K^*(892)^0 e^+ \nu_e$	[d] (1.8 \pm 0.7) $\times 10^{-3}$	
Γ_{29}	$f_0(980) e^+ \nu_e, f_0 \rightarrow \pi^+ \pi^-$	(2.00 \pm 0.32) $\times 10^{-3}$	

Hadronic modes with a $K\bar{K}$ pair

Γ_{30}	$K^+ K_S^0$	(1.48 \pm 0.08) %	
Γ_{31}	$K^+ K^- \pi^+$	[e] (5.49 \pm 0.27) %	
Γ_{32}	$\phi \pi^+$	[d,f] (4.5 \pm 0.4) %	
Γ_{33}	$\phi \pi^+, \phi \rightarrow K^+ K^-$	[f] (2.32 \pm 0.14) %	
Γ_{34}	$K^+ \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+$	(2.60 \pm 0.15) %	
Γ_{35}	$f_0(980) \pi^+, f_0 \rightarrow K^+ K^-$	(1.55 \pm 0.16) %	
Γ_{36}	$f_0(1370) \pi^+, f_0 \rightarrow K^+ K^-$	(2.4 \pm 0.4) $\times 10^{-3}$	
Γ_{37}	$f_0(1710) \pi^+, f_0 \rightarrow K^+ K^-$	(1.87 \pm 0.33) $\times 10^{-3}$	
Γ_{38}	$K^+ \bar{K}_0^*(1430)^0, \bar{K}_0^* \rightarrow K^- \pi^+$	(2.1 \pm 0.4) $\times 10^{-3}$	
Γ_{39}	$K^0 \bar{K}_0^0 \pi^+$	—	
Γ_{40}	$K^*(892)^+ \bar{K}^0$	[d] (5.4 \pm 1.2) %	
Γ_{41}	$K^+ K^- \pi^+ \pi^0$	(5.6 \pm 0.5) %	
Γ_{42}	$\phi \rho^+$	[d] (8.4 \pm 1.9 / -2.3) %	
Γ_{43}	$K_S^0 K^- 2\pi^+$	(1.64 \pm 0.12) %	
Γ_{44}	$K^*(892)^+ \bar{K}^*(892)^0$	[d] (7.2 \pm 2.6) %	
Γ_{45}	$K^+ K_S^0 \pi^+ \pi^-$	(9.6 \pm 1.3) $\times 10^{-3}$	

Γ ₄₆	$K^+ K^- 2\pi^+ \pi^-$	(8.8 ± 1.6) × 10 ⁻³	
Γ ₄₇	$\phi 2\pi^+ \pi^-$	[d] (1.21 ± 0.16) %	
Γ ₄₈	$K^+ K^- \rho^0 \pi^+$ non- ϕ	< 2.6 × 10 ⁻⁴	CL=90%
Γ ₄₉	$\phi \rho^0 \pi^+$, $\phi \rightarrow K^+ K^-$	(6.6 ± 1.3) × 10 ⁻³	
Γ ₅₀	$\phi a_1(1260)^+$, $\phi \rightarrow K^+ K^-$, $a_1^+ \rightarrow \rho^0 \pi^+$	(7.5 ± 1.3) × 10 ⁻³	
Γ ₅₁	$K^+ K^- 2\pi^+ \pi^-$ nonresonant	(9 ± 7) × 10 ⁻⁴	
Γ ₅₂	$2K_S^0 2\pi^+ \pi^-$	(8.3 ± 3.5) × 10 ⁻⁴	

Hadronic modes without K's

Γ ₅₃	$\pi^+ \pi^0$	< 3.4 × 10 ⁻⁴	CL=90%
Γ ₅₄	$2\pi^+ \pi^-$	(1.10 ± 0.06) %	
Γ ₅₅	$\rho^0 \pi^+$	(2.0 ± 1.2) × 10 ⁻⁴	
Γ ₅₆	$\pi^+ (\pi^+ \pi^-)_{S\text{-wave}}$	[g] (9.2 ± 0.6) × 10 ⁻³	
Γ ₅₇	$f_0(980) \pi^+$, $f_0 \rightarrow \pi^+ \pi^-$		
Γ ₅₈	$f_0(1370) \pi^+$, $f_0 \rightarrow \pi^+ \pi^-$		
Γ ₅₉	$f_0(1500) \pi^+$, $f_0 \rightarrow \pi^+ \pi^-$		
Γ ₆₀	$f_2(1270) \pi^+$, $f_2 \rightarrow \pi^+ \pi^-$	(1.11 ± 0.20) × 10 ⁻³	
Γ ₆₁	$\rho(1450)^0 \pi^+$, $\rho^0 \rightarrow \pi^+ \pi^-$	(3.0 ± 2.0) × 10 ⁻⁴	
Γ ₆₂	$\pi^+ 2\pi^0$	(6.5 ± 1.3) × 10 ⁻³	
Γ ₆₃	$2\pi^+ \pi^- \pi^0$	—	
Γ ₆₄	$\eta \pi^+$	[d] (1.83 ± 0.15) %	
Γ ₆₅	$\omega \pi^+$	[d] (2.5 ± 0.7) × 10 ⁻³	
Γ ₆₆	$3\pi^+ 2\pi^-$	(8.0 ± 0.9) × 10 ⁻³	
Γ ₆₇	$2\pi^+ \pi^- 2\pi^0$	—	
Γ ₆₈	$\eta \rho^+$	[d] (8.9 ± 0.8) %	
Γ ₆₉	$\eta \pi^+ \pi^0$ 3-body	[d] < 5 %	CL=90%
Γ ₇₀	$\omega \pi^+ \pi^0$	[d] (2.8 ± 0.7) %	
Γ ₇₁	$3\pi^+ 2\pi^- \pi^0$	(4.9 ± 3.2) %	
Γ ₇₂	$\omega 2\pi^+ \pi^-$	[d] (1.6 ± 0.5) %	
Γ ₇₃	$\eta'(958) \pi^+$	[c,d] (3.94 ± 0.33) %	
Γ ₇₄	$3\pi^+ 2\pi^- 2\pi^0$	—	
Γ ₇₅	$\omega \eta \pi^+$	[d] < 2.13 %	CL=90%
Γ ₇₆	$\eta'(958) \rho^+$	[c,d] (12.5 ± 2.2) %	
Γ ₇₇	$\eta'(958) \pi^+ \pi^0$ 3-body	[d] < 1.8 %	CL=90%

Modes with one or three K's

Γ ₇₈	$K^+ \pi^0$	(6.2 ± 2.1) × 10 ⁻⁴	
Γ ₇₉	$K_S^0 \pi^+$	(1.21 ± 0.08) × 10 ⁻³	
Γ ₈₀	$K^+ \eta$	[d] (1.75 ± 0.35) × 10 ⁻³	
Γ ₈₁	$K^+ \omega$	[d] < 2.4 × 10 ⁻³	CL=90%
Γ ₈₂	$K^+ \eta'(958)$	[d] (1.8 ± 0.6) × 10 ⁻³	
Γ ₈₃	$K^+ \pi^+ \pi^-$	(6.9 ± 0.5) × 10 ⁻³	
Γ ₈₄	$K^+ \rho^0$	(2.7 ± 0.5) × 10 ⁻³	
Γ ₈₅	$K^+ \rho(1450)^0$, $\rho^0 \rightarrow \pi^+ \pi^-$	(7.3 ± 2.6) × 10 ⁻⁴	

Γ_{86}	$K^*(892)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-$	(1.50 ± 0.26) $\times 10^{-3}$	
Γ_{87}	$K^*(1410)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-$	(1.30 ± 0.31) $\times 10^{-3}$	
Γ_{88}	$K^*(1430)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-$	(5 ± 4) $\times 10^{-4}$	
Γ_{89}	$K^+ \pi^+ \pi^-$ nonresonant	(1.1 ± 0.4) $\times 10^{-3}$	
Γ_{90}	$K^0 \pi^+ \pi^0$	(1.00 ± 0.18) %	
Γ_{91}	$K_S^0 2\pi^+ \pi^-$	(2.9 ± 1.1) $\times 10^{-3}$	
Γ_{92}	$K^+ \omega \pi^0$	[d] < 8.2×10^{-3}	CL=90%
Γ_{93}	$K^+ \omega \pi^+ \pi^-$	[d] < 5.4×10^{-3}	CL=90%
Γ_{94}	$K^+ \omega \eta$	[d] < 7.9×10^{-3}	CL=90%
Γ_{95}	$2K^+ K^-$	(4.9 ± 1.7) $\times 10^{-4}$	
Γ_{96}	ϕK^+	[d] < 6×10^{-4}	CL=90%

Doubly Cabibbo-suppressed modes

Γ_{97}	$2K^+ \pi^-$	(1.29 ± 0.18) $\times 10^{-4}$	
---------------	--------------	--------------------------------------	--

Baryon-antibaryon mode

Γ_{98}	$p \bar{n}$	(1.3 ± 0.4) $\times 10^{-3}$	
---------------	-------------	------------------------------------	--

**$\Delta C = 1$ weak neutral current (C1) modes,
Lepton family number (LF), or
Lepton number (L) violating modes**

Γ_{99}	$\pi^+ e^+ e^-$	[h] < 2.2×10^{-5}	CL=90%
Γ_{100}	$\pi^+ \phi, \phi \rightarrow e^+ e^-$	[i] ($6 \begin{smallmatrix} +8 \\ -4 \end{smallmatrix}$) $\times 10^{-6}$	

**$\Delta C = 1$ weak neutral current (C1) modes,
Lepton family number (LF), or
Lepton number (L) violating modes**

Γ_{101}	$\pi^+ \mu^+ \mu^-$	[h] < 2.6×10^{-5}	CL=90%
Γ_{102}	$K^+ e^+ e^-$	C1 < 5.2×10^{-5}	CL=90%
Γ_{103}	$K^+ \mu^+ \mu^-$	C1 < 3.6×10^{-5}	CL=90%
Γ_{104}	$K^*(892)^+ \mu^+ \mu^-$	C1 < 1.4×10^{-3}	CL=90%
Γ_{105}	$\pi^+ e^\pm \mu^\mp$	LF [j] < 6.1×10^{-4}	CL=90%
Γ_{106}	$K^+ e^\pm \mu^\mp$	LF [j] < 6.3×10^{-4}	CL=90%
Γ_{107}	$\pi^- 2e^+$	L < 1.8×10^{-5}	CL=90%
Γ_{108}	$\pi^- 2\mu^+$	L < 2.9×10^{-5}	CL=90%
Γ_{109}	$\pi^- e^+ \mu^+$	L < 7.3×10^{-4}	CL=90%
Γ_{110}	$K^- 2e^+$	L < 1.7×10^{-5}	CL=90%
Γ_{111}	$K^- 2\mu^+$	L < 1.3×10^{-5}	CL=90%
Γ_{112}	$K^- e^+ \mu^+$	L < 6.8×10^{-4}	CL=90%
Γ_{113}	$K^*(892)^- 2\mu^+$	L < 1.4×10^{-3}	CL=90%

Γ_{114}	A dummy mode used by the fit.	(75.7 ± 1.0) %	
----------------	-------------------------------	----------------------	--

- [a] This is the purely e^+ semileptonic branching fraction: the e^+ fraction from τ^+ decays has been subtracted off. The sum of our (non- τ) e^+ exclusive fractions — an $e^+ \nu_e$ with an η , η' , ϕ , K^0 , K^{*0} , or $f_0(980)$ — is 6.9 ± 0.4 %
 - [b] This fraction includes η from η' decays.
 - [c] Two times (to include μ decays) the $\eta' e^+ \nu_e$ branching fraction, plus the $\eta' \pi^+$, $\eta' \rho^+$, and $\eta' K^+$ fractions, is $(18.6 \pm 2.3)\%$, which considerably exceeds the inclusive η' fraction of $(11.7 \pm 1.8)\%$. Our best guess is that the $\eta' \rho^+$ fraction, $(12.5 \pm 2.2)\%$, is too large.
 - [d] This branching fraction includes all the decay modes of the final-state resonance.
 - [e] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers.
 - [f] We decouple the $D_s^+ \rightarrow \phi \pi^+$ branching fraction obtained from mass projections (and used to get some of the other branching fractions) from the $D_s^+ \rightarrow \phi \pi^+$, $\phi \rightarrow K^+ K^-$ branching fraction obtained from the Dalitz-plot analysis of $D_s^+ \rightarrow K^+ K^- \pi^+$. That is, the ratio of these two branching fractions is not exactly the $\phi \rightarrow K^+ K^-$ branching fraction 0.491.
 - [g] This comes from a model-independent and a K -matrix parametrization of the $\pi^+ \pi^-$ S -wave and is a sum over several f_0 mesons.
 - [h] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.
 - [i] This is *not* a test for the $\Delta C=1$ weak neutral current, but leads to the $\pi^+ \ell^+ \ell^-$ final state.
 - [j] The value is for the sum of the charge states or particle/antiparticle states indicated.
-

CONSTRAINED FIT INFORMATION

An overall fit to 16 branching ratios uses 17 measurements and one constraint to determine 12 parameters. The overall fit has a $\chi^2 = 2.4$ for 6 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_{25}	16										
x_{26}	12	2									
x_{30}	0	0	0								
x_{31}	0	0	0	76							
x_{41}	0	0	0	42	48						
x_{43}	0	0	0	51	59	32					
x_{54}	0	0	0	59	74	37	45				
x_{64}	0	0	0	67	51	29	35	40			
x_{65}	0	0	0	11	8	5	6	6	16		
x_{83}	0	0	0	37	45	22	28	33	25	4	
x_{114}	-21	-31	-25	-70	-77	-75	-56	-61	-56	-15	
	x_{23}	x_{25}	x_{26}	x_{30}	x_{31}	x_{41}	x_{43}	x_{54}	x_{64}	x_{65}	
x_{114}	-39										
	x_{83}										

A REVIEW GOES HERE – Check our WWW List of Reviews

D_s^+ BRANCHING RATIOS

A number of older, now obsolete results have been omitted. They may be found in earlier editions.

———— Inclusive modes ————

$\Gamma(e^+ \text{ semileptonic}) / \Gamma_{\text{total}}$

Γ_1 / Γ

This is the purely e^+ semileptonic branching fraction: the e^+ fraction from τ^+ decays has been subtracted off. The sum of our (non- τ) e^+ exclusive fractions — an $e^+ \nu_e$ with an $\eta, \eta', \phi, K^0, K^{*0}$, or $f_0(980)$ — is $6.90 \pm 0.4\%$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
6.52 ± 0.39 ± 0.15	536 ± 29	³ ASNER	10 CLEO	$e^+ e^-$ at 3774 MeV

³ Using the D_s^+ and D^0 lifetimes, ASNER 10 finds that the ratio of the D_s^+ and D^0 semileptonic widths is $0.828 \pm 0.051 \pm 0.025$.

$\Gamma(\pi^+ \text{ anything})/\Gamma_{\text{total}}$ **Γ_2/Γ**

Events with two π^+ 's count twice, etc. But π^+ 's from $K_S^0 \rightarrow \pi^+ \pi^-$ are not included.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
119.3±1.2±0.7	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(\pi^- \text{ anything})/\Gamma_{\text{total}}$ **Γ_3/Γ**

Events with two π^- 's count twice, etc. But π^- 's from $K_S^0 \rightarrow \pi^+ \pi^-$ are not included.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
43.2±0.9±0.3	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(\pi^0 \text{ anything})/\Gamma_{\text{total}}$ **Γ_4/Γ**

Events with two π^0 's count twice, etc. But π^0 's from $K_S^0 \rightarrow 2\pi^0$ are not included.

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
123.4±3.8±5.3	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(K^- \text{ anything})/\Gamma_{\text{total}}$ **Γ_5/Γ**

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
18.7±0.5±0.2	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(K^+ \text{ anything})/\Gamma_{\text{total}}$ **Γ_6/Γ**

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
28.9±0.6±0.3	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(K_S^0 \text{ anything})/\Gamma_{\text{total}}$ **Γ_7/Γ**

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
19.0±1.0±0.4	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(\eta \text{ anything})/\Gamma_{\text{total}}$ **Γ_8/Γ**

This ratio includes η particles from η' decays.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
29.9±2.2±1.7		DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

23.5±3.1±2.0	674 ± 91	HUANG	06B	CLEO See DOBBS 09
--------------	----------	-------	-----	-------------------

$\Gamma(\omega \text{ anything})/\Gamma_{\text{total}}$ **Γ_9/Γ**

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.1±1.4±0.3	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(\eta' \text{ anything})/\Gamma_{\text{total}}$ **Γ_{10}/Γ**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
11.7±1.7±0.7		DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.7±1.9±0.8	68 ± 15	HUANG	06B	CLEO See DOBBS 09
-------------	---------	-------	-----	-------------------

$\Gamma(f_0(980) \text{ anything}, f_0 \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.3	90	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$ Γ_{12}/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$15.7 \pm 0.8 \pm 0.6$		DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$16.1 \pm 1.2 \pm 1.1$	398 ± 27	HUANG	06B	CLEO See DOBBS 09
------------------------	--------------	-------	-----	-------------------

$\Gamma(K^+ K^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{13}/Γ

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$15.8 \pm 0.6 \pm 0.3$	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(K_S^0 K^+ \text{ anything})/\Gamma_{\text{total}}$ Γ_{14}/Γ

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.8 \pm 0.5 \pm 0.1$	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(K_S^0 K^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{15}/Γ

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.9 \pm 0.4 \pm 0.1$	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(2K_S^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{16}/Γ

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.7 \pm 0.3 \pm 0.1$	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(2K^+ \text{ anything})/\Gamma_{\text{total}}$ Γ_{17}/Γ

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.26	90	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(2K^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{18}/Γ

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.06	90	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

———— **Leptonic and semileptonic modes** ————

A REVIEW GOES HERE – Check our WWW List of Reviews

$\Gamma(e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{19}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.2 \times 10^{-4}$	90	ALEXANDER	09	CLEO $e^+ e^-$ at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.3 \times 10^{-4}$	90	DEL-AMO-SA..10J	BABR	$e^+ e^-$, 10.58 GeV
$<1.3 \times 10^{-4}$	90	PEDLAR	07A	CLEO See ALEXANDER 09

$\Gamma(\mu^+ \nu_\mu)/\Gamma_{\text{total}}$ Γ_{20}/Γ

See the note on "Decay Constants of Charged Pseudoscalar Mesons" above.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
5.90±0.33 OUR AVERAGE				
6.02±0.38±0.34	275 ± 17	⁴ DEL-AMO-SA..10J	BABR	$e^+ e^-$, 10.58 GeV
5.65±0.45±0.17	235 ± 14	ALEXANDER 09	CLEO	$e^+ e^-$ at 4170 MeV
6.44±0.76±0.57	169 ± 18	⁵ WIDHALM 08	BELL	$e^+ e^- \approx \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5.94±0.66±0.31	88	⁶ PEDLAR	07A	CLEO See ALEXANDER 09
6.8 ± 1.1 ± 1.8	553	⁷ HEISTER	02I	ALEP Z decays

⁴ DEL-AMO-SANCHEZ 10J uses $\mu^+ \nu_\mu$ and $\tau^+ \nu_\tau$ events together to get $f_{D_s} = (258.6 \pm 6.4 \pm 7.5)$ MeV.

⁵ WIDHALM 08 gets $f_{D_s} = (275 \pm 16 \pm 12)$ MeV from the branching fraction.

⁶ PEDLAR 07A also fits μ^+ and τ^+ events together and gets an effective $\mu^+ \nu_\mu$ branching fraction of $(6.38 \pm 0.59 \pm 0.33) \times 10^{-3}$

⁷ This HEISTER 02I result is not actually an independent measurement of the absolute $\mu^+ \nu_\mu$ branching fraction, but is in fact based on our $\phi\pi^+$ branching fraction of $3.6 \pm 0.9\%$, so it cannot be included in our overall fit. HEISTER 02I combines its $D_s^+ \rightarrow \tau^+ \nu_\tau$ and $\mu^+ \nu_\mu$ branching fractions to get $f_{D_s} = (285 \pm 19 \pm 40)$ MeV.

$\Gamma(\mu^+ \nu_\mu)/\Gamma(\phi\pi^+)$ Γ_{20}/Γ_{32}

See the note on "Decay Constants of Charged Pseudoscalar Mesons" above.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.143±0.018±0.006	489 ± 55	⁸ AUBERT	07V	BABR $e^+ e^- \approx \Upsilon(4S)$
0.23 ± 0.06 ± 0.04	18	⁹ ALEXANDROV 00	BEAT	π^- nucleus, 350 GeV
0.173±0.023±0.035	182	¹⁰ CHADHA	98	CLE2 $e^+ e^- \approx \Upsilon(4S)$
0.245±0.052±0.074	39	¹¹ ACOSTA	94	CLE2 See CHADHA 98

⁸ AUBERT 07V gets $f_{D_s^+} = (283 \pm 17 \pm 16)$ MeV, using $\Gamma(D_s^+ \rightarrow \phi\pi^+)/\Gamma(\text{total}) = (4.71 \pm 0.46)\%$.

⁹ ALEXANDROV 00 uses $f_D^2/f_{D_s}^2 = 0.82 \pm 0.09$ from a lattice-gauge-theory calculation to get the relative numbers of $D^+ \rightarrow \mu^+ \nu_\mu$ and $D_s^+ \rightarrow \mu^+ \nu_\mu$ events. The present result leads to $f_{D_s} = (323 \pm 44 \pm 36)$ MeV.

¹⁰ CHADHA 98 obtains $f_{D_s} = (280 \pm 19 \pm 28 \pm 34)$ MeV from this measurement, using $\Gamma(D_s^+ \rightarrow \phi\pi^+)/\Gamma(\text{total}) = 0.036 \pm 0.009$.

¹¹ ACOSTA 94 obtains $f_{D_s} = (344 \pm 37 \pm 52 \pm 42)$ MeV from this measurement, using $\Gamma(D_s^+ \rightarrow \phi\pi^+)/\Gamma(\text{total}) = 0.037 \pm 0.009$.

$\Gamma(\tau^+ \nu_\tau)/\Gamma_{\text{total}}$ Γ_{21}/Γ

See the note on "Decay Constants of Charged Pseudoscalar Mesons" above.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
5.43±0.31 OUR AVERAGE				
5.00±0.35±0.49	748 ± 53	¹² DEL-AMO-SA..10J	BABR	$e^- \bar{\nu}_e \nu_\tau, \mu^- \bar{\nu}_\mu \nu_\tau$
6.42±0.81±0.18	126 ± 16	¹³ ALEXANDER 09	CLEO	$\tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$
5.52±0.57±0.21	155 ± 17	¹³ NAIK	09A	CLEO $\tau^+ \rightarrow \rho^+ \bar{\nu}_\tau$
5.30±0.47±0.22	181 ± 16	¹³ ONYISI	09	CLEO $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.17±0.71±0.34	102	¹⁴ ECKLUND	08	CLEO	See ONYISI 09
8.0 ±1.3 ±0.4	47	¹⁴ PEDLAR	07A	CLEO	See ALEXANDER 09
5.79±0.77±1.84	881	¹⁵ HEISTER	02I	ALEP	Z decays
7.0 ±2.1 ±2.0	22	¹⁶ ABBIENDI	01L	OPAL	$D_s^{*+} \rightarrow \gamma D_s^+$ from Z's
7.4 ±2.8 ±2.4	16	¹⁷ ACCIARRI	97F	L3	$D_s^{*+} \rightarrow \gamma D_s^+$ from Z's

¹² DEL-AMO-SANCHEZ 10J uses $\mu^+ \nu_\mu$ and $\tau^+ \nu_\tau$ events together to get $f_{D_s} = (258.6 \pm 6.4 \pm 7.5)$ MeV.

¹³ ALEXANDER 09, NAIK 09A, and ONYISI 09 use different τ decay modes and are independent. The three papers combined give $f_{D_s} = (259.7 \pm 7.8 \pm 3.4)$ MeV.

¹⁴ ECKLUND 08 and PEDLAR 07A are independent: ECKLUND 08 uses $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$ events, PEDLAR 07A uses $\tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$ events.

¹⁵ HEISTER 02I combines its $D_s^+ \rightarrow \tau^+ \nu_\tau$ and $\mu^+ \nu_\mu$ branching fractions to get $f_{D_s} = (285 \pm 19 \pm 40)$ MeV.

¹⁶ This ABBIENDI 01L value gives a decay constant f_{D_s} of $(286 \pm 44 \pm 41)$ MeV.

¹⁷ The second ACCIARRI 97F error here combines in quadrature systematic (0.016) and normalization (0.018) errors. The branching fraction gives $f_{D_s} = (309 \pm 58 \pm 33 \pm 38)$ MeV.

$\Gamma(\tau^+ \nu_\tau) / \Gamma(\mu^+ \nu_\mu)$

$\Gamma_{21} / \Gamma_{20}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

11.0±1.4±0.6	102	¹⁸ ECKLUND	08	CLEO	See ONYISI 09
--------------	-----	-----------------------	----	------	---------------

¹⁸ This ECKLUND 08 value also uses results from PEDLAR 07A, and it is not independent of other results in these Listings. Combined with earlier CLEO results, the decay constant f_{D_s} is $274 \pm 10 \pm 5$ MeV.

$\Gamma(K^+ K^- e^+ \nu_e) / \Gamma(K^+ K^- \pi^+)$

$\Gamma_{22} / \Gamma_{31}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.558±0.007±0.016	¹⁹ AUBERT	08AN	BABR	$e^+ e^-$ at $\Upsilon(4S)$
-------------------	----------------------	------	------	-----------------------------

¹⁹ This AUBERT 08AN ratio is only for the $K^+ K^-$ mass in the range 1.01–to–1.03 GeV in the numerator and 1.0095–to–1.0295 GeV in the denominator.

$\Gamma(\phi e^+ \nu_e) / \Gamma_{\text{total}}$

Γ_{23} / Γ

See the end of the D_s^+ Listings for measurements of $D_s^+ \rightarrow \phi e^+ \nu_e$ form factors. Unseen decay modes of the ϕ are included.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

2.49±0.14 OUR FIT

2.54±0.14 OUR AVERAGE

2.36±0.23±0.13	106 ± 10	ECKLUND	09	CLEO	$e^+ e^-$ at 4170 MeV
----------------	----------	---------	----	------	-----------------------

2.61±0.03±0.17	(25 ± 0.5)k	AUBERT	08AN	BABR	$e^+ e^-$ at $\Upsilon(4S)$
----------------	-------------	--------	------	------	-----------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.29±0.37±0.11	45	YELTON	09	CLEO	See ECKLUND 09
----------------	----	--------	----	------	----------------

$\Gamma(\phi e^+ \nu_e)/\Gamma(\phi \pi^+)$

Γ_{23}/Γ_{32}

As noted in the comment column, most of these measurements use $\phi \mu^+ \nu_\mu$ events in addition to or instead of $\phi e^+ \nu_e$ events.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.540 \pm 0.033 \pm 0.048$	793	LINK	02J FOCS	Uses $\phi \mu^+ \nu_\mu$
$0.54 \pm 0.05 \pm 0.04$	367	BUTLER	94 CLE2	Uses $\phi e^+ \nu_e$ and $\phi \mu^+ \nu_\mu$
$0.58 \pm 0.17 \pm 0.07$	97	FRABETTI	93G E687	Uses $\phi \mu^+ \nu_\mu$
$0.57 \pm 0.15 \pm 0.15$	104	ALBRECHT	91 ARG	Uses $\phi e^+ \nu_e$
$0.49 \pm 0.10 \begin{smallmatrix} +0.10 \\ -0.14 \end{smallmatrix}$	54	ALEXANDER	90B CLEO	Uses $\phi e^+ \nu_e$ and $\phi \mu^+ \nu_\mu$

$\Gamma(\eta e^+ \nu_e)/\Gamma_{\text{total}}$

Γ_{25}/Γ

Unseen decay modes of the η are included.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.67 ± 0.29 OUR FIT	Error includes scale factor of 1.1.			
$2.48 \pm 0.29 \pm 0.13$	82	YELTON	09 CLEO	$e^+ e^-$ at 4170 MeV

$\Gamma(\eta e^+ \nu_e)/\Gamma(\phi e^+ \nu_e)$

Γ_{25}/Γ_{23}

Unseen decay modes of the η and the ϕ are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.07 ± 0.12 OUR FIT	Error includes scale factor of 1.1.			
$1.24 \pm 0.12 \pm 0.15$	440	²⁰ BRANDENB...	95 CLE2	$e^+ e^- \approx \gamma(4S)$

²⁰ BRANDENBURG 95 uses both e^+ and μ^+ events and makes a phase-space adjustment to use the μ^+ events as e^+ events.

$\Gamma(\eta'(958) e^+ \nu_e)/\Gamma_{\text{total}}$

Γ_{26}/Γ

Unseen decay modes of the $\eta'(958)$ are included.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.99 ± 0.23 OUR FIT				
$0.91 \pm 0.33 \pm 0.05$	7.5	YELTON	09 CLEO	$e^+ e^-$ at 4170 MeV

$\Gamma(\eta'(958) e^+ \nu_e)/\Gamma(\phi e^+ \nu_e)$

Γ_{26}/Γ_{23}

Unseen decay modes of the resonances are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.40 ± 0.09 OUR FIT				
$0.43 \pm 0.11 \pm 0.07$	29	²¹ BRANDENB...	95 CLE2	$e^+ e^- \approx \gamma(4S)$

²¹ BRANDENBURG 95 uses both e^+ and μ^+ events and makes a phase-space adjustment to use the μ^+ events as e^+ events.

$[\Gamma(\eta e^+ \nu_e) + \Gamma(\eta'(958) e^+ \nu_e)]/\Gamma(\phi e^+ \nu_e)$

$\Gamma_{24}/\Gamma_{23} = (\Gamma_{25} + \Gamma_{26})/\Gamma_{23}$

Unseen decay modes of the resonances are included.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.67 \pm 0.17 \pm 0.17$	²² BRANDENB...	95 CLE2	$e^+ e^- \approx \gamma(4S)$

²² This BRANDENBURG 95 data is redundant with data in previous blocks.

$\Gamma(K^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.37±0.10±0.02	14	YELTON	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(K^*(892)^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{28}/Γ

Unseen decay modes of the $K^*(892)^0$ are included.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.18±0.07±0.01	7.5	YELTON	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(f_0(980) e^+ \nu_e, f_0 \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.20 ±0.03 ±0.01	44 ± 7	ECKLUND	09	CLEO $e^+ e^-$ at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.13 ±0.04 ±0.01	13	YELTON	09	CLEO See ECKLUND 09
------------------	----	--------	----	---------------------

————— Hadronic modes with a $K\bar{K}$ pair. —————

$\Gamma(K^+ K_S^0)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
1.48±0.08 OUR FIT			
1.49±0.07±0.05	²³ ALEXANDER 08	CLEO	$e^+ e^-$ at 4.17 GeV

²³ ALEXANDER 08 uses single- and double-tagged events in an overall fit. The correlation matrix for the branching fractions is used in the fit.

$\Gamma(K^+ K^- \pi^+)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
5.49±0.27 OUR FIT			
5.50±0.23±0.16	²⁴ ALEXANDER 08	CLEO	$e^+ e^-$ at 4.17 GeV

²⁴ ALEXANDER 08 uses single- and double-tagged events in an overall fit. The correlation matrix for the branching fractions is used in the fit.

$\Gamma(\phi\pi^+)/\Gamma_{\text{total}}$ Γ_{32}/Γ

The results here are model-independent. For earlier, model-dependent results, see our PDG 06 edition. We decouple the $D_S^+ \rightarrow \phi\pi^+$ branching fraction obtained from mass projections (and used to get some of the other branching fractions) from the $D_S^+ \rightarrow \phi\pi^+, \phi \rightarrow K^+ K^-$ branching fraction obtained from the Dalitz-plot analysis of $D_S^+ \rightarrow K^+ K^- \pi^+$. That is, the ratio of these two branching fractions is not exactly the $\phi \rightarrow K^+ K^-$ branching fraction 0.491.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.5 ±0.4 OUR AVERAGE				
4.62±0.36±0.51		²⁵ AUBERT	06N BABR	$e^+ e^-$ at $\Upsilon(4S)$
4.81±0.52±0.38	212 ± 19	²⁶ AUBERT	05V BABR	$e^+ e^- \approx \Upsilon(4S)$
3.59±0.77±0.48		²⁷ ARTUSO	96 CLE2	$e^+ e^-$ at $\Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.9 ^{+5.1} _{-1.9} ^{+1.8} _{-1.1}		²⁸ BAI	95C BES	$e^+ e^-$ 4.03 GeV

- ²⁵ This AUBERT 06N measurement uses $\bar{B}^0 \rightarrow D_S^{(*)-} D^{(*)+}$ and $B^- \rightarrow D_S^{(*)-} D^{(*)0}$ decays, including some from other papers. However, the result is independent of AUBERT 05V.
- ²⁶ AUBERT 05V uses the ratio of $B^0 \rightarrow D^{*-} D_S^{*+}$ events seen in two different ways, in both of which the $D^{*-} \rightarrow \bar{D}^0 \pi^-$ decay is fully reconstructed: (1) The $D_S^{*+} \rightarrow D_S^+ \gamma$, $D_S^+ \rightarrow \phi \pi^+$ decay is fully reconstructed. (2) The number of events in the D_S^+ peak in the missing mass spectrum against the $D^{*-} \gamma$ is measured.
- ²⁷ ARTUSO 96 uses partially reconstructed $\bar{B}^0 \rightarrow D^{*+} D_S^{*-}$ decays to get a model-independent value for $\Gamma(D_S^- \rightarrow \phi \pi^-) / \Gamma(D^0 \rightarrow K^- \pi^+)$ of $0.92 \pm 0.20 \pm 0.11$.
- ²⁸ BAI 95C uses $e^+ e^- \rightarrow D_S^+ D_S^-$ events in which one or both of the D_S^\pm are observed to obtain the first model-independent measurement of the $D_S^+ \rightarrow \phi \pi^+$ branching fraction, without assumptions about $\sigma(D_S^\pm)$. However, with only two “doubly-tagged” events, the statistical error is very large.

$\Gamma(\phi \pi^+, \phi \rightarrow K^+ K^-) / \Gamma(K^+ K^- \pi^+)$ $\Gamma_{33} / \Gamma_{31}$

This is the “fit fraction” from the Dalitz-plot analysis. We decouple the $D_S^+ \rightarrow \phi \pi^+$ branching fraction obtained from mass projections (and used to get some of the other branching fractions) from the $D_S^+ \rightarrow \phi \pi^+$, $\phi \rightarrow K^+ K^-$ branching fraction obtained from the Dalitz-plot analysis of $D_S^+ \rightarrow K^+ K^- \pi^+$. That is, the ratio of these two branching fractions is not exactly the $\phi \rightarrow K^+ K^-$ branching fraction 0.491.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.422 ± 0.016 ± 0.003	MITCHELL 09A	CLEO	Dalitz fit, 12k evts
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.396 ± 0.033 ± 0.047	FRABETTI 95B	E687	Dalitz fit, 701 evts

$\Gamma(K^+ \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+) / \Gamma(K^+ K^- \pi^+)$ $\Gamma_{34} / \Gamma_{31}$

This is the “fit fraction” from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.474 ± 0.015 ± 0.004	MITCHELL 09A	CLEO	Dalitz fit, 12k evts
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.478 ± 0.046 ± 0.040	FRABETTI 95B	E687	Dalitz fit, 701 evts

$\Gamma(f_0(980) \pi^+, f_0 \rightarrow K^+ K^-) / \Gamma(K^+ K^- \pi^+)$ $\Gamma_{35} / \Gamma_{31}$

This is the “fit fraction” from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.282 ± 0.019 ± 0.018	MITCHELL 09A	CLEO	Dalitz fit, 12k evts
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.11 ± 0.035 ± 0.026	FRABETTI 95B	E687	Dalitz fit, 701 evts

$\Gamma(f_0(1370) \pi^+, f_0 \rightarrow K^+ K^-) / \Gamma(K^+ K^- \pi^+)$ $\Gamma_{36} / \Gamma_{31}$

This is the “fit fraction” from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.043 ± 0.006 ± 0.005	MITCHELL 09A	CLEO	Dalitz fit, 12k evts

$\Gamma(f_0(1710)\pi^+, f_0 \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+)$ Γ_{37}/Γ_{31}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.034±0.005±0.003	MITCHELL 09A	CLEO	Dalitz fit, 12k evts
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.034±0.023±0.035	FRABETTI 95B	E687	Dalitz fit, 701 evts

$\Gamma(K^+\bar{K}_0^*(1430)^0, \bar{K}_0^* \rightarrow K^-\pi^+)/\Gamma(K^+K^-\pi^+)$ Γ_{38}/Γ_{31}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.039±0.005±0.005	MITCHELL 09A	CLEO	Dalitz fit, 12k evts
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.093±0.032±0.032	FRABETTI 95B	E687	Dalitz fit, 701 evts

$\Gamma(K^*(892)^+\bar{K}^0)/\Gamma(\phi\pi^+)$ Γ_{40}/Γ_{32}

Unseen decay modes of the resonances are included.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.20±0.21±0.13	CHEN 89	CLEO	e^+e^- 10 GeV

$\Gamma(K^+K^-\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{41}/Γ

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.6 ±0.5 OUR FIT			
5.65±0.29±0.40	²⁹ ALEXANDER 08	CLEO	e^+e^- at 4.17 GeV

²⁹ALEXANDER 08 uses single- and double-tagged events in an overall fit. The correlation matrix for the branching fractions is used in the fit.

$\Gamma(\phi\rho^+)/\Gamma(\phi\pi^+)$ Γ_{42}/Γ_{32}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.86±0.26^{+0.29}_{-0.40}	253	AVERY 92	CLE2	$e^+e^- \simeq 10.5$ GeV

$\Gamma(K_S^0K^-2\pi^+)/\Gamma_{\text{total}}$ Γ_{43}/Γ

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.64±0.12 OUR FIT			
1.64±0.10±0.07	³⁰ ALEXANDER 08	CLEO	e^+e^- at 4.17 GeV

³⁰ALEXANDER 08 uses single- and double-tagged events in an overall fit. The correlation matrix for the branching fractions is used in the fit.

$\Gamma(K^*(892)^+\bar{K}^*(892)^0)/\Gamma(\phi\pi^+)$ Γ_{44}/Γ_{32}

Unseen decay modes of the resonances are included.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.6±0.4±0.4	ALBRECHT 92B	ARG	$e^+e^- \simeq 10.4$ GeV

$\Gamma(K^+K_S^0\pi^+\pi^-)/\Gamma(K_S^0K^-2\pi^+)$ Γ_{45}/Γ_{43}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.586±0.052±0.043	476	LINK 01C	FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^+ K^- 2\pi^+ \pi^-)/\Gamma(K^+ K^- \pi^+)$ Γ_{46}/Γ_{31}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.160±0.027 OUR AVERAGE				
0.150±0.019±0.025	240	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
0.188±0.036±0.040	75	FRABETTI	97C E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV

$\Gamma(\phi 2\pi^+ \pi^-)/\Gamma(\phi \pi^+)$ Γ_{47}/Γ_{32}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.269±0.027 OUR AVERAGE				
0.249±0.024±0.021	136	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
0.28 ±0.06 ±0.01	40	FRABETTI	97C E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV
0.58 ±0.21 ±0.10	21	FRABETTI	92 E687	γ Be
0.42 ±0.13 ±0.07	19	ANJOS	88 E691	Photoproduction
1.11 ±0.37 ±0.28	62	ALBRECHT	85D ARG	$e^+ e^-$ 10 GeV

$\Gamma(K^+ K^- \rho^0 \pi^+ \text{non-}\phi)/\Gamma(K^+ K^- 2\pi^+ \pi^-)$ Γ_{48}/Γ_{46}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.03	90	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(\phi \rho^0 \pi^+, \phi \rightarrow K^+ K^-)/\Gamma(K^+ K^- 2\pi^+ \pi^-)$ Γ_{49}/Γ_{46}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.75±0.06±0.04	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(\phi a_1(1260)^+, \phi \rightarrow K^+ K^-, a_1^+ \rightarrow \rho^0 \pi^+)/\Gamma(K^+ K^- \pi^+)$ Γ_{50}/Γ_{31}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.137±0.019±0.011	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^+ K^- 2\pi^+ \pi^- \text{nonresonant})/\Gamma(K^+ K^- 2\pi^+ \pi^-)$ Γ_{51}/Γ_{46}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.10±0.06±0.05	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(2K_S^0 2\pi^+ \pi^-)/\Gamma(K_S^0 K^- 2\pi^+)$ Γ_{52}/Γ_{43}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.051±0.015±0.015	37 ± 10	LINK	04D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

———— Pionic modes ————

$\Gamma(\pi^+ \pi^0)/\Gamma(K^+ K_S^0)$ Γ_{53}/Γ_{30}

<u>VALUE (units 10⁻²)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.3	90	MENDEZ	10 CLEO	$e^+ e^-$ at 4170 MeV
• • •				We do not use the following data for averages, fits, limits, etc. • • •
<4.1	90	ADAMS	07A CLEO	See MENDEZ 10

$\Gamma(2\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{54}/Γ

<u>VALUE (units 10⁻²)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.10±0.06 OUR FIT			
1.11±0.07±0.04	³¹ ALEXANDER 08	CLEO	$e^+ e^-$ at 4.17 GeV

³¹ ALEXANDER 08 uses single- and double-tagged events in an overall fit. The correlation matrix for the branching fractions is used in the fit.

$\Gamma(2\pi^+\pi^-)/\Gamma(K^+K^-\pi^+)$ Γ_{54}/Γ_{31}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.200±0.008 OUR FIT				
0.199±0.004±0.009	≈ 10.5k	AUBERT	090 BABR	$e^+e^- \approx 10.6$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.265±0.041±0.031	98	FRABETTI	97D E687	γ Be ≈ 200 GeV

$\Gamma(\rho^0\pi^+)/\Gamma(2\pi^+\pi^-)$ Γ_{55}/Γ_{54}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.018±0.005±0.010		AUBERT	090 BABR	Dalitz fit, ≈ 10.5k evts
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen		LINK	04 FOCS	Dalitz fit, 1475 ± 50 evts
0.058±0.023±0.037		AITALA	01A E791	Dalitz fit, 848 evts
<0.073	90	FRABETTI	97D E687	γ Be ≈ 200 GeV

$\Gamma(\pi^+(\pi^+\pi^-)_{S\text{-wave}})/\Gamma(2\pi^+\pi^-)$ Γ_{56}/Γ_{54}

This is the “fit fraction” from the Dalitz-plot analysis. See also KLEMPT 08, which uses 568 $D_s^+ \rightarrow 3\pi$ decays (over 280 background events) from FNAL E791 to study various parametrizations of the decay amplitudes. The emphasis there is more on S -wave $\pi\pi$ decay products — 20 different solutions are given — than on D_s^+ fit fractions.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.833 ±0.020 OUR AVERAGE			

0.830 ±0.009 ±0.019	³² AUBERT	090 BABR	Dalitz fit, ≈ 10.5k evts
0.8704±0.0560±0.0438	³³ LINK	04 FOCS	Dalitz fit, 1475 ± 50 evts
³² AUBERT 090 gives the amplitude and phase of the $\pi^+\pi^-$ S -wave in 29 $\pi^+\pi^-$ invariant-mass bins.			
³³ LINK 04 borrows a K-matrix parametrization from ANISOVICH 03 of the full $\pi-\pi$ S -wave isoscalar scattering amplitude to describe the $\pi^+\pi^-$ S -wave component of the $\pi^+\pi^+\pi^-$ state. The fit fraction given above is a sum over five f_0 mesons, the $f_0(980)$, $f_0(1300)$, $f_0(1200-1600)$, $f_0(1500)$, and $f_0(1750)$. See LINK 04 for details and discussion.			

$\Gamma(f_0(980)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{57}/Γ_{54}

This is the “fit fraction” from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.565±0.043±0.047	AITALA	01A E791	Dalitz fit, 848 evts
1.074±0.140±0.043	FRABETTI	97D E687	γ Be ≈ 200 GeV

$\Gamma(f_0(1370)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{58}/Γ_{54}

This is the “fit fraction” from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.324±0.077±0.017	AITALA	01A E791	Dalitz fit, 848 evts

$\Gamma(f_0(1500)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{59}/Γ_{54}

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.274 \pm 0.114 \pm 0.019$	³⁴ FRABETTI	97D E687	γ Be \approx 200 GeV
-----------------------------	------------------------	----------	-------------------------------

³⁴ FRABETTI 97D calls this mode $S(1475)\pi^+$, but finds the mass and width of this $S(1475)$ to be in excellent agreement with those of the $f_0(1500)$.

$\Gamma(f_2(1270)\pi^+, f_2 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{60}/Γ_{54}

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.101 ± 0.018 OUR AVERAGE

$0.101 \pm 0.015 \pm 0.011$	AUBERT	09O BABR	Dalitz fit, \approx 10.5k evts
-----------------------------	--------	----------	----------------------------------

$0.0974 \pm 0.0449 \pm 0.0294$	LINK	04 FOCS	Dalitz fit, 1475 ± 50 evts
--------------------------------	------	---------	--------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.197 \pm 0.033 \pm 0.006$	AITALA	01A E791	Dalitz fit, 848 evts
-----------------------------	--------	----------	----------------------

$0.123 \pm 0.056 \pm 0.018$	FRABETTI	97D E687	γ Be \approx 200 GeV
-----------------------------	----------	----------	-------------------------------

$\Gamma(\rho(1450)^0\pi^+, \rho^0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{61}/Γ_{54}

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.027 ± 0.018 OUR AVERAGE

$0.023 \pm 0.008 \pm 0.017$	AUBERT	09O BABR	Dalitz fit, \approx 10.5k evts
-----------------------------	--------	----------	----------------------------------

$0.0656 \pm 0.0343 \pm 0.0440$	LINK	04 FOCS	Dalitz fit, 1475 ± 50 evts
--------------------------------	------	---------	--------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.044 \pm 0.021 \pm 0.002$	AITALA	01A E791	Dalitz fit, 848 evts
-----------------------------	--------	----------	----------------------

$\Gamma(\pi^+2\pi^0)/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$0.65 \pm 0.13 \pm 0.03$	72 ± 16	NAIK	09A CLEO	e^+e^- at 4170 MeV
--	-------------	------	----------	----------------------

$\Gamma(2\pi^+\pi^-\pi^0)/\Gamma(\phi\pi^+)$ Γ_{63}/Γ_{32}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.3	90	ANJOS	89E E691	Photoproduction
--------	----	-------	----------	-----------------

$\Gamma(\eta\pi^+)/\Gamma_{\text{total}}$ Γ_{64}/Γ

Unseen decay modes of the η are included.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.58 \pm 0.11 \pm 0.18$	³⁵ ALEXANDER	08 CLEO	See MENDEZ 10
--------------------------	-------------------------	---------	---------------

³⁵ ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(\eta\pi^+)/\Gamma(K^+K_S^0)$ Γ_{64}/Γ_{30}

Unseen decay modes of the η are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

1.23 ± 0.08 OUR FIT

$1.236 \pm 0.043 \pm 0.063$	2587 ± 89	MENDEZ	10 CLEO	e^+e^- at 4170 MeV
---	---------------	--------	---------	----------------------

$\Gamma(\eta\pi^+)/\Gamma(\phi\pi^+)$ Γ_{64}/Γ_{32}

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.48 \pm 0.03 \pm 0.04$	920	JESSOP	98	CLE2 $e^+e^- \approx \Upsilon(4S)$
$0.54 \pm 0.09 \pm 0.06$	165	ALEXANDER	92	CLE2 See JESSOP 98

$\Gamma(\omega\pi^+)/\Gamma_{\text{total}}$ Γ_{65}/Γ

Unseen decay modes of the ω are included.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.25 ± 0.07 OUR FIT				
$0.21 \pm 0.09 \pm 0.01$	6 ± 2.4	GE	09A	CLEO e^+e^- at 4170 MeV

$\Gamma(\omega\pi^+)/\Gamma(\eta\pi^+)$ Γ_{65}/Γ_{64}

Unseen decay modes of the resonances are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.14 ± 0.04 OUR FIT			
$0.16 \pm 0.04 \pm 0.03$	BALEST	97	CLE2 $e^+e^- \approx \Upsilon(4S)$

$\Gamma(3\pi^+2\pi^-)/\Gamma(K^+K^-\pi^+)$ Γ_{66}/Γ_{31}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.146 ± 0.014 OUR AVERAGE				
$0.145 \pm 0.011 \pm 0.010$	671	LINK	03D	FOCS $\gamma A, \bar{E}_\gamma \approx 180$ GeV
$0.158 \pm 0.042 \pm 0.031$	37	FRABETTI	97C	E687 $\gamma Be, \bar{E}_\gamma \approx 200$ GeV

$\Gamma(\eta\rho^+)/\Gamma_{\text{total}}$ Γ_{68}/Γ

Unseen decay modes of the η are included.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.9 \pm 0.6 \pm 0.5$	328 ± 22	NAIK	09A	CLEO $\eta \rightarrow 2\gamma$

$\Gamma(\eta\rho^+)/\Gamma(\phi\pi^+)$ Γ_{68}/Γ_{32}

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$2.98 \pm 0.20 \pm 0.39$	447	JESSOP	98	CLE2 $e^+e^- \approx \Upsilon(4S)$
$2.86 \pm 0.38^{+0.36}_{-0.38}$	217	AVERY	92	CLE2 See JESSOP 98

$\Gamma(\eta\pi^+\pi^0\text{3-body})/\Gamma(\phi\pi^+)$ Γ_{69}/Γ_{32}

Unseen decay modes of the resonances are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.1				
<0.82	90	JESSOP	98	CLE2 $e^+e^- \approx \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.82	90	³⁶ DAUDI	92	CLE2 See JESSOP 98

³⁶We use the JESSOP 98 limit, even though the DAUDI 92 limit, from the same experiment but with a much smaller data sample, is more restrictive.

$\Gamma(\omega\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{70}/Γ

Unseen decay modes of the ω are included.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.78 \pm 0.65 \pm 0.25$	34 ± 7.9	GE	09A	CLEO e^+e^- at 4170 MeV

$\Gamma(3\pi^+ 2\pi^- \pi^0)/\Gamma_{\text{total}}$					Γ_{71}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
$0.049^{+0.033}_{-0.030}$	BARLAG	92C	ACCM	π^- 230 GeV	

$\Gamma(\omega 2\pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{72}/Γ
Unseen decay modes of the ω are included.					
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.58 \pm 0.45 \pm 0.09$	29 ± 8.2	GE	09A	CLEO	$e^+ e^-$ at 4170 MeV

$\Gamma(\eta'(958)\pi^+)/\Gamma_{\text{total}}$					Γ_{73}/Γ
Unseen decay modes of the $\eta'(958)$ are included.					
VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$3.77 \pm 0.25 \pm 0.30$	³⁷ ALEXANDER 08	CLEO	See MENDEZ 10		
³⁷ ALEXANDER 08 uses single- and double-tagged events in an overall fit.					

$\Gamma(\eta'(958)\pi^+)/\Gamma(K^+ K_S^0)$					Γ_{73}/Γ_{30}
Unseen decay modes of the $\eta'(958)$ are included.					
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.654 \pm 0.088 \pm 0.139$	1436 ± 47	MENDEZ	10	CLEO	$e^+ e^-$ at 4170 MeV

$\Gamma(\eta'(958)\pi^+)/\Gamma(\phi\pi^+)$					Γ_{73}/Γ_{32}
Unseen decay modes of the resonances are included.					
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$1.03 \pm 0.06 \pm 0.07$	537	JESSOP	98	CLE2	$e^+ e^- \approx \Upsilon(4S)$
$1.20 \pm 0.15 \pm 0.11$	281	ALEXANDER	92	CLE2	See JESSOP 98
$2.5 \pm 1.0 \pm 1.5_{-0.4}$	22	ALVAREZ	91	NA14	Photoproduction
$2.5 \pm 0.5 \pm 0.3$	215	ALBRECHT	90D	ARG	$e^+ e^- \approx 10.4$ GeV

$\Gamma(\omega\eta\pi^+)/\Gamma_{\text{total}}$					Γ_{75}/Γ
Unseen decay modes of the ω and η are included.					
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 2.13 \times 10^{-2}$	90	GE	09A	CLEO	$e^+ e^-$ at 4170 MeV

$\Gamma(\eta'(958)\rho^+)/\Gamma(\phi\pi^+)$					Γ_{76}/Γ_{32}
Unseen decay modes of the resonances are included.					
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.78 \pm 0.28 \pm 0.30$	137	JESSOP	98	CLE2	$e^+ e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$3.44 \pm 0.62 \pm 0.44_{-0.46}$	68	AVERY	92	CLE2	See JESSOP 98

$\Gamma(\eta'(958)\pi^+\pi^0\text{-body})/\Gamma(\phi\pi^+)$					Γ_{77}/Γ_{32}
Unseen decay modes of the resonances are included.					
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
< 0.4	90	JESSOP	98	CLE2	$e^+ e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.85	90	DAOUDI	92	CLE2	See JESSOP 98

———— Modes with one or three *K*'s ————

$\Gamma(K^+\pi^0)/\Gamma(K^+K_S^0)$ Γ_{78}/Γ_{30}

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.2±1.4±0.2	202 ± 70	MENDEZ	10	CLEO e^+e^- at 4170 MeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5.5±1.3±0.7	141 ± 34	ADAMS	07A	CLEO See MENDEZ 10

$\Gamma(K_S^0\pi^+)/\Gamma(K^+K_S^0)$ Γ_{79}/Γ_{30}

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.12±0.28 OUR AVERAGE				
8.5 ± 0.7 ± 0.2	393 ± 33	MENDEZ	10	CLEO e^+e^- at 4170 MeV
8.03±0.24±0.19	17.6k±481	WON	09	BELL e^+e^- at $\Upsilon(4S)$
10.4 ± 2.4 ± 1.4	113 ± 26	LINK	08	FOCS γ A, $\bar{E}_\gamma \approx 180$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
8.2 ± 0.9 ± 0.2	206 ± 22	ADAMS	07A	CLEO See MENDEZ 10

$\Gamma(K^+\eta)/\Gamma(K^+K_S^0)$ Γ_{80}/Γ_{30}

Unseen decay modes of the η are included.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
11.8±2.2±0.6	222 ± 41	MENDEZ	10	CLEO e^+e^- at 4170 MeV

$\Gamma(K^+\eta)/\Gamma(\eta\pi^+)$ Γ_{80}/Γ_{64}

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
8.9±1.5±0.4	113 ± 18	ADAMS	07A	CLEO See MENDEZ 10

$\Gamma(K^+\omega)/\Gamma_{\text{total}}$ Γ_{81}/Γ

Unseen decay modes of the ω are included.

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.24	90	GE	09A	CLEO e^+e^- at 4170 MeV

$\Gamma(K^+\eta'(958))/\Gamma(K^+K_S^0)$ Γ_{82}/Γ_{30}

Unseen decay modes of the $\eta'(958)$ are included.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
11.8±3.6±0.7	56 ± 17	MENDEZ	10	CLEO e^+e^- at 4170 MeV

$\Gamma(K^+\eta'(958))/\Gamma(\eta'(958)\pi^+)$ Γ_{82}/Γ_{73}

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.2±1.3±0.3	28 ± 9	ADAMS	07A	CLEO See MENDEZ 10

$\Gamma(K^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{83}/Γ

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.69±0.05 OUR FIT			
0.69±0.05±0.03	³⁸ ALEXANDER	08	CLEO e^+e^- at 4.17 GeV

³⁸ ALEXANDER 08 uses single- and double-tagged events in an overall fit. The correlation matrix for the branching fractions is used in the fit.

$\Gamma(K^+ \pi^+ \pi^-) / \Gamma(K^+ K^- \pi^+)$ $\Gamma_{83} / \Gamma_{31}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.126 ± 0.009 OUR FIT				
0.127 ± 0.007 ± 0.014	567 ± 31	LINK	04F FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^+ \rho^0) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{84} / \Gamma_{83}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.3883 ± 0.0531 ± 0.0261	LINK	04F FOCS	Dalitz fit, 567 evts

$\Gamma(K^+ \rho(1450)^0, \rho^0 \rightarrow \pi^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{85} / \Gamma_{83}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.1062 ± 0.0351 ± 0.0104	LINK	04F FOCS	Dalitz fit, 567 evts

$\Gamma(K^*(892)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{86} / \Gamma_{83}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.2164 ± 0.0321 ± 0.0114	LINK	04F FOCS	Dalitz fit, 567 evts

$\Gamma(K^*(1410)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{87} / \Gamma_{83}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.1882 ± 0.0403 ± 0.0122	LINK	04F FOCS	Dalitz fit, 567 evts

$\Gamma(K^*(1430)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{88} / \Gamma_{83}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.0765 ± 0.0500 ± 0.0170	LINK	04F FOCS	Dalitz fit, 567 evts

$\Gamma(K^+ \pi^+ \pi^- \text{ nonresonant}) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{89} / \Gamma_{83}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.1588 ± 0.0492 ± 0.0153	LINK	04F FOCS	Dalitz fit, 567 evts

$\Gamma(K^0 \pi^+ \pi^0) / \Gamma_{\text{total}}$ Γ_{90} / Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.00 ± 0.18 ± 0.04	44 ± 8	NAIK	09A CLEO	$e^+ e^-$ at 4170 MeV

$\Gamma(K_S^0 2\pi^+ \pi^-) / \Gamma(K_S^0 K^- 2\pi^+)$ $\Gamma_{91} / \Gamma_{43}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.18 ± 0.04 ± 0.05	179 ± 36	LINK	08 FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^+ \omega \pi^0) / \Gamma_{\text{total}}$ Γ_{92} / Γ

Unseen decay modes of the ω are included.

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.82	90	GE	09A CLEO	$e^+ e^-$ at 4170 MeV

$\Gamma(K^+ \omega \pi^+ \pi^-) / \Gamma_{\text{total}}$ **Γ_{93} / Γ**

Unseen decay modes of the ω are included.

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.54	90	GE	09A CLEO	$e^+ e^-$ at 4170 MeV

$\Gamma(K^+ \omega \eta) / \Gamma_{\text{total}}$ **Γ_{94} / Γ**

Unseen decay modes of the ω and η are included.

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.79	90	GE	09A CLEO	$e^+ e^-$ at 4170 MeV

$\Gamma(2K^+ K^-) / \Gamma(K^+ K^- \pi^+)$ **$\Gamma_{95} / \Gamma_{31}$**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.95 \pm 2.12^{+2.24}_{-2.31}$	31	LINK	02I FOCS	γ nucleus, ≈ 180 GeV

$\Gamma(\phi K^+) / \Gamma(\phi \pi^+)$ **$\Gamma_{96} / \Gamma_{32}$**

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.013	90	FRABETTI	95F E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.071	90	ANJOS	92D E691	γ Be, $\bar{E}_\gamma = 145$ GeV
--------	----	-------	----------	---

————— **Doubly Cabibbo-suppressed modes** —————

$\Gamma(2K^+ \pi^-) / \Gamma(K^+ K^- \pi^+)$ **$\Gamma_{97} / \Gamma_{31}$**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.35 ± 0.30 OUR AVERAGE				
$2.29 \pm 0.28 \pm 0.12$	281 ± 34	KO	09 BELL	$e^+ e^-$ at $\Upsilon(4S)$
$5.2 \pm 1.7 \pm 1.1$	27 ± 9	LINK	05K FOCS	<0.78%, CL = 90%

————— **Baryon-antibaryon mode** —————

$\Gamma(p \bar{n}) / \Gamma_{\text{total}}$ **Γ_{98} / Γ**

This is the only baryonic mode allowed kinematically.

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.30 \pm 0.36^{+0.12}_{-0.16}$	13.0 ± 3.6	ATHAR	08 CLEO	$e^+ e^-$, $E_{\text{cm}} \approx 4170$ MeV

————— **Rare or forbidden modes** —————

$\Gamma(\pi^+ e^+ e^-) / \Gamma_{\text{total}}$ **Γ_{99} / Γ**

This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 2.2×10^{-5}	90	³⁹ RUBIN	10 CLEO	$e^+ e^-$ at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 27×10^{-5}	90	AITALA	99G E791	$\pi^- N$ 500 GeV
-----------------------	----	--------	----------	-------------------

³⁹This RUBIN 10 limit is for the $e^+ e^-$ mass in the continuum away from the $\phi(1020)$.
See the next data block.

$\Gamma(\pi^+ \phi, \phi \rightarrow e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{100}/Γ**

This is *not* a test for the $\Delta C = 1$ weak neutral current, but leads to the $\pi^+ e^+ e^-$ final state.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$(6_{-4}^{+8} \pm 1) \times 10^{-6}$	3	RUBIN	10	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(\pi^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{101}/Γ**

This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 2.6 \times 10^{-5}$	90		LINK	03F	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.4 \times 10^{-4}$	90		AITALA	99G	E791 $\pi^- N$ 500 GeV
$< 4.3 \times 10^{-4}$	90	0	KODAMA	95	E653 π^- emulsion 600 GeV

$\Gamma(K^+ e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{102}/Γ**

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 5.2 \times 10^{-5}$	90		RUBIN	10	CLEO $e^+ e^-$ at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.6 \times 10^{-3}$	90		AITALA	99G	E791 $\pi^- N$ 500 GeV
------------------------	----	--	--------	-----	------------------------

$\Gamma(K^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{103}/Γ**

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 3.6 \times 10^{-5}$	90		LINK	03F	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.4 \times 10^{-4}$	90		AITALA	99G	E791 $\pi^- N$ 500 GeV
$< 5.9 \times 10^{-4}$	90	0	KODAMA	95	E653 π^- emulsion 600 GeV

$\Gamma(K^*(892)^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{104}/Γ**

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.4 \times 10^{-3}$	90	0	KODAMA	95	E653 π^- emulsion 600 GeV

$\Gamma(\pi^+ e^\pm \mu^\mp)/\Gamma_{\text{total}}$ **Γ_{105}/Γ**

A test of lepton-family-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 6.1 \times 10^{-4}$	90		AITALA	99G	E791 $\pi^- N$ 500 GeV

$\Gamma(K^+ e^\pm \mu^\mp)/\Gamma_{\text{total}}$ **Γ_{106}/Γ**

A test of lepton-family-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 6.3 \times 10^{-4}$	90		AITALA	99G	E791 $\pi^- N$ 500 GeV

$\Gamma(\pi^- 2e^+)/\Gamma_{\text{total}}$ **Γ_{107}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.8 \times 10^{-5}$	90	RUBIN	10	CLEO $e^+ e^-$ at 4170 MeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 69 \times 10^{-5}$	90	AITALA	99G	E791 $\pi^- N$ 500 GeV

$\Gamma(\pi^- 2\mu^+)/\Gamma_{\text{total}}$ **Γ_{108}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 2.9 \times 10^{-5}$	90		LINK	03F	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 8.2 \times 10^{-5}$	90		AITALA	99G	E791 $\pi^- N$ 500 GeV
$< 4.3 \times 10^{-4}$	90	0	KODAMA	95	E653 π^- emulsion 600 GeV

$\Gamma(\pi^- e^+ \mu^+)/\Gamma_{\text{total}}$ **Γ_{109}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 7.3 \times 10^{-4}$	90	AITALA	99G	E791 $\pi^- N$ 500 GeV

$\Gamma(K^- 2e^+)/\Gamma_{\text{total}}$ **Γ_{110}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.7 \times 10^{-5}$	90	RUBIN	10	CLEO $e^+ e^-$ at 4170 MeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 63 \times 10^{-5}$	90	AITALA	99G	E791 $\pi^- N$ 500 GeV

$\Gamma(K^- 2\mu^+)/\Gamma_{\text{total}}$ **Γ_{111}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.3 \times 10^{-5}$	90		LINK	03F	FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 1.8 \times 10^{-4}$	90		AITALA	99G	E791 $\pi^- N$ 500 GeV
$< 5.9 \times 10^{-4}$	90	0	KODAMA	95	E653 π^- emulsion 600 GeV

$\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$ **Γ_{112}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 6.8 \times 10^{-4}$	90	AITALA	99G	E791 $\pi^- N$ 500 GeV

$\Gamma(K^*(892)^- 2\mu^+)/\Gamma_{\text{total}}$ **Γ_{113}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.4 \times 10^{-3}$	90	0	KODAMA	95	E653 π^- emulsion 600 GeV

$D_s^+ - D_s^-$ CP-VIOLATING DECAY-RATE ASYMMETRIES

This is the difference of the D_s^+ and D_s^- partial widths divided by the sum of the widths.

$A_{CP}(\mu^\pm \nu)$ in $D_s^+ \rightarrow \mu^+ \nu, D_s^- \rightarrow \mu^- \bar{\nu}_\mu$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
+4.8±6.1	ALEXANDER 09	CLEO	$e^+ e^-$ at 4170 MeV

$A_{CP}(K^\pm K_S^0)$ in $D_s^\pm \rightarrow K^\pm K_S^0$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.3 ± 0.4 OUR AVERAGE				
+0.12±0.36±0.22		KO 10	BELL	$e^+ e^- \approx \Upsilon(4S)$
+4.7 ± 1.8 ± 0.9	4.0k	MENDEZ 10	CLEO	$e^+ e^-$ at 4170 MeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+4.9 ± 2.1 ± 0.9		ALEXANDER 08	CLEO	See MENDEZ 10

$A_{CP}(K^+ K^- \pi^\pm)$ in $D_s^\pm \rightarrow K^+ K^- \pi^\pm$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
+0.3±1.1±0.8	ALEXANDER 08	CLEO	$e^+ e^-$ at 4.17 GeV

$A_{CP}(K^+ K^- \pi^\pm \pi^0)$ in $D_s^\pm \rightarrow K^+ K^- \pi^\pm \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-5.9±4.2±1.2	ALEXANDER 08	CLEO	$e^+ e^-$ at 4.17 GeV

$A_{CP}(K_S^0 K^\mp 2\pi^\pm)$ in $D_s^+ \rightarrow K_S^0 K^\mp 2\pi^\pm$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-0.7±3.6±1.1	ALEXANDER 08	CLEO	$e^+ e^-$ at 4.17 GeV

$A_{CP}(\pi^+ \pi^- \pi^\pm)$ in $D_s^\pm \rightarrow \pi^+ \pi^- \pi^\pm$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
+2.0±4.6±0.7	ALEXANDER 08	CLEO	$e^+ e^-$ at 4.17 GeV

$A_{CP}(\pi^\pm \eta)$ in $D_s^\pm \rightarrow \pi^\pm \eta$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-4.6±2.9±0.3	2.5k	MENDEZ 10	CLEO	$e^+ e^-$ at 4170 MeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-8.2±5.2±0.8		ALEXANDER 08	CLEO	See MENDEZ 10

$A_{CP}(\pi^\pm \eta')$ in $D_s^\pm \rightarrow \pi^\pm \eta'$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-6.1±3.0±0.3	1.4k	MENDEZ 10	CLEO	$e^+ e^-$ at 4170 MeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-5.5±3.7±1.2		ALEXANDER 08	CLEO	See MENDEZ 10

$A_{CP}(K^\pm \pi^0)$ in $D_s^\pm \rightarrow K^\pm \pi^0$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$-26.6 \pm 23.8 \pm 0.9$	202 ± 70	MENDEZ	10 CLEO	e^+e^- at 4170 MeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$+2 \pm 29$		ADAMS	07A CLEO	See MENDEZ 10

$A_{CP}(K_S^0 \pi^\pm)$ in $D_s^\pm \rightarrow K_S^0 \pi^\pm$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
6.6 ± 3.3 OUR AVERAGE	Error includes scale factor of 1.4.			
$+5.45 \pm 2.50 \pm 0.33$		KO	10 BELL	$e^+e^- \approx \Upsilon(4S)$
$+16.3 \pm 7.3 \pm 0.3$	393 ± 33	MENDEZ	10 CLEO	e^+e^- at 4170 MeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$+27 \pm 11$		ADAMS	07A CLEO	See MENDEZ 10

$A_{CP}(K^\pm \pi^+ \pi^-)$ in $D_s^\pm \rightarrow K^\pm \pi^+ \pi^-$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$+11.2 \pm 7.0 \pm 0.9$	ALEXANDER 08	CLEO	e^+e^- at 4.17 GeV

$A_{CP}(K^\pm \eta)$ in $D_s^\pm \rightarrow K^\pm \eta$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$+9.3 \pm 15.2 \pm 0.9$	222 ± 41	MENDEZ	10 CLEO	e^+e^- at 4170 MeV
••• We do not use the following data for averages, fits, limits, etc. •••				
-20 ± 18		ADAMS	07A CLEO	See MENDEZ 10

$A_{CP}(K^\pm \eta'(958))$ in $D_s^\pm \rightarrow K^\pm \eta'(958)$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$+6.0 \pm 18.9 \pm 0.9$	56 ± 17	MENDEZ	10 CLEO	e^+e^- at 4170 MeV
••• We do not use the following data for averages, fits, limits, etc. •••				
-17 ± 37		ADAMS	07A CLEO	See MENDEZ 10

$D_s^+ - D_s^-$ T-VIOLATING DECAY-RATE ASYMMETRIES

$A_{Tviol}(K_S^0 K^\pm \pi^+ \pi^-)$ in $D_s^\pm \rightarrow K_S^0 K^\pm \pi^+ \pi^-$

$C_T \equiv \vec{p}_{K^+} \cdot (\vec{p}_{\pi^+} \times \vec{p}_{\pi^-})$ is a T -odd correlation of the K^+ , π^+ , and π^- momenta for the D_s^+ . $\bar{C}_T \equiv \vec{p}_{K^-} \cdot (\vec{p}_{\pi^-} \times \vec{p}_{\pi^+})$ is the corresponding quantity for the D_s^- . $A_T \equiv [\Gamma(C_T > 0) - \Gamma(C_T < 0)] / [\Gamma(C_T > 0) + \Gamma(C_T < 0)]$ would, in the absence of strong phases, test for T violation in D_s^+ decays (the Γ 's are partial widths). With $\bar{A}_T \equiv [\Gamma(-\bar{C}_T > 0) - \Gamma(-\bar{C}_T < 0)] / [\Gamma(-\bar{C}_T > 0) + \Gamma(-\bar{C}_T < 0)]$, the asymmetry $A_{Tviol} \equiv \frac{1}{2}(A_T - \bar{A}_T)$ tests for T violation even with nonzero strong phases.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$-0.036 \pm 0.067 \pm 0.023$	508 ± 34	LINK	05E FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$D_s^+ \rightarrow \phi \ell^+ \nu_\ell$ FORM FACTORS

$r_2 \equiv A_2(0)/A_1(0)$ in $D_s^+ \rightarrow \phi \ell^+ \nu_\ell$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.84 ± 0.11 OUR AVERAGE Error includes scale factor of 2.4.				
0.816 ± 0.036 ± 0.030	25 ± 0.5k	⁴⁰ AUBERT	08AN BABR	$\phi e^+ \nu_e$
0.713 ± 0.202 ± 0.284	793	LINK	04C FOCS	$\phi \mu^+ \nu_\mu$
1.57 ± 0.25 ± 0.19	271	AITALA	99D E791	$\phi e^+ \nu_e, \phi \mu^+ \nu_\mu$
1.4 ± 0.5 ± 0.3	308	AVERY	94B CLE2	$\phi e^+ \nu_e$
1.1 ± 0.8 ± 0.1	90	FRABETTI	94F E687	$\phi \mu^+ \nu_\mu$
2.1 ^{+0.6} _{-0.5} ± 0.2	19	KODAMA	93 E653	$\phi \mu^+ \nu_\mu$

⁴⁰ To compare with previous measurements, this AUBERT 08AN value is from a fit that fixes the pole masses at $m_A = 2.5 \text{ GeV}/c^2$ and $m_V = 2.1 \text{ GeV}/c^2$. A simultaneous fit to r_2 , r_V , r_0 (a significant s -wave contribution) and m_A , gives $r_2 = 0.763 \pm 0.071 \pm 0.065$.

$r_V \equiv V(0)/A_1(0)$ in $D_s^+ \rightarrow \phi \ell^+ \nu_\ell$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.80 ± 0.08 OUR AVERAGE				
1.807 ± 0.046 ± 0.065	25 ± 0.5k	⁴¹ AUBERT	08AN BABR	$\phi e^+ \nu_e$
1.549 ± 0.250 ± 0.148	793	LINK	04C FOCS	$\phi \mu^+ \nu_\mu$
2.27 ± 0.35 ± 0.22	271	AITALA	99D E791	$\phi e^+ \nu_e, \phi \mu^+ \nu_\mu$
0.9 ± 0.6 ± 0.3	308	AVERY	94B CLE2	$\phi e^+ \nu_e$
1.8 ± 0.9 ± 0.2	90	FRABETTI	94F E687	$\phi \mu^+ \nu_\mu$
2.3 ^{+1.1} _{-0.9} ± 0.4	19	KODAMA	93 E653	$\phi \mu^+ \nu_\mu$

⁴¹ To compare with previous measurements, this AUBERT 08AN value is from a fit that fixes the pole masses at $m_A = 2.5 \text{ GeV}/c^2$ and $m_V = 2.1 \text{ GeV}/c^2$. A simultaneous fit to r_2 , r_V , r_0 (a significant s -wave contribution) and m_A , gives $r_V = 1.849 \pm 0.060 \pm 0.095$.

Γ_L/Γ_T in $D_s^+ \rightarrow \phi \ell^+ \nu_\ell$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.72 ± 0.18 OUR AVERAGE				
1.0 ± 0.3 ± 0.2	308	AVERY	94B CLE2	$\phi e^+ \nu_e$
1.0 ± 0.5 ± 0.1	90	⁴² FRABETTI	94F E687	$\phi \mu^+ \nu_\mu$
0.54 ± 0.21 ± 0.10	19	⁴² KODAMA	93 E653	$\phi \mu^+ \nu_\mu$

⁴² FRABETTI 94F and KODAMA 93 evaluate Γ_L/Γ_T for a lepton mass of zero.

D_s^\pm REFERENCES

ASNER	10	PR D81 052007	D.M. Asner <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA...	10J	PR D82 091103R	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
KO	10	PRL 104 181602	B.R. Ko <i>et al.</i>	(BELLE Collab.)
MENDEZ	10	PR D81 052013	H. Mendez <i>et al.</i>	(CLEO Collab.)
RUBIN	10	PR D82 092007	P. Rubin <i>et al.</i>	(CLEO Collab.)
ALEXANDER	09	PR D79 052001	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
AUBERT	09O	PR D79 032003	B. Aubert <i>et al.</i>	(BABAR Collab.)
DOBBS	09	PR D79 112008	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ECKLUND	09	PR D80 052009	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
GE	09A	PR D80 051102R	J.Y. Ge <i>et al.</i>	(CLEO Collab.)
KO	09	PRL 102 221802	B.R. Ko <i>et al.</i>	(BELLE Collab.)

MITCHELL	09A	PR D79 072008	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
NAIK	09A	PR D80 112004	P. Naik <i>et al.</i>	(CLEO Collab.)
ONYISI	09	PR D79 052002	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)
WON	09	PR D80 111101R	E. Won <i>et al.</i>	(BELLE Collab.)
YELTON	09	PR D80 052007	J. Yelton <i>et al.</i>	(CLEO Collab.)
ALEXANDER	08	PRL 100 161804	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ATHAR	08	PRL 100 181802	S.B. Athar <i>et al.</i>	(CLEO Collab.)
AUBERT	08AN	PR D78 051101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
ECKLUND	08	PRL 100 161801	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
KLEMPPT	08	EPJ C55 39	E. Klempt, M. Matveev, A.V. Sarantsev	(BONN+)
LINK	08	PL B660 147	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
WIDHALM	08	PRL 100 241801	L. Widhalm <i>et al.</i>	(BELLE Collab.)
ADAMS	07A	PRL 99 191805	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	07V	PRL 98 141801	B. Aubert <i>et al.</i>	(BABAR Collab.)
PEDLAR	07A	PR D76 072002	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
Also		PRL 99 071802	M. Artuso <i>et al.</i>	(CLEO Collab.)
AUBERT	06N	PR D74 031103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
HUANG	06B	PR D74 112005	G.S. Huang <i>et al.</i>	(CLEO Collab.)
PDG	06	JPG 33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
AUBERT	05V	PR D71 091104R	B. Aubert <i>et al.</i>	(BABAR Collab.)
LINK	05E	PL B622 239	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05J	PRL 95 052003	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05K	PL B624 166	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04	PL B585 200	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04C	PL B586 183	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04D	PL B586 191	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04F	PL B601 10	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ACOSTA	03D	PR D68 072004	D. Acosta <i>et al.</i>	(FNAL CDF-II Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
LINK	03D	PL B561 225	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	03F	PL B572 21	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
AUBERT	02G	PR D65 091104R	B. Aubert <i>et al.</i>	(BaBar Collab.)
HEISTER	02I	PL B528 1	A. Heister <i>et al.</i>	(ALEPH Collab.)
LINK	02I	PL B541 227	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02J	PL B541 243	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABBIENDI	01L	PL B516 236	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
IORI	01	PL B523 22	M. Iori <i>et al.</i>	(FNAL SELEX Collab.)
LINK	01C	PRL 87 162001	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ALEXANDROV	00	PL B478 31	Y. Alexandrov <i>et al.</i>	(CERN BEATRICE Collab.)
AITALA	99	PL B445 449	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	99D	PL B450 294	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	99G	PL B462 401	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BONVICINI	99	PRL 82 4586	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
CHADHA	98	PR D58 032002	M. Chada <i>et al.</i>	(CLEO Collab.)
JESSOP	98	PR D58 052002	C.P. Jessop <i>et al.</i>	(CLEO Collab.)
ACCIARRI	97F	PL B396 327	M. Acciarri <i>et al.</i>	(L3 Collab.)
BALEST	97	PRL 79 1436	R. Balest <i>et al.</i>	(CLEO Collab.)
FRABETTI	97C	PL B401 131	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	97D	PL B407 79	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ARTUSO	96	PL B378 364	M. Artuso <i>et al.</i>	(CLEO Collab.)
BAI	95C	PR D52 3781	J.Z. Bai <i>et al.</i>	(BES Collab.)
BRANDENB...	95	PRL 75 3804	G.W. Brandenburg <i>et al.</i>	(CLEO Collab.)
FRABETTI	95B	PL B351 591	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	95F	PL B363 259	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KODAMA	95	PL B345 85	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ACOSTA	94	PR D49 5690	D. Acosta <i>et al.</i>	(CLEO Collab.)
AVERY	94B	PL B337 405	P. Avery <i>et al.</i>	(CLEO Collab.)
BROWN	94	PR D50 1884	D. Brown <i>et al.</i>	(CLEO Collab.)
BUTLER	94	PL B324 255	F. Butler <i>et al.</i>	(CLEO Collab.)
FRABETTI	94F	PL B328 187	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	93F	PRL 71 827	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	93G	PL B313 253	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KODAMA	93	PL B309 483	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ALBRECHT	92B	ZPHY C53 361	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	92	PRL 68 1275	J. Alexander <i>et al.</i>	(CLEO Collab.)
ANJOS	92D	PRL 69 2892	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
AVERY	92	PRL 68 1279	P. Avery <i>et al.</i>	(CLEO Collab.)
BARLAG	92C	ZPHY C55 383	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
Also		ZPHY C48 29	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
DAOUDI	92	PR D45 3965	M. Daoudi <i>et al.</i>	(CLEO Collab.)

FRABETTI	92	PL B281 167	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ALBRECHT	91	PL B255 634	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALVAREZ	91	PL B255 639	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
ALBRECHT	90D	PL B245 315	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	90B	PRL 65 1531	J. Alexander <i>et al.</i>	(CLEO Collab.)
BARLAG	90C	ZPHY C46 563	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
FRABETTI	90	PL B251 639	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ANJOS	89E	PL B223 267	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
CHEN	89	PL B226 192	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
ALBRECHT	88	PL B207 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANJOS	88	PRL 60 897	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
RAAB	88	PR D37 2391	J.R. Raab <i>et al.</i>	(FNAL E691 Collab.)
BECKER	87B	PL B184 277	H. Becker <i>et al.</i>	(NA11 and NA32 Collab.)
BLAYLOCK	87	PRL 58 2171	G.T. Blaylock <i>et al.</i>	(Mark III Collab.)
USHIDA	86	PRL 56 1767	N. Ushida <i>et al.</i>	(FNAL E531 Collab.)
ALBRECHT	85D	PL 153B 343	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
DERRICK	85B	PRL 54 2568	M. Derrick <i>et al.</i>	(HRS Collab.)
AIHARA	84D	PRL 53 2465	H. Aihara <i>et al.</i>	(TPC Collab.)
ALTHOFF	84	PL 136B 130	M. Althoff <i>et al.</i>	(TASSO Collab.)
BAILEY	84	PL 139B 320	R. Bailey <i>et al.</i>	(ACCMOR Collab.)
CHEN	83C	PRL 51 634	A. Chen <i>et al.</i>	(CLEO Collab.)

OTHER RELATED PAPERS

RICHMAN	95	RMP 67 893	J.D. Richman, P.R. Burchat	(UCSB, STAN)
---------	----	------------	----------------------------	--------------