

$\psi(2S)$

$$J^{PC} = 0^{--}(1^{--})$$


See the Review on “ $\psi(2S)$ and χ_c branching ratios” before the $\chi_{c0}(1P)$ Listings.

$\psi(2S)$ MASS

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3686.09 ± 0.04 OUR FIT				Error includes scale factor of 1.6.
3686.093 ± 0.034 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
3686.111 ± 0.025 ± 0.009		AULCHENKO 03	KEDR	$e^+e^- \rightarrow$ hadrons
3685.95 ± 0.10	413	¹ ARTAMONOV 00	OLYA	$e^+e^- \rightarrow$ hadrons
3685.98 ± 0.09 ± 0.04		² ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3686.00 ± 0.10	413	³ ZHOLENTZ 80	OLYA	e^+e^-

- ¹ Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).
² Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $J/\psi(1S)$ mass from AULCHENKO 03.
³ Superseded by ARTAMONOV 00.



$m_{\psi(2S)} - m_{J/\psi(1S)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
589.188 ± 0.028 OUR AVERAGE			
589.194 ± 0.027 ± 0.011	⁴ AULCHENKO 03	KEDR	$e^+ e^- \rightarrow \text{hadrons}$
589.7 ± 1.2	LEMOIGNE 82	GOLI	185 $\pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- \text{A}$
589.07 ± 0.13	⁴ ZHOLENTZ 80	OLYA	$e^+ e^-$
588.7 ± 0.8	LUTH 75	MRK1	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
588 ± 1	⁵ BAI 98E	BES	$e^+ e^-$
⁴ Redundant with data in mass above.			
⁵ Systematic errors not evaluated.			

$\psi(2S)$ WIDTH

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
304 ± 9 OUR FIT				
286 ± 16 OUR AVERAGE				
358 ± 88 ± 4		ABLIKIM 08B	BES2	$e^+ e^- \rightarrow \text{hadrons}$
290 ± 25 ± 4	2.7k	ANDREOTTI 07	E835	$p\bar{p} \rightarrow e^+ e^-, J/\psi X$
331 ± 58 ± 2		ABLIKIM 06L	BES2	$e^+ e^- \rightarrow \text{hadrons}$
264 ± 27		⁶ BAI 02B	BES2	$e^+ e^-$
287 ± 37 ± 16		⁷ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+ e^-$
⁶ From a simultaneous fit to the hadronic and $\mu^+ \mu^-$ cross section, assuming $\Gamma = \Gamma_h + \Gamma_e + \Gamma_\mu + \Gamma_\tau$ and lepton universality. Does not include vacuum polarization correction.				
⁷ The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].				

$\psi(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(97.85 ± 0.13) %	
Γ_2 virtual $\gamma \rightarrow \text{hadrons}$	(1.73 ± 0.14) %	S=1.5
Γ_3 ggg	(10.6 ± 1.6) %	
Γ_4 γgg	(1.03 ± 0.29) %	
Γ_5 light hadrons	(15.4 ± 1.5) %	
Γ_6 $e^+ e^-$	(7.73 ± 0.17) × 10 ⁻³	
Γ_7 $\mu^+ \mu^-$	(7.7 ± 0.8) × 10 ⁻³	
Γ_8 $\tau^+ \tau^-$	(3.0 ± 0.4) × 10 ⁻³	

Decays into $J/\psi(1S)$ and anything

Γ_9 $J/\psi(1S)$ anything	(59.5 ± 0.8) %	
Γ_{10} $J/\psi(1S)$ neutrals	(24.6 ± 0.4) %	
Γ_{11} $J/\psi(1S) \pi^+ \pi^-$	(33.6 ± 0.4) %	
Γ_{12} $J/\psi(1S) \pi^0 \pi^0$	(17.76 ± 0.34) %	
Γ_{13} $J/\psi(1S) \eta$	(3.28 ± 0.07) %	
Γ_{14} $J/\psi(1S) \pi^0$	(1.30 ± 0.10) × 10 ⁻³	S=1.4

Hadronic decays

Γ_{15}	$\pi^0 h_c(1P)$	$(8.4 \pm 1.6) \times 10^{-4}$	
Γ_{16}	$3(\pi^+ \pi^-) \pi^0$	$(3.5 \pm 1.6) \times 10^{-3}$	
Γ_{17}	$2(\pi^+ \pi^-) \pi^0$	$(2.9 \pm 1.0) \times 10^{-3}$	S=4.6
Γ_{18}	$\rho a_2(1320)$	$(2.6 \pm 0.9) \times 10^{-4}$	
Γ_{19}	$\rho \bar{p}$	$(2.76 \pm 0.12) \times 10^{-4}$	
Γ_{20}	$\Delta^{++} \bar{\Delta}^{--}$	$(1.28 \pm 0.35) \times 10^{-4}$	
Γ_{21}	$\Lambda \bar{\Lambda} \pi^0$	$< 1.2 \times 10^{-4}$	CL=90%
Γ_{22}	$\Lambda \bar{\Lambda} \eta$	$< 4.9 \times 10^{-5}$	CL=90%
Γ_{23}	$\Lambda \bar{p} K^+$	$(1.00 \pm 0.14) \times 10^{-4}$	
Γ_{24}	$\Lambda \bar{p} K^+ \pi^+ \pi^-$	$(1.8 \pm 0.4) \times 10^{-4}$	
Γ_{25}	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	$(2.8 \pm 0.6) \times 10^{-4}$	
Γ_{26}	$\Lambda \bar{\Lambda}$	$(2.8 \pm 0.5) \times 10^{-4}$	S=2.6
Γ_{27}	$\Sigma^+ \bar{\Sigma}^-$	$(2.6 \pm 0.8) \times 10^{-4}$	
Γ_{28}	$\Sigma^0 \bar{\Sigma}^0$	$(2.2 \pm 0.4) \times 10^{-4}$	S=1.5
Γ_{29}	$\Sigma(1385)^+ \bar{\Sigma}(1385)^-$	$(1.1 \pm 0.4) \times 10^{-4}$	
Γ_{30}	$\Xi^- \bar{\Xi}^+$	$(1.8 \pm 0.6) \times 10^{-4}$	S=2.8
Γ_{31}	$\Xi^0 \bar{\Xi}^0$	$(2.8 \pm 0.9) \times 10^{-4}$	
Γ_{32}	$\Xi(1530)^0 \bar{\Xi}(1530)^0$	$< 8.1 \times 10^{-5}$	CL=90%
Γ_{33}	$\Omega^- \bar{\Omega}^+$	$< 7.3 \times 10^{-5}$	CL=90%
Γ_{34}	$\pi^0 \rho \bar{p}$	$(1.50 \pm 0.08) \times 10^{-4}$	S=1.1
Γ_{35}	$N_1^*(1440) \bar{p} \rightarrow \pi^0 \rho \bar{p}$	$(8.1 \pm 0.8) \times 10^{-5}$	
Γ_{36}	$\pi^0 f_0(2100) \rightarrow \pi^0 \rho \bar{p}$	$(1.1 \pm 0.4) \times 10^{-5}$	
Γ_{37}	$\eta \rho \bar{p}$	$(5.7 \pm 0.6) \times 10^{-5}$	
Γ_{38}	$\eta f_0(2100) \rightarrow \eta \rho \bar{p}$	$(1.2 \pm 0.4) \times 10^{-5}$	
Γ_{39}	$N^*(1535) \bar{p} \rightarrow \eta \rho \bar{p}$	$(4.4 \pm 0.7) \times 10^{-5}$	
Γ_{40}	$\omega \rho \bar{p}$	$(6.9 \pm 2.1) \times 10^{-5}$	
Γ_{41}	$\phi \rho \bar{p}$	$< 2.4 \times 10^{-5}$	CL=90%
Γ_{42}	$\pi^+ \pi^- \rho \bar{p}$	$(6.0 \pm 0.4) \times 10^{-4}$	
Γ_{43}	$\rho \bar{n} \pi^-$ or c.c.	$(2.48 \pm 0.17) \times 10^{-4}$	
Γ_{44}	$\rho \bar{n} \pi^- \pi^0$	$(3.2 \pm 0.7) \times 10^{-4}$	
Γ_{45}	$2(\pi^+ \pi^- \pi^0)$	$(4.8 \pm 1.5) \times 10^{-3}$	
Γ_{46}	$\eta \pi^+ \pi^-$	$< 1.6 \times 10^{-4}$	CL=90%
Γ_{47}	$\eta \pi^+ \pi^- \pi^0$	$(9.5 \pm 1.7) \times 10^{-4}$	
Γ_{48}	$2(\pi^+ \pi^-) \eta$	$(1.2 \pm 0.6) \times 10^{-3}$	
Γ_{49}	$\eta' \pi^+ \pi^- \pi^0$	$(4.5 \pm 2.1) \times 10^{-4}$	
Γ_{50}	$\omega \pi^+ \pi^-$	$(7.3 \pm 1.2) \times 10^{-4}$	S=2.1
Γ_{51}	$b_1^\pm \pi^\mp$	$(4.0 \pm 0.6) \times 10^{-4}$	S=1.1
Γ_{52}	$b_1^0 \pi^0$	$(2.4 \pm 0.6) \times 10^{-4}$	
Γ_{53}	$\omega f_2(1270)$	$(2.2 \pm 0.4) \times 10^{-4}$	
Γ_{54}	$\pi^+ \pi^- K^+ K^-$	$(7.5 \pm 0.9) \times 10^{-4}$	S=1.9
Γ_{55}	$\rho^0 K^+ K^-$	$(2.2 \pm 0.4) \times 10^{-4}$	
Γ_{56}	$K^*(892)^0 \bar{K}_2^*(1430)^0$	$(1.9 \pm 0.5) \times 10^{-4}$	
Γ_{57}	$K^+ K^- \pi^+ \pi^- \eta$	$(1.3 \pm 0.7) \times 10^{-3}$	

Γ_{58}	$K^+ K^- 2(\pi^+ \pi^-) \pi^0$	$(1.00 \pm 0.31) \times 10^{-3}$	
Γ_{59}	$K^+ K^- 2(\pi^+ \pi^-)$	$(1.9 \pm 0.9) \times 10^{-3}$	
Γ_{60}	$K_1(1270)^\pm K^\mp$	$(1.00 \pm 0.28) \times 10^{-3}$	
Γ_{61}	$K_S^0 K_S^0 \pi^+ \pi^-$	$(2.2 \pm 0.4) \times 10^{-4}$	
Γ_{62}	$\rho^0 p \bar{p}$	$(5.0 \pm 2.2) \times 10^{-5}$	
Γ_{63}	$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	$(6.7 \pm 2.5) \times 10^{-4}$	
Γ_{64}	$2(\pi^+ \pi^-)$	$(2.4 \pm 0.6) \times 10^{-4}$	S=2.2
Γ_{65}	$\rho^0 \pi^+ \pi^-$	$(2.2 \pm 0.6) \times 10^{-4}$	S=1.4
Γ_{66}	$K^+ K^- \pi^+ \pi^- \pi^0$	$(1.26 \pm 0.09) \times 10^{-3}$	
Γ_{67}	$\omega f_0(1710) \rightarrow \omega K^+ K^-$	$(5.9 \pm 2.2) \times 10^{-5}$	
Γ_{68}	$K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.}$	$(8.6 \pm 2.2) \times 10^{-4}$	
Γ_{69}	$K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.}$	$(9.6 \pm 2.8) \times 10^{-4}$	
Γ_{70}	$K^*(892)^+ K^- \rho^0 + \text{c.c.}$	$(7.3 \pm 2.6) \times 10^{-4}$	
Γ_{71}	$K^*(892)^0 K^- \rho^+ + \text{c.c.}$	$(6.1 \pm 1.8) \times 10^{-4}$	
Γ_{72}	$\eta K^+ K^-$	$< 1.3 \times 10^{-4}$	CL=90%
Γ_{73}	$\omega K^+ K^-$	$(1.85 \pm 0.25) \times 10^{-4}$	S=1.1
Γ_{74}	$3(\pi^+ \pi^-)$	$(3.5 \pm 2.0) \times 10^{-4}$	S=2.8
Γ_{75}	$\rho \bar{p} \pi^+ \pi^- \pi^0$	$(7.3 \pm 0.7) \times 10^{-4}$	
Γ_{76}	$K^+ K^-$	$(6.3 \pm 0.7) \times 10^{-5}$	
Γ_{77}	$K_S^0 K_L^0$	$(5.4 \pm 0.5) \times 10^{-5}$	
Γ_{78}	$\pi^+ \pi^- \pi^0$	$(1.68 \pm 0.26) \times 10^{-4}$	S=1.4
Γ_{79}	$\rho(2150) \pi \rightarrow \pi^+ \pi^- \pi^0$	$(1.9 \pm_{-0.4}^{+1.2}) \times 10^{-4}$	
Γ_{80}	$\rho(770) \pi \rightarrow \pi^+ \pi^- \pi^0$	$(3.2 \pm 1.2) \times 10^{-5}$	S=1.8
Γ_{81}	$\pi^+ \pi^-$	$(8 \pm 5) \times 10^{-5}$	
Γ_{82}	$K_1(1400)^\pm K^\mp$	$< 3.1 \times 10^{-4}$	CL=90%
Γ_{83}	$K^+ K^- \pi^0$	$< 2.96 \times 10^{-5}$	CL=90%
Γ_{84}	$K^+ \bar{K}^*(892)^- + \text{c.c.}$	$(1.7 \pm_{-0.7}^{+0.8}) \times 10^{-5}$	
Γ_{85}	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$(1.09 \pm 0.20) \times 10^{-4}$	
Γ_{86}	$\phi \pi^+ \pi^-$	$(1.17 \pm 0.29) \times 10^{-4}$	S=1.7
Γ_{87}	$\phi f_0(980) \rightarrow \pi^+ \pi^-$	$(6.8 \pm 2.5) \times 10^{-5}$	S=1.1
Γ_{88}	$2(K^+ K^-)$	$(6.0 \pm 1.4) \times 10^{-5}$	
Γ_{89}	$\phi K^+ K^-$	$(7.0 \pm 1.6) \times 10^{-5}$	
Γ_{90}	$2(K^+ K^-) \pi^0$	$(1.10 \pm 0.28) \times 10^{-4}$	
Γ_{91}	$\phi \eta$	$(2.8 \pm_{-0.8}^{+1.0}) \times 10^{-5}$	
Γ_{92}	$\phi \eta'$	$(3.1 \pm 1.6) \times 10^{-5}$	
Γ_{93}	$\omega \eta'$	$(3.2 \pm_{-2.1}^{+2.5}) \times 10^{-5}$	
Γ_{94}	$\omega \pi^0$	$(2.1 \pm 0.6) \times 10^{-5}$	
Γ_{95}	$\rho \eta'$	$(1.9 \pm_{-1.2}^{+1.7}) \times 10^{-5}$	
Γ_{96}	$\rho \eta$	$(2.2 \pm 0.6) \times 10^{-5}$	S=1.1
Γ_{97}	$\omega \eta$	$< 1.1 \times 10^{-5}$	CL=90%
Γ_{98}	$\phi \pi^0$	$< 4 \times 10^{-6}$	CL=90%

Γ_{99}	$\eta_c \pi^+ \pi^- \pi^0$	$< 1.0 \times 10^{-3}$	CL=90%
Γ_{100}	$p \bar{p} K^+ K^-$	$(2.7 \pm 0.7) \times 10^{-5}$	
Γ_{101}	$\bar{\Lambda} n K_S^0 + \text{c.c.}$	$(8.1 \pm 1.8) \times 10^{-5}$	
Γ_{102}	$\phi f_2'(1525)$	$(4.4 \pm 1.6) \times 10^{-5}$	
Γ_{103}	$\Theta(1540) \bar{\Theta}(1540) \rightarrow$ $K_S^0 p K^- \bar{n} + \text{c.c.}$	$< 8.8 \times 10^{-6}$	CL=90%
Γ_{104}	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	$< 1.0 \times 10^{-5}$	CL=90%
Γ_{105}	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	$< 7.0 \times 10^{-6}$	CL=90%
Γ_{106}	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	$< 2.6 \times 10^{-5}$	CL=90%
Γ_{107}	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	$< 6.0 \times 10^{-6}$	CL=90%
Γ_{108}	$K_S^0 K_S^0$	$< 4.6 \times 10^{-6}$	

Radiative decays

Γ_{109}	$\gamma \chi_{c0}(1P)$	$(9.68 \pm 0.31) \%$	
Γ_{110}	$\gamma \chi_{c1}(1P)$	$(9.2 \pm 0.4) \%$	
Γ_{111}	$\gamma \chi_{c2}(1P)$	$(8.75 \pm 0.35) \%$	
Γ_{112}	$\gamma \eta_c(1S)$	$(3.4 \pm 0.5) \times 10^{-3}$	S=1.3
Γ_{113}	$\gamma \eta_c(2S)$	$< 8 \times 10^{-4}$	CL=90%
Γ_{114}	$\gamma \pi^0$	$(1.6 \pm 0.4) \times 10^{-6}$	
Γ_{115}	$\gamma \eta'(958)$	$(1.23 \pm 0.06) \times 10^{-4}$	
Γ_{116}	$\gamma f_2(1270)$	$(2.1 \pm 0.4) \times 10^{-4}$	
Γ_{117}	$\gamma f_0(1710)$		
Γ_{118}	$\gamma f_0(1710) \rightarrow \gamma \pi \pi$	$(3.0 \pm 1.3) \times 10^{-5}$	
Γ_{119}	$\gamma f_0(1710) \rightarrow \gamma K \bar{K}$	$(6.0 \pm 1.6) \times 10^{-5}$	
Γ_{120}	$\gamma \gamma$	$< 1.4 \times 10^{-4}$	CL=90%
Γ_{121}	$\gamma \eta$	$(1.4 \pm 0.5) \times 10^{-6}$	
Γ_{122}	$\gamma \eta \pi^+ \pi^-$	$(8.7 \pm 2.1) \times 10^{-4}$	
Γ_{123}	$\gamma \eta(1405)$		
Γ_{124}	$\gamma \eta(1405) \rightarrow \gamma K \bar{K} \pi$	$< 9 \times 10^{-5}$	CL=90%
Γ_{125}	$\gamma \eta(1405) \rightarrow \eta \pi^+ \pi^-$	$(3.6 \pm 2.5) \times 10^{-5}$	
Γ_{126}	$\gamma \eta(1475)$		
Γ_{127}	$\gamma \eta(1475) \rightarrow K \bar{K} \pi$	$< 1.4 \times 10^{-4}$	CL=90%
Γ_{128}	$\gamma \eta(1475) \rightarrow \eta \pi^+ \pi^-$	$< 8.8 \times 10^{-5}$	CL=90%
Γ_{129}	$\gamma 2(\pi^+ \pi^-)$	$(4.0 \pm 0.6) \times 10^{-4}$	
Γ_{130}	$\gamma K^{*0} K^+ \pi^- + \text{c.c.}$	$(3.7 \pm 0.9) \times 10^{-4}$	
Γ_{131}	$\gamma K^{*0} \bar{K}^{*0}$	$(2.4 \pm 0.7) \times 10^{-4}$	
Γ_{132}	$\gamma K_S^0 K^+ \pi^- + \text{c.c.}$	$(2.6 \pm 0.5) \times 10^{-4}$	
Γ_{133}	$\gamma K^+ K^- \pi^+ \pi^-$	$(1.9 \pm 0.5) \times 10^{-4}$	
Γ_{134}	$\gamma p \bar{p}$	$(3.9 \pm 0.5) \times 10^{-5}$	S=2.0
Γ_{135}	$\gamma f_2(1950) \rightarrow \gamma p \bar{p}$	$(1.20 \pm 0.22) \times 10^{-5}$	
Γ_{136}	$\gamma f_2(2150) \rightarrow \gamma p \bar{p}$	$(7.2 \pm 1.8) \times 10^{-6}$	
Γ_{137}	$\gamma X(1835) \rightarrow \gamma p \bar{p}$	$< 1.6 \times 10^{-6}$	CL=90%
Γ_{138}	$\gamma X \rightarrow \gamma p \bar{p}$	[a] $< 2 \times 10^{-6}$	CL=90%

Γ_{139}	$\gamma\pi^+\pi^-\rho\bar{p}$	$(2.8 \pm 1.4) \times 10^{-5}$	
Γ_{140}	$\gamma 2(\pi^+\pi^-)K^+K^-$	$< 2.2 \times 10^{-4}$	CL=90%
Γ_{141}	$\gamma 3(\pi^+\pi^-)$	$< 1.7 \times 10^{-4}$	CL=90%
Γ_{142}	$\gamma K^+K^-K^+K^-$	$< 4 \times 10^{-5}$	CL=90%

[a] For a narrow resonance in the range $2.2 < M(X) < 2.8$ GeV.

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 24 combinations of partial widths obtained from integrated cross section, and 83 branching ratios uses 218 measurements to determine 48 parameters. The overall fit has a $\chi^2 = 307.7$ for 170 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_7	5									
x_8	1	0								
x_{11}	44	12	3							
x_{12}	39	8	2	64						
x_{13}	27	7	2	57	35					
x_{19}	2	1	0	7	5	4				
x_{109}	2	1	0	4	3	2	0			
x_{110}	2	1	0	5	2	3	0	0		
x_{111}	3	1	0	6	4	4	0	0	0	
Γ	-79	-6	-2	-52	-46	-32	-10	-2	-3	-3
	x_6	x_7	x_8	x_{11}	x_{12}	x_{13}	x_{19}	x_{109}	x_{110}	x_{111}

$\psi(2S)$ PARTIAL WIDTHS

$\Gamma(\text{hadrons})$

Γ_1

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
258 ± 26	BAI	02B	BES2 e^+e^-
224 ± 56	LUTH	75	MRK1 e^+e^-

$\Gamma(e^+e^-)$

Γ_6

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.35 ± 0.04 OUR FIT			
2.33 ± 0.07 OUR AVERAGE			
2.338 ± 0.037 ± 0.096	ABLIKIM	08B BES2	$e^+e^- \rightarrow$ hadrons
2.330 ± 0.036 ± 0.110	ABLIKIM	06L BES2	$e^+e^- \rightarrow$ hadrons
2.44 ± 0.21	⁸ BAI	02B BES2	e^+e^-
2.14 ± 0.21	ALEXANDER	89 RVUE	See Υ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.0 ± 0.3	BRANDELIK	79C DASP	e^+e^-
2.1 ± 0.3	⁹ LUTH	75 MRK1	e^+e^-

⁸From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channel, assuming $\Gamma_e = \Gamma_\mu = \Gamma_\tau/0.38847$.

⁹From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

$\Gamma(\gamma\gamma)$

Γ_{120}

<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<43	90	BRANDELIK	79C DASP	e^+e^-

$\psi(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel(i) in the e^+e^- annihilation. We list only data that have not been used to determine the partial width $\Gamma(i)$ or the branching ratio $\Gamma(i)/\text{total}$.

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$\Gamma_1\Gamma_6/\Gamma$

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.2 ± 0.4	ABRAMS	75 MRK1	e^+e^-

$\Gamma(\tau^+\tau^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$\Gamma_8\Gamma_6/\Gamma$


<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9.0 ± 2.6	79	¹⁰ ANASHIN	07 KEDR	$e^+e^- \rightarrow \psi(2S) \rightarrow \tau^+\tau^-$
¹⁰ Using $\psi(2S)$ total width of 337 ± 13 keV. Systematic errors not evaluated.				

$\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$\Gamma_{11}\Gamma_6/\Gamma$

<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.789 ± 0.015 OUR FIT				
0.82 ± 0.04 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
0.852 ± 0.010 ± 0.026	19.5k ± 243	ADAM	06 CLEO	3.773 $e^+e^- \rightarrow \gamma\psi(2S)$
0.76 ± 0.05 ± 0.01	544	¹¹ AUBERT	05D BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-\gamma$
0.68 ± 0.09		¹² BAI	98E BES	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.90 ± 0.08 ± 0.05	256	¹³ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$

- ¹¹ AUBERT 05D reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = 0.0450 \pm 0.0018 \pm 0.0022$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.93 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ¹² The value of $\Gamma(e^+e^-)$ quoted in BAI 98E is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6) \times 10^{-2}$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1203 \pm 0.0038$. Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.
- ¹³ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0186 \pm 0.0012 \pm 0.0011$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.07 \pm 0.12) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(J/\psi(1S)\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{12}\Gamma_6/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.417±0.010 OUR FIT					
0.411±0.008±0.018	3.6k±96	ADAM	06	CLEO	3.773 $e^+e^- \rightarrow \gamma\psi(2S)$

$\Gamma(J/\psi(1S)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{13}\Gamma_6/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
77.0± 1.9 OUR FIT					
87 ± 9 OUR AVERAGE					
83 ±25 ±5	14	¹⁴ AUBERT	07AU	BABR	10.6 $e^+e^- \rightarrow J/\psi\pi^+\pi^-\pi^0\gamma$
88 ± 6 ±7	291 ± 24	ADAM	06	CLEO	3.773 $e^+e^- \rightarrow \gamma\psi(2S)$

¹⁴ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow J/\psi\eta) \cdot B(J/\psi \rightarrow \mu^+\mu^-) \cdot B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.11 \pm 0.33 \pm 0.07$ eV.

$\Gamma(J/\psi(1S)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{14}\Gamma_6/\Gamma$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<8	90	<37	ADAM	06	CLEO 3.773 $e^+e^- \rightarrow \gamma\psi(2S)$

$\Gamma(\rho\bar{\rho}) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{19}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.647±0.028 OUR FIT				
0.59 ±0.05 OUR AVERAGE				
0.579±0.038±0.036	2.7k	ANDREOTTI 07	E835	$\rho\bar{\rho} \rightarrow e^+e^-$, $J/\psi X$
0.70 ±0.17 ±0.03	22	AUBERT 06B		$e^+e^- \rightarrow \rho\bar{\rho}\gamma$

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{26}\Gamma_6/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
1.5±0.4±0.1	AUBERT 07BD	BABR	10.6 $e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$

$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{45}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.2±3.3±1.3	43	AUBERT 06D	BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$

$\Gamma(K^+K^-2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{59}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.4±2.1±0.3	26	AUBERT 06D	BABR	10.6 $e^+e^- \rightarrow K^+K^-2(\pi^+\pi^-)\gamma$

$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{54}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.56±0.42±0.16	85	AUBERT 07AK	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

$\Gamma(\phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{87}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.347±0.169±0.003	6 ± 3	¹⁵ AUBERT 07AK	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

¹⁵ AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)]/\Gamma_{total} \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.17 \pm 0.08 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{86}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.57±0.23±0.01	10	¹⁶ AUBERT,BE 06D	BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

¹⁶ AUBERT,BE 06D reports $[\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)]/\Gamma_{total} \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.28 \pm 0.11 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-))/\Gamma_{\text{total}}$					$\Gamma_{17}\Gamma_6/\Gamma$
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
29.7±2.2±1.8	410	AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)π ⁰ γ	

$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{50}\Gamma_6/\Gamma$
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
3.01±0.84±0.02	37	17 AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → ωπ ⁺ π ⁻ γ	
¹⁷ AUBERT 07AU reports [$\Gamma(\psi(2S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(ω(782) → π ⁺ π ⁻ π ⁰)] = 2.69 ± 0.73 ± 0.16 eV which we divide by our best value B(ω(782) → π ⁺ π ⁻ π ⁰) = (89.2 ± 0.7) × 10 ⁻² . Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(2(\pi^+\pi^-\eta) \times \Gamma(e^+e^-))/\Gamma_{\text{total}}$					$\Gamma_{48}\Gamma_6/\Gamma$
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2.87±1.41±0.01	16	18 AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)ηγ	
¹⁸ AUBERT 07AU reports [$\Gamma(\psi(2S) \rightarrow 2(\pi^+\pi^-\eta) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(η → 2γ)] = 1.13 ± 0.55 ± 0.08 eV which we divide by our best value B(η → 2γ) = (39.31 ± 0.20) × 10 ⁻² . Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{66}\Gamma_6/\Gamma$
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
4.4±1.3±0.3	32	AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ π ⁰ γ	

$\Gamma(K^+K^-\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{57}\Gamma_6/\Gamma$
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
3.05±1.80±0.02	7	19 AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ ηγ	
¹⁹ AUBERT 07AU reports [$\Gamma(\psi(2S) \rightarrow K^+K^-\pi^+\pi^-\eta) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(η → 2γ)] = 1.2 ± 0.7 ± 0.1 eV which we divide by our best value B(η → 2γ) = (39.31 ± 0.20) × 10 ⁻² . Our first error is their experiment's error and our second error is the systematic error from using our best value.					

ψ(2S) BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$					Γ_1/Γ
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.9785±0.0013 OUR AVERAGE					
0.9779±0.0015		²⁰ BAI	02B	BES2 e ⁺ e ⁻	
0.981 ±0.003		²⁰ LUTH	75	MRK1 e ⁺ e ⁻	
²⁰ Includes cascade decay into J/ψ(1S).					

$\Gamma(\text{virtual}\gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$					Γ_2/Γ
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.0173±0.0014 OUR AVERAGE	Error includes scale factor of 1.5.				
0.0166±0.0010		^{21,22} SETH	04	RVUE e ⁺ e ⁻	
0.0199±0.0019		²¹ BAI	02B	BES2 e ⁺ e ⁻	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.029 ±0.004		²¹ LUTH	75	MRK1 e ⁺ e ⁻	

²¹ Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.

²² Using $B(\psi(2S) \rightarrow \ell^+ \ell^-) = (0.73 \pm 0.04)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C.

$\Gamma(g g g)/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.58 ± 1.62	2.9 M	²³ LIBBY	09	CLEO $\psi(2S) \rightarrow \text{hadrons}$

²³ Calculated using $\Gamma(\gamma g g)/\Gamma(g g g) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09, $B(\psi(2S) \rightarrow X J/\psi)$ relative and absolute branching fractions from MENDEZ 08, $B(\psi(2S) \rightarrow \gamma \eta_c)$ from MITCHELL 09, and $B(\psi(2S) \rightarrow \text{virtual } \gamma \rightarrow \text{hadrons})$, $B(\psi(2S) \rightarrow \gamma \chi_{cJ})$, and $B(\psi(2S) \rightarrow \ell^+ \ell^-)$ from PDG 08. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(\gamma g g)/\Gamma_{\text{total}}$ LIBBY 09 measurement.

$\Gamma(\gamma g g)/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.025 ± 0.288	200 k	²⁴ LIBBY	09	CLEO $\psi(2S) \rightarrow \gamma + \text{hadrons}$

²⁴ Calculated using $\Gamma(\gamma g g)/\Gamma(g g g) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(g g g)/\Gamma_{\text{total}}$ LIBBY 09 measurement.

$\Gamma(\gamma g g)/\Gamma(g g g)$ Γ_4/Γ_3

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.7 ± 2.6 ± 1.6	2.9 M	LIBBY	09	CLEO $\psi(2S) \rightarrow (\gamma +) \text{hadrons}$

$\Gamma(\text{light hadrons})/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.154 ± 0.015	²⁵ MENDEZ	08	CLEO $e^+ e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.169 ± 0.026	²⁶ ADAM	05A	CLEO $e^+ e^- \rightarrow \psi(2S)$
---------------	--------------------	-----	-------------------------------------

²⁵ Uses $B(\psi(2S) \rightarrow J/\psi X)$ from MENDEZ 08 and other branching fractions from PDG 07.

²⁶ Uses $B(J/\psi X)$ from ADAM 05A, $B(\chi_{cJ} \gamma)$, $B(\eta_c \gamma)$ from ATHAR 04 and $B(\ell^+ \ell^-)$ from PDG 04. Superseded by MENDEZ 08.

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
77.3 ± 1.7 OUR FIT			

• • • We do not use the following data for averages, fits, limits, etc. • • •

88 ± 13	²⁷ FELDMAN	77	RVUE $e^+ e^-$
---------	-----------------------	----	----------------

²⁷ From an overall fit assuming equal partial widths for $e^+ e^-$ and $\mu^+ \mu^-$. For a measurement of the ratio see the entry $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$ below. Includes LUTH 75, HILGER 75, BURMESTER 77.

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
77 ± 8 OUR FIT	

$\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$ Γ_7/Γ_6

VALUE DOCUMENT ID TECN COMMENT

1.00 ± 0.11 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.89 ± 0.16 BOYARSKI 75C MRK1 $e^+ e^-$

$\Gamma(\tau^+ \tau^-)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

30 ± 4 OUR FIT

30.8 ± 2.1 ± 3.8 ²⁸ ABLIKIM 06W BES $e^+ e^- \rightarrow \psi(2S)$

²⁸ Computed using PDG 02 value of $B(\psi(2S) \rightarrow \text{hadrons}) = 0.9810 \pm 0.0030$ to estimate the total number of $\psi(2S)$ events.

————— **DECAYS INTO $J/\psi(1S)$ AND ANYTHING** —————

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE EVTS DOCUMENT ID TECN COMMENT

0.595 ± 0.008 OUR FIT

0.55 ± 0.07 OUR AVERAGE

0.51 ± 0.12 BRANDELIK 79C DASP $e^+ e^- \rightarrow \mu^+ \mu^- X$

0.57 ± 0.08 ABRAMS 75B MRK1 $e^+ e^- \rightarrow \mu^+ \mu^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6254 ± 0.0016 ± 0.0155 1.1M ²⁹ MENDEZ 08 CLEO $\psi(2S) \rightarrow \ell^+ \ell^- X$

0.5950 ± 0.0015 ± 0.0190 151k ADAM 05A CLEO Repl. by MENDEZ 08

²⁹ Not independent from other measurements of MENDEZ 08.

$\Gamma(e^+ e^-)/\Gamma(J/\psi(1S)\text{anything})$
 $\Gamma_6/\Gamma_9 = \Gamma_6/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.344\Gamma_{110} + 0.195\Gamma_{111})$

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

1.299 ± 0.026 OUR FIT

1.28 ± 0.04 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.


1.22 ± 0.02 ± 0.05 5097 ± 73 ³⁰ ANDREOTTI 05 E835 $p\bar{p} \rightarrow \psi(2S) \rightarrow e^+ e^-$

1.28 ± 0.03 ± 0.02 ³⁰ AMBROGIANI 00A E835 $p\bar{p} \rightarrow \psi(2S)$

1.44 ± 0.08 ± 0.02 ³⁰ ARMSTRONG 97 E760 $\bar{p}p \rightarrow \psi(2S)$

³⁰ Using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

WEIGHTED AVERAGE
 1.28 ± 0.04 (Error scaled by 1.6)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

			χ^2
ANDREOTTI	05	E835	1.3
AMBROGIANI	00A	E835	0.0
ARMSTRONG	97	E760	3.7
			5.0

(Confidence Level = 0.082)

$\Gamma(e^+ e^-) / \Gamma(J/\psi(1S)\text{anything})$ (units 10^{-2})

$\Gamma(\mu^+ \mu^-) / \Gamma(J/\psi(1S)\text{anything})$

$$\Gamma_7 / \Gamma_9 = \Gamma_7 / (\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.344\Gamma_{110} + 0.195\Gamma_{111})$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0130 ± 0.0014 OUR FIT			
0.014 ± 0.003	HILGER	75	SPEC $e^+ e^-$

$\Gamma(J/\psi(1S)\text{neutrals}) / \Gamma_{\text{total}}$

Γ_{10} / Γ

VALUE	DOCUMENT ID
0.246 ± 0.004 OUR FIT	

$\Gamma(J/\psi(1S)\pi^+ \pi^-) / \Gamma_{\text{total}}$

Γ_{11} / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.336 ± 0.004 OUR FIT				
0.343 ± 0.011 OUR AVERAGE				Error includes scale factor of 1.7.
$0.3504 \pm 0.0007 \pm 0.0077$	565k	MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+ \ell^- \pi^+ \pi^-$
0.323 ± 0.014		BAI	02B	BES2 $e^+ e^-$
0.32 ± 0.04		ABRAMS	75B	MRK1 $e^+ e^- \rightarrow J/\psi \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.3354 \pm 0.0014 \pm 0.0110$	60k	³¹ ADAM	05A	CLEO Repl. by MENDEZ 08
--------------------------------	-----	--------------------	-----	-------------------------

³¹Not independent from other values reported by ADAM 05A.

$\Gamma(e^+ e^-) / \Gamma(J/\psi(1S)\pi^+ \pi^-)$

Γ_6 / Γ_{11}

VALUE	DOCUMENT ID	TECN	COMMENT
0.0230 ± 0.0005 OUR FIT			
$0.0252 \pm 0.0028 \pm 0.0011$	³² AUBERT	02B	BABR $e^+ e^-$

³²Using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

$\Gamma(\mu^+ \mu^-) / \Gamma(J/\psi(1S) \pi^+ \pi^-)$ Γ_7 / Γ_{11}

VALUE	DOCUMENT ID	TECN	COMMENT
0.0229 ± 0.0025 OUR FIT			
0.0224 ± 0.0029 OUR AVERAGE			
0.0216 ± 0.0026 ± 0.0014	³³ AUBERT	02B	BABR $e^+ e^-$
0.0327 ± 0.0077 ± 0.0072	³³ GRIBUSHIN	96	FMPs 515 $\pi^- \text{Be} \rightarrow 2\mu X$

³³ Using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.


$\Gamma(\tau^+ \tau^-) / \Gamma(J/\psi(1S) \pi^+ \pi^-)$ Γ_8 / Γ_{11}

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
9.0 ± 1.1 OUR FIT			
8.73 ± 1.39 ± 1.57	BAI	02	BES $e^+ e^-$

$\Gamma(J/\psi(1S) \pi^+ \pi^-) / \Gamma(J/\psi(1S) \text{anything})$ Γ_{11} / Γ_9

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.5647 ± 0.0026 OUR FIT				
0.554 ± 0.008 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
0.5604 ± 0.0009 ± 0.0062	565k	MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+ \ell^- \pi^+ \pi^-$
0.525 ± 0.009 ± 0.022	4k	ANDREOTTI	05	E835 $\psi(2S) \rightarrow J/\psi X$
0.536 ± 0.007 ± 0.016	20k	^{34,35} ABLIKIM	04B	BES $\psi(2S) \rightarrow J/\psi X$
0.496 ± 0.037		ARMSTRONG	97	E760 $\bar{p} p \rightarrow \psi(2S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.5637 ± 0.0027 ± 0.0046	60k	ADAM	05A	CLEO Repl. by MENDEZ 08

³⁴ From a fit to the J/ψ recoil mass spectra.
³⁵ ABLIKIM 04B quotes $B(\psi(2S) \rightarrow J/\psi X) / B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)$.



$$\frac{\Gamma(J/\psi(1S)\text{ neutrals})/\Gamma(J/\psi(1S)\pi^+\pi^-)}{\Gamma_{10}/\Gamma_{11} = (0.9761\Gamma_{12} + 0.719\Gamma_{13} + 0.344\Gamma_{110} + 0.195\Gamma_{111})/\Gamma_{11}}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.731 ± 0.008 OUR FIT			
0.73 ± 0.09	TANENBAUM 76	MRK1	e^+e^-

$$\frac{\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma_{\text{total}}}{\Gamma_{12}/\Gamma}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.1776 ± 0.0034 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.1769 ± 0.0008 ± 0.0053	61k	³⁶ MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+\ell^-2\pi^0$
0.1652 ± 0.0014 ± 0.0058	13.4k	³⁷ ADAM	05A	CLEO	Repl. by MENDEZ 08

³⁶ Not independent from other measurements of MENDEZ 08.

³⁷ Not independent from other values reported by ADAM 05A.

$$\frac{\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\text{ anything})}{\Gamma_{12}/\Gamma_9}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.2982 ± 0.0032 OUR FIT				

0.320 ± 0.012 OUR AVERAGE

0.300 ± 0.008 ± 0.022	1655 ± 44	ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$
0.328 ± 0.013 ± 0.008		AMBROGIANI 00A	E835	$p\bar{p} \rightarrow \psi(2S)$
0.323 ± 0.033		ARMSTRONG 97	E760	$\bar{p}p \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.2829 ± 0.0012 ± 0.0056	61k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+\ell^-2\pi^0$
0.2776 ± 0.0025 ± 0.0043	13.4k	ADAM	05A	CLEO	Repl. by MENDEZ 08

$$\frac{\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)}{\Gamma_{12}/\Gamma_{11}}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.528 ± 0.008 OUR FIT				

0.513 ± 0.022 OUR AVERAGE Error includes scale factor of 2.2.

0.5047 ± 0.0022 ± 0.0102	61k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+\ell^-2\pi^0$
0.570 ± 0.009 ± 0.026	14k	³⁸ ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.4924 ± 0.0047 ± 0.0086	73k	^{39,40} ADAM	05A	CLEO	Repl. by MENDEZ 08
0.571 ± 0.018 ± 0.044		⁴¹ ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$	
0.53 ± 0.06		TANENBAUM 76	MRK1	e^+e^-	
0.64 ± 0.15		⁴² HILGER 75	SPEC	e^+e^-	

³⁸ From a fit to the J/ψ recoil mass spectra.

³⁹ Not independent from other values reported by ADAM 05A.

⁴⁰ Using 13,217 $J/\psi\pi^0\pi^0$ and 60,010 $J/\psi\pi^+\pi^-$ events.

⁴¹ Not independent from other values reported by ANDREOTTI 05.

⁴² Ignoring the $J/\psi(1S)\eta$ and $J/\psi(1S)\gamma\gamma$ decays.


$\Gamma(J/\psi(1S)\eta)/\Gamma_{total}$

Γ_{13}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0328 ± 0.0007 OUR FIT				
0.0296 ± 0.0031 OUR AVERAGE				Error includes scale factor of 1.8. See the ideogram below.
0.0298 ± 0.0009 ± 0.0023	5.7k	BAI	04I	BES2 $\psi(2S) \rightarrow J/\psi\gamma\gamma$
0.0255 ± 0.0029	386	43 OREGLIA	80	CBAL $e^+e^- \rightarrow J/\psi 2\gamma$
0.045 ± 0.012	17	44 BRANDELIK	79B	DASP $e^+e^- \rightarrow J/\psi 2\gamma$
0.042 ± 0.006	164	44 BARTEL	78B	CNTR e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.0343 ± 0.0004 ± 0.0009	18.4k	45 MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+\ell^-\eta$
0.0325 ± 0.0006 ± 0.0011	2.8k	46 ADAM	05A	CLEO Repl. by MENDEZ 08
0.043 ± 0.008	44	TANENBAUM	76	MRK1 e^+e^-

- 43 Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.
- 44 Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.
- 45 Not independent from other measurements of MENDEZ 08.
- 46 Not independent from other values reported by ADAM 05A.

WEIGHTED AVERAGE
0.0296 ± 0.0031 (Error scaled by 1.8)




$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\text{anything})$

Γ_{13}/Γ_9

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0551 ± 0.0009 OUR FIT				
0.058 ± 0.007 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
0.050 ± 0.006 ± 0.003	298 ± 20	ANDREOTTI	05	E835 $\psi(2S) \rightarrow J/\psi X$
0.072 ± 0.009		AMBROGIANI	00A	E835 $p\bar{p} \rightarrow \psi(2S)$
0.061 ± 0.015		ARMSTRONG	97	E760 $\bar{p}p \rightarrow \psi(2S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				

0.0549 ± 0.0006 ± 0.0009 18.4k ⁴⁷ MENDEZ 08 CLEO $\psi(2S) \rightarrow \ell^+ \ell^- \eta$
 0.0546 ± 0.0010 ± 0.0007 2.8k ADAM 05A CLEO Repl. by MENDEZ 08

⁴⁷ Not independent from other measurements of MENDEZ 08.



$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{13}/Γ_{11}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.0976 ± 0.0016 OUR FIT
0.0979 ± 0.0018 OUR AVERAGE

0.0979 ± 0.0010 ± 0.0015	18.4k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
0.098 ± 0.005 ± 0.010	2k	⁴⁸ ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$
0.091 ± 0.021		⁴⁹ HIMEL	80	MRK2	$e^+ e^- \rightarrow \psi(2S) X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0968 ± 0.0019 ± 0.0013	2.8k	⁵⁰ ADAM	05A	CLEO	Repl. by MENDEZ 08
0.095 ± 0.007 ± 0.007		⁵¹ ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$

⁴⁸ From a fit to the J/ψ recoil mass spectra.

⁴⁹ The value for $B(\psi(2S) \rightarrow J/\psi(1S)\eta)$ reported in HIMEL 80 is derived using $B(\psi(2S)) \rightarrow J/\psi(1S)\pi^+\pi^- = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (0.1181 \pm 0.0020)$.

⁵⁰ Not independent from other values reported by ADAM 05A.

⁵¹ Not independent from other values reported by ANDREOTTI 05.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$

Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

13.0±1.0 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.


13.3±0.8±0.3	530	MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+ \ell^- 2\gamma$
14.3±1.4±1.2	280	BAI	04I	BES2 $\psi(2S) \rightarrow J/\psi\gamma\gamma$
14 ±6	7	HIMEL	80	MRK2 e^+e^-
9 ±2 ±1	23	⁵² OREGLIA	80	CBAL $\psi(2S) \rightarrow J/\psi 2\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

13 ±1 ±1	88	ADAM	05A	CLEO Repl. by MENDEZ 08
----------	----	------	-----	-------------------------

⁵² Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

WEIGHTED AVERAGE
13.0±1.0 (Error scaled by 1.4)



$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\text{anything})$

$$\Gamma_{14}/\Gamma_9 = \Gamma_{14}/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.344\Gamma_{110} + 0.195\Gamma_{111})$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.213±0.012±0.003	527	⁵³ MENDEZ	08	CLEO $e^+e^- \rightarrow J/\psi\gamma\gamma$
0.22 ±0.02 ±0.01		⁵⁴ ADAM	05A	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow J/\psi\gamma\gamma$

⁵³ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

⁵⁴ Not independent from other values reported by ADAM 05A.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{14}/Γ_{11}

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.380 \pm 0.022 \pm 0.005$	527	⁵⁵ MENDEZ	08	CLEO $e^+e^- \rightarrow J/\psi\gamma\gamma$
$0.39 \pm 0.04 \pm 0.01$		⁵⁶ ADAM	05A	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow J/\psi\gamma\gamma$

⁵⁵ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

⁵⁶ Not independent from other values reported by ADAM 05A.

————— **HADRONIC DECAYS** —————

$\Gamma(\pi^0 h_c(1P))/\Gamma_{total}$ Γ_{15}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

$8.4 \pm 1.3 \pm 1.0$	11k	ABLIKIM	10B	BES3 $\psi(2S) \rightarrow \pi^0 h_c$
---	-----	---------	-----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	92^{+23}_{-22}	ADAMS	09	CLEO $\psi(2S) \rightarrow 2\pi^+ 2\pi^- 2\pi^0$
seen	1282	DOBBS	08A	CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
seen	168 ± 40	ROSNER	05	CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$

$\Gamma(3(\pi^+\pi^-)\pi^0)/\Gamma_{total}$ Γ_{16}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

35 ± 16	6	FRANKLIN	83	MRK2 $e^+e^- \rightarrow$ hadrons
-------------------------------	---	----------	----	-----------------------------------

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{total}$ Γ_{17}/Γ


<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

28 ± 10 OUR AVERAGE		Error includes scale factor of 4.6. See the ideogram below.		
---	--	---	--	--

$24.9 \pm 0.7 \pm 3.6$	2173	ABLIKIM	07D	BES2 $e^+e^- \rightarrow \psi(2S)$
$127 \pm 12 \pm 2$	410	⁵⁷ AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0\gamma$
$26.1 \pm 0.7 \pm 3.0$	1703	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$

30 ± 8	42	FRANKLIN	83	MRK2 e^+e^-
------------	----	----------	----	---------------

⁵⁷ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma_{total}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (297 \pm 22 \pm 18) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.35 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(\rho a_2(1320)) / \Gamma_{\text{total}}$


Γ_{18} / Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.55 \pm 0.73 \pm 0.47$		112 ± 31	BAI	04C BES2	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.3	90		BAI	98J BES	$e^+ e^-$

$\Gamma(\rho \bar{\rho}) / \Gamma_{\text{total}}$

Γ_{19} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
(2.76 ± 0.12) OUR FIT				
(2.95 ± 0.23) OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
$3.36 \pm 0.09 \pm 0.25$	1618	ABLIKIM	07C BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \rho \bar{\rho}$
$2.87 \pm 0.12 \pm 0.15$	557	PEDLAR	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \rho \bar{\rho}$
1.4 ± 0.8	4	BRANDELIK	79C DASP	$e^+ e^- \rightarrow \psi(2S) \rightarrow \rho \bar{\rho}$
2.3 ± 0.7		FELDMAN	77 MRK1	$e^+ e^- \rightarrow \psi(2S) \rightarrow \rho \bar{\rho}$



$\Gamma(p\bar{p})/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_{19}/Γ_{11}

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
(8.2±0.4) OUR FIT			
6.98±0.49±0.97	BAI	01	BES $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$

$\Gamma(\Delta^{++}\bar{\Delta}^{--})/\Gamma_{\text{total}}$

Γ_{20}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
12.8±1.0±3.4	157	⁵⁸ BAI	01	BES $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

⁵⁸ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$

Γ_{21}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	⁵⁹ ABLIKIM	07H	BES2 $e^+e^- \rightarrow \psi(2S)$

⁵⁹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$

Γ_{22}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.49	90	⁶⁰ ABLIKIM	07H	BES2 $e^+e^- \rightarrow \psi(2S)$

⁶⁰ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

$\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$

Γ_{23}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.0±0.1±0.1	74.0	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ $p\bar{p}K^+\pi^-$

$\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$			Γ_{24}/Γ		
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.8±0.3±0.3	45.8	BRIERE 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+\pi^+\pi^-\pi^-$	

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$			Γ_{25}/Γ		
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.8±0.4±0.5	73.4	BRIERE 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}2(\pi^+\pi^-)$	

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$			Γ_{26}/Γ		
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT

(2.8±0.5) OUR AVERAGE Error includes scale factor of 2.6. See the ideogram below.


3.39±0.20±0.32	337	ABLIKIM	07C	BES	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
6.4 ±1.8 ±0.1		⁶¹ AUBERT	07BD	BABR	10.6 $e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$
3.28±0.23±0.25	208	PEDLAR	05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
1.81±0.20±0.27	80	⁶² BAI	01	BES	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4	90	FELDMAN	77	MRK1	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
-----	----	---------	----	------	---

⁶¹AUBERT 07BD reports $[\Gamma(\psi(2S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (15 \pm 4 \pm 1) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.35 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶²Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.




$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ **Γ_{27}/Γ**

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$25.7 \pm 4.4 \pm 6.8$	35	PEDLAR	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ **Γ_{28}/Γ**

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
22 ± 4 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
$23.5 \pm 3.6 \pm 3.2$	59	ABLIKIM	07C	BES $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
$26.3 \pm 3.5 \pm 2.1$	58	PEDLAR	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
$12 \pm 4 \pm 4$	8	⁶³ BAI	01	BES $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

⁶³ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.



$\Gamma(\Sigma(1385)^+\bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$ **Γ_{29}/Γ**

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$11 \pm 3 \pm 3$	14	⁶⁴ BAI	01	BES $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons


⁶⁴ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Xi^- \Xi^+)/\Gamma_{\text{total}}$

Γ_{30}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
18 ±6 OUR AVERAGE			Error includes scale factor of 2.8. See the ideogram below.		
30.3 ± 4.0 ± 3.2		67	ABLIKIM	07C	BES $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
23.8 ± 3.0 ± 2.1		63	PEDLAR	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
9.4 ± 2.7 ± 1.5		12	⁶⁵ BAI	01	BES $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<20	90		FELDMAN	77	MRK1 $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
⁶⁵ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.					

WEIGHTED AVERAGE
18±6 (Error scaled by 2.8)



$\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$

Γ_{31}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
27.5 ± 6.4 ± 6.1	19	PEDLAR	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

$\Gamma(\Xi(1530)^0 \Xi(1530)^0)/\Gamma_{\text{total}}$

Γ_{32}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 8.1	90	⁶⁶ BAI	01	BES $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<32	90	PEDLAR	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
⁶⁶ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.				

$\Gamma(\Omega^- \bar{\Omega}^+)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 7.3	90	⁶⁷ BAI	01 BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<16	90	PEDLAR	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
⁶⁷ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.				

$\Gamma(\pi^0 p \bar{p})/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
(1.50±0.08) OUR AVERAGE Error includes scale factor of 1.1.				
1.54±0.06±0.06	948	ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \pi^0 p \bar{p}$
1.32±0.10±0.15	256 ± 18	⁶⁸ ABLIKIM	05E BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow p \bar{p} \gamma \gamma$
1.4 ± 0.5	9	FRANKLIN	83 MRK2	$e^+ e^-$
⁶⁸ Computed using $B(\pi^0 \rightarrow \gamma \gamma) = (98.80 \pm 0.03)\%$.				

$\Gamma(N_1^*(1440) \bar{p} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
8.1±0.7±0.3	474	⁶⁹ ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \pi^0 p \bar{p}$
⁶⁹ From a fit of the $p \bar{p}$ and $p \pi^0$ mass distributions to a combination of $N_1^*(1440) \bar{p}$, $\pi^0 f_0(2100)$, and two other broad, unestablished resonances.				

$\Gamma(\pi^0 f_0(2100) \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.1±0.4±0.1	76	⁷⁰ ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \pi^0 p \bar{p}$
⁷⁰ From a fit of the $p \bar{p}$ and $p \pi^0$ mass distributions to a combination of $N_1^*(1440) \bar{p}$, $\pi^0 f_0(2100)$, and two other broad, unestablished resonances.				

$\Gamma(\eta p \bar{p})/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
(5.7±0.6) OUR AVERAGE				
5.6±0.6±0.3	154	ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \eta p \bar{p}$
5.8±1.1±0.7	44.8 ± 8.5	⁷¹ ABLIKIM	05E BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow p \bar{p} \gamma \gamma$
8 ± 3 ± 3	9.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p \bar{p} \pi^+ \pi^- \pi^0$
⁷¹ Computed using $B(\eta \rightarrow \gamma \gamma) = (39.43 \pm 0.26)\%$.				

$\Gamma(\eta f_0(2100) \rightarrow \eta p \bar{p})/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.2±0.4±0.1	31	⁷² ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \eta p \bar{p}$
⁷² From a fit of the $p \bar{p}$ and $p \eta$ distributions to a combination of $N^*(1535) \bar{p}$ and $\eta f_0(2100)$.				

$\Gamma(N^*(1535)\bar{p} \rightarrow \eta p \bar{p})/\Gamma_{\text{total}}$ Γ_{39}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.4 \pm 0.6 \pm 0.3$	123	⁷³ ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \eta p \bar{p}$

⁷³ From a fit of the $p\bar{p}$ and $p\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and $\eta f_0(2100)$.

$\Gamma(\omega p \bar{p})/\Gamma_{\text{total}}$ Γ_{40}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.69 ± 0.21 OUR AVERAGE				
$0.6 \pm 0.2 \pm 0.2$	21.2	BRIERE 05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$
$0.8 \pm 0.3 \pm 0.1$	14.9 ± 0.1	⁷⁴ BAI 03B	BES	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

⁷⁴ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(\phi p \bar{p})/\Gamma_{\text{total}}$ Γ_{41}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 0.24	90	BRIERE 05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.26	90	⁷⁵ BAI 03B	BES	$\psi(2S) \rightarrow K^+ K^- p \bar{p}$
----------	----	-----------------------	-----	--

⁷⁵ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(\pi^+ \pi^- p \bar{p})/\Gamma_{\text{total}}$ Γ_{42}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
(6.0 ± 0.4) OUR AVERAGE				
$5.9 \pm 0.2 \pm 0.4$	904.5	BRIERE 05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-$
8 ± 2		⁷⁶ TANENBAUM 78	MRK1	$e^+ e^-$

⁷⁶ Assuming entirely strong decay.

$\Gamma(p\bar{n}\pi^- \text{ or c.c.})/\Gamma_{\text{total}}$ Γ_{43}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
(2.48 ± 0.17) OUR AVERAGE				
$2.45 \pm 0.11 \pm 0.21$	851	ABLIKIM 06i	BES2	$e^+ e^- \rightarrow p\pi^- X$
$2.52 \pm 0.12 \pm 0.22$	849	ABLIKIM 06i	BES2	$e^+ e^- \rightarrow \bar{p}\pi^+ X$

$\Gamma(p\bar{n}\pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{44}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.18 \pm 0.50 \pm 0.50$	135 ± 21	ABLIKIM 06i	BES2	$e^+ e^- \rightarrow p\pi^- \pi^0 X$

$\Gamma(\eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{45}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.6	90	BRIERE 05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

$\Gamma(\eta\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ **Γ_{47}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$9.5 \pm 0.7 \pm 1.5$		77 BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadr
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$10.3 \pm 0.8 \pm 1.4$	201.7	78 BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ $\eta 3\pi(\eta \rightarrow \gamma\gamma)$
$8.1 \pm 1.4 \pm 1.6$	50.0	78 BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ $\eta 3\pi(\eta \rightarrow 3\pi)$

⁷⁷ Average of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow 3\pi$.

⁷⁸ Not independent from other values reported by BRIERE 05.

$\Gamma(2(\pi^+\pi^-\eta))/\Gamma_{\text{total}}$ **Γ_{48}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.2 \pm 0.6 \pm 0.1$	16	79 AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\eta)\eta\gamma$
⁷⁹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+\pi^-\eta)) \cdot B(\eta \rightarrow \gamma\gamma) = 1.2 \pm 0.7 \pm 0.1$ eV.				


$\Gamma(\eta'\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ **Γ_{49}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.5 \pm 1.6 \pm 1.3$	12.8	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadr

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{50}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
(7.3 ± 1.2)	OUR AVERAGE	Error includes scale factor of 2.1. See the ideogram below.		
$8.4 \pm 0.5 \pm 1.2$	386	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
$12.2 \pm 2.2 \pm 0.7$	37	80 AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
$8.2 \pm 0.5 \pm 0.7$	391	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ $2(\pi^+\pi^-\pi^0)$
$4.8 \pm 0.6 \pm 0.7$	100 ± 22	81 BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-\pi^0)$
⁸⁰ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow \omega\pi^+\pi^-) \cdot B(\omega \rightarrow 3\pi) = 2.69 \pm 0.73 \pm 0.16$ eV.				
⁸¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.				

WEIGHTED AVERAGE
7.3±1.2 (Error scaled by 2.1)



$\Gamma(b_1^\pm \pi^\mp)/\Gamma_{\text{total}}$

Γ_{51}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
(4.0±0.6) OUR AVERAGE Error includes scale factor of 1.1.				
5.1 ± 0.6 ± 0.8	202	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
4.18 ^{+0.43} _{-0.42} ± 0.92	170	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
3.2 ± 0.6 ± 0.5	61 ± 11	^{82,83} BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.2 ± 0.8 ± 1.0		⁸² BAI	99C BES	Repl. by BAI 03B
⁸² Assuming $B(b_1 \rightarrow \omega\pi)=1$.				
⁸³ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.				

$\Gamma(b_1^0 \pi^0)/\Gamma_{\text{total}}$

Γ_{52}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.35^{+0.47}_{-0.42} ± 0.40	45	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$

Γ_{53}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
(2.2±0.4) OUR AVERAGE					
2.3 ± 0.5 ± 0.4		57	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
2.05±0.41±0.38		62 ± 12	BAI	04C BES2	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.5	90		⁸⁴ BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
<1.7	90		BAI	98J BES	Repl. by BAI 03B
⁸⁴ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.					

$\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ **Γ_{54}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
(7.5±0.9) OUR AVERAGE Error includes scale factor of 1.9.				
10.9±1.9±0.2	85	⁸⁵ AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
7.1±0.3±0.4	817.2	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow K^+K^-\pi^+\pi^-$
16 ±4		⁸⁶ TANENBAUM	78 MRK1	e^+e^-
⁸⁵ AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (2.56 \pm 0.42 \pm 0.16) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.35 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
⁸⁶ Assuming entirely strong decay.				

$\Gamma(\rho^0K^+K^-)/\Gamma_{\text{total}}$ **Γ_{55}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.2±0.2±0.4	223.8	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow K^+K^-\pi^+\pi^-$

$\Gamma(K^*(892)^0\bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$ **Γ_{56}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.86±0.32±0.43	93 ± 16	BAI	04C		$\psi(2S) \rightarrow K^+K^-\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.2	90	BAI	98J BES		e^+e^-

$\Gamma(K^+K^-\pi^+\pi^-\eta)/\Gamma_{\text{total}}$ **Γ_{57}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.3±0.7±0.1	7	⁸⁷ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\eta\gamma$
⁸⁷ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+\pi)\eta) \cdot B(\eta \rightarrow \gamma\gamma) = 1.2 \pm 0.7 \pm 0.1$ eV.				

$\Gamma(K^+K^-2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ **Γ_{58}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.0±2.5±1.8	65	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$ **Γ_{60}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.0±1.8±2.1	⁸⁸ BAI	99C BES	e^+e^-
⁸⁸ Assuming $B(K_1(1270) \rightarrow K\rho) = 0.42 \pm 0.06$			

$\Gamma(K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ **Γ_{61}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.20±0.25±0.37	83 ± 9	ABLIKIM	050 BES2	$e^+e^- \rightarrow \psi(2S)$

$\Gamma(\rho^0 p\bar{p})/\Gamma_{\text{total}}$ **Γ_{62}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.5 ±0.1 ±0.2	61.1	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-$

$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{63}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.7 ± 2.5	TANENBAUM 78	MRK1	$e^+ e^-$

$\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ **Γ_{64}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
(2.4+0.6) OUR AVERAGE Error includes scale factor of 2.2.				
$2.2 \pm 0.2 \pm 0.2$	308	BRIERE 05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$
4.5 ± 1.0		TANENBAUM 78	MRK1	$e^+ e^-$

$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ **Γ_{65}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
(2.2+0.6) OUR AVERAGE Error includes scale factor of 1.4.				
$2.0 \pm 0.2 \pm 0.4$	285.5	BRIERE 05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$
4.2 ± 1.5		TANENBAUM 78	MRK1	$e^+ e^-$

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ **Γ_{66}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
12.6 ± 0.9 OUR AVERAGE				
$18.7 \pm 5.7 \pm 0.3$	32	⁸⁹ AUBERT 07AU	BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$
$11.7 \pm 1.0 \pm 1.5$	597	ABLIKIM 06G	BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
$12.7 \pm 0.5 \pm 1.0$	711.6	BRIERE 05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

⁸⁹AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (44 \pm 13 \pm 3) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.35 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega f_0(1710) \rightarrow \omega K^+ K^-)/\Gamma_{\text{total}}$ **Γ_{67}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.9 \pm 2.0 \pm 0.9$	19	ABLIKIM 06G	BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{68}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.6 \pm 1.3 \pm 1.8$	238	ABLIKIM 06G	BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{69}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$9.6 \pm 2.2 \pm 1.7$	133	ABLIKIM 06G	BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(K^*(892)^+ K^- \rho^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{70}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$7.3 \pm 2.2 \pm 1.4$	78	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(K^*(892)^0 K^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{71}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$6.1 \pm 1.3 \pm 1.2$	125	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(\eta K^+ K^-)/\Gamma_{\text{total}}$ Γ_{72}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.3	90	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{73}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
(1.85 ± 0.25) OUR AVERAGE		Error includes scale factor of 1.1.		
$2.38 \pm 0.37 \pm 0.29$	78	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
$1.9 \pm 0.3 \pm 0.3$	76.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
$1.5 \pm 0.3 \pm 0.2$	23.0 ± 5.2	⁹⁰ BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

⁹⁰ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{74}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
(3.5 ± 2.0) OUR AVERAGE		Error includes scale factor of 2.8.		
$5.45 \pm 0.42 \pm 0.87$	671	ABLIKIM	05H BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow 3(\pi^+ \pi^-)$
1.5 ± 1.0		⁹¹ TANENBAUM	78 MRK1	$e^+ e^-$

⁹¹ Assuming entirely strong decay.

$\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{75}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$7.3 \pm 0.4 \pm 0.6$	434.9	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_{76}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
(6.3 ± 0.7) OUR AVERAGE				
$6.3 \pm 0.6 \pm 0.3$		DOBBS	06A CLEO	$e^+ e^-$
10 ± 7		BRANDELIK	79C DASP	$e^+ e^-$
< 5	90	FELDMAN	77 MRK1	$e^+ e^-$

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$

Γ_{77}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

(5.4±0.5) OUR AVERAGE

5.8 ± 0.8 ± 0.4		DOBBS	06A CLEO	$e^+ e^-$
5.24 ± 0.47 ± 0.48	156 ± 14	⁹² BAI	04B BES2	$\psi(2S) \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$

⁹² Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6860 \pm 0.0027$.

$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$


Γ_{78}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

(1.68±0.26) OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

1.81 ± 0.18 ± 0.19	260 ± 19	⁹³ ABLIKIM	05J BES2	$e^+ e^- \rightarrow \psi(2S)$
1.88 ^{+0.16} _{-0.15} ± 0.28	194	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
0.85 ± 0.46	4	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow \text{hadrons}$

⁹³ From a PW analysis of $\psi(2S) \rightarrow \pi^+ \pi^- \pi^0$.



$\Gamma(\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

Γ_{79}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.94 ± 0.25 ^{+1.15} _{-0.34}	⁹⁴ ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0$
---	-----------------------	----------	--

⁹⁴ From a PW analysis of $\psi(2S) \rightarrow \pi^+ \pi^- \pi^0$.

$\Gamma(\rho(770)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

0.32±0.12 OUR AVERAGE Error includes scale factor of 1.8.

0.51±0.07±0.11			⁹⁵ ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(770)\pi \rightarrow \pi^+\pi^-\pi^0$
----------------	--	--	-----------------------	----------	---

0.24 ^{+0.08} _{-0.07} ±0.02		22	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
--	--	----	------	---------	-------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.83	90	1	FRANKLIN	83 MRK2	e^+e^-
-------	----	---	----------	---------	----------

<10	90		BARTEL	76 CNTR	e^+e^-
-----	----	--	--------	---------	----------

<10	90		⁹⁶ ABRAMS	75 MRK1	e^+e^-
-----	----	--	----------------------	---------	----------

⁹⁵ From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

⁹⁶ Final state $\rho^0\pi^0$.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

8 ±5 BRANDELIK 79C DASP e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.1	90	DOBBS	06A CLEO	$e^+e^- \rightarrow \psi(2S)$
------	----	-------	----------	-------------------------------

<5	90	FELDMAN	77 MRK1	e^+e^-
----	----	---------	---------	----------

$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<3.1 ⁹⁷ BAI 99C BES e^+e^-

⁹⁷ Assuming $B(K_1(1400) \rightarrow K^*\pi)=0.94 \pm 0.06$

$\Gamma(K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<2.96 90 1 FRANKLIN 83 MRK2 $e^+e^- \rightarrow$ hadrons

$\Gamma(K^+\bar{K}^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{84}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

(1.7±0.8-0.7) OUR AVERAGE

2.9 ^{+1.3} _{-1.7} ±0.4		9.6 ± 4.2	ABLIKIM	05I BES2	$e^+e^- \rightarrow \psi(2S)$
--	--	-----------	---------	----------	-------------------------------

1.3 ^{+1.0} _{-0.7} ±0.3		7	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
--	--	---	------	---------	-------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5.4	90	FRANKLIN	83 MRK2	$e^+e^- \rightarrow$ hadrons
------	----	----------	---------	------------------------------

$\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{85}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

10.9±2.0 OUR AVERAGE

13.3 ^{+2.4} _{-2.8} ±1.7	65.6 ± 9.0	ABLIKIM	05I BES2	$e^+e^- \rightarrow \psi(2S)$
---	------------	---------	----------	-------------------------------

9.2 ^{+2.7} _{-2.2} ±0.9	25	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
--	----	------	---------	-------------------------------

$\Gamma(K^+ \bar{K}^*(892)^- + \text{c.c.}) / \Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})$ $\Gamma_{84} / \Gamma_{85}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.16 ± 0.06 OUR AVERAGE			
0.22 ^{+0.10} _{-0.14}	ABLIKIM	05I	BES2 $e^+ e^- \rightarrow \psi(2S)$
0.14 ^{+0.08} _{-0.06}	ADAM	05	CLEO $e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\phi \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{86} / Γ

<u>VALUE (units 10⁻⁴)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
(1.17 ± 0.29) OUR AVERAGE				Error includes scale factor of 1.7.

2.43 ± 0.95 ± 0.04	10 ± 4	^{98,99} AUBERT	07AK	BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
0.9 ± 0.2 ± 0.1	47.6	BRIERE	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
1.5 ± 0.2 ± 0.2	51.5 ± 8.3	¹⁰⁰ BAI	03B	BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$

⁹⁸ AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.57 \pm 0.22 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.35 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁹⁹ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

¹⁰⁰ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(\phi f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{87} / Γ

<u>VALUE (units 10⁻⁴)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.68 ± 0.25 OUR AVERAGE				Error includes scale factor of 1.1.

1.45 ± 0.70 ± 0.03	6 ± 3	^{101,102} AUBERT	07AK	BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
0.6 ± 0.2 ± 0.1	18.4 ± 6.4	¹⁰³ BAI	03B	BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$

¹⁰¹ AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.34 \pm 0.16 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.35 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁰² Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

¹⁰³ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(2(K^+ K^-)) / \Gamma_{\text{total}}$ Γ_{88} / Γ

<u>VALUE (units 10⁻⁴)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.6 ± 0.1 ± 0.1	59.2	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$

$\Gamma(\phi K^+ K^-) / \Gamma_{\text{total}}$ Γ_{89} / Γ

<u>VALUE (units 10⁻⁴)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.70 ± 0.16 OUR AVERAGE				

0.8 ± 0.2 ± 0.1	36.8	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$
0.6 ± 0.2 ± 0.1	16.1 ± 5.0	¹⁰⁴ BAI	03B	BES $\psi(2S) \rightarrow 2(K^+ K^-)$

¹⁰⁴ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(2(K^+ K^-)\pi^0)/\Gamma_{\text{total}}$					Γ_{90}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.1±0.2±0.2	44.7	BRIERE	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)\pi^0$

$\Gamma(\phi\eta)/\Gamma_{\text{total}}$					Γ_{91}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
(2.8+1.0-0.8) OUR AVERAGE					
$2.0^{+1.5}_{-1.1} \pm 0.4$	6	ADAM	05	CLEO	$e^+ e^- \rightarrow \psi(2S)$
$3.3 \pm 1.1 \pm 0.5$	17	ABLIKIM	04k	BES	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\phi\eta')/\Gamma_{\text{total}}$					Γ_{92}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
3.1±1.4±0.7	8	¹⁰⁵ ABLIKIM	04k	BES	$e^+ e^- \rightarrow \psi(2S)$
¹⁰⁵ Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels.					

$\Gamma(\omega\eta')/\Gamma_{\text{total}}$					Γ_{93}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
3.2^{+2.4}_{-2.0}±0.7	4	¹⁰⁶ ABLIKIM	04k	BES	$e^+ e^- \rightarrow \psi(2S)$
¹⁰⁶ Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels.					

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$					Γ_{94}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
(2.1+0.6) OUR AVERAGE					
$2.5^{+1.2}_{-1.0} \pm 0.2$	14	ADAM	05	CLEO	$e^+ e^- \rightarrow \psi(2S)$
$1.87^{+0.68}_{-0.62} \pm 0.28$	14	ABLIKIM	04L	BES	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\rho\eta')/\Gamma_{\text{total}}$					Γ_{95}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.87^{+1.64}_{-1.11}±0.33	2	ABLIKIM	04L	BES	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\rho\eta)/\Gamma_{\text{total}}$					Γ_{96}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
(2.2+0.6) OUR AVERAGE Error includes scale factor of 1.1.					
$3.0^{+1.1}_{-0.9} \pm 0.2$	18	ADAM	05	CLEO	$e^+ e^- \rightarrow \psi(2S)$
$1.78^{+0.67}_{-0.62} \pm 0.17$	13	ABLIKIM	04L	BES	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\omega\eta)/\Gamma_{\text{total}}$					Γ_{97}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<1.1	90	ADAM	05	CLEO	$e^+ e^- \rightarrow \psi(2S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<3.1	90	ABLIKIM	04k	BES	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$					Γ_{98}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<0.4	90	ABLIKIM	04K	BES	$e^+e^- \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.7	90	ADAM	05	CLEO	$e^+e^- \rightarrow \psi(2S)$
$\Gamma(\eta_c\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{99}/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
<1.0	90	PEDLAR	07	CLEO	$e^+e^- \rightarrow \psi(2S)$
$\Gamma(p\bar{p}K^+K^-)/\Gamma_{\text{total}}$					Γ_{100}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.7 \pm 0.6 \pm 0.4$	30.1	BRIERE	05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$
$\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{101}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.81 \pm 0.11 \pm 0.14$	50	¹⁰⁷ ABLIKIM	08C	BES2	$e^+e^- \rightarrow J/\psi$
¹⁰⁷ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.					
$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$					Γ_{102}/Γ
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$0.44 \pm 0.12 \pm 0.11$	20 ± 6	BAI	04C		$\psi(2S) \rightarrow 2(K^+K^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.45	90	BAI	98J	BES	$e^+e^- \rightarrow 2(K^+K^-)$
$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{103}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<0.88	90	BAI	04G	BES2	e^+e^-
$\Gamma(\Theta(1540)K^-\bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$					Γ_{104}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<1.0	90	BAI	04G	BES2	e^+e^-
$\Gamma(\Theta(1540)K_S^0\bar{p} \rightarrow K_S^0\bar{p}K^+n)/\Gamma_{\text{total}}$					Γ_{105}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<0.70	90	BAI	04G	BES2	e^+e^-
$\Gamma(\bar{\Theta}(1540)K^+n \rightarrow K_S^0\bar{p}K^+n)/\Gamma_{\text{total}}$					Γ_{106}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<2.6	90	BAI	04G	BES2	e^+e^-
$\Gamma(\bar{\Theta}(1540)K_S^0p \rightarrow K_S^0pK^-\bar{n})/\Gamma_{\text{total}}$					Γ_{107}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<0.60	90	BAI	04G	BES2	e^+e^-

$\Gamma(K_S^0 \bar{K}_S^0)/\Gamma_{\text{total}}$					Γ_{108}/Γ
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT		
<0.046	¹⁰⁸ BAI	04D	BES	$e^+ e^-$	

¹⁰⁸ Forbidden by CP.

————— **RADIATIVE DECAYS** —————

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$					Γ_{109}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
9.68±0.31 OUR FIT					
9.2 ±0.4 OUR AVERAGE					
9.22±0.11±0.46	72600	ATHAR	04	CLEO	$e^+ e^- \rightarrow \gamma X$
9.9 ±0.5 ±0.8		¹⁰⁹ GAISER	86	CBAL	$e^+ e^- \rightarrow \gamma X$
7.2 ±2.3		¹⁰⁹ BIDDICK	77	CNTR	$e^+ e^- \rightarrow \gamma X$
7.5 ±2.6		¹⁰⁹ WHITAKER	76	MRK1	$e^+ e^-$

¹⁰⁹ Angular distribution $(1+\cos^2\theta)$ assumed.

$\Gamma(\gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$					Γ_{110}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
9.2 ±0.4 OUR FIT					
8.9 ±0.5 OUR AVERAGE					
9.07±0.11±0.54	76700	ATHAR	04	CLEO	$e^+ e^- \rightarrow \gamma X$
9.0 ±0.5 ±0.7		¹¹⁰ GAISER	86	CBAL	$e^+ e^- \rightarrow \gamma X$
7.1 ±1.9		¹¹¹ BIDDICK	77	CNTR	$e^+ e^- \rightarrow \gamma X$

¹¹⁰ Angular distribution $(1-0.189 \cos^2\theta)$ assumed.
¹¹¹ Valid for isotropic distribution of the photon.

$\Gamma(\gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$					Γ_{111}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
8.75±0.35 OUR FIT					
8.8 ±0.5 OUR AVERAGE					Error includes scale factor of 1.1.
9.33±0.14±0.61	79300	ATHAR	04	CLEO	$e^+ e^- \rightarrow \gamma X$
8.0 ±0.5 ±0.7		¹¹² GAISER	86	CBAL	$e^+ e^- \rightarrow \gamma X$
7.0 ±2.0		¹¹³ BIDDICK	77	CNTR	$e^+ e^- \rightarrow \gamma X$

¹¹² Angular distribution $(1-0.052 \cos^2\theta)$ assumed.
¹¹³ Valid for isotropic distribution of the photon.

$[\Gamma(\gamma\chi_{c0}(1P)) + \Gamma(\gamma\chi_{c1}(1P)) + \Gamma(\gamma\chi_{c2}(1P))]/\Gamma_{\text{total}}$					$(\Gamma_{109}+\Gamma_{110}+\Gamma_{111})/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
27.6±0.3±2.0	¹¹⁴ ATHAR	04	CLEO	$e^+ e^- \rightarrow \gamma X$	

¹¹⁴ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c1}(1P))$					$\Gamma_{109}/\Gamma_{110}$
VALUE	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.02±0.01±0.07	¹¹⁵ ATHAR	04	CLEO	$e^+ e^- \rightarrow \gamma X$	

¹¹⁵ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$\Gamma(\gamma\chi_{c2}(1P))/\Gamma(\gamma\chi_{c1}(1P))$

$\Gamma_{111}/\Gamma_{110}$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.03 \pm 0.02 \pm 0.03$ ¹¹⁶ ATHAR 04 CLEO $e^+e^- \rightarrow \gamma X$

¹¹⁶ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c2}(1P))$

$\Gamma_{109}/\Gamma_{111}$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.99 \pm 0.02 \pm 0.08$ ¹¹⁷ ATHAR 04 CLEO $e^+e^- \rightarrow \gamma X$

¹¹⁷ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$

Γ_{112}/Γ

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

0.34 ± 0.05 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.


$0.432 \pm 0.016 \pm 0.060$ MITCHELL 09 CLEO $e^+e^- \rightarrow \gamma X$

$0.32 \pm 0.04 \pm 0.06$ 2560 ¹¹⁸ ATHAR 04 CLEO $e^+e^- \rightarrow \gamma X$

0.28 ± 0.06 ¹¹⁹ GAISER 86 CBAL $e^+e^- \rightarrow \gamma X$

¹¹⁸ ATHAR 04 used $\Gamma_{\eta_c(1S)} = 24.8 \pm 4.9$ MeV to obtain this result.

¹¹⁹ GAISER 86 used $\Gamma_{\eta_c(1S)} = 11.5 \pm 4.5$ MeV to obtain this result.



$\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$ **Γ_{113}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 8 \times 10^{-4}$	90	120 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K \bar{K} \pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 2 \times 10^{-3}$	90	ATHAR	04	CLEO $e^+ e^- \rightarrow \gamma X$
$0.2-1.3 \times 10^{-2}$	95	EDWARDS	82C	CBAL $e^+ e^- \rightarrow \gamma X$

¹²⁰ CRONIN-HENNESSY 10 reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K \bar{K} \pi)] < 14.5 \times 10^{-6}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K \bar{K} \pi) = 1.9 \times 10^{-2}$. This measurement assumes $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$ **Γ_{114}/Γ**

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

$1.58 \pm 0.40 \pm 0.13$		37	ABLIKIM	10F	BES3 $\psi(2S) \rightarrow \gamma\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 5	90		PEDLAR	09	CLE3 $\psi(2S) \rightarrow \gamma X$
< 5400	95	121	LIBERMAN	75	SPEC $e^+ e^-$
$< 1 \times 10^4$	90		WIIK	75	DASP $e^+ e^-$

¹²¹ Restated by us using $B(\psi(2S) \rightarrow \mu^+ \mu^-) = 0.0077$.

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ **Γ_{115}/Γ**

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

(1.23 ± 0.06) OUR AVERAGE					
$1.26 \pm 0.03 \pm 0.08$		2226	122 ABLIKIM	10F	BES3 $\psi(2S) \rightarrow 3\gamma\pi^+\pi^-$, $2\gamma\pi^+\pi^-$
$1.19 \pm 0.08 \pm 0.03$			PEDLAR	09	CLE3 $\psi(2S) \rightarrow \gamma X$
$1.24 \pm 0.27 \pm 0.15$		23	ABLIKIM	06R	BES2 $e^+ e^- \rightarrow \psi(2S)$
$1.54 \pm 0.31 \pm 0.20$		~ 43	BAI	98F	BES $\psi(2S) \rightarrow \pi^+\pi^-2\gamma$, $\pi^+\pi^-3\gamma$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

< 60	90	123	BRAUNSCH...	77	DASP $e^+ e^-$
< 11	90	124	BARTEL	76	CNTR $e^+ e^-$

¹²² Combining the results from $\eta' \rightarrow \pi^+\pi^-\eta$ and $\eta' \rightarrow \pi^+\pi^-\gamma$ decay modes.

¹²³ Restated by us using total decay width 228 keV.

¹²⁴ The value is normalized to the branching ratio for $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$.

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ **Γ_{116}/Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$2.12 \pm 0.19 \pm 0.32$		125,126	BAI	03C BES $\psi(2S) \rightarrow \gamma\pi\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$2.08 \pm 0.19 \pm 0.33$	200.6 ± 18.8	125	BAI	03C BES $\psi(2S) \rightarrow \gamma\pi^+\pi^-$
$2.90 \pm 1.08 \pm 1.07$	29.9 ± 11.1	125	BAI	03C BES $\psi(2S) \rightarrow \gamma\pi^0\pi^0$

¹²⁵ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

¹²⁶ Combining the results from $\pi^+\pi^-$ and $\pi^0\pi^0$ decay modes.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{118} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.301 ± 0.041 ± 0.124	35.6 ± 4.8	127 BAI	03C BES	$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
127 Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.				

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{119} / Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.604 ± 0.090 ± 0.132		39.6 ± 5.9 ^{128,129}	BAI	03C BES	$\psi(2S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.56		90 6.8 ± 3.1 ^{128,129}	BAI	03C BES	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
128 Includes unknown branching fractions to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied the $K^+ K^-$ result by a factor of 2 and the $K_S^0 K_S^0$ result by a factor of 4 to obtain the $K \bar{K}$ result.					
129 Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.					

$\Gamma(\gamma \eta) / \Gamma_{\text{total}}$ Γ_{121} / Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.38 ± 0.48 ± 0.09		13 130	ABLIKIM	10F BES3	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0, \gamma 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 2		90	PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$
< 90		90	BAI	98F BES	$\psi(2S) \rightarrow \pi^+ \pi^- 3\gamma$
< 200		90	YAMADA	77 DASP	$e^+ e^- \rightarrow 3\gamma$
130 Combining the results from $\eta \rightarrow \pi^+ \pi^- \pi^0$ and $\eta \rightarrow 3\pi^0$ decay modes.					

$\Gamma(\gamma \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{122} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.71 ± 1.25 ± 1.64	418	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

$\Gamma(\gamma \eta(1405) \rightarrow \gamma K \bar{K} \pi) / \Gamma_{\text{total}}$ Γ_{124} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.9		90 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 1.3		90 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
< 1.2		90 131 SCHARRE	80 MRK1	$e^+ e^-$
131 Includes unknown branching fraction $\eta(1405) \rightarrow K \bar{K} \pi$.				

$\Gamma(\gamma \eta(1405) \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{125} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.36 ± 0.25 ± 0.05	10	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

$\Gamma(\gamma \eta(1475) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}$ Γ_{127} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.4		90 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 1.5		90 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$

$\Gamma(\gamma\eta(1475) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{128}/Γ

<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.88	90	ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

$\Gamma(\gamma 2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{129}/Γ

<u>VALUE (units 10⁻⁵)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
39.6±2.8±5.0	583	ABLIKIM	07D	BES2 $e^+e^- \rightarrow \psi(2S)$

$\Gamma(\gamma K^{*0}K^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{130}/Γ

<u>VALUE (units 10⁻⁵)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
37.0±6.1±7.2	237	ABLIKIM	07D	BES2 $e^+e^- \rightarrow \psi(2S)$

$\Gamma(\gamma K^{*0}\bar{K}^{*0})/\Gamma_{\text{total}}$ Γ_{131}/Γ

<u>VALUE (units 10⁻⁵)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
24.0±4.5±5.0	41	ABLIKIM	07D	BES2 $e^+e^- \rightarrow \psi(2S)$

$\Gamma(\gamma K_S^0 K^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{132}/Γ

<u>VALUE (units 10⁻⁵)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
25.6±3.6±3.6	115	ABLIKIM	07D	BES2 $e^+e^- \rightarrow \psi(2S)$

$\Gamma(\gamma K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{133}/Γ

<u>VALUE (units 10⁻⁵)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
19.1±2.7±4.3	132	ABLIKIM	07D	BES2 $e^+e^- \rightarrow \psi(2S)$

$\Gamma(\gamma\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{134}/Γ

<u>VALUE (units 10⁻⁵)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
(3.9±0.5) OUR AVERAGE				Error includes scale factor of 2.0.
4.18±0.26±0.18	348	¹³² ALEXANDER	10	CLEO $\psi(2S) \rightarrow \gamma\rho\bar{\rho}$
2.9 ±0.4 ±0.4	142	ABLIKIM	07D	BES2 $e^+e^- \rightarrow \psi(2S)$

¹³² From a fit of the $\rho\bar{\rho}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma\rho\bar{\rho}$ phase space, for $M(\rho\bar{\rho}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0\rho\bar{\rho}$ and continuum.

$\Gamma(\gamma f_2(1950) \rightarrow \gamma\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{135}/Γ

<u>VALUE (units 10⁻⁵)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.2±0.2±0.1	111	¹³³ ALEXANDER	10	CLEO $\psi(2S) \rightarrow \gamma\rho\bar{\rho}$

¹³³ From a fit of the $\rho\bar{\rho}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma\rho\bar{\rho}$ phase space, for $M(\rho\bar{\rho}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0\rho\bar{\rho}$ and continuum.

$\Gamma(\gamma f_2(2150) \rightarrow \gamma\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{136}/Γ

<u>VALUE (units 10⁻⁵)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.72±0.18±0.03	73	¹³⁴ ALEXANDER	10	CLEO $\psi(2S) \rightarrow \gamma\rho\bar{\rho}$

¹³⁴ From a fit of the $\rho\bar{\rho}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma\rho\bar{\rho}$ phase space, for $M(\rho\bar{\rho}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0\rho\bar{\rho}$ and continuum.

$\Gamma(\gamma X(1835) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{137}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<5.4	90	ABLIKIM 07D	BES	$\psi(2S) \rightarrow \gamma p \bar{p}$

$\Gamma(\gamma X \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{138}/Γ

For a narrow resonance in the range $2.2 < M(X) < 2.8$ GeV.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$

$\Gamma(\gamma \pi^+ \pi^- p \bar{p})/\Gamma_{\text{total}}$ Γ_{139}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.8 \pm 1.2 \pm 0.7$	17	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\gamma 2(\pi^+ \pi^-) K^+ K^-)/\Gamma_{\text{total}}$ Γ_{140}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<22	90	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{141}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<17	90	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\gamma K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{142}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

$\psi(2S)$ CROSS-PARTICLE BRANCHING RATIOS

For measurements involving $B(\psi(2S) \rightarrow \gamma \chi_{cJ}(1P)) \times B(\chi_{cJ}(1P) \rightarrow X)$ see the corresponding entries in the $\chi_{cJ}(1P)$ sections.

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

$\psi(2S) \rightarrow \gamma \chi_{cJ}(1P)$ and $\chi_{cJ} \rightarrow \gamma J/\psi(1S)$

$a_2(\chi_{c1})/a_2(\chi_{c2})$ Magnetic quadrupole transition amplitude ratio

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$\frac{67^{+19}}{-13}$	59k	¹³⁵ ARTUSO 09	CLEO	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

¹³⁵ Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $a_2(\chi_{c1}(1P))$ and $a_2(\chi_{c2}(1P))$ from ARTUSO 09.

$b_2(\chi_{c2})/b_2(\chi_{c1})$ Magnetic quadrupole transition amplitude ratio

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
37^{+53}_{-47}	59k	¹³⁶ ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

¹³⁶ Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $b_2(\chi_{c1}(1P))$ and $b_2(\chi_{c2}(1P))$ from ARTUSO 09.

 $\psi(2S)$ REFERENCES

ABLIKIM	10B	PRL 104 132002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	10F	PRL 105 261801	M. Ablikim <i>et al.</i>	(BES III Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN...	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)
ADAMS	09	PR D80 051106	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)
LIBBY	09	PR D80 072002	J. Libby <i>et al.</i>	(CLEO Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)
DOBBS	08A	PRL 101 182003	S. Dobbs <i>et al.</i>	(CLEO Collab.)
MENDEZ	08	PR D78 011102R	H. Mendez <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ABLIKIM	07C	PL B648 149	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07D	PRL 99 011802	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ANASHIN	07	JETPL 85 347	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
		Translated from ZETFP 85 429.		
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)
PDG	07	Unofficial 2007 WWW edition		(PDG Collab.)
PEDLAR	07	PR D75 011102R	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	06G	PR D73 052004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06W	PR D74 112003	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT, BE	06D	PR D74 091103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
DOBBS	06A	PR D74 011105R	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ABLIKIM	05E	PR D71 072006	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05I	PL B614 37	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05J	PL B619 247	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	05	PRL 94 012005	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05	PR D71 032006	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
BRIERE	05	PRL 95 062001	R.A. Briere <i>et al.</i>	(CLEO Collab.)
PEDLAR	05	PR D72 051108R	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ROSNER	05	PRL 95 102003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04K	PR D70 112003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04L	PR D70 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BAI	04B	PRL 92 052001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04C	PR D69 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)

BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
SETH	04	PR D69 097503	K.K. Seth	
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03B	PR D67 052002	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)
AUBERT	02B	PR D65 031101R	B. Aubert <i>et al.</i>	(BaBar Collab.)
BAI	02	PR D65 052004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	02B	PL B550 24	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
PDG	02	PR D66 010001	K. Hagiwara <i>et al.</i>	
BAI	01	PR D63 032002	J.Z. Bai <i>et al.</i>	(BES Collab.)
AMBROGIANI	00A	PR D62 032004	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98F	PR D58 097101	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98J	PRL 81 5080	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARMSTRONG	97	PR D55 1153	T.A. Armstrong <i>et al.</i>	(E760 Collab.)
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 Collab., E706 Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)
OREGLIA	80	PRL 45 959	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)
BRAUNSCH...	77	PL 67B 249	W. Braunschweig <i>et al.</i>	(DASP Collab.)
BURMESTER	77	PL 66B 395	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	76	PRL 36 402	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL) IG
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)
ABRAMS	75	Stanford Symp. 25	G.S. Abrams	(LBL)
ABRAMS	75B	PRL 34 1181	G.S. Abrams <i>et al.</i>	(LBL, SLAC)
BOYARSKI	75C	Palermo Conf. 54	A.M. Boyarski <i>et al.</i>	(SLAC, LBL)
HILGER	75	PRL 35 625	E. Hilger <i>et al.</i>	(STAN, PENN)
LIBERMAN	75	Stanford Symp. 55	A.D. Liberman	(STAN)
LUTH	75	PRL 35 1124	V. Luth <i>et al.</i>	(SLAC, LBL) JPC
WIJK	75	Stanford Symp. 69	B.H. Wiik	(DESY)