

ψ(3770)

$$I^G(J^{PC}) = 0^-(1^{--})$$

ψ(3770) MASS (MeV)

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

3772.92 ± 0.35 OUR FIT Error includes scale factor of 1.1.


3775.2 ± 1.7 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

3772.0 ± 1.9		¹ ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
3775.5 ± 2.4 ± 0.5	57	AUBERT	08B BABR	$B \rightarrow D\bar{D}K$
3776 ± 5 ± 4	68	BRODZICKA	08 BELL	$B^+ \rightarrow D^0\bar{D}^0 K^+$
3778.8 ± 1.9 ± 0.9		AUBERT	07BE BABR	$e^+e^- \rightarrow D\bar{D}\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3778.4 ± 3.0 ± 1.3	34	CHISTOV	04 BELL	Sup. by BRODZICKA 08
--------------------	----	---------	---------	----------------------

¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.



$m_{\psi(3770)} - m_{\psi(2S)}$


OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
86.83 ± 0.35 OUR FIT	Error includes scale factor of 1.1.		
86.6 ± 0.7 OUR AVERAGE	Error includes scale factor of 2.0. See the ideogram below.		
86.9 ± 0.4	² ABLIKIM	07E	BES2 $e^+e^- \rightarrow$ hadrons
86.7 ± 0.7	ABLIKIM	06L	BES2 $e^+e^- \rightarrow$ hadrons
80 ± 2	SCHINDLER	80	MRK2 e^+e^-
86 ± 2	³ BACINO	78	DLCO e^+e^-
88 ± 3	RAPIDIS	77	LGW e^+e^-

² BES-II $\psi(2S)$ mass subtracted (see ABLIKIM 06L).

³ SPEAR $\psi(2S)$ mass subtracted (see SCHINDLER 80).

WEIGHTED AVERAGE
86.6 ± 0.7 (Error scaled by 2.0)



$\psi(3770)$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
27.3 ± 1.0 OUR FIT				
27.6 ± 1.0 OUR AVERAGE				
30.4 ± 8.5		⁴ ABLIKIM	08D	BES2 $e^+e^- \rightarrow$ hadrons
27 ± 10 ± 5	68	BRODZICKA	08	BELL $B^+ \rightarrow D^0 \bar{D}^0 K^+$
28.5 ± 1.2 ± 0.2		ABLIKIM	07E	BES2 $e^+e^- \rightarrow$ hadrons
23.5 ± 3.7 ± 0.9		AUBERT	07BE	BABR $e^+e^- \rightarrow D \bar{D} \gamma$

$26.9 \pm 2.4 \pm 0.3$	ABLIKIM	06L	BES2	$e^+ e^- \rightarrow \text{hadrons}$
24 ± 5	SCHINDLER	80	MRK2	$e^+ e^-$
24 ± 5	BACINO	78	DLCO	$e^+ e^-$
28 ± 5	RAPIDIS	77	LGW	$e^+ e^-$

⁴ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.

$\psi(3770)$ DECAY MODES

In addition to the dominant decay mode to $D\bar{D}$, $\psi(3770)$ was found to decay into the final states containing the J/ψ (BAI 05, ADAM 06). ADAMS 06 and HUANG 06A searched for various decay modes with light hadrons and found a statistically significant signal for the decay to $\phi\eta$ only (ADAMS 06).

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $D\bar{D}$	$(93 \pm 8 \text{ } ^{+8}_{-9})\%$	S=2.0
Γ_2 $D^0\bar{D}^0$	$(52 \pm 5)\%$	S=2.0
Γ_3 D^+D^-	$(41 \pm 4)\%$	S=2.0
Γ_4 $J/\psi\pi^+\pi^-$	$(1.93 \pm 0.28) \times 10^{-3}$	
Γ_5 $J/\psi\pi^0\pi^0$	$(8.0 \pm 3.0) \times 10^{-4}$	
Γ_6 $J/\psi\eta$	$(9 \pm 4) \times 10^{-4}$	
Γ_7 $J/\psi\pi^0$	$< 2.8 \times 10^{-4}$	CL=90%
Γ_8 e^+e^-	$(9.7 \pm 0.7) \times 10^{-6}$	S=1.2

Decays to light hadrons

Γ_9 $b_1(1235)\pi$	$< 1.4 \times 10^{-5}$	CL=90%
Γ_{10} $\phi\eta'$	$< 7 \times 10^{-4}$	CL=90%
Γ_{11} $\omega\eta'$	$< 4 \times 10^{-4}$	CL=90%
Γ_{12} $\rho^0\eta'$	$< 6 \times 10^{-4}$	CL=90%
Γ_{13} $\phi\eta$	$(3.1 \pm 0.7) \times 10^{-4}$	
Γ_{14} $\omega\eta$	$< 1.4 \times 10^{-5}$	CL=90%
Γ_{15} $\rho^0\eta$	$< 5 \times 10^{-4}$	CL=90%
Γ_{16} $\phi\pi^0$	$< 3 \times 10^{-5}$	CL=90%
Γ_{17} $\omega\pi^0$	$< 6 \times 10^{-4}$	CL=90%
Γ_{18} $\pi^+\pi^-\pi^0$	$< 5 \times 10^{-6}$	CL=90%
Γ_{19} $\rho\pi$	$< 5 \times 10^{-6}$	CL=90%
Γ_{20} $K^*(892)^+K^- + \text{c.c.}$	$< 1.4 \times 10^{-5}$	CL=90%
Γ_{21} $K^*(892)^0\bar{K}^0 + \text{c.c.}$	$< 1.2 \times 10^{-3}$	CL=90%
Γ_{22} $K_S^0 K_L^0$	$< 1.2 \times 10^{-5}$	CL=90%
Γ_{23} $2(\pi^+\pi^-)$	$< 1.12 \times 10^{-3}$	CL=90%
Γ_{24} $2(\pi^+\pi^-\pi^0)$	$< 1.06 \times 10^{-3}$	CL=90%
Γ_{25} $2(\pi^+\pi^-\pi^0)$	$< 5.85\%$	CL=90%
Γ_{26} $\omega\pi^+\pi^-$	$< 6.0 \times 10^{-4}$	CL=90%

Γ_{27}	$3(\pi^+\pi^-)$	< 9.1	$\times 10^{-3}$	
Γ_{28}	$3(\pi^+\pi^-)\pi^0$	< 1.37	%	
Γ_{29}	$3(\pi^+\pi^-)2\pi^0$	< 11.74	%	CL=90%
Γ_{30}	$\eta\pi^+\pi^-$	< 1.24	$\times 10^{-3}$	CL=90%
Γ_{31}	$\pi^+\pi^-2\pi^0$	< 8.9	$\times 10^{-3}$	CL=90%
Γ_{32}	$\rho^0\pi^+\pi^-$	< 6.9	$\times 10^{-3}$	CL=90%
Γ_{33}	$\eta3\pi$	< 1.34	$\times 10^{-3}$	CL=90%
Γ_{34}	$\eta2(\pi^+\pi^-)$	< 2.43	%	
Γ_{35}	$\eta\rho^0\pi^+\pi^-$	< 1.45	%	CL=90%
Γ_{36}	$\eta'3\pi$	< 2.44	$\times 10^{-3}$	CL=90%
Γ_{37}	$K^+K^-\pi^+\pi^-$	< 9.0	$\times 10^{-4}$	CL=90%
Γ_{38}	$\phi\pi^+\pi^-$	< 4.1	$\times 10^{-4}$	CL=90%
Γ_{39}	$K^+K^-2\pi^0$	< 4.2	$\times 10^{-3}$	CL=90%
Γ_{40}	$4(\pi^+\pi^-)$	< 1.67	%	CL=90%
Γ_{41}	$4(\pi^+\pi^-)\pi^0$	< 3.06	%	CL=90%
Γ_{42}	$\phi f_0(980)$	< 4.5	$\times 10^{-4}$	CL=90%
Γ_{43}	$K^+K^-\pi^+\pi^-\pi^0$	< 2.36	$\times 10^{-3}$	CL=90%
Γ_{44}	$K^+K^-\rho^0\pi^0$	< 8	$\times 10^{-4}$	CL=90%
Γ_{45}	$K^+K^-\rho^+\pi^-$	< 1.46	%	CL=90%
Γ_{46}	ωK^+K^-	< 3.4	$\times 10^{-4}$	CL=90%
Γ_{47}	$\phi\pi^+\pi^-\pi^0$	< 3.8	$\times 10^{-3}$	CL=90%
Γ_{48}	$K^{*0}K^-\pi^+\pi^0 + \text{c.c.}$	< 1.62	%	CL=90%
Γ_{49}	$K^{*+}K^-\pi^+\pi^- + \text{c.c.}$	< 3.23	%	CL=90%
Γ_{50}	$K^+K^-\pi^+\pi^-2\pi^0$	< 2.67	%	CL=90%
Γ_{51}	$K^+K^-2(\pi^+\pi^-)$	< 1.03	%	CL=90%
Γ_{52}	$K^+K^-2(\pi^+\pi^-)\pi^0$	< 3.60	%	CL=90%
Γ_{53}	ηK^+K^-	< 4.1	$\times 10^{-4}$	CL=90%
Γ_{54}	$\eta K^+K^-\pi^+\pi^-$	< 1.24	%	CL=90%
Γ_{55}	$\rho^0 K^+K^-$	< 5.0	$\times 10^{-3}$	CL=90%
Γ_{56}	$2(K^+K^-)$	< 6.0	$\times 10^{-4}$	CL=90%
Γ_{57}	ϕK^+K^-	< 7.5	$\times 10^{-4}$	CL=90%
Γ_{58}	$2(K^+K^-)\pi^0$	< 2.9	$\times 10^{-4}$	CL=90%
Γ_{59}	$2(K^+K^-)\pi^+\pi^-$	< 3.2	$\times 10^{-3}$	CL=90%
Γ_{60}	$K_S^0 K^-\pi^+$	< 3.2	$\times 10^{-3}$	CL=90%
Γ_{61}	$K_S^0 K^-\pi^+\pi^0$	< 1.33	%	CL=90%
Γ_{62}	$K_S^0 K^-\rho^+$	< 6.6	$\times 10^{-3}$	CL=90%
Γ_{63}	$K_S^0 K^-\pi^+\pi^-$	< 8.7	$\times 10^{-3}$	CL=90%
Γ_{64}	$K_S^0 K^-\pi^+\rho^0$	< 1.6	%	CL=90%
Γ_{65}	$K_S^0 K^-\pi^+\eta$	< 1.3	%	CL=90%
Γ_{66}	$K_S^0 K^-\pi^+\pi^-\pi^0$	< 4.18	%	CL=90%
Γ_{67}	$K_S^0 K^-\pi^+\pi^-\eta$	< 4.8	%	CL=90%
Γ_{68}	$K_S^0 K^-\pi^+2(\pi^+\pi^-)$	< 1.22	%	CL=90%
Γ_{69}	$K_S^0 K^-\pi^+2\pi^0$	< 2.65	%	CL=90%

Γ_{70}	$K_S^0 K^- K^+ K^- \pi^+$	< 4.9	$\times 10^{-3}$	CL=90%
Γ_{71}	$K_S^0 K^- K^+ K^- \pi^+ \pi^0$	< 3.0	%	CL=90%
Γ_{72}	$K_S^0 K^- K^+ K^- \pi^+ \eta$	< 2.2	%	CL=90%
Γ_{73}	$K^{*0} K^- \pi^+ + \text{c.c.}$	< 9.7	$\times 10^{-3}$	CL=90%
Γ_{74}	$\rho \bar{p} \pi^0$	< 1.2	$\times 10^{-3}$	
Γ_{75}	$\rho \bar{p} \pi^+ \pi^-$	< 5.8	$\times 10^{-4}$	CL=90%
Γ_{76}	$\Lambda \bar{\Lambda}$	< 1.2	$\times 10^{-4}$	CL=90%
Γ_{77}	$\rho \bar{p} \pi^+ \pi^- \pi^0$	< 1.85	$\times 10^{-3}$	CL=90%
Γ_{78}	$\omega \rho \bar{p}$	< 2.9	$\times 10^{-4}$	CL=90%
Γ_{79}	$\Lambda \bar{\Lambda} \pi^0$	< 1.2	$\times 10^{-3}$	CL=90%
Γ_{80}	$\rho \bar{p} 2(\pi^+ \pi^-)$	< 2.6	$\times 10^{-3}$	CL=90%
Γ_{81}	$\eta \rho \bar{p}$	< 5.4	$\times 10^{-4}$	CL=90%
Γ_{82}	$\eta \rho \bar{p} \pi^+ \pi^-$	< 3.3	$\times 10^{-3}$	CL=90%
Γ_{83}	$\rho^0 \rho \bar{p}$	< 1.7	$\times 10^{-3}$	CL=90%
Γ_{84}	$\rho \bar{p} K^+ K^-$	< 3.2	$\times 10^{-4}$	CL=90%
Γ_{85}	$\eta \rho \bar{p} K^+ K^-$	< 6.9	$\times 10^{-3}$	CL=90%
Γ_{86}	$\pi^0 \rho \bar{p} K^+ K^-$	< 1.2	$\times 10^{-3}$	CL=90%
Γ_{87}	$\phi \rho \bar{p}$	< 1.3	$\times 10^{-4}$	CL=90%
Γ_{88}	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	< 2.5	$\times 10^{-4}$	CL=90%
Γ_{89}	$\Lambda \bar{p} K^+$	< 2.8	$\times 10^{-4}$	CL=90%
Γ_{90}	$\Lambda \bar{p} K^+ \pi^+ \pi^-$	< 6.3	$\times 10^{-4}$	CL=90%

Radiative decays

Γ_{91}	$\gamma \chi_{c2}$	< 9	$\times 10^{-4}$	CL=90%
Γ_{92}	$\gamma \chi_{c1}$	(2.9 ± 0.6)	$\times 10^{-3}$	
Γ_{93}	$\gamma \chi_{c0}$	(7.3 ± 0.9)	$\times 10^{-3}$	
Γ_{94}	$\gamma \eta'$	< 1.8	$\times 10^{-4}$	CL=90%
Γ_{95}	$\gamma \eta$	< 1.5	$\times 10^{-4}$	CL=90%
Γ_{96}	$\gamma \pi^0$	< 2	$\times 10^{-4}$	CL=90%

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 3 branching ratios uses 21 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 17.9$ for 17 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_3	98		
x_8	0	0	
Γ	0	0	-46
	x_2	x_3	x_8

Mode	Rate (MeV)	Scale factor
Γ_2 $D^0 \bar{D}^0$	14.1 ± 1.4	1.7
Γ_3 $D^+ D^-$	11.2 ± 1.1	1.7
Γ_8 $e^+ e^-$	$(2.65 \pm 0.18) \times 10^{-4}$	1.3

$\psi(3770)$ PARTIAL WIDTHS

$\Gamma(e^+ e^-)$	Γ_8
VALUE (keV)	EVTS DOCUMENT ID TECN COMMENT
0.265 ± 0.018 OUR FIT	Error includes scale factor of 1.3.
0.259 ± 0.016 OUR AVERAGE	Error includes scale factor of 1.2.
0.22 ± 0.05	⁵ ABLIKIM 08D BES2 $e^+ e^- \rightarrow$ hadrons
$0.277 \pm 0.011 \pm 0.013$	ABLIKIM 07E BES2 $e^+ e^- \rightarrow$ hadrons
$0.203 \pm 0.003^{+0.041}_{-0.027}$ 1.427M	⁶ BESSON 06 CLEO $e^+ e^- \rightarrow$ hadrons
0.276 ± 0.050	SCHINDLER 80 MRK2 $e^+ e^-$
0.18 ± 0.06	BACINO 78 DLCO $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.37 ± 0.09	⁷ RAPIDIS 77 LGW $e^+ e^-$
⁵ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.	
⁶ BESSON 06 (as corrected in BESSON 10) measure $\sigma(e^+ e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = 6.36 \pm 0.08^{+0.41}_{-0.30}$ nb at $\sqrt{s} = 3773 \pm 1$ MeV, and obtain Γ_{ee} from the Born-level cross section calculated using $\psi(3770)$ mass and width from our 2004 edition, PDG 04.	
⁷ See also $\Gamma(e^+ e^-) / \Gamma_{\text{total}}$ below.	

$\psi(3770)$ BRANCHING RATIOS

$\Gamma(D\bar{D})/\Gamma_{\text{total}}$ $\Gamma_1/\Gamma = (\Gamma_2+\Gamma_3)/\Gamma$
VALUE EVTS DOCUMENT ID TECN COMMENT

0.93 $^{+0.08}_{-0.09}$ OUR FIT Error includes scale factor of 2.0.

0.93 $^{+0.08}_{-0.09}$ OUR AVERAGE Error includes scale factor of 2.1.

0.849 ± 0.056 ± 0.018		⁸ ABLIKIM	08B	BES2	$e^+e^- \rightarrow \text{non-}D\bar{D}$
1.033 ± 0.014 $^{+0.048}_{-0.066}$	1.427M	⁹ BESSON	06	CLEO	$e^+e^- \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.866 ± 0.050 ± 0.036		^{10,11} ABLIKIM	07K	BES2	$e^+e^- \rightarrow \text{non-}D\bar{D}$
0.836 ± 0.073 ± 0.042		¹¹ ABLIKIM	06L	BES2	$e^+e^- \rightarrow D\bar{D}$
0.855 ± 0.017 ± 0.058		^{11,12} ABLIKIM	06N	BES2	$e^+e^- \rightarrow D\bar{D}$

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$ Γ_2/Γ
VALUE DOCUMENT ID TECN COMMENT

0.52 ± 0.05 OUR FIT Error includes scale factor of 2.0.

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.467 ± 0.047 ± 0.023		ABLIKIM	06L	BES2	$e^+e^- \rightarrow D^0\bar{D}^0$
0.499 ± 0.013 ± 0.038		¹² ABLIKIM	06N	BES2	$e^+e^- \rightarrow D^0\bar{D}^0$

$\Gamma(D^+D^-)/\Gamma_{\text{total}}$ Γ_3/Γ
VALUE DOCUMENT ID TECN COMMENT

0.41 ± 0.04 OUR FIT Error includes scale factor of 2.0.

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.369 ± 0.037 ± 0.028		ABLIKIM	06L	BES2	$e^+e^- \rightarrow D^+D^-$
0.357 ± 0.011 ± 0.034		¹² ABLIKIM	06N	BES2	$e^+e^- \rightarrow D^+D^-$

$\Gamma(D^0\bar{D}^0)/\Gamma(D^+D^-)$ Γ_2/Γ_3
VALUE EVTS DOCUMENT ID TECN COMMENT

1.260 ± 0.021 OUR FIT

1.260 ± 0.021 OUR AVERAGE

1.39 ± 0.31 ± 0.12		PAKHLOVA	08	BELL	10.6 $e^+e^- \rightarrow D\bar{D}\gamma$
1.78 ± 0.33 ± 0.24		AUBERT	07BE	BABR	$e^+e^- \rightarrow D\bar{D}\gamma$
1.258 ± 0.016 ± 0.014		DOBBS	07	CLEO	$e^+e^- \rightarrow D\bar{D}$
1.27 ± 0.12 ± 0.08		ABLIKIM	06L	BES2	$e^+e^- \rightarrow D\bar{D}$
2.43 ± 1.50 ± 0.43	34	¹³ CHISTOV	04	BELL	$B^+ \rightarrow \psi(3770)K^+$

$\Gamma(J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_4/Γ
VALUE (units 10⁻³) EVTS DOCUMENT ID TECN COMMENT

1.93 ± 0.28 OUR AVERAGE

1.89 ± 0.20 ± 0.20	231 ± 33	ADAM	06	CLEO	$e^+e^- \rightarrow \psi(3770)$
3.4 ± 1.4 ± 0.9	17.8 ± 4.8	BAI	05	BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(J/\psi\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ
VALUE (units 10⁻²) EVTS DOCUMENT ID TECN COMMENT

0.080 ± 0.025 ± 0.016	39 ± 14	ADAM	06	CLEO	$e^+e^- \rightarrow \psi(3770)$
------------------------------	---------	------	----	------	---------------------------------

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
$87 \pm 33 \pm 22$	22 ± 10	ADAM	06	CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(J/\psi\pi^0)/\Gamma_{\text{total}}$					Γ_7/Γ	
VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
<28	90	<10	ADAM	06	CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$					Γ_8/Γ	
VALUE (units 10^{-5})			DOCUMENT ID	TECN	COMMENT	
0.97 ± 0.07 OUR FIT	Error includes scale factor of 1.2.					
1.3 ± 0.2			RAPIDIS	77	LGW	e^+e^-

⁸ Neglecting interference.

⁹ Obtained by comparing a measurement of the total cross section (corrected in BESSON 10) with that of $D\bar{D}$ reported by CLEO in DOBBS 07.

¹⁰ Using $\sigma^{obs} = 7.07 \pm 0.58$ nb and neglecting interference.

¹¹ Not independent of ABLIKIM 08B.

¹² From a measurement of $\sigma(e^+e^- \rightarrow D\bar{D})$ at $\sqrt{s} = 3773$ MeV, using the $\psi(3770)$ resonance parameters measured by ABLIKIM 06L.

¹³ See ADLER 88C for older measurements of this quantity.

———— DECAYS TO LIGHT HADRONS ————

$\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$					Γ_9/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<1.4	90	¹⁴ ADAMS	06	CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\phi\eta')/\Gamma_{\text{total}}$					Γ_{10}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
<7	90	¹⁴ ADAMS	06	CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\omega\eta')/\Gamma_{\text{total}}$					Γ_{11}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
<4	90	¹⁴ ADAMS	06	CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\rho^0\eta')/\Gamma_{\text{total}}$					Γ_{12}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
<6	90	¹⁴ ADAMS	06	CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\phi\eta)/\Gamma_{\text{total}}$					Γ_{13}/Γ	
VALUE (units 10^{-4})			DOCUMENT ID	TECN	COMMENT	
$3.1 \pm 0.6 \pm 0.3$			¹⁴ ADAMS	06	CLEO	$3.773 e^+e^- \rightarrow \phi\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19			¹⁵ ABLIKIM	07B	BES2	$e^+e^- \rightarrow \psi(3770)$
-------	--	--	-----------------------	-----	------	---------------------------------

$\Gamma(\omega\eta)/\Gamma_{\text{total}}$					Γ_{14}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<1.4	90	¹⁴ ADAMS	06	CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\rho^0\eta)/\Gamma_{\text{total}}$ Γ_{15}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	14 ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$ Γ_{16}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 3	90	14 ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<50		15 ABLIKIM	07B	BES2 $e^+e^- \rightarrow \psi(3770)$
-----	--	------------	-----	--------------------------------------

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6	90	14 ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{18}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	14,16 ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_{19}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	14,16 ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(K^*(892)^+K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{20}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.4	90	14 ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{21}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.2	90	14 ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ Γ_{22}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.2	90	17 CRONIN-HEN..06		CLEO $e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90	18 ABLIKIM	04F	BES $e^+e^- \rightarrow \psi(3770)$
-----	----	------------	-----	-------------------------------------

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{23}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<11.2	90	19 HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<48		15 ABLIKIM	07B	BES2 $e^+e^- \rightarrow \psi(3770)$
-----	--	------------	-----	--------------------------------------

$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$					Γ_{24}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<10.6	90	19 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<62		15 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$					Γ_{25}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<58.5	90	305	ABLIKIM	08N BES2	$e^+e^- \rightarrow \psi(3770)$
$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{26}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 6.0	90	19 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<55	90	15 ABLIKIM	07I BES2	$3.77 e^+e^-$	
$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$					Γ_{27}/Γ
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<91	15 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$					Γ_{28}/Γ
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<137	15 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(3(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$					Γ_{29}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<117.4	90	59	ABLIKIM	08N BES2	$e^+e^- \rightarrow \psi(3770)$
$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{30}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<1.24	90	19 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<2.3	90	15 ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$					Γ_{31}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8.9	90	218	ABLIKIM	08N BES2	$e^+e^- \rightarrow \psi(3770)$
$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{32}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<6.9	90	15 ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(\eta3\pi)/\Gamma_{\text{total}}$					Γ_{33}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<13.4	90	19 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$	

$\Gamma(\eta 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$			Γ_{34}/Γ		
<u>VALUE (units 10^{-4})</u>			<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<243		15	ABLIKIM	07B	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(\eta \rho^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$			Γ_{35}/Γ		
<u>VALUE (units 10^{-2})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.45	90	15	ABLIKIM	10D	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(\eta' 3\pi)/\Gamma_{\text{total}}$			Γ_{36}/Γ		
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<24.4	90	19	HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$			Γ_{37}/Γ		
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 9.0	90	19	HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<48		15	ABLIKIM	07B	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(\phi \pi^+ \pi^-)/\Gamma_{\text{total}}$			Γ_{38}/Γ		
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 4.1	90	19	HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<16		15	ABLIKIM	07B	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$			Γ_{39}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.2	90	14	ABLIKIM	08N	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(4(\pi^+ \pi^-))/\Gamma_{\text{total}}$			Γ_{40}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<16.7	90	15	ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(4(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$			Γ_{41}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<30.6	90	15	ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$			Γ_{42}/Γ		
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.5	90	19	HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$			Γ_{43}/Γ		
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 23.6	90	19	HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<111		15	ABLIKIM	07B	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K^+ K^- \rho^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<8	90	15 ABLIKIM 07I	BES2	$3.77 e^+ e^-$

$\Gamma(K^+ K^- \rho^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<146	90	15 ABLIKIM 07I	BES2	$3.77 e^+ e^-$

$\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.4	90	19 HUANG 06A	CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<66	90	15 ABLIKIM 07I	BES2	$3.77 e^+ e^-$
-----	----	----------------	------	----------------

$\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<38	90	15 ABLIKIM 07I	BES2	$3.77 e^+ e^-$

$\Gamma(K^{*0} K^- \pi^+ \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<162	90	15 ABLIKIM 07I	BES2	$3.77 e^+ e^-$

$\Gamma(K^{*+} K^- \pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<323	90	15 ABLIKIM 07I	BES2	$3.77 e^+ e^-$

$\Gamma(K^+ K^- \pi^+ \pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<26.7	90	24	ABLIKIM 08N	BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<10.3	90	15 ABLIKIM 07F	BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<36.0	90	15 ABLIKIM 07F	BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\eta K^+ K^-)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.1	90	19 HUANG 06A	CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<31	90	15 ABLIKIM 10D	BES2	$e^+ e^- \rightarrow \psi(3770)$
-----	----	----------------	------	----------------------------------

$\Gamma(\eta K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$			Γ_{54}/Γ		
<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<1.24	90	15 ABLIKIM	10D	BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\rho^0 K^+ K^-)/\Gamma_{\text{total}}$			Γ_{55}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<5.0	90	15 ABLIKIM	07F	BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$			Γ_{56}/Γ		
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 6.0	90	19 HUANG	06A	CLEO	$e^+ e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<17		15 ABLIKIM	07B	BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$			Γ_{57}/Γ		
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 7.5	90	19 HUANG	06A	CLEO	$e^+ e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<24		15 ABLIKIM	07B	BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(2(K^+ K^-)\pi^0)/\Gamma_{\text{total}}$			Γ_{58}/Γ		
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 2.9	90	19 HUANG	06A	CLEO	$e^+ e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<46		15 ABLIKIM	07B	BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(2(K^+ K^-)\pi^+ \pi^-)/\Gamma_{\text{total}}$			Γ_{59}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<3.2	90	15 ABLIKIM	07F	BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \pi^+)/\Gamma_{\text{total}}$			Γ_{60}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.2	90	18	ABLIKIM	08M	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$			Γ_{61}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<13.3	90	40	ABLIKIM	08M	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \rho^+)/\Gamma_{\text{total}}$			Γ_{62}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6.6	90		ABLIKIM	09C	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- 2\pi^+ \pi^-)/\Gamma_{\text{total}}$			Γ_{63}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8.7	90	39	ABLIKIM	08M	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \pi^+ \rho^0)/\Gamma_{\text{total}}$			Γ_{64}/Γ		
<u>VALUE (units 10^{-2})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.6	90		ABLIKIM	09C	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(K_S^0 K^- \pi^+ \eta)/\Gamma_{\text{total}}$			Γ_{65}/Γ		
<u>VALUE (units 10^{-2})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.3	90		ABLIKIM	09C	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(K_S^0 K^- 2\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$			Γ_{66}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<41.8	90	23	ABLIKIM	08M	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(K_S^0 K^- 2\pi^+ \pi^- \eta)/\Gamma_{\text{total}}$			Γ_{67}/Γ		
<u>VALUE (units 10^{-2})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.8	90		ABLIKIM	09C	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(K_S^0 K^- \pi^+ 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$			Γ_{68}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<12.2	90	4	ABLIKIM	08M	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(K_S^0 K^- \pi^+ 2\pi^0)/\Gamma_{\text{total}}$			Γ_{69}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<26.5	90	17	ABLIKIM	08M	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(K_S^0 K^- K^+ K^- \pi^+)/\Gamma_{\text{total}}$			Γ_{70}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.9	90		ABLIKIM	09C	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(K_S^0 K^- K^+ K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$			Γ_{71}/Γ		
<u>VALUE (units 10^{-2})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.0	90		ABLIKIM	09C	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(K_S^0 K^- K^+ K^- \pi^+ \eta)/\Gamma_{\text{total}}$			Γ_{72}/Γ		
<u>VALUE (units 10^{-2})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.2	90		ABLIKIM	09C	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(K^{*0} K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$			Γ_{73}/Γ		
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<9.7	90	15	ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$
$\Gamma(\rho\bar{\rho}\pi^0)/\Gamma_{\text{total}}$			Γ_{74}/Γ		
<u>VALUE (units 10^{-4})</u>			<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<12			15 ABLIKIM	07B	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\rho\bar{\rho}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{75}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 5.8	90	19 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<16		15 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{76}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.2	90	19 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<4	90	15 ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\rho\bar{\rho}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{77}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<18.5	90	19 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<73		15 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\omega\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{78}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 2.9	90	19 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<30	90	20 ABLIKIM	07I BES2	$3.77 e^+e^-$

$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{79}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<12	90	15 ABLIKIM	07I BES2	$3.77 e^+e^-$

$\Gamma(\rho\bar{\rho}2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{80}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.6	90	15 ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\eta\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{81}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 5.4	90	19 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<11	90	15 ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\eta\rho\bar{\rho}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{82}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.3	90	15 ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\rho^0\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{83}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.7	90	15 ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\rho\bar{\rho}K^+K^-)/\Gamma_{\text{total}}$ Γ_{84}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.2	90	¹⁹ HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<11		¹⁵ ABLIKIM	07B	BES2 $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\eta\rho\bar{\rho}K^+K^-)/\Gamma_{\text{total}}$ Γ_{85}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<6.9	90	¹⁵ ABLIKIM	10D	BES2 $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\pi^0\rho\bar{\rho}K^+K^-)/\Gamma_{\text{total}}$ Γ_{86}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	¹⁵ ABLIKIM	10D	BES2 $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\phi\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{87}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	¹⁹ HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<9		¹⁵ ABLIKIM	07B	BES2 $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{88}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.5	90	¹⁹ HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<39	90	¹⁵ ABLIKIM	07F	BES2 $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\Lambda\bar{\rho}K^+)/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.8	90	¹⁹ HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(\Lambda\bar{\rho}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{90}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6.3	90	¹⁹ HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$

¹⁴ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

¹⁵ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

¹⁶ Data suggest possible destructive interference with continuum.

¹⁷ Using $\sigma(e^+e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = (6.38 \pm 0.08^{+0.41}_{-0.30})$ nb from BESSON 06 and $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6895 \pm 0.0014$.

¹⁸ Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6860 \pm 0.0027$.

¹⁹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

²⁰ Using $\sigma^{obs} = 7.15 \pm 0.27 \pm 0.27$ nb and neglecting interference.

RADIATIVE DECAYS

$\Gamma(\gamma\chi_{c2})/\Gamma_{\text{total}}$ Γ_{91}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.9	90	21 COAN	06A CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.0	90	22 BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$
------	----	-----------	---------	---

$\Gamma(\gamma\chi_{c1})/\Gamma_{\text{total}}$ Γ_{92}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.9 \pm 0.5 \pm 0.4$		23 BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}, \gamma\gamma J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.9 \pm 1.4 \pm 0.6$	54 ± 17	24 BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$
-----------------------	-------------	-----------	---------	---

$2.8 \pm 0.5 \pm 0.4$	53 ± 10	21 COAN	06A CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$
-----------------------	-------------	---------	----------	---

$\Gamma(\gamma\chi_{c1})/\Gamma(J/\psi\pi^+\pi^-)$ Γ_{92}/Γ_4

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.49 \pm 0.31 \pm 0.26$	53 ± 10	25 COAN	06A CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$

$\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$ Γ_{93}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$7.3 \pm 0.7 \pm 0.6$		274 ± 27	26 BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 44	90	21 COAN	06A CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$
------	----	---------	----------	---

$\Gamma(\gamma\chi_{c0})/\Gamma(\gamma\chi_{c2})$ Γ_{93}/Γ_{91}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>8	90	27 BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\gamma\chi_{c0})/\Gamma(\gamma\chi_{c1})$ Γ_{93}/Γ_{92}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.5 ± 0.6	27 BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

$\Gamma(\gamma\eta')/\Gamma_{\text{total}}$ Γ_{94}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.8	90	28 PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$

$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$ Γ_{95}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.5	90	28 PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$					Γ_{96}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
<2	90	PEDLAR	09	CLE3	$\psi(2S) \rightarrow \gamma X$
<p>²¹ Using $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$ keV from ADAM 06 and taking $\sigma(e^+e^- \rightarrow D\bar{D})$ from HE 05 for $\sigma(e^+e^- \rightarrow \psi(3770))$.</p> <p>²² Uses $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = 9.22 \pm 0.11 \pm 0.46\%$ from ATHAR 04, $\psi(2S)$ mass and width from PDG 04, and $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$ keV from ADAM 06.</p> <p>²³ Averages the two measurements from COAN 06A and BRIERE 06.</p> <p>²⁴ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = 9.07 \pm 0.11 \pm 0.54\%$ from ATHAR 04, $\psi(2S)$ mass and width from PDG 04, and $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$ keV from ADAM 06.</p> <p>²⁵ Using $B(\psi(3770) \rightarrow J/\psi\pi^+\pi^-) = (1.89 \pm 0.20 \pm 0.20) \times 10^{-3}$ from ADAM 06.</p> <p>²⁶ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = 9.33 \pm 0.14 \pm 0.61\%$ from ATHAR 04, $\psi(2S)$ mass and width from PDG 04, and $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$ keV from ADAM 06.</p> <p>²⁷ Not independent of other results in BRIERE 06.</p> <p>²⁸ Assuming maximal destructive interference between $\psi(3770)$ and continuum sources.</p>					

$\psi(3770)$ REFERENCES

ABLIKIM	10D	EPJ C66 11	M. Ablikim <i>et al.</i>	(BES II Collab.)
BESSION	10	PRL 104 159901E	D. Besson <i>et al.</i>	(CLEO Collab.)
ABLIKIM	09C	EPJ C64 243	M. Ablikim <i>et al.</i>	(BES Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08M	PL B670 179	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08N	PL B670 184	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	08B	PR D77 011102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
BRODZICKA	08	PRL 100 092001	J. Brodzicka <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08	PR D77 011103R	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	07B	PL B650 111	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07E	PL B652 238	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07F	PL B656 30	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07I	EPJ C52 805	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07K	PR D76 122002	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	07BE	PR D76 111105R	B. Aubert <i>et al.</i>	(BABAR Collab.)
DOBBS	07	PR D76 112001	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06N	PL B641 145	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ADAMS	06	PR D73 012002	G.S. Adams <i>et al.</i>	(CLEO Collab.)
BESSION	06	PRL 96 092002	D. Besson <i>et al.</i>	(CLEO Collab.)
Also		PRL 104 159901E	D. Besson <i>et al.</i>	(CLEO Collab.)
BRIERE	06	PR D74 031106R	R.A. Briere <i>et al.</i>	(CLEO Collab.)
COAN	06A	PRL 96 182002	T.E. Coan <i>et al.</i>	(CLEO Collab.)
CRONIN-HENNESSY	06	PR D74 012005	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
HUANG	06A	PRL 96 032003	G.S. Huang <i>et al.</i>	(CLEO Collab.)
BAI	05	PL B605 63	J.Z. Bai <i>et al.</i>	(BES Collab.)
HE	05	PRL 95 121801	Q. He <i>et al.</i>	(CLEO Collab.)
Also		PRL 96 199903 (err.)	Q. He <i>et al.</i>	(CLEO Collab.)
ABLIKIM	04F	PR D70 077101	M. Ablikim <i>et al.</i>	(BES Collab.)
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
ADLER	88C	PRL 60 89	J. Adler <i>et al.</i>	(Mark III Collab.)
SCHINDLER	80	PR D21 2716	R.H. Schindler <i>et al.</i>	(Mark II Collab.)
BACINO	78	PRL 40 671	W.J. Bacino <i>et al.</i>	(SLAC, UCLA, UCI)
RAPIDIS	77	PRL 39 526	P.A. Rapidis <i>et al.</i>	(LGW Collab.)