

$\Delta(1900) 1/2^-$ $I(J^P) = \frac{3}{2}(\frac{1}{2}^-)$ Status: **

OMITTED FROM SUMMARY TABLE

Some obsolete results published before 1980 were last included in our 2006 edition, Journal of Physics, G **33** 1 (2006). Some further obsolete results published before 1984 were last included in our 2006 edition, Journal of Physics, G **33** 1 (2006).

The latest GWU analysis (ARNDT 06) finds no evidence for this resonance.

 $\Delta(1900)$ BREIT-WIGNER MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1840 to 1920 (≈ 1860) OUR ESTIMATE			
1840 ± 30	ANISOVICH	12A DPWA	Multichannel
1920 ± 24	MANLEY	92 IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
1890 ± 50	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$
1908 ± 30	HOEHLER	79 IPWA	$\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1802 ± 87	VRANA	00 DPWA	Multichannel
1918.5 ± 23.0	CHEW	80 BPWA	$\pi^+ p \rightarrow \pi^+ p$

 $\Delta(1900)$ BREIT-WIGNER WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
300 ± 45	ANISOVICH	12A DPWA	Multichannel
263 ± 39	MANLEY	92 IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
170 ± 50	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$
140 ± 40	HOEHLER	79 IPWA	$\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
48 ± 45	VRANA	00 DPWA	Multichannel
93.5 ± 54.0	CHEW	80 BPWA	$\pi^+ p \rightarrow \pi^+ p$

 $\Delta(1900)$ POLE POSITION**REAL PART**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1845 ± 25	ANISOVICH	12A DPWA	Multichannel
1780	¹ HOEHLER	93 SPED	$\pi N \rightarrow \pi N$
1870 ± 40	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1795	VRANA	00 DPWA	Multichannel
not seen	ARNDT	91 DPWA	$\pi N \rightarrow \pi N$ Soln SM90
2029 or 2025	² LONGACRE	78 IPWA	$\pi N \rightarrow N\pi\pi$

–2×IMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
300±45	ANISOVICH 12A	DPWA	Multichannel
180±50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
58	VRANA 00	DPWA	Multichannel
not seen	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
164 or 163	² LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$

Δ(1900) ELASTIC POLE RESIDUE

MODULUS |r|

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10±3	ANISOVICH 12A	DPWA	Multichannel
10±3	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
–125±20	ANISOVICH 12A	DPWA	Multichannel
+ 20±40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

Δ(1900) INELASTIC POLE RESIDUE

The “normalized residue” is the residue divided by Γ_{pole} .

Normalized residue in $N\pi \rightarrow \Delta(1900) \rightarrow \Sigma K$

MODULUS (%)	PHASE (°)	DOCUMENT ID	TECN	COMMENT
7±2	–50 ± 30	ANISOVICH 12A	DPWA	Multichannel

Normalized residue in $N\pi \rightarrow \Delta(1900) \rightarrow \Delta\pi, D\text{-wave}$

MODULUS (%)	PHASE (°)	DOCUMENT ID	TECN	COMMENT
12 ⁺⁸ _{–5}	110 ± 20	ANISOVICH 12A	DPWA	Multichannel

Δ(1900) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	10–30 %
Γ_2 ΣK	
Γ_3 $N\pi\pi$	
Γ_4 $\Delta\pi$	
Γ_5 $\Delta(1232)\pi, D\text{-wave}$	
Γ_6 $N\rho$	
Γ_7 $N\rho, S=1/2, S\text{-wave}$	
Γ_8 $N\rho, S=3/2, D\text{-wave}$	
Γ_9 $N(1440)\pi, S\text{-wave}$	
Γ_{10} $N\gamma, \text{helicity}=1/2$	

Δ(1900) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7 ± 3	ANISOVICH	12A	DPWA Multichannel
41 ± 4	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
10 ± 3	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
8 ± 4	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
33 ± 10	VRANA	00	DPWA Multichannel
28	CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow \Sigma K$ $(\Gamma_1\Gamma_2)^{1/2}/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.03	CANDLIN	84	DPWA $\pi^+ p \rightarrow \Sigma^+ K^+$

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow \Delta(1232)\pi$, *D-wave* $(\Gamma_1\Gamma_5)^{1/2}/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
+0.25 ± 0.07	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$

$\Gamma(\Delta(1232)\pi, D\text{-wave})/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
15 ⁺⁵⁰ ₋₁₀	ANISOVICH	12A	DPWA Multichannel
28 ± 1	VRANA	00	DPWA Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow N\rho, S=1/2, S\text{-wave}$ $(\Gamma_1\Gamma_7)^{1/2}/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.14 ± 0.11	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$

$\Gamma(N\rho, S=1/2, S\text{-wave})/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
30 ± 2	VRANA	00	DPWA Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow N\rho, S=3/2, D\text{-wave}$ $(\Gamma_1\Gamma_8)^{1/2}/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.37 ± 0.07	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$

$\Gamma(N\rho, S=3/2, D\text{-wave})/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5 ± 1	VRANA	00	DPWA Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow N(1440)\pi, S\text{-wave}$ $(\Gamma_1\Gamma_9)^{1/2}/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.16 ± 0.11	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$

$\Gamma(N(1440)\pi, S\text{-wave})/\Gamma_{\text{total}}$	Γ_9/Γ		
VALUE (%)	DOCUMENT ID	TECN	COMMENT
4±1	VRANA	00	DPWA Multichannel

$\Delta(1900)$ PHOTON DECAY AMPLITUDES

Papers on γN amplitudes predating 1981 may be found in our 2006 edition, Journal of Physics, G **33** 1 (2006).

$\Delta(1900) \rightarrow N\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE ($\text{GeV}^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
0.059±0.016	³ ANISOVICH 12A	DPWA	Phase = $(60 \pm 25)^\circ$
-0.004±0.016	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
0.029±0.008	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$

$\Delta(1900)$ FOOTNOTES

¹ See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

² LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.

³ This ANISOVICH 12A value is the complex helicity amplitude at the pole position.

$\Delta(1900)$ REFERENCES

For early references, see Physics Letters **111B** 1 (1982).

ANISOVICH 12A	EPJ A48 15	A.V. Anisovich <i>et al.</i>	(BONN, PNPI)
ARNDT 06	PR C74 045205	R.A. Arndt <i>et al.</i>	(GWU)
PDG 06	JPG 33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
VRANA 00	PRPL 328 181	T.P. Vrana, S.A. Dytman,, T.-S.H. Lee	(PITT+)
HOEHLER 93	πN Newsletter 9 1	G. Hohler	(KARL)
MANLEY 92	PR D45 4002	D.M. Manley, E.M. Saleski	(KENT) IJP
Also	PR D30 904	D.M. Manley <i>et al.</i>	(VPI)
ARNDT 91	PR D43 2131	R.A. Arndt <i>et al.</i>	(VPI, TELE) IJP
CANDLIN 84	NP B238 477	D.J. Candlin <i>et al.</i>	(EDIN, RAL, LOWC)
CRAWFORD 83	NP B211 1	R.L. Crawford, W.T. Morton	(GLAS)
AWAJI 81	Bonn Conf. 352	N. Awaji, R. Kajikawa	(NAGO)
Also	NP B197 365	K. Fujii <i>et al.</i>	(NAGO)
CHEW 80	Toronto Conf. 123	D.M. Chew	(LBL) IJP
CUTKOSKY 80	Toronto Conf. 19	R.E. Cutkosky <i>et al.</i>	(CMU, LBL) IJP
Also	PR D20 2839	R.E. Cutkosky <i>et al.</i>	(CMU, LBL) IJP
HOEHLER 79	PDAT 12-1	G. Hohler <i>et al.</i>	(KARLT) IJP
Also	Toronto Conf. 3	R. Koch	(KARLT) IJP
LONGACRE 78	PR D17 1795	R.S. Longacre <i>et al.</i>	(LBL, SLAC)