

WIMPs and Other Particles Searches for

OMITTED FROM SUMMARY TABLE
 A REVIEW GOES HERE – Check our WWW List of Reviews

GALACTIC WIMP SEARCHES

Cross-Section Limits for Dark Matter Particles (X^0) on Nuclei

These limits are for weakly-interacting stable particles that may constitute the invisible mass in the galaxy. Unless otherwise noted, a local mass density of 0.3 GeV/cm^3 is assumed; see each paper for velocity distribution assumptions. In the papers the limit is given as a function of the X^0 mass. Here we list limits only for typical mass values of 20 GeV, 100 GeV, and 1 TeV. Specific limits on supersymmetric dark matter particles may be found in the Supersymmetry section.

For $m_{X^0} = 20 \text{ GeV}$

<u>VALUE (nb)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1		AALSETH 11	CGNT	Ge
2		AALSETH 11A	CGNT	Ge
3		AHLEN 11	DMTP	CF ₄
4		AHMED 11	CDM2	Ge, inelastic
5		AHMED 11A	RVUE	Ge
6		AHMED 11B	CDM2	Ge, low threshold
7		ANGLE 11	XE10	Xe
8		APRILE 11	X100	Xe
9		APRILE 11B	X100	Xe
10		BEHNKE 11	COUP	CF ₃ I
11		HORN 11	ZEP3	Xe
12		TANAKA 11	SKAM	H, solar ν
13		AKERIB 10	CDM2	Si, Ge, low threshold
14		APRILE 10	X100	Xe
15		ARMENGAUD 10	EDE2	Ge
16		FELIZARDO 10	SMPL	C ₂ ClF ₃
17		AHMED 09	CDM2	Ge
18		ARCHAMBAUD 09	PICA	F
19		LEBEDENKO 09A	ZEP3	Xe
20		LIN 09	TEXO	Ge
21		AALSETH 08	CGNT	Ge
22		ANGLE 08A	XE10	Xe
23		ALNER 07	ZEP2	Xe
24		LEE 07A	KIMS	CsI
25		AKERIB 06	CDMS	⁷³ Ge, ²⁹ Si
26		SHIMIZU 06A	CNTR	F (CaF ₂)
27		ALNER 05	NAIA	NaI
28		BARNABE-HELY 05	PICA	F (C ₄ F ₁₀)
29		BENOIT 05	EDEL	⁷³ Ge

		30	GIRARD	05	SMPL	F (C ₂ ClF ₅)
		31	KLAPDOR-K...	05	HDMS	⁷³ Ge (enriched)
		32	MIUCHI	03	BOLO	LiF
		33	TAKEDA	03	BOLO	NaF
< 0.08	90	34	ANGLOHER	02	CRES	Al
		35	BENOIT	00	EDEL	Ge
< 0.04	95	36	KLIMENKO	98	CNTR	⁷³ Ge, incl.
< 0.8			ALESSAND...	96	CNTR	O
< 6			ALESSAND...	96	CNTR	Te
< 0.02	90	37	BELLI	96	CNTR	¹²⁹ Xe, incl.
		38	BELLI	96C	CNTR	¹²⁹ Xe
< 0.004	90	39	BERNABEI	96	CNTR	Na
< 0.3	90	39	BERNABEI	96	CNTR	I
< 0.2	95	40	SARSA	96	CNTR	Na
< 0.015	90	41	SMITH	96	CNTR	Na
< 0.05	95	42	GARCIA	95	CNTR	Natural Ge
< 0.1	95		QUENBY	95	CNTR	Na
< 90	90	43	SNOWDEN-...	95	MICA	¹⁶ O
< 4 × 10 ³	90	43	SNOWDEN-...	95	MICA	³⁹ K
< 0.7	90		BACCI	92	CNTR	Na
< 0.12	90	44	REUSSER	91	CNTR	Natural Ge
< 0.06	95		CALDWELL	88	CNTR	Natural Ge

¹ AALSETH 11 give $\sigma < 5 \times 10^{-5}$ pb (90% CL) for spin-independent X^0 -nucleon cross section. See their Fig. 4 for limits extending to $m_{X^0} = 3.5$ GeV.

² AALSETH 11A find indications of annual modulation of the data, the energy spectrum being compatible with X^0 mass around 8 GeV.

³ AHLEN 11 give $\sigma < 1 \times 10^5$ pb (90% CL) for spin-dependent X^0 -proton cross section.

⁴ AHMED 11 search for X^0 inelastic scattering. See their Figs. 8–10 for limits.

⁵ AHMED 11A combine CDMS and EDELWEISS data and give $\sigma < 2.7 \times 10^{-7}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.

⁶ AHMED 11B give limits on spin-independent X^0 -nucleon and spin-dependent X^0 -neutron cross section for $m_{X^0} = 4$ –12 GeV in the range 10^{-5} – 10^{-3} pb and 10 – 10^3 pb, respectively. See their Fig. 3.

⁷ ANGLE 11 give $\sigma < 5 \times 10^{-7}$ pb (90% CL) for spin-independent X^0 -nucleon cross section. See their Fig. 3 for limits down to $m_{X^0} = 4$ GeV.

⁸ APRILE 11 reanalyze APRILE 10 data and give $\sigma < 7 \times 10^{-8}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.

⁹ APRILE 11B give $\sigma < 2 \times 10^{-8}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.

¹⁰ BEHNKE 11 give $\sigma < 0.1$ pb (90% CL) for spin-dependent X^0 -proton cross section with a direction sensitive detector.

¹¹ HORN 11 perform detector calibration by neutrons. Earlier results are only marginally affected.

¹² TANAKA 11 search for neutrinos produced by X^0 annihilation in the Sun and give $\sigma < 1.5 \times 10^{-2}$ pb (90% CL) (for $m_{X^0} = 10$ GeV) for spin-dependent X^0 -proton cross section, assuming that X^0 pairs annihilate to $b\bar{b}$.

¹³ AKERIB 10 give $\sigma < 1 \times 10^{-5}$ pb (90% CL) for spin-independent X^0 -nucleon cross section. See their Figs. 10 and 12 for limits extending to X^0 mass of 1 GeV.

¹⁴ APRILE 10 give $\sigma < 1 \times 10^{-7}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.

- 15 ARMENGAUD 10 give $\sigma < 2 \times 10^{-6}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.
- 16 FELIZARDO 10 give $\sigma < 4 \times 10^{-5}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.
- 17 AHMED 09 give $\sigma < 0.06$ pb (90% CL) for spin-dependent X^0 -neutron cross section.
- 18 ARCHAMBAULT 09 give $\sigma < 0.2$ pb (90% CL) for spin-dependent X^0 -proton cross section.
- 19 LEBEDENKO 09A give $\sigma < 4$ (0.04) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 20 See their Fig. 6(b) for limits on spin-dependent X^0 -neutron cross section for m_{X^0} between 2 and 10 GeV.
- 21 See their Fig. 2 for cross section limits for m_{X^0} between 4 and 10 GeV.
- 22 ANGLE 08A give $\sigma < 0.6$ (0.007) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 23 ALNER 07 give $\sigma < 100$ (0.5) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 24 LEE 07A give $\sigma < 1$ (25) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 25 AKERIB 06 give $\sigma < 20$ (0.3) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section. See also AKERIB 05.
- 26 SHIMIZU 06A give $\sigma < 2$ (30) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 27 ALNER 05 give $\sigma < 0.5$ (60) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 28 BARNABE-HEIDER 05 give $\sigma < 1.5$ (20) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 29 BENOIT 05 give $\sigma < 10$ pb (90% CL) for spin-dependent X^0 -neutron cross section.
- 30 GIRARD 05 give $\sigma < 1.5$ pb (90% CL) for spin-dependent X^0 -proton cross section.
- 31 KLAPDOR-KLEINGROTHAUS 05 give $\sigma < 4$ pb (90% CL) for spin-dependent X^0 -neutron cross section.
- 32 MIUCHI 03 give model-independent limit $\sigma < 35$ pb (90% CL) for spin-dependent X^0 -proton cross section.
- 33 TAKEDA 03 give model-independent limit $\sigma < 0.03$ (0.6) nb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 34 ANGLOHER 02 limit is for spin-dependent WIMP-Aluminum cross section.
- 35 BENOIT 00 find four event categories in Ge detectors and suggest that low-energy surface nuclear recoils can explain anomalous events reported by UKDMC and Saclay NaI experiments.
- 36 KLIMENKO 98 limit is for inelastic scattering $X^0 \text{ } ^{73}\text{Ge} \rightarrow X^0 \text{ } ^{73}\text{Ge}^*$ (13.26 keV).
- 37 BELLI 96 limit for inelastic scattering $X^0 \text{ } ^{129}\text{Xe} \rightarrow X^0 \text{ } ^{129}\text{Xe}^*$ (39.58 keV).
- 38 BELLI 96C use background subtraction and obtain $\sigma < 150$ pb (< 1.5 fb) (90% CL) for spin-dependent (independent) X^0 -proton cross section. The confidence level is from R. Bernabei, private communication, May 20, 1999.
- 39 BERNABEI 96 use pulse shape discrimination to enhance the possible signal. The limit here is from R. Bernabei, private communication, September 19, 1997.
- 40 SARSA 96 search for annual modulation of WIMP signal. See SARSA 97 for details of the analysis. The limit here is from M.L. Sarsa, private communication, May 26, 1997.
- 41 SMITH 96 use pulse shape discrimination to enhance the possible signal. A dark matter density of 0.4 GeV cm^{-3} is assumed.
- 42 GARCIA 95 limit is from the event rate. A weaker limit is obtained from searches for diurnal and annual modulation.
- 43 SNOWDEN-IFFT 95 look for recoil tracks in an ancient mica crystal. Similar limits are also given for ^{27}Al and ^{28}Si . See COLLAR 96 and SNOWDEN-IFFT 96 for discussion on potential backgrounds.
- 44 REUSSER 91 limit here is changed from published (0.04) after reanalysis by authors. J.L. Vuilleumier, private communication, March 29, 1996.

For $m_{\chi_0} = 100$ GeV

<u>VALUE (nb)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		1 AALSETH	11	CGNT Ge
		2 AHLEN	11	DMTP CF ₄
		3 AHMED	11	CDM2 Ge, inelastic
		4 AHMED	11A	RVUE Ge
		5 AJELLO	11	FLAT
		6 APRILE	11	X100 Xe
		7 APRILE	11A	X100 Xe, inelastic
		8 APRILE	11B	X100 Xe
		9 ARMENGAUD	11	EDE2 Ge
		10 BEHNKE	11	COUP CF ₃ I
		11 HORN	11	ZEP3 Xe
		12 TANAKA	11	SKAM H, solar ν
		13 AKERIB	10	CDM2 Si, Ge, low threshold
		14 AKIMOV	10	ZEP3 Xe, inelastic scattering
		15 APRILE	10	X100 Xe
		16 ARMENGAUD	10	EDE2 Ge
		17 FELIZARDO	10	SMPL C ₂ F ₃
		18 MIUCHI	10	NAGE CF ₄
		19 AHMED	09	CDM2 Ge
		20 ANGLE	09	XE10 Xe, inelastic
		21 ARCHAMBAU..	09	PICA F
		22 LEBEDENKO	09A	ZEP3 Xe
		23 ANGLE	08A	XE10 Xe
		24 BEDNYAKOV	08	RVUE Ge
		25 ALNER	07	ZEP2 Xe
		26 LEE	07A	KIMS CsI
		27 MIUCHI	07	CNTR F (CF ₄)
		28 AKERIB	06	CDMS ⁷³ Ge, ²⁹ Si
		29 SHIMIZU	06A	CNTR F (CaF ₂)
		30 ALNER	05	NAIA NaI
		31 BARNABE-HE..	05	PICA F (C ₄ F ₁₀)
		32 BENOIT	05	EDEL ⁷³ Ge
		33 GIRARD	05	SMPL F (C ₂ F ₅)
		34 GIULIANI	05	RVUE
		35 GIULIANI	05A	RVUE
		36 KLAPDOR-K...	05	HDMS ⁷³ Ge (enriched)
		37 GIULIANI	04	RVUE
		38 GIULIANI	04A	RVUE
		39 MIUCHI	03	BOLO LiF
		40 MIUCHI	03	BOLO LiF
		41 TAKEDA	03	BOLO NaF
< 0.3	90	42 ANGLOHER	02	CRES Al
		43 BELLI	02	RVUE
		44 BERNABEI	02C	DAMA
		45 GREEN	02	RVUE
		46 ULLIO	01	RVUE
		47 BENOIT	00	EDEL Ge

< 0.004	90	48 BERNABEI	00D		^{129}Xe , inel.
		49 AMBROSIO	99	MCRO	
		50 BRHLIK	99	RVUE	
< 0.008	95	51 KLIMENKO	98	CNTR	^{73}Ge , inel.
< 0.08	95	52 KLIMENKO	98	CNTR	^{73}Ge , inel.
< 4		ALESSAND...	96	CNTR	O
< 25		ALESSAND...	96	CNTR	Te
< 0.006	90	53 BELLI	96	CNTR	^{129}Xe , inel.
		54 BELLI	96C	CNTR	^{129}Xe
< 0.001	90	55 BERNABEI	96	CNTR	Na
< 0.3	90	55 BERNABEI	96	CNTR	I
< 0.7	95	56 SARSA	96	CNTR	Na
< 0.03	90	57 SMITH	96	CNTR	Na
< 0.8	90	57 SMITH	96	CNTR	I
< 0.35	95	58 GARCIA	95	CNTR	Natural Ge
< 0.6	95	QUENBY	95	CNTR	Na
< 3	95	QUENBY	95	CNTR	I
< 1.5×10^2	90	59 SNOWDEN-...	95	MICA	^{16}O
< 4×10^2	90	59 SNOWDEN-...	95	MICA	^{39}K
< 0.08	90	60 BECK	94	CNTR	^{76}Ge
< 2.5	90	BACCI	92	CNTR	Na
< 3	90	BACCI	92	CNTR	I
< 0.9	90	61 REUSSER	91	CNTR	Natural Ge
< 0.7	95	CALDWELL	88	CNTR	Natural Ge

¹ AALSETH 11 give $\sigma < 2 \times 10^{-4}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.

² AHLEN 11 give $\sigma < 2 \times 10^3$ pb (90% CL) for spin-dependent X^0 -proton cross section.

³ AHMED 11 search for X^0 inelastic scattering. See their Figs. 8–10 for limits.

⁴ AHMED 11A combine CDMS and EDELWEISS data and give $\sigma < 3.3 \times 10^{-8}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.

⁵ AJELLO 11 search for e^\pm flux from X^0 annihilations in the Sun. Models in which X^0 annihilates into an intermediate long-lived weakly interacting particles or X^0 scatters inelastically are constrained. See their Figs. 6–8 for limits.

⁶ APRILE 11 reanalyze APRILE 10 data and give $\sigma < 3 \times 10^{-8}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.

⁷ APRILE 11A search for X^0 inelastic scattering. See their Figs. 2 and 3 for limits.

⁸ APRILE 11B give $\sigma < 1 \times 10^{-8}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.

⁹ ARMENGAUD 11 give $\sigma < 5 \times 10^{-8}$ pb (90% CL) for spin-independent X^0 -nucleon cross section. Supersedes ARMENGAUD 10. A limit on inelastic cross section is also given.

¹⁰ BEHNKE 11 give $\sigma < 0.07$ pb (90% CL) for spin-dependent X^0 -proton cross section with a direction sensitive detector.

¹¹ HORN 11 perform detector calibration by neutrons. Earlier results are only marginally affected.

¹² TANAKA 11 search for neutrinos produced by X^0 annihilation in the Sun and give $\sigma < 2.7 \times 10^{-4}$ (4.5×10^{-3}) pb (90% CL) for spin-dependent X^0 -proton cross section, if X^0 pairs annihilate to $W^+ W^- (b\bar{b})$.

¹³ AKERIB 10 give $\sigma < 9 \times 10^{-6}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.

¹⁴ AKIMOV 10 give cross section limits for inelastically scattering dark matter. See their Fig. 4.

- 15 APRILE 10 give $\sigma < 5 \times 10^{-8}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.
- 16 ARMENGAUD 10 give $\sigma < 1 \times 10^{-7}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.
- 17 FELIZARDO 10 give $\sigma < 3 \times 10^{-5}$ pb (90% CL) for spin-independent X^0 -nucleon cross section. See their Fig. 3 for limits on spin-dependent proton/neutron couplings for X^0 mass of 50 GeV.
- 18 MIUCHI 10 give $\sigma < 6 \times 10^3$ pb (90% CL) for spin-dependent X^0 -proton cross section with a direction sensitive detector.
- 19 AHMED 09 give $\sigma < 0.02$ pb (90% CL) for spin-dependent X^0 -neutron cross section.
- 20 ANGLE 09 search for X^0 inelastic scattering. See their Fig. 4 for limits.
- 21 ARCHAMBAULT 09 give $\sigma < 0.4$ pb (90% CL) for spin-dependent X^0 -proton cross section.
- 22 LEBEDENKO 09A give $\sigma < 0.8$ (0.01) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 23 ANGLE 08A give $\sigma < 0.9$ (0.01) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 24 BEDNYAKOV 08 reanalyze KLAPDOR-KLEINGROTHAUS 05 and BAUDIS 01 data and give $\sigma < 0.05$ pb (90% CL) for spin-dependent X^0 -neutron cross section.
- 25 ALNER 07 give $\sigma < 15$ (0.08) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 26 LEE 07A give $\sigma < 0.2$ (6) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 27 MIUCHI 07 give $\sigma < 1 \times 10^4$ pb (90% CL) for spin-dependent X^0 -proton cross section with a direction-sensitive detector.
- 28 AKERIB 06 give $\sigma < 5$ (0.07) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section. See also AKERIB 05.
- 29 SHIMIZU 06A give $\sigma < 2$ (30) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 30 ALNER 05 give $\sigma < 0.3$ (10) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 31 BARNABE-HEIDER 05 give $\sigma < 2$ (30) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 32 BENOIT 05 give $\sigma < 100$ (0.7) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 33 GIRARD 05 give $\sigma < 1.5$ pb (90% CL) for spin-dependent X^0 -proton cross section.
- 34 GIULIANI 05 analyzes the spin-independent X^0 -nucleon cross section limits with both isoscalar and isovector couplings. See Figs. 3 and 4 for limits on the couplings.
- 35 GIULIANI 05A analyze available data and give combined limits $\sigma < 0.7$ (0.2) pb for spin-dependent X^0 -proton (neutron) cross section.
- 36 KLAPDOR-KLEINGROTHAUS 05 give $\sigma < 1.5$ pb (90% CL) for spin-dependent X^0 -neutron cross section.
- 37 GIULIANI 04 reanalyze COLLAR 00 data and give limits for spin-dependent X^0 -proton and neutron couplings.
- 38 GIULIANI 04A gives limits for spin-dependent X^0 -proton and neutron couplings from existing data.
- 39 MIUCHI 03 give model-independent limit for spin-dependent X^0 -proton and neutron cross sections. See their Fig. 5.
- 40 MIUCHI 03 give model-independent limit $\sigma < 35$ pb (90% CL) for spin-dependent X^0 -proton cross section.
- 41 TAKEDA 03 give model-independent limit $\sigma < 0.04$ (0.8) nb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 42 ANGLOHER 02 limit is for spin-dependent WIMP-Aluminum cross section.
- 43 BELLI 02 discuss dependence of the extracted WIMP cross section on the assumptions of the galactic halo structure.

- 44 BERNABEI 02C analyze the DAMA data in the scenario in which χ^0 scatters into a slightly heavier state as discussed by SMITH 01.
- 45 GREEN 02 discusses dependence of extracted WIMP cross section limits on the assumptions of the galactic halo structure.
- 46 ULLIO 01 disfavor the possibility that the BERNABEI 99 signal is due to spin-dependent WIMP coupling.
- 47 BENOIT 00 find four event categories in Ge detectors and suggest that low-energy surface nuclear recoils can explain anomalous events reported by UKDMC and Saclay NaI experiments.
- 48 BERNABEI 00D limit is for inelastic scattering $\chi^0 129\text{Xe} \rightarrow \chi^0 129\text{Xe}$ (39.58 keV).
- 49 AMBROSIO 99 search for upgoing muon events induced by neutrinos originating from WIMP annihilations in the Sun and Earth.
- 50 BRHLIK 99 discuss the effect of astrophysical uncertainties on the WIMP interpretation of the BERNABEI 99 signal.
- 51 KLIMENKO 98 limit is for inelastic scattering $\chi^0 73\text{Ge} \rightarrow \chi^0 73\text{Ge}^*$ (13.26 keV).
- 52 KLIMENKO 98 limit is for inelastic scattering $\chi^0 73\text{Ge} \rightarrow \chi^0 73\text{Ge}^*$ (66.73 keV).
- 53 BELLI 96 limit for inelastic scattering $\chi^0 129\text{Xe} \rightarrow \chi^0 129\text{Xe}^*$ (39.58 keV).
- 54 BELLI 96C use background subtraction and obtain $\sigma < 0.35 \text{ pb}$ ($< 0.15 \text{ fb}$) (90% CL) for spin-dependent (independent) χ^0 -proton cross section. The confidence level is from R. Bernabei, private communication, May 20, 1999.
- 55 BERNABEI 96 use pulse shape discrimination to enhance the possible signal. The limit here is from R. Bernabei, private communication, September 19, 1997.
- 56 SARSA 96 search for annual modulation of WIMP signal. See SARSA 97 for details of the analysis. The limit here is from M.L. Sarsa, private communication, May 26, 1997.
- 57 SMITH 96 use pulse shape discrimination to enhance the possible signal. A dark matter density of 0.4 GeV cm^{-3} is assumed.
- 58 GARCIA 95 limit is from the event rate. A weaker limit is obtained from searches for diurnal and annual modulation.
- 59 SNOWDEN-IFFT 95 look for recoil tracks in an ancient mica crystal. Similar limits are also given for ^{27}Al and ^{28}Si . See COLLAR 96 and SNOWDEN-IFFT 96 for discussion on potential backgrounds.
- 60 BECK 94 uses enriched ^{76}Ge (86% purity).
- 61 REUSSER 91 limit here is changed from published (0.3) after reanalysis by authors. J.L. Vuilleumier, private communication, March 29, 1996.

For $m_{\chi^0} = 1 \text{ TeV}$

VALUE (nb)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1		AHLEN 11	DMTP	CF ₄
2		AHMED 11	CDM2	Ge, inelastic
3		AHMED 11A	RVUE	Ge
4		APRILE 11	X100	Xe
5		APRILE 11A	X100	Xe, inelastic
6		APRILE 11B	X100	Xe
7		ARMENGAUD 11	EDE2	Ge
8		BEHNKE 11	COUP	CF ₃ I
9		HORN 11	ZEP3	Xe
10		TANAKA 11	SKAM	H, solar ν
11		ABBASI 10	ICCB	KK dark matter
12		APRILE 10	X100	Xe
13		ARMENGAUD 10	EDE2	Ge

		14	MIUCHI	10	NAGE	CF ₄
		15	ABBASI	09B	ICCB	H, solar ν
		16	AHMED	09	CDM2	Ge
		17	ARCHAMBAU..	09	PICA	F
		18	LEBEDENKO	09A	ZEP3	Xe
		19	ANGLE	08A	XE10	Xe
		20	BEDNYAKOV	08	RVUE	Ge
		21	ALNER	07	ZEP2	Xe
		22	LEE	07A	KIMS	CsI
		23	MIUCHI	07	CNTR	F (CF ₄)
		24	AKERIB	06	CDMS	⁷³ Ge, ²⁹ Si
		25	ALNER	05	NAIA	NaI
		26	BARNABE-HE..	05	PICA	F (C ₄ F ₁₀)
		27	BENOIT	05	EDEL	⁷³ Ge
		28	GIRARD	05	SMPL	F (C ₂ ClF ₅)
		29	KLAPDOR-K...	05	HDMS	⁷³ Ge (enriched)
		30	MIUCHI	03	BOLO	LiF
		31	TAKEDA	03	BOLO	NaF
< 3	90	32	ANGLOHER	02	CRES	Al
		33	BENOIT	00	EDEL	Ge
		34	BERNABEI	99D	CNTR	SIMP
		35	DERBIN	99	CNTR	SIMP
< 0.06	95	36	KLIMENKO	98	CNTR	⁷³ Ge, inel.
< 0.4	95	37	KLIMENKO	98	CNTR	⁷³ Ge, inel.
< 40			ALESSAND...	96	CNTR	O
< 700			ALESSAND...	96	CNTR	Te
< 0.05	90	38	BELLI	96	CNTR	¹²⁹ Xe, inel.
< 1.5	90	39	BELLI	96	CNTR	¹²⁹ Xe, inel.
		40	BELLI	96C	CNTR	¹²⁹ Xe
< 0.01	90	41	BERNABEI	96	CNTR	Na
< 9	90	41	BERNABEI	96	CNTR	I
< 7	95	42	SARSA	96	CNTR	Na
< 0.3	90	43	SMITH	96	CNTR	Na
< 6	90	43	SMITH	96	CNTR	I
< 6	95	44	GARCIA	95	CNTR	Natural Ge
< 8	95		QUENBY	95	CNTR	Na
< 50	95		QUENBY	95	CNTR	I
< 7 × 10 ²	90	45	SNOWDEN-...	95	MICA	¹⁶ O
< 1 × 10 ³	90	45	SNOWDEN-...	95	MICA	³⁹ K
< 0.8	90	46	BECK	94	CNTR	⁷⁶ Ge
< 30	90		BACCI	92	CNTR	Na
< 30	90		BACCI	92	CNTR	I
< 15	90	47	REUSSER	91	CNTR	Natural Ge
< 6	95		CALDWELL	88	CNTR	Natural Ge

¹ AHLEN 11 give $\sigma < 8 \times 10^3$ pb (90% CL) for spin-dependent X^0 -proton cross section.

² AHMED 11 search for X^0 inelastic scattering. See their Figs. 8–10 for limits.

³ AHMED 11A combine CDMS and EDELWEISS data and give $\sigma < 1.5 \times 10^{-7}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.

- 4 APRILE 11 reanalyze APRILE 10 data and give $\sigma < 2 \times 10^{-7}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.
- 5 APRILE 11A search for X^0 inelastic scattering. See their Figs. 2 and 3 for limits.
- 6 APRILE 11B give $\sigma < 8 \times 10^{-8}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.
- 7 ARMENGAUD 11 give $\sigma < 2 \times 10^{-7}$ pb (90% CL) for spin-independent X^0 -nucleon cross section. Supersedes ARMENGAUD 10. A limit on inelastic cross section is also given.
- 8 BEHNKE 11 give $\sigma < 0.4$ pb (90% CL) for spin-dependent X^0 -proton cross section with a direction sensitive detector.
- 9 HORN 11 perform detector calibration by neutrons. Earlier results are only marginally affected.
- 10 TANAKA 11 search for neutrinos produced by X^0 annihilation in the Sun and give $\sigma < 2 \times 10^{-3}$ (2×10^{-2}) pb (90% CL) for spin-dependent X^0 -proton cross section, if X^0 pairs annihilate to $W^+ W^- (b\bar{b})$.
- 11 ABBASI 10 search for ν_μ from annihilations of Kaluza-Klein photon dark matter in the Sun and give $\sigma < 1 \times 10^{-3}$ pb (90% CL) for spin-dependent X^0 -proton cross section.
- 12 APRILE 10 give $\sigma < 4 \times 10^{-7}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.
- 13 ARMENGAUD 10 give $\sigma < 6 \times 10^{-7}$ pb (90% CL) for spin-independent X^0 -nucleon cross section.
- 14 MIUCHI 10 give $\sigma < 2 \times 10^4$ pb (90% CL) for spin-dependent X^0 -proton cross section with a direction sensitive detector.
- 15 ABBASI 09B search for neutrinos produced by X^0 annihilation in the Sun and give $\sigma < 8.7 \times 10^{-4}$ (2.2×10^{-2}) pb (90% CL) for spin-dependent X^0 -proton cross section, if X^0 pairs annihilate to $W^+ W^- (b\bar{b})$.
- 16 AHMED 09 give $\sigma < 0.2$ pb (90% CL) for spin-dependent X^0 -neutron cross section. Superseded by AHMED 10.
- 17 ARCHAMBAULT 09 give $\sigma < 3$ pb (90% CL) for spin-dependent X^0 -proton cross section.
- 18 LEBEDENKO 09A give $\sigma < 6$ (0.1) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 19 ANGLE 08A give $\sigma < 8$ (0.1) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 20 BEDNYAKOV 08 reanalyze KLAPDOR-KLEINGROTHAUS 05 and BAUDIS 01 data and give $\sigma < 0.25$ pb (90% CL) for spin-dependent X^0 -neutron cross section.
- 21 ALNER 07 give $\sigma < 100$ (0.6) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 22 LEE 07A give $\sigma < 0.8$ (30) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 23 MIUCHI 07 give $\sigma < 4 \times 10^4$ pb (90% CL) for spin-dependent X^0 -proton cross section with a direction-sensitive detector.
- 24 AKERIB 06 give $\sigma < 30$ (0.5) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section. See also AKERIB 05.
- 25 ALNER 05 give $\sigma < 1.5$ (40) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 26 BARNABE-HEIDER 05 give $\sigma < 15$ (200) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 27 BENOIT 05 give $\sigma < 600$ (4) pb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 28 GIRARD 05 give $\sigma < 10$ pb (90% CL) for spin-dependent X^0 -proton cross section.
- 29 KLAPDOR-KLEINGROTHAUS 05 give $\sigma < 10$ pb (90% CL) for spin-dependent X^0 -neutron cross section.
- 30 MIUCHI 03 give model-independent limit $\sigma < 260$ pb (90% CL) for spin-dependent X^0 -proton cross section.

- 31 TAKEDA 03 give model-independent limit $\sigma < 0.15$ (4) nb (90% CL) for spin-dependent X^0 -proton (neutron) cross section.
- 32 ANGLOHER 02 limit is for spin-dependent WIMP-Aluminum cross section.
- 33 BENOIT 00 find four event categories in Ge detectors and suggest that low-energy surface nuclear recoils can explain anomalous events reported by UKDMC and Saclay NaI experiments.
- 34 BERNABEI 99D search for SIMPs (Strongly Interacting Massive Particles) in the mass range 10^3 – 10^{16} GeV. See their Fig. 3 for cross-section limits.
- 35 DERBIN 99 search for SIMPs (Strongly Interacting Massive Particles) in the mass range 10^2 – 10^{14} GeV. See their Fig. 3 for cross-section limits.
- 36 KLIMENKO 98 limit is for inelastic scattering $X^0 \text{ } ^{73}\text{Ge} \rightarrow X^0 \text{ } ^{73}\text{Ge}^*$ (13.26 keV).
- 37 KLIMENKO 98 limit is for inelastic scattering $X^0 \text{ } ^{73}\text{Ge} \rightarrow X^0 \text{ } ^{73}\text{Ge}^*$ (66.73 keV).
- 38 BELLI 96 limit for inelastic scattering $X^0 \text{ } ^{129}\text{Xe} \rightarrow X^0 \text{ } ^{129}\text{Xe}^*$ (39.58 keV).
- 39 BELLI 96 limit for inelastic scattering $X^0 \text{ } ^{129}\text{Xe} \rightarrow X^0 \text{ } ^{129}\text{Xe}^*$ (236.14 keV).
- 40 BELLI 96C use background subtraction and obtain $\sigma < 0.7$ pb (< 0.7 fb) (90% CL) for spin-dependent (independent) X^0 -proton cross section. The confidence level is from R. Bernabei, private communication, May 20, 1999.
- 41 BERNABEI 96 use pulse shape discrimination to enhance the possible signal. The limit here is from R. Bernabei, private communication, September 19, 1997.
- 42 SARSA 96 search for annual modulation of WIMP signal. See SARSA 97 for details of the analysis. The limit here is from M.L. Sarsa, private communication, May 26, 1997.
- 43 SMITH 96 use pulse shape discrimination to enhance the possible signal. A dark matter density of 0.4 GeV cm^{-3} is assumed.
- 44 GARCIA 95 limit is from the event rate. A weaker limit is obtained from searches for diurnal and annual modulation.
- 45 SNOWDEN-IFFT 95 look for recoil tracks in an ancient mica crystal. Similar limits are also given for ^{27}Al and ^{28}Si . See COLLAR 96 and SNOWDEN-IFFT 96 for discussion on potential backgrounds.
- 46 BECK 94 uses enriched ^{76}Ge (86% purity).
- 47 REUSSER 91 limit here is changed from published (5) after reanalysis by authors. J.L. Vuilleumier, private communication, March 29, 1996.

————— X^0 Annihilation Cross Section —————

Limits are on σv for X^0 pair annihilation at threshold.

VALUE (cm^3s^{-1})	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<10^{-22}$	90	¹ ABBASI 11C	ICCB	Galctic halo, $m=1$ TeV
$<3 \times 10^{-25}$	95	² ABRAMOWSKI11	HESS	Near Galactic center, $m=1$ TeV
$<10^{-26}$	95	³ ACKERMANN 11	FLAT	Satellite galaxy, $m=10$ GeV
$<10^{-25}$	95	³ ACKERMANN 11	FLAT	Satellite galaxy, $m=100$ GeV
$<10^{-24}$	95	³ ACKERMANN 11	FLAT	Satellite galaxy, $m=1$ TeV

¹ ABBASI 11C search for ν_μ from X^0 annihilation in the outer halo of the Milky Way. The limit assumes annihilation into $\nu\nu$. See their Fig. 9 for limits with other annihilation channels.

² ABRAMOWSKI 11 search for γ from X^0 annihilation near the Galactic center. The limit assumes Einasto DM density profile.

³ ACKERMANN 11 search for γ from X^0 annihilation in ten dwarf spheroidal satellite galaxies of the Milky Way. The limit for $m = 10$ GeV assumes annihilation into $b\bar{b}$, the others W^+W^- . See their Fig. 2 for limits with other final states. See also GERINGER-SAMETH 11 for a different analysis of the same data.

CONCENTRATION OF STABLE PARTICLES IN MATTER**Concentration of Heavy (Charge +1) Stable Particles in Matter**

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<4 \times 10^{-17}$	95	¹ YAMAGATA	93	SPEC Deep sea water, $M=5-1600m_p$
$<6 \times 10^{-15}$	95	² VERKERK	92	SPEC Water, $M= 10^5$ to 3×10^7 GeV
$<7 \times 10^{-15}$	95	² VERKERK	92	SPEC Water, $M= 10^4$, 6×10^7 GeV
$<9 \times 10^{-15}$	95	² VERKERK	92	SPEC Water, $M= 10^8$ GeV
$<3 \times 10^{-23}$	90	³ HEMMICK	90	SPEC Water, $M = 1000m_p$
$<2 \times 10^{-21}$	90	³ HEMMICK	90	SPEC Water, $M = 5000m_p$
$<3 \times 10^{-20}$	90	³ HEMMICK	90	SPEC Water, $M = 10000m_p$
$<1. \times 10^{-29}$		SMITH	82B	SPEC Water, $M=30-400m_p$
$<2. \times 10^{-28}$		SMITH	82B	SPEC Water, $M=12-1000m_p$
$<1. \times 10^{-14}$		SMITH	82B	SPEC Water, $M >1000 m_p$
$<(0.2-1.) \times 10^{-21}$		SMITH	79	SPEC Water, $M=6-350 m_p$

¹ YAMAGATA 93 used deep sea water at 4000 m since the concentration is enhanced in deep sea due to gravity.

² VERKERK 92 looked for heavy isotopes in sea water and put a bound on concentration of stable charged massive particle in sea water. The above bound can be translated into into a bound on charged dark matter particle (5×10^6 GeV), assuming the local density, $\rho=0.3$ GeV/cm³, and the mean velocity $\langle v \rangle=300$ km/s.

³ See HEMMICK 90 Fig. 7 for other masses 100–10000 m_p .

Concentration of Heavy Stable Particles Bound to Nuclei

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<1.2 \times 10^{-11}$	95	¹ JAVORSEK	01	SPEC Au, $M= 3$ GeV
$<6.9 \times 10^{-10}$	95	¹ JAVORSEK	01	SPEC Au, $M= 144$ GeV
$<1 \times 10^{-11}$	95	² JAVORSEK	01B	SPEC Au, $M= 188$ GeV
$<1 \times 10^{-8}$	95	² JAVORSEK	01B	SPEC Au, $M= 1669$ GeV
$<6 \times 10^{-9}$	95	² JAVORSEK	01B	SPEC Fe, $M= 188$ GeV
$<1 \times 10^{-8}$	95	² JAVORSEK	01B	SPEC Fe, $M= 647$ GeV
$<4 \times 10^{-20}$	90	³ HEMMICK	90	SPEC C, $M = 100m_p$
$<8 \times 10^{-20}$	90	³ HEMMICK	90	SPEC C, $M = 1000m_p$
$<2 \times 10^{-16}$	90	³ HEMMICK	90	SPEC C, $M = 10000m_p$
$<6 \times 10^{-13}$	90	³ HEMMICK	90	SPEC Li, $M = 1000m_p$
$<1 \times 10^{-11}$	90	³ HEMMICK	90	SPEC Be, $M = 1000m_p$
$<6 \times 10^{-14}$	90	³ HEMMICK	90	SPEC B, $M = 1000m_p$
$<4 \times 10^{-17}$	90	³ HEMMICK	90	SPEC O, $M = 1000m_p$
$<4 \times 10^{-15}$	90	³ HEMMICK	90	SPEC F, $M = 1000m_p$
$< 1.5 \times 10^{-13}$ /nucleon	68	⁴ NORMAN	89	SPEC $^{206}\text{Pb}X^-$
$< 1.2 \times 10^{-12}$ /nucleon	68	⁴ NORMAN	87	SPEC $^{56,58}\text{Fe}X^-$

- ¹ JAVORSEK 01 search for (neutral) SIMPs (strongly interacting massive particles) bound to Au nuclei. Here M is the effective SIMP mass.
- ² JAVORSEK 01B search for (neutral) SIMPs (strongly interacting massive particles) bound to Au and Fe nuclei from various origins with exposures on the earth's surface, in a satellite, heavy ion collisions, etc. Here M is the mass of the anomalous nucleus. See also JAVORSEK 02.
- ³ See HEMMICK 90 Fig. 7 for other masses 100–10000 m_p .
- ⁴ Bound valid up to $m_{X^-} \sim 100$ TeV.

GENERAL NEW PHYSICS SEARCHES

This subsection lists some of the search experiments which look for general signatures characteristic of new physics, independent of the framework of a specific model.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
¹	AAD	11S ATLS	jet + \cancel{E}_T
²	AALTONEN	11AF CDF	$\ell^\pm \ell^\pm$
³	CHATRCHYAN	11C CMS	$\ell^+ \ell^- + \text{jets} + \cancel{E}_T$
⁴	CHATRCHYAN	11U CMS	jet + \cancel{E}_T
⁵	AALTONEN	10AF CDF	$\gamma\gamma + \ell, \cancel{E}_T$
⁶	AALTONEN	09AF CDF	$\ell\gamma b \cancel{E}_T$
⁷	AALTONEN	09G CDF	$\ell\ell\ell \cancel{E}_T$

- ¹ AAD 11S search for events with one jet and missing E_T in pp collisions at $E_{\text{cm}} = 7$ TeV with $L = 33 \text{ pb}^{-1}$. The observed events are compatible with Standard Model expectation.
 - ² AALTONEN 11AF search for high- p_T like-sign dileptons in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV with $L = 6.1 \text{ fb}^{-1}$. The observed events are compatible with Standard Model expectation.
 - ³ CHATRCHYAN 11C search for events with an opposite-sign lepton pair, jets, and missing E_T in pp collisions at $E_{\text{cm}} = 7$ TeV with $L = 34 \text{ pb}^{-1}$. The observed events are compatible with Standard Model expectation.
 - ⁴ CHATRCHYAN 11U search for events with one jet and missing E_T in pp collisions at $E_{\text{cm}} = 7$ TeV with $L = 36 \text{ pb}^{-1}$. The observed events are compatible with Standard Model expectation.
 - ⁵ AALTONEN 10AF search for $\gamma\gamma$ events with e, μ, τ , or missing E_T in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV with $L = 1.1\text{--}2.0 \text{ fb}^{-1}$. The observed events are compatible with Standard Model expectation.
 - ⁶ AALTONEN 09AF search for $\ell\gamma b$ events with missing E_T in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV with $L = 1.9 \text{ fb}^{-1}$. The observed events are compatible with Standard Model expectation including $t\bar{t}\gamma$ production.
 - ⁷ AALTONEN 09G search for $\mu\mu\mu$ and $\mu\mu e$ events with missing E_T in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV with $L = 976 \text{ pb}^{-1}$. The observed events are compatible with Standard Model expectation.
-

LIMITS ON JET-JET RESONANCES

Heavy Particle Production Cross Section

Limits are for a particle decaying to two hadronic jets.

Units(pb)	CL%	Mass(GeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
			¹ AAD	11AG ATLS	7 TeV $pp \rightarrow 2$ jets
			² AALTONEN	11M CDF	1.96 TeV $p\bar{p} \rightarrow W + 2$ jets
			³ ABAZOV	11i D0	1.96 TeV $p\bar{p} \rightarrow W + 2$ jets
			⁴ AAD	10 ATLS	7 TeV $pp \rightarrow 2$ jets
			⁵ KHACHATRYAN	10 CMS	7 TeV $pp \rightarrow 2$ jets
			⁶ ABE	99F CDF	1.8 TeV $p\bar{p} \rightarrow b\bar{b} + \text{anything}$
			⁷ ABE	97G CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets
<2603	95	200	⁸ ABE	93G CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets
< 44	95	400	⁸ ABE	93G CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets
< 7	95	600	⁸ ABE	93G CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets

¹ AAD 11AG search for dijet resonances in pp collisions at $E_{\text{cm}} = 7$ TeV with $L = 36 \text{ pb}^{-1}$. Limits on number of events for $m = 0.6\text{--}4$ TeV are given in their Table 3.

² AALTONEN 11M find a peak in two jet invariant mass distribution around 140 GeV in $W + 2$ jet events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV with $L = 4.3 \text{ fb}^{-1}$.

³ ABAZOV 11i search for two-jet resonances in $W + 2$ jet events in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV with $L = 4.3 \text{ fb}^{-1}$ and give limits $\sigma < (2.6\text{--}1.3) \text{ pb}$ (95% CL) for $m = 110\text{--}170$ GeV. The result is incompatible with AALTONEN 11M.

⁴ AAD 10 search for narrow dijet resonances in pp collisions at $E_{\text{cm}} = 7$ TeV with $L = 315 \text{ nb}^{-1}$. Limits on the cross section in the range $10\text{--}10^3 \text{ pb}$ is given for $m = 0.3\text{--}1.7$ TeV.

⁵ KHACHATRYAN 10 search for narrow dijet resonances in pp collisions at $E_{\text{cm}} = 7$ TeV with $L = 2.9 \text{ pb}^{-1}$. Limits on the cross section in the range $1\text{--}300 \text{ pb}$ is given for $m = 0.5\text{--}2.6$ TeV separately in the final states qq , qg , and gg .

⁶ ABE 99F search for narrow $b\bar{b}$ resonances in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.8$ TeV. Limits on $\sigma(p\bar{p} \rightarrow X + \text{anything}) \times B(X \rightarrow b\bar{b})$ in the range $3\text{--}10^3 \text{ pb}$ (95%CL) are given for $m_X = 200\text{--}750$ GeV. See their Table I.

⁷ ABE 97G search for narrow dijet resonances in $p\bar{p}$ collisions with 106 pb^{-1} of data at $E_{\text{cm}} = 1.8$ TeV. Limits on $\sigma(p\bar{p} \rightarrow X + \text{anything}) \cdot B(X \rightarrow jj)$ in the range $10^4\text{--}10^{-1} \text{ pb}$ (95%CL) are given for dijet mass $m = 200\text{--}1150$ GeV with both jets having $|\eta| < 2.0$ and the dijet system having $|\cos\theta^*| < 0.67$. See their Table I for the list of limits. Supersedes ABE 93G.

⁸ ABE 93G gives cross section times branching ratio into light (d , u , s , c , b) quarks for $\Gamma = 0.02 M$. Their Table II gives limits for $M = 200\text{--}900$ GeV and $\Gamma = (0.02\text{--}0.2) M$.

LIMITS ON NEUTRAL PARTICLE PRODUCTION

Production Cross Section of Radiatively-Decaying Neutral Particle

VALUE (pb)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<(0.043–0.17)	95	¹ ABBIENDI	00D OPAL	$e^+e^- \rightarrow X^0 Y^0$, $X^0 \rightarrow Y^0 \gamma$
<(0.05–0.8)	95	² ABBIENDI	00D OPAL	$e^+e^- \rightarrow X^0 X^0$, $X^0 \rightarrow Y^0 \gamma$
<(2.5–0.5)	95	³ ACKERSTAFF	97B OPAL	$e^+e^- \rightarrow X^0 Y^0$, $X^0 \rightarrow Y^0 \gamma$
<(1.6–0.9)	95	⁴ ACKERSTAFF	97B OPAL	$e^+e^- \rightarrow X^0 X^0$, $X^0 \rightarrow Y^0 \gamma$

- ¹ ABBIENDI 00D associated production limit is for $m_{\chi^0} = 90\text{--}188$ GeV, $m_{\gamma^0} = 0$ at $E_{\text{cm}} = 189$ GeV. See also their Fig. 9.
- ² ABBIENDI 00D pair production limit is for $m_{\chi^0} = 45\text{--}94$ GeV, $m_{\gamma^0} = 0$ at $E_{\text{cm}} = 189$ GeV. See also their Fig. 12.
- ³ ACKERSTAFF 97B associated production limit is for $m_{\chi^0} = 80\text{--}160$ GeV, $m_{\gamma^0} = 0$ from 10.0 pb^{-1} at $E_{\text{cm}} = 161$ GeV. See their Fig. 3(a).
- ⁴ ACKERSTAFF 97B pair production limit is for $m_{\chi^0} = 40\text{--}80$ GeV, $m_{\gamma^0} = 0$ from 10.0 pb^{-1} at $E_{\text{cm}} = 161$ GeV. See their Fig. 3(b).

Heavy Particle Production Cross Section

VALUE (cm^2/N)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

			¹ ADAMS	97B	KTEV	$m = 1.2\text{--}5$ GeV
$< 10^{-36}\text{--}10^{-33}$	90		² GALLAS	95	TOF	$m = 0.5\text{--}20$ GeV
$< (4\text{--}0.3) \times 10^{-31}$	95		³ AKESSON	91	CNTR	$m = 0\text{--}5$ GeV
$< 2 \times 10^{-36}$	90	0	⁴ BADIER	86	BDMP	$\tau = (0.05\text{--}1.) \times 10^{-8}$ s
$< 2.5 \times 10^{-35}$		0	⁵ GUSTAFSON	76	CNTR	$\tau > 10^{-7}$ s

- ¹ ADAMS 97B search for a hadron-like neutral particle produced in pN interactions, which decays into a ρ^0 and a weakly interacting massive particle. Upper limits are given for the ratio to K_L production for the mass range 1.2–5 GeV and lifetime $10^{-9}\text{--}10^{-4}$ s. See also our Light Gluino Section.
- ² GALLAS 95 limit is for a weakly interacting neutral particle produced in 800 GeV/c pN interactions decaying with a lifetime of $10^{-4}\text{--}10^{-8}$ s. See their Figs. 8 and 9. Similar limits are obtained for a stable particle with interaction cross section $10^{-29}\text{--}10^{-33} \text{ cm}^2$. See Fig. 10.
- ³ AKESSON 91 limit is from weakly interacting neutral long-lived particles produced in pN reaction at 450 GeV/c performed at CERN SPS. Bourquin-Gaillard formula is used as the production model. The above limit is for $\tau > 10^{-7}$ s. For $\tau > 10^{-9}$ s, $\sigma < 10^{-30} \text{ cm}^2/\text{nucleon}$ is obtained.
- ⁴ BADIER 86 looked for long-lived particles at 300 GeV π^- beam dump. The limit applies for nonstrongly interacting neutral or charged particles with mass > 2 GeV. The limit applies for particle modes, $\mu^+ \pi^-$, $\mu^+ \mu^-$, $\pi^+ \pi^- X$, $\pi^+ \pi^- \pi^\pm$ etc. See their figure 5 for the contours of limits in the mass- τ plane for each mode.
- ⁵ GUSTAFSON 76 is a 300 GeV FNAL experiment looking for heavy ($m > 2$ GeV) long-lived neutral hadrons in the M4 neutral beam. The above typical value is for $m = 3$ GeV and assumes an interaction cross section of 1 mb. Values as a function of mass and interaction cross section are given in figure 2.

Production of New Penetrating Non- ν Like States in Beam Dump

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ LOSECCO 81 CALO 28 GeV protons

- ¹ No excess neutral-current events leads to $\sigma(\text{production}) \times \sigma(\text{interaction}) \times \text{acceptance} < 2.26 \times 10^{-71} \text{ cm}^4/\text{nucleon}^2$ (CL = 90%) for light neutrals. Acceptance depends on models (0.1 to $4. \times 10^{-4}$).

LIMITS ON CHARGED PARTICLES IN e^+e^- **Heavy Particle Production Cross Section in e^+e^-**

Ratio to $\sigma(e^+e^- \rightarrow \mu^+\mu^-)$ unless noted. See also entries in Free Quark Search and Magnetic Monopole Searches.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • •					We do not use the following data for averages, fits, limits, etc. • • •
			¹ ACKERSTAFF 98P	OPAL	$Q=1,2/3, m=45-89.5$ GeV
			² ABREU 97D	DLPH	$Q=1,2/3, m=45-84$ GeV
			³ BARATE 97K	ALEP	$Q=1, m=45-85$ GeV
$<2 \times 10^{-5}$	95		⁴ AKERS 95R	OPAL	$Q=1, m=5-45$ GeV
$<1 \times 10^{-5}$	95		⁴ AKERS 95R	OPAL	$Q=2, m=5-45$ GeV
$<2 \times 10^{-3}$	90		⁵ BUSKULIC 93C	ALEP	$Q=1, m=32-72$ GeV
$<(10^{-2}-1)$	95		⁶ ADACHI 90C	TOPZ	$Q=1, m=1-16,$ 18-27 GeV
$<7 \times 10^{-2}$	90		⁷ ADACHI 90E	TOPZ	$Q=1, m=5-25$ GeV
$<1.6 \times 10^{-2}$	95	0	⁸ KINOSHITA 82	PLAS	$Q=3-180, m < 14.5$ GeV
$<5.0 \times 10^{-2}$	90	0	⁹ BARTEL 80	JADE	$Q=(3,4,5)/3$ 2-12 GeV

¹ ACKERSTAFF 98P search for pair production of long-lived charged particles at E_{cm} between 130 and 183 GeV and give limits $\sigma < (0.05-0.2)$ pb (95%CL) for spin-0 and spin-1/2 particles with $m=45-89.5$ GeV, charge 1 and 2/3. The limit is translated to the cross section at $E_{cm}=183$ GeV with the s dependence described in the paper. See their Figs. 2-4.

² ABREU 97D search for pair production of long-lived particles and give limits $\sigma < (0.4-2.3)$ pb (95%CL) for various center-of-mass energies $E_{cm}=130-136, 161,$ and 172 GeV, assuming an almost flat production distribution in $\cos\theta$.

³ BARATE 97K search for pair production of long-lived charged particles at $E_{cm} = 130, 136, 161,$ and 172 GeV and give limits $\sigma < (0.2-0.4)$ pb (95%CL) for spin-0 and spin-1/2 particles with $m=45-85$ GeV. The limit is translated to the cross section at $E_{cm}=172$ GeV with the E_{cm} dependence described in the paper. See their Figs. 2 and 3 for limits on $J = 1/2$ and $J = 0$ cases.

⁴ AKERS 95R is a CERN-LEP experiment with $W_{cm} \sim m_Z$. The limit is for the production of a stable particle in multihadron events normalized to $\sigma(e^+e^- \rightarrow \text{hadrons})$. Constant phase space distribution is assumed. See their Fig. 3 for bounds for $Q = \pm 2/3, \pm 4/3$.

⁵ BUSKULIC 93C is a CERN-LEP experiment with $W_{cm} = m_Z$. The limit is for a pair or single production of heavy particles with unusual ionization loss in TPC. See their Fig. 5 and Table 1.

⁶ ADACHI 90C is a KEK-TRISTAN experiment with $W_{cm} = 52-60$ GeV. The limit is for pair production of a scalar or spin-1/2 particle. See Figs. 3 and 4.

⁷ ADACHI 90E is KEK-TRISTAN experiment with $W_{cm} = 52-61.4$ GeV. The above limit is for inclusive production cross section normalized to $\sigma(e^+e^- \rightarrow \mu^+\mu^-) \cdot \beta(3-\beta^2)/2$, where $\beta = (1 - 4m^2/W_{cm}^2)^{1/2}$. See the paper for the assumption about the production mechanism.

⁸ KINOSHITA 82 is SLAC PEP experiment at $W_{cm} = 29$ GeV using lexan and ³⁹Cr plastic sheets sensitive to highly ionizing particles.

⁹ BARTEL 80 is DESY-PETRA experiment with $W_{cm} = 27-35$ GeV. Above limit is for inclusive pair production and ranges between $1. \times 10^{-1}$ and $1. \times 10^{-2}$ depending on mass and production momentum distributions. (See their figures 9, 10, 11).

Branching Fraction of Z^0 to a Pair of Stable Charged Heavy Fermions

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<5 \times 10^{-6}$	95	¹ AKERS	95R OPAL	$m = 40.4\text{--}45.6$ GeV
$<1 \times 10^{-3}$	95	AKRAWY	90O OPAL	$m = 29\text{--}40$ GeV
¹ AKERS 95R give the 95% CL limit $\sigma(X\bar{X})/\sigma(\mu\mu) < 1.8 \times 10^{-4}$ for the pair production of singly- or doubly-charged stable particles. The limit applies for the mass range 40.4–45.6 GeV for X^\pm and < 45.6 GeV for $X^{\pm\pm}$. See the paper for bounds for $Q = \pm 2/3, \pm 4/3$.				

LIMITS ON CHARGED PARTICLES IN HADRONIC REACTIONS**Heavy Particle Production Cross Section**

<u>VALUE (nb)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<1.2 \times 10^{-3}$	95	¹ AAD	11I ATLS	$ q =10e, m=0.2\text{--}1$ TeV
$<1.0 \times 10^{-5}$	95	^{2,3} AALTONEN	09Z CDF	$m > 100$ GeV, noncolored
$<4.8 \times 10^{-5}$	95	^{2,4} AALTONEN	09Z CDF	$m > 100$ GeV, colored
$<0.31\text{--}0.04 \times 10^{-3}$	95	⁵ ABAZOV	09M D0	pair production
<0.19	95	⁶ AKTAS	04C H1	$m = 3\text{--}10$ GeV
<0.05	95	⁷ ABE	92J CDF	$m = 50\text{--}200$ GeV
$<30\text{--}130$		⁸ CARROLL	78 SPEC	$m = 2\text{--}2.5$ GeV
<100		⁹ LEIPUNER	73 CNTR	$m = 3\text{--}11$ GeV

¹ AAD 11I search for production of highly ionizing massive particles in pp collisions at $E_{\text{cm}} = 7$ TeV with $L = 3.1 \text{ pb}^{-1}$. See their Table 5 for similar limits for $|q| = 6e$ and $17e$, Table 6 for limits on pair production cross section.

² AALTONEN 09Z search for long-lived charged particles in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV with $L = 1.0 \text{ fb}^{-1}$. The limits are on production cross section for a particle of mass above 100 GeV in the region $|\eta| \lesssim 0.7, p_T > 40$ GeV, and $0.4 < \beta < 1.0$.

³ Limit for weakly interacting charge-1 particle.

⁴ Limit for up-quark like particle.

⁵ ABAZOV 09M search for pair production of long-lived charged particles in $p\bar{p}$ collisions at $E_{\text{cm}} = 1.96$ TeV with $L = 1.1 \text{ fb}^{-1}$. Limit on the cross section of $(0.31\text{--}0.04)$ pb (95% CL) is given for the mass range of 60–300 GeV, assuming the kinematics of stau pair production.

⁶ AKTAS 04C look for charged particle photoproduction at HERA with mean c.m. energy of 200 GeV.

⁷ ABE 92J look for pair production of unit-charged particles which leave detector before decaying. Limit shown here is for $m = 50$ GeV. See their Fig. 5 for different charges and stronger limits for higher mass.

⁸ CARROLL 78 look for neutral, $S = -2$ dihyperon resonance in $pp \rightarrow 2K^+ X$. Cross section varies within above limits over mass range and $p_{\text{lab}} = 5.1\text{--}5.9$ GeV/ c .

⁹ LEIPUNER 73 is an NAL 300 GeV p experiment. Would have detected particles with lifetime greater than 200 ns.

Heavy Particle Production Differential Cross Section

<u>VALUE</u> ($\text{cm}^2\text{sr}^{-1}\text{GeV}^{-1}$)	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
$<2.6 \times 10^{-36}$	90	0	¹ BALDIN	76	CNTR	– $Q=1$, $m=2.1\text{--}9.4$ GeV
$<2.2 \times 10^{-33}$	90	0	² ALBROW	75	SPEC	\pm $Q= \pm 1$, $m=4\text{--}15$ GeV
$<1.1 \times 10^{-33}$	90	0	² ALBROW	75	SPEC	\pm $Q= \pm 2$, $m=6\text{--}27$ GeV
$<8. \times 10^{-35}$	90	0	³ JOVANO...	75	CNTR	\pm $m=15\text{--}26$ GeV
$<1.5 \times 10^{-34}$	90	0	³ JOVANO...	75	CNTR	\pm $Q= \pm 2$, $m=3\text{--}10$ GeV
$<6. \times 10^{-35}$	90	0	³ JOVANO...	75	CNTR	\pm $Q= \pm 2$, $m=10\text{--}26$ GeV
$<1. \times 10^{-31}$	90	0	⁴ APPEL	74	CNTR	\pm $m=3.2\text{--}7.2$ GeV
$<5.8 \times 10^{-34}$	90	0	⁵ ALPER	73	SPEC	\pm $m=1.5\text{--}24$ GeV
$<1.2 \times 10^{-35}$	90	0	⁶ ANTIPOV	71B	CNTR	– $Q=-$, $m=2.2\text{--}2.8$
$<2.4 \times 10^{-35}$	90	0	⁷ ANTIPOV	71C	CNTR	– $Q=-$, $m=1.2\text{--}1.7$, $2.1\text{--}4$
$<2.4 \times 10^{-35}$	90	0	BINON	69	CNTR	– $Q=-$, $m=1\text{--}1.8$ GeV
$<1.5 \times 10^{-36}$		0	⁸ DORFAN	65	CNTR	Be target $m=3\text{--}7$ GeV
$<3.0 \times 10^{-36}$		0	⁸ DORFAN	65	CNTR	Fe target $m=3\text{--}7$ GeV

¹ BALDIN 76 is a 70 GeV Serpukhov experiment. Value is per Al nucleus at $\theta = 0$. For other charges in range -0.5 to -3.0 , CL = 90% limit is $(2.6 \times 10^{-36})/|(\text{charge})|$ for mass range $(2.1\text{--}9.4 \text{ GeV}) \times |(\text{charge})|$. Assumes stable particle interacting with matter as do antiprotons.

² ALBROW 75 is a CERN ISR experiment with $E_{\text{cm}} = 53$ GeV. $\theta = 40$ mr. See figure 5 for mass ranges up to 35 GeV.

³ JOVANOVIICH 75 is a CERN ISR 26+26 and 15+15 GeV pp experiment. Figure 4 covers ranges $Q = 1/3$ to 2 and $m = 3$ to 26 GeV. Value is per GeV momentum.

⁴ APPEL 74 is NAL 300 GeV pW experiment. Studies forward production of heavy (up to 24 GeV) charged particles with momenta 24–200 GeV ($-$ charge) and 40–150 GeV ($+$ charge). Above typical value is for 75 GeV and is per GeV momentum per nucleon.

⁵ ALPER 73 is CERN ISR 26+26 GeV pp experiment. $p > 0.9$ GeV, $0.2 < \beta < 0.65$.

⁶ ANTIPOV 71B is from same 70 GeV p experiment as ANTIPOV 71C and BINON 69.

⁷ ANTIPOV 71C limit inferred from flux ratio. 70 GeV p experiment.

⁸ DORFAN 65 is a 30 GeV/ c p experiment at BNL. Units are per GeV momentum per nucleus.

Long-Lived Heavy Particle Invariant Cross Section

<u>VALUE</u> ($\text{cm}^2/\text{GeV}^2/N$)	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$<5\text{--}700 \times 10^{-35}$	90	¹ BERNSTEIN	88	CNTR	
$<5\text{--}700 \times 10^{-37}$	90	¹ BERNSTEIN	88	CNTR	
$<2.5 \times 10^{-36}$	90	² THRON	85	CNTR	– $Q=1$, $m=4\text{--}12$ GeV
$<1. \times 10^{-35}$	90	² THRON	85	CNTR	+ $Q=1$, $m=4\text{--}12$ GeV
$<6. \times 10^{-33}$	90	³ ARMITAGE	79	SPEC	$m=1.87$ GeV

$<1.5 \times 10^{-33}$	90	³ ARMITAGE	79	SPEC	$m=1.5-3.0$ GeV
		⁴ BOZZOLI	79	CNTR \pm	$Q = (2/3, 1, 4/3, 2)$
$<1.1 \times 10^{-37}$	90	⁵ CUTTS	78	CNTR	$m=4-10$ GeV
$<3.0 \times 10^{-37}$	90	⁶ VIDAL	78	CNTR	$m=4.5-6$ GeV

¹BERNSTEIN 88 limits apply at $x = 0.2$ and $p_T = 0$. Mass and lifetime dependence of limits are shown in the regions: $m = 1.5-7.5$ GeV and $\tau = 10^{-8}-2 \times 10^{-6}$ s. First number is for hadrons; second is for weakly interacting particles.

²THRON 85 is FNAL 400 GeV proton experiment. Mass determined from measured velocity and momentum. Limits are for $\tau > 3 \times 10^{-9}$ s.

³ARMITAGE 79 is CERN-ISR experiment at $E_{cm} = 53$ GeV. Value is for $x = 0.1$ and $p_T = 0.15$. Observed particles at $m = 1.87$ GeV are found all consistent with being antideuterons.

⁴BOZZOLI 79 is CERN-SPS 200 GeV pN experiment. Looks for particle with τ larger than 10^{-8} s. See their figure 11-18 for production cross-section upper limits vs mass.

⁵CUTTS 78 is p Be experiment at FNAL sensitive to particles of $\tau > 5 \times 10^{-8}$ s. Value is for $-0.3 < x < 0$ and $p_T = 0.175$.

⁶VIDAL 78 is FNAL 400 GeV proton experiment. Value is for $x = 0$ and $p_T = 0$. Puts lifetime limit of $< 5 \times 10^{-8}$ s on particle in this mass range.

Long-Lived Heavy Particle Production ($\sigma(\text{Heavy Particle}) / \sigma(\pi)$)

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<10^{-8}$		¹ NAKAMURA	89	SPEC \pm	$Q = (-5/3, \pm 2)$
	0	² BUSSIERE	80	CNTR \pm	$Q = (2/3, 1, 4/3, 2)$

¹NAKAMURA 89 is KEK experiment with 12 GeV protons on Pt target. The limit applies for mass $\lesssim 1.6$ GeV and lifetime $\gtrsim 10^{-7}$ s.

²BUSSIERE 80 is CERN-SPS experiment with 200-240 GeV protons on Be and Al target. See their figures 6 and 7 for cross-section ratio vs mass.

Production and Capture of Long-Lived Massive Particles

<u>VALUE (10^{-36} cm²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<20 to 800	0	¹ ALEKSEEV	76	ELEC	$\tau=5$ ms to 1 day
<200 to 2000	0	¹ ALEKSEEV	76B	ELEC	$\tau=100$ ms to 1 day
<1.4 to 9	0	² FRANKEL	75	CNTR	$\tau=50$ ms to 10 hours
<0.1 to 9	0	³ FRANKEL	74	CNTR	$\tau=1$ to 1000 hours

¹ALEKSEEV 76 and ALEKSEEV 76B are 61-70 GeV p Serpukhov experiment. Cross section is per Pb nucleus.

²FRANKEL 75 is extension of FRANKEL 74.

³FRANKEL 74 looks for particles produced in thick Al targets by 300-400 GeV/c protons.

Long-Lived Particle Search at Hadron Collisions

Limits are for cross section times branching ratio.

<u>VALUE (pb/nucleon)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------	------------	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2	90	0	¹ BADIER	86	BDMP $\tau = (0.05-1.) \times 10^{-8}$ s
------	----	---	---------------------	----	--

¹BADIER 86 looked for long-lived particles at 300 GeV π^- beam dump. The limit applies for nonstrongly interacting neutral or charged particles with mass >2 GeV. The limit applies for particle modes, $\mu^+ \pi^-$, $\mu^+ \mu^-$, $\pi^+ \pi^- X$, $\pi^+ \pi^- \pi^\pm$ etc. See their figure 5 for the contours of limits in the mass- τ plane for each mode.

Long-Lived Heavy Particle Cross Section

<u>VALUE (pb/sr)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<34	95	¹ RAM	94	SPEC 1015 < $m_{X^{++}} < 1085$ MeV
<75	95	¹ RAM	94	SPEC 920 < $m_{X^{++}} < 1025$ MeV

¹RAM 94 search for a long-lived doubly-charged fermion X^{++} with mass between m_N and $m_N + m_\pi$ and baryon number +1 in the reaction $pp \rightarrow X^{++}n$. No candidate is found. The limit is for the cross section at 15° scattering angle at 460 MeV incident energy and applies for $\tau(X^{++}) \gg 0.1 \mu\text{s}$.

LIMITS ON CHARGED PARTICLES IN COSMIC RAYS

Heavy Particle Flux in Cosmic Rays

<u>VALUE (cm⁻²sr⁻¹s⁻¹)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
~ 6 × 10 ⁻⁹		2	¹ SAITO	90		$Q \simeq 14, m \simeq 370m_p$
< 1.4 × 10 ⁻¹²	90	0	² MINCER	85	CALO	$m \geq 1 \text{ TeV}$
< 1.7 × 10 ⁻¹¹	99	0	³ SAKUYAMA	83B	PLAS	$m \sim 1 \text{ TeV}$
< 1. × 10 ⁻⁹	90	0	⁴ BHAT	82	CC	
< 1. × 10 ⁻⁹		0	⁵ MARINI	82	CNTR ±	$Q=1, m \sim 4.5m_p$
2. × 10 ⁻⁹		3	⁶ YOCK	81	SPRK ±	$Q=1, m \sim 4.5m_p$
		3	⁶ YOCK	81	SPRK	Fractionally charged
3.0 × 10 ⁻⁹		3	⁷ YOCK	80	SPRK	$m \sim 4.5 m_p$
(4 ± 1) × 10 ⁻¹¹		3	GOODMAN	79	ELEC	$m \geq 5 \text{ GeV}$
< 1.3 × 10 ⁻⁹	90	0	⁸ BHAT	78	CNTR ±	$m > 1 \text{ GeV}$
< 1.0 × 10 ⁻⁹		0	BRIATORE	76	ELEC	
< 7. × 10 ⁻¹⁰	90	0	YOCK	75	ELEC ±	$Q > 7e \text{ or } < -7e$
> 6. × 10 ⁻⁹		5	⁹ YOCK	74	CNTR	$m > 6 \text{ GeV}$
< 3.0 × 10 ⁻⁸		0	DARDO	72	CNTR	
< 1.5 × 10 ⁻⁹		0	TONWAR	72	CNTR	$m > 10 \text{ GeV}$
< 3.0 × 10 ⁻¹⁰		0	BJORNBOE	68	CNTR	$m > 5 \text{ GeV}$
< 5.0 × 10 ⁻¹¹	90	0	JONES	67	ELEC	$m=5-15 \text{ GeV}$

¹SAITO 90 candidates carry about 450 MeV/nucleon. Cannot be accounted for by conventional backgrounds. Consistent with strange quark matter hypothesis.

²MINCER 85 is high statistics study of calorimeter signals delayed by 20–200 ns. Calibration with AGS beam shows they can be accounted for by rare fluctuations in signals from low-energy hadrons in the shower. Claim that previous delayed signals including BJORNBOE 68, DARDO 72, BHAT 82, SAKUYAMA 83B below may be due to this fake effect.

³SAKUYAMA 83B analyzed 6000 extended air shower events. Increase of delayed particles and change of lateral distribution above 10^{17} eV may indicate production of very heavy parent at top of atmosphere.

- ⁴ BHAT 82 observed 12 events with delay $> 2. \times 10^{-8}$ s and with more than 40 particles. 1 eV has good hadron shower. However all events are delayed in only one of two detectors in cloud chamber, and could not be due to strongly interacting massive particle.
- ⁵ MARINI 82 applied PEP-counter for TOF. Above limit is for velocity = 0.54 of light. Limit is inconsistent with YOCC 80 YOCC 81 events if isotropic dependence on zenith angle is assumed.
- ⁶ YOCC 81 saw another 3 events with $Q = \pm 1$ and m about $4.5m_p$ as well as 2 events with $m > 5.3m_p$, $Q = \pm 0.75 \pm 0.05$ and $m > 2.8m_p$, $Q = \pm 0.70 \pm 0.05$ and 1 event with $m = (9.3 \pm 3.)m_p$, $Q = \pm 0.89 \pm 0.06$ as possible heavy candidates.
- ⁷ YOCC 80 events are with charge exactly or approximately equal to unity.
- ⁸ BHAT 78 is at Kolar gold fields. Limit is for $\tau > 10^{-6}$ s.
- ⁹ YOCC 74 events could be tritons.

Superheavy Particle (Quark Matter) Flux in Cosmic Rays

$\frac{VALUE}{(cm^{-2}sr^{-1}s^{-1})}$	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 5 \times 10^{-16}$	90		1 AMBROSIO	00B	MCRO $m > 5 \times 10^{14}$ GeV
$< 1.8 \times 10^{-12}$	90		2 ASTONE	93	CNTR $m \geq 1.5 \times 10^{-13}$ gram
$< 1.1 \times 10^{-14}$	90		3 AHLEN	92	MCRO $10^{-10} < m < 0.1$ gram
$< 2.2 \times 10^{-14}$	90	0	4 NAKAMURA	91	PLAS $m > 10^{11}$ GeV
$< 6.4 \times 10^{-16}$	90	0	5 ORITO	91	PLAS $m > 10^{12}$ GeV
$< 2.0 \times 10^{-11}$	90		6 LIU	88	BOLO $m > 1.5 \times 10^{-13}$ gram
$< 4.7 \times 10^{-12}$	90		7 BARISH	87	CNTR $1.4 \times 10^8 < m < 10^{12}$ GeV
$< 3.2 \times 10^{-11}$	90	0	8 NAKAMURA	85	CNTR $m > 1.5 \times 10^{-13}$ gram
$< 3.5 \times 10^{-11}$	90	0	9 ULLMAN	81	CNTR Planck-mass 10^{19} GeV
$< 7. \times 10^{-11}$	90	0	9 ULLMAN	81	CNTR $m \leq 10^{16}$ GeV

- ¹ AMBROSIO 00B searched for quark matter ("nuclearites") in the velocity range $(10^{-5}-1) c$. The listed limit is for $2 \times 10^{-3} c$.
- ² ASTONE 93 searched for quark matter ("nuclearites") in the velocity range $(10^{-3}-1) c$. Their Table 1 gives a compilation of searches for nuclearites.
- ³ AHLEN 92 searched for quark matter ("nuclearites"). The bound applies to velocity $< 2.5 \times 10^{-3} c$. See their Fig. 3 for other velocity/ c and heavier mass range.
- ⁴ NAKAMURA 91 searched for quark matter in the velocity range $(4 \times 10^{-5}-1) c$.
- ⁵ ORITO 91 searched for quark matter. The limit is for the velocity range $(10^{-4}-10^{-3}) c$.
- ⁶ LIU 88 searched for quark matter ("nuclearites") in the velocity range $(2.5 \times 10^{-3}-1) c$. A less stringent limit of 5.8×10^{-11} applies for $(1-2.5) \times 10^{-3} c$.
- ⁷ BARISH 87 searched for quark matter ("nuclearites") in the velocity range $(2.7 \times 10^{-4}-5 \times 10^{-3}) c$.
- ⁸ NAKAMURA 85 at KEK searched for quark-matter. These might be lumps of strange quark matter with roughly equal numbers of u , d , s quarks. These lumps or nuclearites were assumed to have velocity of $(10^{-4}-10^{-3}) c$.
- ⁹ ULLMAN 81 is sensitive for heavy slow singly charge particle reaching earth with vertical velocity 100–350 km/s.

Highly Ionizing Particle Flux

$\frac{VALUE}{(m^{-2}yr^{-1})}$	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.4	95	0	KINOSHITA	81B	PLAS Z/β 30–100
---------	----	---	-----------	-----	-----------------------

REFERENCES FOR Searches for WIMPs and Other Particles

AAD	11AG	NJP 13 053044	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	11I	PL B698 353	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	11S	PL B705 294	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALSETH	11	PRL 106 131301	C.E. Aalseth <i>et al.</i>	(CoGeNT Collab.)
AALSETH	11A	PRL 107 141301	C.E. Aalseth <i>et al.</i>	(CoGeNT Collab.)
AALTONEN	11AF	PRL 107 181801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	11M	PRL 106 171801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	11I	PRL 107 011804	V. M. Abazov <i>et al.</i>	(D0 Collab.)
ABBASI	11C	PR D84 022004	R. Abbasi <i>et al.</i>	(IceCube Collab.)
ABRAMOWSKI	11	PRL 106 161301	A. Abramowski <i>et al.</i>	(H.E.S.S. Collab.)
ACKERMANN	11	PRL 107 241302	M. Ackermann <i>et al.</i>	(Fermi-LAT Collab.)
AHLEN	11	PL B695 124	S. Ahlen <i>et al.</i>	(DMTPC Collab.)
AHMED	11	PR D83 112002	Z. Ahmed <i>et al.</i>	(CDMS Collab.)
AHMED	11A	PR D84 011102	Z. Ahmed <i>et al.</i>	(CDMS and EDELWEISS Collabs.)
AHMED	11B	PRL 106 131302	Z. Ahmed <i>et al.</i>	(CDMS Collab.)
AJELLO	11	PR D84 032007	M. Ajello <i>et al.</i>	(Fermi-LAT)
ANGLE	11	PRL 107 051301	J. Angle <i>et al.</i>	(XENON10 Collab.)
APRILE	11	PR D84 052003	E. Aprile <i>et al.</i>	(XENON100 Collab.)
APRILE	11A	PR D84 061101	E. Aprile <i>et al.</i>	(XENON100 Collab.)
APRILE	11B	PRL 107 131302	E. Aprile <i>et al.</i>	(XENON100 Collab.)
ARMENGAUD	11	PL B702 329	E. Armengaud <i>et al.</i>	(EDELWEISS II Collab.)
BEHNKE	11	PRL 106 021303	E. Behnke <i>et al.</i>	(COUPP Collab.)
CHATRCHYAN	11C	JHEP 1106 026	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	11U	PRL 107 201804	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
GERINGER-SA...	11	PRL 107 241303	A. Geringer-Sameth, S.M. Koushiappas	
HORN	11	PL B705 471	M. Horn <i>et al.</i>	(ZEPLIN-III Collab.)
TANAKA	11	APJ 742 78	T. Tanaka <i>et al.</i>	(Super-Kamiokande Collab.)
AAD	10	PRL 105 161801	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	10AF	PR D82 052005	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABBASI	10	PR D81 057101	R. Abbasi <i>et al.</i>	(IceCube Collab.)
AHMED	10	SCI 327 1619	Z. Ahmed <i>et al.</i>	(CDMS II Collab.)
AKERIB	10	PR D82 122004	D.S. Akerib <i>et al.</i>	(CDMS-II Collab.)
AKIMOV	10	PL B692 180	D.Yu. Akimov <i>et al.</i>	(ZEPLIN-III Collab.)
APRILE	10	PRL 105 131302	E. Aprile <i>et al.</i>	(XENON100 Collab.)
ARMENGAUD	10	PL B687 294	E. Armengaud <i>et al.</i>	(EDELWEISS II Collab.)
FELIZARDO	10	PRL 105 211301	M. Felizardo <i>et al.</i>	(The SIMPLE Collab.)
KHACHATRY...	10	PRL 105 211801	V. Khachatryan <i>et al.</i>	(CMS Collab.)
Also		PRL 106 029902E	V. Khachatryan <i>et al.</i>	(CMS Collab.)
MIUCHI	10	PL B686 11	K. Miuchi <i>et al.</i>	(NEWAGE Collab.)
AALTONEN	09AF	PR D80 011102R	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09G	PR D79 052004	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09Z	PRL 103 021802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	09M	PRL 102 161802	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABBASI	09B	PRL 102 201302	R. Abbasi <i>et al.</i>	(IceCube Collab.)
AHMED	09	PRL 102 011301	Z. Ahmed <i>et al.</i>	(CDMS Collab.)
ANGLE	09	PR D80 115005	J. Angle <i>et al.</i>	(XENON10 Collab.)
ARCHAMBAU...	09	PL B682 185	S. Archambault <i>et al.</i>	(PICASSO Collab.)
LEBEDENKO	09A	PRL 103 151302	V.N. Lebedenko <i>et al.</i>	(ZEPLIN-III Collab.)
LIN	09	PR D79 061101R	S.T. Lin <i>et al.</i>	(TEXONO Collab.)
AALSETH	08	PRL 101 251301	C.E. Aalseth <i>et al.</i>	(CoGeNT Collab.)
Also		PRL 102 109903 (errat)	C.E. Aalseth <i>et al.</i>	(CoGeNT Collab.)
ANGLE	08A	PRL 101 091301	J. Angle <i>et al.</i>	(XENON10 Collab.)
BEDNYAKOV	08	PAN 71 111	V.A. Bednyakov, H.P. Klapdor-Kleingrothaus, I.V. Krivosheina	
		Translated from YAF 71 112.		
ALNER	07	PL B653 161	G.J. Alner <i>et al.</i>	(ZEPLIN-II Collab.)
LEE	07A	PRL 99 091301	H.S. Lee <i>et al.</i>	(KIMS Collab.)
MIUCHI	07	PL B654 58	K. Miuchi <i>et al.</i>	
AKERIB	06	PR D73 011102R	D.S. Akerib <i>et al.</i>	(CDMS Collab.)
SHIMIZU	06A	PL B633 195	Y. Shimizu <i>et al.</i>	
AKERIB	05	PR D72 052009	D.S. Akerib <i>et al.</i>	(CDMS Collab.)
ALNER	05	PL B616 17	G.J. Alner <i>et al.</i>	(UK Dark Matter Collab.)
BARNABE-HE...	05	PL B624 186	M. Barnabe-Heider <i>et al.</i>	(PICASSO Collab.)
BENOIT	05	PL B616 25	A. Benoit <i>et al.</i>	(EDELWEISS Collab.)
GIRARD	05	PL B621 233	T.A. Girard <i>et al.</i>	(SIMPLE Collab.)
GIULIANI	05	PRL 95 101301	F. Giuliani	
GIULIANI	05A	PR D71 123503	F. Giuliani, T.A. Girard	
KLAPDOR-K...	05	PL B609 226	H.V. Klapdor-Kleingrothaus, I.V. Krivosheina, C. Tomei	
AKTAS	04C	EPJ C36 413	A. Aktas <i>et al.</i>	(H1 Collab.)
GIULIANI	04	PL B588 151	F. Giuliani, T.A. Girard	

GIULIANI	04A	PRL 93 161301	F. Giuliani	
MIUCHI	03	ASP 19 135	K. Miuchi <i>et al.</i>	
TAKEDA	03	PL B572 145	A. Takeda <i>et al.</i>	
ANGLOHER	02	ASP 18 43	G. Angloher <i>et al.</i>	(CRESST Collab.)
BELLI	02	PR D66 043503	P. Belli <i>et al.</i>	
BERNABEI	02C	EPJ C23 61	R. Bernabei <i>et al.</i>	(DAMA Collab.)
GREEN	02	PR D66 083003	A.M. Green	
JAVORSEK	02	PR D65 072003	D. Javorsek II <i>et al.</i>	
BAUDIS	01	PR D63 022001	L. Baudis <i>et al.</i>	(Heidelberg-Moscow Collab.)
JAVORSEK	01	PR D64 012005	D. Javorsek II <i>et al.</i>	
JAVORSEK	01B	PRL 87 231804	D. Javorsek II <i>et al.</i>	
SMITH	01	PR D64 043502	D. Smith, N. Weiner	
ULLIO	01	JHEP 0107 044	P. Ullio, M. Kamionkowski, P. Vogel	
ABBIENDI	00D	EPJ C13 197	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AMBROSIO	00B	EPJ C13 453	M. Ambrosio <i>et al.</i>	(MACRO Collab.)
BENOIT	00	PL B479 8	A. Benoit <i>et al.</i>	(EDELWEISS Collab.)
BERNABEI	00D	NJP 2 15	R. Bernabei <i>et al.</i>	(DAMA Collab.)
COLLAR	00	PRL 85 3083	J.I. Collar <i>et al.</i>	(SIMPLE Collab.)
ABE	99F	PRL 82 2038	F. Abe <i>et al.</i>	(CDF Collab.)
AMBROSIO	99	PR D60 082002	M. Ambrosio <i>et al.</i>	(Macro Collab.)
BERNABEI	99	PL B450 448	R. Bernabei <i>et al.</i>	(DAMA Collab.)
BERNABEI	99D	PRL 83 4918	R. Bernabei <i>et al.</i>	(DAMA Collab.)
BRHLIK	99	PL B464 303	M. Brhlik, L. Roszkowski	
DERBIN	99	PAN 62 1886	A.V. Derbin <i>et al.</i>	
ACKERSTAFF	98P	Translated from YAF 62 2034. PL B433 195	K. Akerstaff <i>et al.</i>	(OPAL Collab.)
KLIMENKO	98	JETPL 67 875	A.A. Klimenko <i>et al.</i>	
		Translated from ZETFP 67 835.		
ABE	97G	PR D55 R5263	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	97D	PL B396 315	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACKERSTAFF	97B	PL B391 210	K. Akerstaff <i>et al.</i>	(OPAL Collab.)
ADAMS	97B	PRL 79 4083	J. Adams <i>et al.</i>	(FNAL KTeV Collab.)
BARATE	97K	PL B405 379	R. Barate <i>et al.</i>	(ALEPH Collab.)
SARSA	97	PR D56 1856	M.L. Sarsa <i>et al.</i>	(ZARA)
ALESSAND...	96	PL B384 316	A. Alessandrello <i>et al.</i>	(MILA, MILAI, SASSO)
BELLI	96	PL B387 222	P. Belli <i>et al.</i>	(DAMA Collab.)
Also		PL B389 783 (erratum)	P. Belli <i>et al.</i>	(DAMA Collab.)
BELLI	96C	NC 19C 537	P. Belli <i>et al.</i>	(DAMA Collab.)
BERNABEI	96	PL B389 757	R. Bernabei <i>et al.</i>	(DAMA Collab.)
COLLAR	96	PRL 76 331	J.I. Collar	(SCUC)
SARSA	96	PL B386 458	M.L. Sarsa <i>et al.</i>	(ZARA)
Also		PR D56 1856	M.L. Sarsa <i>et al.</i>	(ZARA)
SMITH	96	PL B379 299	P.F. Smith <i>et al.</i>	(RAL, SHEF, LOIC+)
SNOWDEN-...	96	PRL 76 332	D.P. Snowden-Ifft, E.S. Freeman, P.B. Price	(UCB)
AKERS	95R	ZPHY C67 203	R. Akers <i>et al.</i>	(OPAL Collab.)
GALLAS	95	PR D52 6	E. Gallas <i>et al.</i>	(MSU, FNAL, MIT, FLOR)
GARCIA	95	PR D51 1458	E. Garcia <i>et al.</i>	(ZARA, SCUC, PNL)
QUENBY	95	PL B351 70	J.J. Quenby <i>et al.</i>	(LOIC, RAL, SHEF+)
SNOWDEN-...	95	PRL 74 4133	D.P. Snowden-Ifft, E.S. Freeman, P.B. Price	(UCB)
Also		PRL 76 331	J.I. Collar	(SCUC)
Also		PRL 76 332	D.P. Snowden-Ifft, E.S. Freeman, P.B. Price	(UCB)
BECK	94	PL B336 141	M. Beck <i>et al.</i>	(MPIH, KIAE, SASSO)
RAM	94	PR D49 3120	S. Ram <i>et al.</i>	(TELA, TRIU)
ABE	93G	PRL 71 2542	F. Abe <i>et al.</i>	(CDF Collab.)
ASTONE	93	PR D47 4770	P. Astone <i>et al.</i>	(ROMA, ROMAI, CATA, FRAS)
BUSKULIC	93C	PL B303 198	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
YAMAGATA	93	PR D47 1231	T. Yamagata, Y. Takamori, H. Utsunomiya	(KONAN)
ABE	92J	PR D46 R1889	F. Abe <i>et al.</i>	(CDF Collab.)
AHLEN	92	PRL 69 1860	S.P. Ahlen <i>et al.</i>	(MACRO Collab.)
BACCI	92	PL B293 460	C. Bacci <i>et al.</i>	(Beijing-Roma-Saclay Collab.)
VERKERK	92	PRL 68 1116	P. Verkerk <i>et al.</i>	(ENSP, SACL, PAST)
AKESSON	91	ZPHY C52 219	T. Akesson <i>et al.</i>	(HELIOS Collab.)
NAKAMURA	91	PL B263 529	S. Nakamura <i>et al.</i>	
ORITO	91	PRL 66 1951	S. Orito <i>et al.</i>	(ICEPP, WASC, NIHO, ICRR)
REUSSER	91	PL B255 143	D. Reusser <i>et al.</i>	(NEUC, CIT, PSI)
ADACHI	90C	PL B244 352	I. Adachi <i>et al.</i>	(TOPAZ Collab.)
ADACHI	90E	PL B249 336	I. Adachi <i>et al.</i>	(TOPAZ Collab.)
AKRAWY	90O	PL B252 290	M.Z. Akrawy <i>et al.</i>	(OPAL Collab.)
HEMMICK	90	PR D41 2074	T.K. Hemmick <i>et al.</i>	(ROCH, MICH, OHIO+)
SAITO	90	PRL 65 2094	T. Saito <i>et al.</i>	(ICRR, KOBE)
NAKAMURA	89	PR D39 1261	T.T. Nakamura <i>et al.</i>	(KYOT, TMTC)

NORMAN	89	PR D39 2499	E.B. Norman <i>et al.</i>	(LBL)
BERNSTEIN	88	PR D37 3103	R.M. Bernstein <i>et al.</i>	(STAN, WISC)
CALDWELL	88	PRL 61 510	D.O. Caldwell <i>et al.</i>	(UCSB, UCB, LBL)
LIU	88	PRL 61 271	G. Liu, B. Barish	
BARISH	87	PR D36 2641	B.C. Barish, G. Liu, C. Lane	(CIT)
NORMAN	87	PRL 58 1403	E.B. Norman, S.B. Gazes, D.A. Bennett	(LBL)
BADIER	86	ZPHY C31 21	J. Badier <i>et al.</i>	(NA3 Collab.)
MINCER	85	PR D32 541	A. Mincer <i>et al.</i>	(UMD, GMAS, NSF)
NAKAMURA	85	PL 161B 417	K. Nakamura <i>et al.</i>	(KEK, INUS)
THRON	85	PR D31 451	J.L. Thron <i>et al.</i>	(YALE, FNAL, IOWA)
SAKUYAMA	83B	LNC 37 17	H. Sakuyama, N. Suzuki	(MEIS)
Also		LNC 36 389	H. Sakuyama, K. Watanabe	(MEIS)
Also		NC 78A 147	H. Sakuyama, K. Watanabe	(MEIS)
Also		NC 6C 371	H. Sakuyama, K. Watanabe	(MEIS)
BHAT	82	PR D25 2820	P.N. Bhat <i>et al.</i>	(TATA)
KINOSHITA	82	PRL 48 77	K. Kinoshita, P.B. Price, D. Fryberger	(UCB+)
MARINI	82	PR D26 1777	A. Marini <i>et al.</i>	(FRAS, LBL, NWES, STAN+)
SMITH	82B	NP B206 333	P.F. Smith <i>et al.</i>	(RAL)
KINOSHITA	81B	PR D24 1707	K. Kinoshita, P.B. Price	(UCB)
LOSECCO	81	PL 102B 209	J.M. LoSecco <i>et al.</i>	(MICH, PENN, BNL)
ULLMAN	81	PRL 47 289	J.D. Ullman	(LEHM, BNL)
YOCK	81	PR D23 1207	P.C.M. Yock	(AUCK)
BARTEL	80	ZPHY C6 295	W. Bartel <i>et al.</i>	(JADE Collab.)
BUSSIERE	80	NP B174 1	A. Bussiere <i>et al.</i>	(BGNA, SACL, LAPP)
YOCK	80	PR D22 61	P.C.M. Yock	(AUCK)
ARMITAGE	79	NP B150 87	J.C.M. Armitage <i>et al.</i>	(CERN, DARE, FOM+)
BOZZOLI	79	NP B159 363	W. Bozzoli <i>et al.</i>	(BGNA, LAPP, SACL+)
GOODMAN	79	PR D19 2572	J.A. Goodman <i>et al.</i>	(UMD)
SMITH	79	NP B149 525	P.F. Smith, J.R.J. Bennett	(RHEL)
BHAT	78	PRAM 10 115	P.N. Bhat, P.V. Ramana Murthy	(TATA)
CARROLL	78	PRL 41 777	A.S. Carroll <i>et al.</i>	(BNL, PRIN)
CUTTS	78	PRL 41 363	D. Cutts <i>et al.</i>	(BROW, FNAL, ILL, BARI+)
VIDAL	78	PL 77B 344	R.A. Vidal <i>et al.</i>	(COLU, FNAL, STON+)
ALEKSEEV	76	SJNP 22 531	G.D. Alekseev <i>et al.</i>	(JINR)
		Translated from YAF 22	1021.	
ALEKSEEV	76B	SJNP 23 633	G.D. Alekseev <i>et al.</i>	(JINR)
		Translated from YAF 23	1190.	
BALDIN	76	SJNP 22 264	B.Y. Baldin <i>et al.</i>	(JINR)
		Translated from YAF 22	512.	
BRIATORE	76	NC 31A 553	L. Briatore <i>et al.</i>	(LCGT, FRAS, FREIB)
GUSTAFSON	76	PRL 37 474	H.R. Gustafson <i>et al.</i>	(MICH)
ALBROW	75	NP B97 189	M.G. Albrow <i>et al.</i>	(CERN, DARE, FOM+)
FRANKEL	75	PR D12 2561	S. Frankel <i>et al.</i>	(PENN, FNAL)
JOVANOVI...	75	PL 56B 105	J.V. Jovanovich <i>et al.</i>	(MANI, AACH, CERN+)
YOCK	75	NP B86 216	P.C.M. Yock	(AUCK, SLAC)
APPEL	74	PRL 32 428	J.A. Appel <i>et al.</i>	(COLU, FNAL)
FRANKEL	74	PR D9 1932	S. Frankel <i>et al.</i>	(PENN, FNAL)
YOCK	74	NP B76 175	P.C.M. Yock	(AUCK)
ALPER	73	PL 46B 265	B. Alper <i>et al.</i>	(CERN, LIVP, LUND, BOHR+)
LEIPUNER	73	PRL 31 1226	L.B. Leipuner <i>et al.</i>	(BNL, YALE)
DARDO	72	NC 9A 319	M. Dardo <i>et al.</i>	(TORI)
TONWAR	72	JPA 5 569	S.C. Tonwar, S. Naranan, B.V. Sreekantan	(TATA)
ANTIPOV	71B	NP B31 235	Y.M. Antipov <i>et al.</i>	(SERP)
ANTIPOV	71C	PL 34B 164	Y.M. Antipov <i>et al.</i>	(SERP)
BINON	69	PL 30B 510	F.G. Binon <i>et al.</i>	(SERP)
BJORNBOE	68	NC B53 241	J. Bjornboe <i>et al.</i>	(BOHR, TATA, BERN+)
JONES	67	PR 164 1584	L.W. Jones	(MICH, WISC, LBL, UCLA, MINN+)
DORFAN	65	PRL 14 999	D.E. Dorfan <i>et al.</i>	(COLU)