

CHARMED MESONS

($C = \pm 1$)

$$D^+ = c\bar{d}, D^0 = c\bar{u}, \bar{D}^0 = \bar{c}u, D^- = \bar{c}d, \quad \text{similarly for } D^{*'}\text{'s}$$

D^\pm

$$I(J^P) = \frac{1}{2}(0^-)$$

$$\text{Mass } m = 1869.62 \pm 0.15 \text{ MeV} \quad (S = 1.1)$$

$$\text{Mean life } \tau = (1040 \pm 7) \times 10^{-15} \text{ s}$$

$$c\tau = 311.8 \text{ } \mu\text{m}$$

c-quark decays

$$\Gamma(c \rightarrow \ell^+ \text{ anything}) / \Gamma(c \rightarrow \text{ anything}) = 0.096 \pm 0.004 \text{ [a]}$$

$$\Gamma(c \rightarrow D^{*(2010)^+} \text{ anything}) / \Gamma(c \rightarrow \text{ anything}) = 0.255 \pm 0.017$$

CP-violation decay-rate asymmetries

$$A_{CP}(\mu^\pm \nu) = (8 \pm 8)\%$$

$$A_{CP}(K_S^0 \pi^\pm) = (-0.54 \pm 0.14)\%$$

$$A_{CP}(K^\mp 2\pi^\pm) = (-0.1 \pm 1.0)\%$$

$$A_{CP}(K^\mp \pi^\pm \pi^\pm \pi^0) = (1.0 \pm 1.3)\%$$

$$A_{CP}(K_S^0 \pi^\pm \pi^0) = (0.3 \pm 0.9)\%$$

$$A_{CP}(K_S^0 \pi^\pm \pi^+ \pi^-) = (0.1 \pm 1.3)\%$$

$$A_{CP}(\pi^\pm \pi^0) = (2.9 \pm 2.9)\%$$

$$A_{CP}(\pi^\pm \eta) = (1.0 \pm 1.5)\% \quad (S = 1.4)$$

$$A_{CP}(\pi^\pm \eta'(958)) = (-0.5 \pm 1.2)\% \quad (S = 1.1)$$

$$A_{CP}(K_S^0 K^\pm) = (-0.1 \pm 0.6)\%$$

$$A_{CP}(K^+ K^- \pi^\pm) = (0.3 \pm 0.6)\%$$

$$A_{CP}(K^\pm K^{*0}) = (0.1 \pm 1.3)\%$$

$$A_{CP}(\phi \pi^\pm) = (0.42 \pm 0.28)\%$$

$$A_{CP}(K^\pm K_0^*(1430)^0) = (8_{-6}^{+7})\%$$

$$A_{CP}(K^\pm K_2^*(1430)^0) = (43_{-26}^{+20})\%$$

$$A_{CP}(K^\pm K_0^*(800)) = (-12_{-13}^{+18})\%$$

$$A_{CP}(a_0(1450)^0 \pi^\pm) = (-19_{-16}^{+14})\%$$

$$A_{CP}(\phi(1680) \pi^\pm) = (-9 \pm 26)\%$$

$$A_{CP}(\pi^+ \pi^- \pi^\pm) = (-2 \pm 4)\%$$

$$A_{CP}(K_S^0 K^\pm \pi^+ \pi^-) = (-4 \pm 7)\%$$

$$A_{CP}(K^\pm \pi^0) = (-4 \pm 11)\%$$

T-violation decay-rate asymmetry

$$A_T(K_S^0 K^\pm \pi^+ \pi^-) = (-12 \pm 11) \times 10^{-3} \text{ [b]}$$

D^+ form factors

$$\begin{aligned}
 f_+(0)|V_{cs}| \text{ in } \bar{K}^0 \ell^+ \nu_\ell &= 0.707 \pm 0.013 \\
 r_1 \equiv a_1/a_0 \text{ in } \bar{K}^0 \ell^+ \nu_\ell &= -1.7 \pm 0.5 \\
 r_2 \equiv a_2/a_0 \text{ in } \bar{K}^0 \ell^+ \nu_\ell &= -14 \pm 11 \\
 f_+(0)|V_{cd}| \text{ in } \pi^0 \ell^+ \nu_\ell &= 0.146 \pm 0.007 \\
 r_1 \equiv a_1/a_0 \text{ in } \pi^0 \ell^+ \nu_\ell &= -1.4 \pm 0.9 \\
 r_2 \equiv a_2/a_0 \text{ in } \pi^0 \ell^+ \nu_\ell &= -4 \pm 5 \\
 f_+(0)|V_{cd}| \text{ in } D^+ \rightarrow \eta e^+ \nu_e &= 0.086 \pm 0.006 \\
 r_1 \equiv a_1/a_0 \text{ in } D^+ \rightarrow \eta e^+ \nu_e &= -1.8 \pm 2.2 \\
 r_v \equiv V(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 1.51 \pm 0.07 \quad (S = 2.2) \\
 r_2 \equiv A_2(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 0.807 \pm 0.025 \\
 r_3 \equiv A_3(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 0.0 \pm 0.4 \\
 \Gamma_L/\Gamma_T \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 1.13 \pm 0.08 \\
 \Gamma_+/\Gamma_- \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 0.22 \pm 0.06 \quad (S = 1.6)
 \end{aligned}$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as \bar{K}^0 modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Inclusive modes			
e^+ semileptonic	(16.07±0.30) %		—
μ^+ anything	(17.6 ±3.2) %		—
K^- anything	(25.7 ±1.4) %		—
\bar{K}^0 anything + K^0 anything	(61 ±5) %		—
K^+ anything	(5.9 ±0.8) %		—
$K^*(892)^-$ anything	(6 ±5) %		—
$\bar{K}^*(892)^0$ anything	(23 ±5) %		—
$K^*(892)^0$ anything	< 6.6 %	CL=90%	—
η anything	(6.3 ±0.7) %		—
η' anything	(1.04±0.18) %		—
ϕ anything	(1.03±0.12) %		—
Leptonic and semileptonic modes			
$e^+ \nu_e$	< 8.8 × 10 ⁻⁶	CL=90%	935
$\mu^+ \nu_\mu$	(3.82±0.33) × 10 ⁻⁴		932
$\tau^+ \nu_\tau$	< 1.2 × 10 ⁻³	CL=90%	91
$\bar{K}^0 e^+ \nu_e$	(8.83±0.22) %		869
$\bar{K}^0 \mu^+ \nu_\mu$	(9.2 ±0.6) %		865
$K^- \pi^+ e^+ \nu_e$	(4.00±0.10) %		864

$\bar{K}^*(892)^0 e^+ \nu_e, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(3.68±0.10) %		722
$(K^- \pi^+)_{S\text{-wave}} e^+ \nu_e$	(2.32±0.10) × 10 ⁻³		—
$\bar{K}^*(1410)^0 e^+ \nu_e,$ $\bar{K}^*(1410)^0 \rightarrow K^- \pi^+$	< 6 × 10 ⁻³	CL=90%	—
$\bar{K}_2^*(1430)^0 e^+ \nu_e,$ $\bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	< 5 × 10 ⁻⁴	CL=90%	—
$K^- \pi^+ e^+ \nu_e$ nonresonant	< 7 × 10 ⁻³	CL=90%	864
$K^- \pi^+ \mu^+ \nu_\mu$	(3.8 ±0.4) %		851
$\bar{K}^*(892)^0 \mu^+ \nu_\mu,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(3.52±0.10) %		717
$K^- \pi^+ \mu^+ \nu_\mu$ nonresonant	(2.0 ±0.5) × 10 ⁻³		851
$K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	< 1.6 × 10 ⁻³	CL=90%	825
$\pi^0 e^+ \nu_e$	(4.05±0.18) × 10 ⁻³		930
$\eta e^+ \nu_e$	(1.14±0.10) × 10 ⁻³		855
$\rho^0 e^+ \nu_e$	(2.2 ±0.4) × 10 ⁻³		774
$\rho^0 \mu^+ \nu_\mu$	(2.4 ±0.4) × 10 ⁻³		770
$\omega e^+ \nu_e$	(1.6 ^{+0.7} / _{-0.6}) × 10 ⁻³		771
$\eta'(958) e^+ \nu_e$	(2.2 ±0.5) × 10 ⁻⁴		689
$\phi e^+ \nu_e$	< 9 × 10 ⁻⁵	CL=90%	657

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$\bar{K}^*(892)^0 e^+ \nu_e$	(5.52±0.15) %		722
$\bar{K}^*(892)^0 \mu^+ \nu_\mu$	(5.28±0.15) %		717
$\bar{K}_0^*(1430)^0 \mu^+ \nu_\mu$	< 2.4 × 10 ⁻⁴		380
$\bar{K}^*(1680)^0 \mu^+ \nu_\mu$	< 1.5 × 10 ⁻³		105

Hadronic modes with a \bar{K} or $\bar{K}K\bar{K}$

$K_S^0 \pi^+$	(1.47±0.07) %	S=2.0	863
$K_L^0 \pi^+$	(1.46±0.05) %		863
$K^- 2\pi^+$	[c] (9.13±0.19) %		846
$(K^- \pi^+)_{S\text{-wave}} \pi^+$	(7.32±0.19) %		846
$\bar{K}_0^*(1430)^0 \pi^+,$ $\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$	[d] (1.21±0.06) %		382
$\bar{K}^*(892)^0 \pi^+,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(1.01±0.11) %		714
$\bar{K}^*(1410)^0 \pi^+, \bar{K}^{*0} \rightarrow K^- \pi^+$	not seen		381
$\bar{K}_2^*(1430)^0 \pi^+,$ $\bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	[d] (2.2 ±0.7) × 10 ⁻⁴		371
$\bar{K}^*(1680)^0 \pi^+,$ $\bar{K}^*(1680)^0 \rightarrow K^- \pi^+$	[d] (2.1 ±1.1) × 10 ⁻⁴		58

$K^-(2\pi^+)_{I=2}$	(1.41±0.26) %		—
$K_S^0 \pi^+ \pi^0$	[c] (6.99±0.27) %		845
$K_S^0 \rho^+$	(4.8 ±1.0) %		677
$\overline{K}^*(892)^0 \pi^+$,	(1.3 ±0.6) %		714
$\overline{K}^*(892)^0 \rightarrow K_S^0 \pi^0$			
$K_S^0 \pi^+ \pi^0$ nonresonant	(9 ±7) × 10 ⁻³		845
$K^- 2\pi^+ \pi^0$	[e] (5.99±0.18) %		816
$K_S^0 2\pi^+ \pi^-$	[e] (3.12±0.11) %		814
$K^- 3\pi^+ \pi^-$	[c] (5.6 ±0.5) × 10 ⁻³	S=1.1	772
$\overline{K}^*(892)^0 2\pi^+ \pi^-$,	(1.2 ±0.4) × 10 ⁻³		645
$\overline{K}^*(892)^0 \rightarrow K^- \pi^+$			
$\overline{K}^*(892)^0 \rho^0 \pi^+$,	(2.2 ±0.4) × 10 ⁻³		239
$\overline{K}^*(892)^0 \rightarrow K^- \pi^+$			
$\overline{K}^*(892)^0 a_1(1260)^+$	[f] (9.0 ±1.8) × 10 ⁻³		†
$K^- \rho^0 2\pi^+$	(1.68±0.27) × 10 ⁻³		524
$K^- 3\pi^+ \pi^-$ nonresonant	(3.9 ±2.9) × 10 ⁻⁴		772
$K^+ 2K_S^0$	(4.5 ±2.0) × 10 ⁻³		545
$K^+ K^- K_S^0 \pi^+$	(2.4 ±0.6) × 10 ⁻⁴		436

Pionic modes

$\pi^+ \pi^0$	(1.19±0.06) × 10 ⁻³		925
$2\pi^+ \pi^-$	(3.18±0.18) × 10 ⁻³		909
$\rho^0 \pi^+$	(8.1 ±1.5) × 10 ⁻⁴		767
$\pi^+(\pi^+ \pi^-)_{S\text{-wave}}$	(1.78±0.16) × 10 ⁻³		909
$\sigma \pi^+$, $\sigma \rightarrow \pi^+ \pi^-$	(1.34±0.12) × 10 ⁻³		—
$f_0(980) \pi^+$,	(1.52±0.33) × 10 ⁻⁴		669
$f_0(980) \rightarrow \pi^+ \pi^-$			
$f_0(1370) \pi^+$,	(8 ±4) × 10 ⁻⁵		—
$f_0(1370) \rightarrow \pi^+ \pi^-$			
$f_2(1270) \pi^+$,	(4.9 ±0.9) × 10 ⁻⁴		485
$f_2(1270) \rightarrow \pi^+ \pi^-$			
$\rho(1450)^0 \pi^+$,	< 8 × 10 ⁻⁵	CL=95%	338
$\rho(1450)^0 \rightarrow \pi^+ \pi^-$			
$f_0(1500) \pi^+$,	(1.1 ±0.4) × 10 ⁻⁴		—
$f_0(1500) \rightarrow \pi^+ \pi^-$			
$f_0(1710) \pi^+$,	< 5 × 10 ⁻⁵	CL=95%	—
$f_0(1710) \rightarrow \pi^+ \pi^-$			
$f_0(1790) \pi^+$,	< 6 × 10 ⁻⁵	CL=95%	—
$f_0(1790) \rightarrow \pi^+ \pi^-$			
$(\pi^+ \pi^+)_{S\text{-wave}} \pi^-$	< 1.2 × 10 ⁻⁴	CL=95%	909
$2\pi^+ \pi^-$ nonresonant	< 1.1 × 10 ⁻⁴	CL=95%	909
$\pi^+ 2\pi^0$	(4.6 ±0.4) × 10 ⁻³		910
$2\pi^+ \pi^- \pi^0$	(1.13±0.08) %		883

$\eta\pi^+, \eta \rightarrow \pi^+\pi^-\pi^0$	$(8.0 \pm 0.5) \times 10^{-4}$		848
$\omega\pi^+, \omega \rightarrow \pi^+\pi^-\pi^0$	$< 3 \times 10^{-4}$	CL=90%	763
$3\pi^+2\pi^-$	$(1.61 \pm 0.16) \times 10^{-3}$		845

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$\eta\pi^+$	$(3.53 \pm 0.21) \times 10^{-3}$		848
$\eta\pi^+\pi^0$	$(1.38 \pm 0.35) \times 10^{-3}$		830
$\omega\pi^+$	$< 3.4 \times 10^{-4}$	CL=90%	764
$\eta'(958)\pi^+$	$(4.67 \pm 0.29) \times 10^{-3}$		681
$\eta'(958)\pi^+\pi^0$	$(1.6 \pm 0.5) \times 10^{-3}$		654

Hadronic modes with a $K\bar{K}$ pair

$K^+K_S^0$	$(2.83 \pm 0.16) \times 10^{-3}$	S=2.2	793
$K^+K^-\pi^+$	[c] $(9.54 \pm 0.26) \times 10^{-3}$	S=1.1	744
$\phi\pi^+, \phi \rightarrow K^+K^-$	$(2.65_{-0.09}^{+0.08}) \times 10^{-3}$		647
$K^+\bar{K}^*(892)^0, \bar{K}^*(892)^0 \rightarrow K^-\pi^+$	$(2.45_{-0.14}^{+0.09}) \times 10^{-3}$		613
$K^+\bar{K}_0^*(1430)^0, \bar{K}_0^*(1430)^0 \rightarrow K^-\pi^+$	$(1.79 \pm 0.34) \times 10^{-3}$		—
$K^+\bar{K}_2^*(1430)^0, \bar{K}_2^* \rightarrow K^-\pi^+$	$(1.6_{-0.8}^{+1.2}) \times 10^{-4}$		—
$K^+\bar{K}_0^*(800), \bar{K}_0^* \rightarrow K^-\pi^+$	$(6.7_{-2.1}^{+3.4}) \times 10^{-4}$		—
$a_0(1450)^0\pi^+, a_0^0 \rightarrow K^+K^-$	$(4.4_{-1.8}^{+7.0}) \times 10^{-4}$		—
$\phi(1680)\pi^+, \phi \rightarrow K^+K^-$	$(4.9_{-1.9}^{+4.0}) \times 10^{-5}$		—
$K^+K^-\pi^+$ nonresonant	not seen		744
$K^+K_S^0\pi^+\pi^-$	$(1.75 \pm 0.18) \times 10^{-3}$		678
$K_S^0K^-2\pi^+$	$(2.40 \pm 0.18) \times 10^{-3}$		678
$K^+K^-2\pi^+\pi^-$	$(2.2 \pm 1.2) \times 10^{-4}$		600

A few poorly measured branching fractions:

$\phi\pi^+\pi^0$	$(2.3 \pm 1.0) \%$		619
$\phi\rho^+$	$< 1.5 \%$	CL=90%	259
$K^+K^-\pi^+\pi^0$ non- ϕ	$(1.5_{-0.6}^{+0.7}) \%$		682
$K^*(892)^+K_S^0$	$(1.6 \pm 0.7) \%$		612

Doubly Cabibbo-suppressed modes

$K^+ \pi^0$	$(1.83 \pm 0.26) \times 10^{-4}$	S=1.4	864
$K^+ \eta$	$(1.08 \pm 0.17) \times 10^{-4}$		776
$K^+ \eta'(958)$	$(1.76 \pm 0.22) \times 10^{-4}$		571
$K^+ \pi^+ \pi^-$	$(5.27 \pm 0.23) \times 10^{-4}$		846
$K^+ \rho^0$	$(2.0 \pm 0.5) \times 10^{-4}$		679
$K^*(892)^0 \pi^+, K^*(892)^0 \rightarrow$	$(2.5 \pm 0.4) \times 10^{-4}$		714
$K^+ \pi^-$			
$K^+ f_0(980), f_0(980) \rightarrow$	$(4.7 \pm 2.8) \times 10^{-5}$		—
$\pi^+ \pi^-$			
$K_2^*(1430)^0 \pi^+, K_2^*(1430)^0 \rightarrow$	$(4.2 \pm 2.9) \times 10^{-5}$		—
$K^+ \pi^-$			
$K^+ \pi^+ \pi^-$ nonresonant	not seen		846
$2K^+ K^-$	$(8.7 \pm 2.0) \times 10^{-5}$		550

**$\Delta C = 1$ weak neutral current (C1) modes, or
Lepton Family number (LF) or Lepton number (L) violating modes**

$\pi^+ e^+ e^-$	C1	< 1.1	$\times 10^{-6}$	CL=90%	930
$\pi^+ \phi, \phi \rightarrow e^+ e^-$	[g]	$(1.7 \pm_{-0.9}^{+1.4})$	$\times 10^{-6}$		—
$\pi^+ \mu^+ \mu^-$	C1	< 3.9	$\times 10^{-6}$	CL=90%	918
$\pi^+ \phi, \phi \rightarrow \mu^+ \mu^-$	[g]	(1.8 ± 0.8)	$\times 10^{-6}$		—
$\rho^+ \mu^+ \mu^-$	C1	< 5.6	$\times 10^{-4}$	CL=90%	757
$K^+ e^+ e^-$	[h]	< 1.0	$\times 10^{-6}$	CL=90%	870
$K^+ \mu^+ \mu^-$	[h]	< 4.3	$\times 10^{-6}$	CL=90%	856
$\pi^+ e^+ \mu^-$	LF	< 2.9	$\times 10^{-6}$	CL=90%	927
$\pi^+ e^- \mu^+$	LF	< 3.6	$\times 10^{-6}$	CL=90%	927
$K^+ e^+ \mu^-$	LF	< 1.2	$\times 10^{-6}$	CL=90%	866
$K^+ e^- \mu^+$	LF	< 2.8	$\times 10^{-6}$	CL=90%	866
$\pi^- 2e^+$	L	< 1.1	$\times 10^{-6}$	CL=90%	930
$\pi^- 2\mu^+$	L	< 2.0	$\times 10^{-6}$	CL=90%	918
$\pi^- e^+ \mu^+$	L	< 2.0	$\times 10^{-6}$	CL=90%	927
$\rho^- 2\mu^+$	L	< 5.6	$\times 10^{-4}$	CL=90%	757
$K^- 2e^+$	L	< 9	$\times 10^{-7}$	CL=90%	870
$K^- 2\mu^+$	L	< 1.0	$\times 10^{-5}$	CL=90%	856
$K^- e^+ \mu^+$	L	< 1.9	$\times 10^{-6}$	CL=90%	866
$K^*(892)^- 2\mu^+$	L	< 8.5	$\times 10^{-4}$	CL=90%	703

D⁰

$$I(J^P) = \frac{1}{2}(0^-)$$

Mass $m = 1864.86 \pm 0.13$ MeV

$m_{D^\pm} - m_{D^0} = 4.76 \pm 0.10$ MeV (S = 1.1)

Mean life $\tau = (410.1 \pm 1.5) \times 10^{-15}$ s

$c\tau = 122.9$ μm

$|m_{D_1^0} - m_{D_2^0}| = (1.44_{-0.50}^{+0.48}) \times 10^{10} \hbar \text{ s}^{-1}$

$(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma = 2y = (1.60_{-0.26}^{+0.25}) \times 10^{-2}$

$|q/p| = 0.88_{-0.15}^{+0.16}$

$A_\Gamma = (0.26 \pm 2.31) \times 10^{-3}$

$K^+ \pi^-$ relative strong phase: $\cos \delta = 1.03_{-0.18}^{+0.32}$

$K^- \pi^+ \pi^0$ coherence factor $R_{K \pi \pi^0} = 0.78_{-0.25}^{+0.11}$

$K^- \pi^+ \pi^0$ average relative strong phase $\delta^{K \pi \pi^0} = (239_{-28}^{+32})^\circ$

$K^- \pi^- 2\pi^+$ coherence factor $R_{K 3\pi} = 0.36_{-0.30}^{+0.24}$

$K^- \pi^- 2\pi^+$ average relative strong phase $\delta^{K 3\pi} = (118_{-50}^{+60})^\circ$

CP-violation decay-rate asymmetries (labeled by the D⁰ decay)

$A_{CP}(K^+ K^-) = (-0.21 \pm 0.17)\%$

$A_{CP}(2K_S^0) = (-23 \pm 19)\%$

$A_{CP}(\pi^+ \pi^-) = (0.22 \pm 0.21)\%$

$A_{CP}(2\pi^0) = (0 \pm 5)\%$

$A_{CP}(\pi^+ \pi^- \pi^0) = (0.3 \pm 0.4)\%$

$A_{CP}(\rho(770)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) = (1.2 \pm 0.9)\% [i]$

$A_{CP}(\rho(770)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (-3.1 \pm 3.0)\% [i]$

$A_{CP}(\rho(770)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) = (-1.0 \pm 1.7)\% [i]$

$A_{CP}(\rho(1450)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) = (0 \pm 70)\% [i]$

$A_{CP}(\rho(1450)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (-20 \pm 40)\% [i]$

$A_{CP}(\rho(1450)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) = (6 \pm 9)\% [i]$

$A_{CP}(\rho(1700)^+ \pi^- \rightarrow \pi^+ \pi^- \pi^0) = (-5 \pm 14)\% [i]$

$A_{CP}(\rho(1700)^0 \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (13 \pm 9)\% [i]$

$A_{CP}(\rho(1700)^- \pi^+ \rightarrow \pi^+ \pi^- \pi^0) = (8 \pm 11)\% [i]$

$A_{CP}(f_0(980) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (0 \pm 35)\% [i]$

$A_{CP}(f_0(1370) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (25 \pm 18)\% [i]$

$A_{CP}(f_0(1500) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (0 \pm 18)\% [i]$

$A_{CP}(f_0(1710) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (0 \pm 24)\% [i]$

$A_{CP}(f_2(1270) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (-4 \pm 6)\% [i]$

$A_{CP}(\sigma(400) \pi^0 \rightarrow \pi^+ \pi^- \pi^0) = (6 \pm 8)\% [i]$

$A_{CP}(\text{nonresonant } \pi^+ \pi^- \pi^0) = (-13 \pm 23)\% [i]$

$A_{CP}(K^+ K^- \pi^0) = (-1.0 \pm 1.7)\%$

$A_{CP}(K^*(892)^+ K^- \rightarrow K^+ K^- \pi^0) = (-0.9 \pm 1.3)\% [i]$

$$\begin{aligned}
 A_{CP}(K^*(1410)^+ K^- \rightarrow K^+ K^- \pi^0) &= (-21 \pm 24)\% [i] \\
 A_{CP}((K^+ \pi^0)_S K^- \rightarrow K^+ K^- \pi^0) &= (7 \pm 15)\% [i] \\
 A_{CP}(\phi(1020) \pi^0 \rightarrow K^+ K^- \pi^0) &= (1.1 \pm 2.2)\% [i] \\
 A_{CP}(f_0(980) \pi^0 \rightarrow K^+ K^- \pi^0) &= (-3 \pm 19)\% [i] \\
 A_{CP}(a_0(980)^0 \pi^0 \rightarrow K^+ K^- \pi^0) &= (-5 \pm 16)\% [i] \\
 A_{CP}(f'_2(1525) \pi^0 \rightarrow K^+ K^- \pi^0) &= (0 \pm 160)\% [i] \\
 A_{CP}(K^*(892)^- K^+ \rightarrow K^+ K^- \pi^0) &= (-5 \pm 4)\% [i] \\
 A_{CP}(K^*(1410)^- K^+ \rightarrow K^+ K^- \pi^0) &= (-17 \pm 29)\% [i] \\
 A_{CP}((K^- \pi^0)_{S-wave} K^+ \rightarrow K^+ K^- \pi^0) &= (-10 \pm 40)\% [i] \\
 A_{CP}(K_S^0 \pi^0) &= (-0.27 \pm 0.21)\% \\
 A_{CP}(K_S^0 \eta) &= (0.5 \pm 0.5)\% \\
 A_{CP}(K_S^0 \eta') &= (1.0 \pm 0.7)\% \\
 A_{CP}(K_S^0 \phi) &= (-3 \pm 9)\% \\
 A_{CP}(K^- \pi^+) &= (0.1 \pm 0.7)\% \\
 A_{CP}(K^+ \pi^-) &= (2.2 \pm 3.2)\% \\
 A_{CP}(K^- \pi^+ \pi^0) &= (0.2 \pm 0.9)\% \\
 A_{CP}(K^+ \pi^- \pi^0) &= (0 \pm 5)\% \\
 A_{CP}(K_S^0 \pi^+ \pi^-) &= (-0.9^{+2.6}_{-6.0})\% \\
 A_{CP}(K^*(892)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &< 3.5 \times 10^{-4}, \text{ CL} = 95\% \\
 A_{CP}(K^*(892)^+ \pi^- \rightarrow K_S^0 \pi^+ \pi^-) &< 7.8 \times 10^{-4}, \text{ CL} = 95\% \\
 A_{CP}(\bar{K}^0 \rho^0 \rightarrow K_S^0 \pi^+ \pi^-) &< 4.8 \times 10^{-4}, \text{ CL} = 95\% \\
 A_{CP}(\bar{K}^0 \omega \rightarrow K_S^0 \pi^+ \pi^-) &< 9.2 \times 10^{-4}, \text{ CL} = 95\% \\
 A_{CP}(\bar{K}^0 f_0(980) \rightarrow K_S^0 \pi^+ \pi^-) &< 6.8 \times 10^{-4}, \text{ CL} = 95\% \\
 A_{CP}(\bar{K}^0 f_2(1270) \rightarrow K_S^0 \pi^+ \pi^-) &< 13.5 \times 10^{-4}, \text{ CL} = 95\% \\
 A_{CP}(\bar{K}^0 f_0(1370) \rightarrow K_S^0 \pi^+ \pi^-) &< 25.5 \times 10^{-4}, \text{ CL} = 95\% \\
 A_{CP}(K_0^*(1430)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &< 9.0 \times 10^{-4}, \text{ CL} = 95\% \\
 A_{CP}(K_2^*(1430)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &< 6.5 \times 10^{-4}, \text{ CL} = 95\% \\
 A_{CP}(K^*(1680)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &< 28.4 \times 10^{-4}, \text{ CL} = 95\% \\
 A_{CP}(K^- \pi^+ \pi^+ \pi^-) &= (0.7 \pm 1.0)\% \\
 A_{CP}(K^+ \pi^- \pi^+ \pi^-) &= (-2 \pm 4)\% \\
 A_{CP}(K^+ K^- \pi^+ \pi^-) &= (-8 \pm 7)\% \\
 \Delta A_{CP} = A_{CP}(K^+ K^-) - A_{CP}(\pi^+ \pi^-) &= (-0.65 \pm 0.18)\%
 \end{aligned}$$

T-violation decay-rate asymmetry

$$A_T(K^+ K^- \pi^+ \pi^-) = (1 \pm 7) \times 10^{-3} [b]$$

CPT-violation decay-rate asymmetry

$$A_{CPT}(K^\mp \pi^\pm) = 0.008 \pm 0.008$$

Form factors

$$\begin{aligned}
 r_V &\equiv V(0)/A_1(0) \text{ in } D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell = 1.7 \pm 0.8 \\
 r_2 &\equiv A_2(0)/A_1(0) \text{ in } D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell = 0.9 \pm 0.4 \\
 f_+(0) &\text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = 0.727 \pm 0.011 \\
 f_+(0)|V_{cs}| &\text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = 0.726 \pm 0.009 \\
 r_1 &\equiv a_1/a_0 \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = -2.65 \pm 0.35 \\
 r_2 &\equiv a_1/a_0 \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell = 13 \pm 9 \\
 f_+(0)|V_{cd}| &\text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell = 0.152 \pm 0.005 \\
 r_1 &\equiv a_1/a_0 \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell = -2.8 \pm 0.5 \\
 r_2 &\equiv a_1/a_0 \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell = 6 \pm 3.0
 \end{aligned}$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as \bar{K}^0 modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D^0 DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Topological modes			
0-prongs	[j] (15 ± 6) %		—
2-prongs	(70 ± 6) %		—
4-prongs	[k] (14.5 ± 0.5) %		—
6-prongs	[l] (6.4 ± 1.3) × 10 ⁻⁴		—
Inclusive modes			
e ⁺ anything	[m] (6.49 ± 0.11) %		—
μ ⁺ anything	(6.7 ± 0.6) %		—
K ⁻ anything	(54.7 ± 2.8) %	S=1.3	—
\bar{K}^0 anything + K ⁰ anything	(47 ± 4) %		—
K ⁺ anything	(3.4 ± 0.4) %		—
K*(892) ⁻ anything	(15 ± 9) %		—
$\bar{K}^*(892)^0$ anything	(9 ± 4) %		—
K*(892) ⁺ anything	< 3.6 %	CL=90%	—
K*(892) ⁰ anything	(2.8 ± 1.3) %		—
η anything	(9.5 ± 0.9) %		—
η' anything	(2.48 ± 0.27) %		—
φ anything	(1.05 ± 0.11) %		—

Semileptonic modes

$K^- e^+ \nu_e$	(3.55 ± 0.04) %	S=1.2	867
$K^- \mu^+ \nu_\mu$	(3.30 ± 0.13) %		864
$K^*(892)^- e^+ \nu_e$	(2.16 ± 0.16) %		719
$K^*(892)^- \mu^+ \nu_\mu$	(1.90 ± 0.24) %		714
$K^- \pi^0 e^+ \nu_e$	(1.6 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 1.3 \\ 0.5 \end{smallmatrix}$) %		861
$\bar{K}^0 \pi^- e^+ \nu_e$	(2.7 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 0.9 \\ 0.7 \end{smallmatrix}$) %		860
$K^- \pi^+ \pi^- e^+ \nu_e$	(2.8 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 1.4 \\ 1.1 \end{smallmatrix}$) × 10 ⁻⁴		843
$K_1(1270)^- e^+ \nu_e$	(7.6 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 4.0 \\ 3.1 \end{smallmatrix}$) × 10 ⁻⁴		498
$K^- \pi^+ \pi^- \mu^+ \nu_\mu$	< 1.2 × 10 ⁻³	CL=90%	821
$(\bar{K}^*(892)\pi)^- \mu^+ \nu_\mu$	< 1.4 × 10 ⁻³	CL=90%	692
$\pi^- e^+ \nu_e$	(2.89 ± 0.08) × 10 ⁻³	S=1.1	927
$\pi^- \mu^+ \nu_\mu$	(2.37 ± 0.24) × 10 ⁻³		924
$\rho^- e^+ \nu_e$	(1.9 ± 0.4) × 10 ⁻³		771

Hadronic modes with one \bar{K}

$K^- \pi^+$	(3.88 ± 0.05) %	S=1.2	861
$K_S^0 \pi^0$	(1.19 ± 0.04) %		860
$K_L^0 \pi^0$	(10.0 ± 0.7) × 10 ⁻³		860
$K_S^0 \pi^+ \pi^-$	[c] (2.82 ± 0.19) %	S=1.1	842
$K_S^0 \rho^0$	(6.3 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 0.7 \\ 0.8 \end{smallmatrix}$) × 10 ⁻³		674
$K_S^0 \omega, \omega \rightarrow \pi^+ \pi^-$	(2.0 ± 0.6) × 10 ⁻⁴		670
$K_S^0 (\pi^+ \pi^-)_{S\text{-wave}}$	(3.4 ± 0.8) × 10 ⁻³		842
$K_S^0 f_0(980),$ $f_0(980) \rightarrow \pi^+ \pi^-$	(1.21 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 0.40 \\ 0.24 \end{smallmatrix}$) × 10 ⁻³		549
$K_S^0 f_0(1370),$ $f_0(1370) \rightarrow \pi^+ \pi^-$	(2.8 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 0.9 \\ 1.3 \end{smallmatrix}$) × 10 ⁻³		†
$K_S^0 f_2(1270),$ $f_2(1270) \rightarrow \pi^+ \pi^-$	(9 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 10 \\ 6 \end{smallmatrix}$) × 10 ⁻⁵		262
$K^*(892)^- \pi^+,$ $K^*(892)^- \rightarrow K_S^0 \pi^-$	(1.66 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 0.15 \\ 0.17 \end{smallmatrix}$) %		711
$K_0^*(1430)^- \pi^+,$ $K_0^*(1430)^- \rightarrow K_S^0 \pi^-$	(2.69 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 0.40 \\ 0.33 \end{smallmatrix}$) × 10 ⁻³		378
$K_2^*(1430)^- \pi^+,$ $K_2^*(1430)^- \rightarrow K_S^0 \pi^-$	(3.4 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 1.9 \\ 1.0 \end{smallmatrix}$) × 10 ⁻⁴		367
$K^*(1680)^- \pi^+,$ $K^*(1680)^- \rightarrow K_S^0 \pi^-$	(4 ± 4) × 10 ⁻⁴		46

$K^*(892)^+ \pi^-$, $K^*(892)^+ \rightarrow K_S^0 \pi^+$	[η] (1.13 \pm 0.60 \pm 0.34) $\times 10^{-4}$		711
$K_0^*(1430)^+ \pi^-$, $K_0^*(1430)^+ \rightarrow K_S^0 \pi^+$	[η] < 1.4 $\times 10^{-5}$	CL=95%	–
$K_2^*(1430)^+ \pi^-$, $K_2^*(1430)^+ \rightarrow K_S^0 \pi^+$	[η] < 3.4 $\times 10^{-5}$	CL=95%	–
$K_S^0 \pi^+ \pi^-$ nonresonant	(2.5 \pm 6.0 \pm 1.6) $\times 10^{-4}$		842
$K^- \pi^+ \pi^0$	[c] (13.9 \pm 0.5) %	S=1.7	844
$K^- \rho^+$	(10.8 \pm 0.7) %		675
$K^- \rho(1700)^+$, $\rho(1700)^+ \rightarrow \pi^+ \pi^0$	(7.9 \pm 1.7) $\times 10^{-3}$		†
$K^*(892)^- \pi^+$, $K^*(892)^- \rightarrow K^- \pi^0$	(2.22 \pm 0.40 \pm 0.19) %		711
$\bar{K}^*(892)^0 \pi^0$, $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(1.88 \pm 0.23) %		711
$K_0^*(1430)^- \pi^+$, $K_0^*(1430)^- \rightarrow K^- \pi^0$	(4.6 \pm 2.1) $\times 10^{-3}$		378
$\bar{K}_0^*(1430)^0 \pi^0$, $\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$	(5.7 \pm 5.0 \pm 1.5) $\times 10^{-3}$		379
$K^*(1680)^- \pi^+$, $K^*(1680)^- \rightarrow K^- \pi^0$	(1.8 \pm 0.7) $\times 10^{-3}$		46
$K^- \pi^+ \pi^0$ nonresonant	(1.11 \pm 0.50 \pm 0.19) %		844
$K_S^0 2\pi^0$	(9.1 \pm 1.1) $\times 10^{-3}$	S=2.2	843
$K_S^0(2\pi^0)$ -S-wave	(2.6 \pm 0.7) $\times 10^{-3}$		–
$\bar{K}^*(892)^0 \pi^0$, $\bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$	(7.8 \pm 0.7) $\times 10^{-3}$		711
$\bar{K}^*(1430)^0 \pi^0, \bar{K}^{*0} \rightarrow$ $K_S^0 \pi^0$	(4 \pm 23) $\times 10^{-5}$		–
$\bar{K}^*(1680)^0 \pi^0, \bar{K}^{*0} \rightarrow$ $K_S^0 \pi^0$	(1.0 \pm 0.4) $\times 10^{-3}$		–
$K_S^0 f_2(1270), f_2 \rightarrow 2\pi^0$	(2.3 \pm 1.1) $\times 10^{-4}$		–
$2K_S^0, \text{one } K_S^0 \rightarrow 2\pi^0$	(3.2 \pm 1.1) $\times 10^{-4}$		–
$K^- 2\pi^+ \pi^-$	[c] (8.07 \pm 0.21 \pm 0.19) %	S=1.3	813
$K^- \pi^+ \rho^0$ total	(6.74 \pm 0.33) %		609
$K^- \pi^+ \rho^0$ 3-body	(5.1 \pm 2.3) $\times 10^{-3}$		609
$\bar{K}^*(892)^0 \rho^0$, $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(1.05 \pm 0.23) %		416
$K^- a_1(1260)^+$, $a_1(1260)^+ \rightarrow 2\pi^+ \pi^-$	(3.6 \pm 0.6) %		327

$\bar{K}^*(892)^0 \pi^+ \pi^-$ total,	(1.6 ± 0.4) %	685
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$		
$\bar{K}^*(892)^0 \pi^+ \pi^-$ 3-body,	(9.9 ± 2.3) × 10 ⁻³	685
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$		
$K_1(1270)^- \pi^+$,	[o] (2.9 ± 0.3) × 10 ⁻³	484
$K_1(1270)^- \rightarrow K^- \pi^+ \pi^-$		
$K^- 2\pi^+ \pi^-$ nonresonant	(1.88 ± 0.26) %	813
$K_S^0 \pi^+ \pi^- \pi^0$	[p] (5.2 ± 0.6) %	813
$K_S^0 \eta, \eta \rightarrow \pi^+ \pi^- \pi^0$	(1.02 ± 0.09) × 10 ⁻³	772
$K_S^0 \omega, \omega \rightarrow \pi^+ \pi^- \pi^0$	(9.9 ± 0.5) × 10 ⁻³	670
$K^- 2\pi^+ \pi^- \pi^0$	(4.2 ± 0.4) %	771
$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0,$	(1.3 ± 0.6) %	643
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$		
$K^- \pi^+ \omega, \omega \rightarrow \pi^+ \pi^- \pi^0$	(2.7 ± 0.5) %	605
$\bar{K}^*(892)^0 \omega,$	(6.5 ± 3.0) × 10 ⁻³	410
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+,$		
$\omega \rightarrow \pi^+ \pi^- \pi^0$		
$K_S^0 \eta \pi^0$	(5.5 ± 1.1) × 10 ⁻³	721
$K_S^0 a_0(980), a_0(980) \rightarrow \eta \pi^0$	(6.5 ± 2.0) × 10 ⁻³	—
$\bar{K}^*(892)^0 \eta,$	(1.6 ± 0.5) × 10 ⁻³	—
$\bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$		
$K_S^0 2\pi^+ 2\pi^-$	(2.68 ± 0.30) × 10 ⁻³	768
$K_S^0 \rho^0 \pi^+ \pi^-$, no $K^*(892)^-$	(1.1 ± 0.7) × 10 ⁻³	—
$K^*(892)^- 2\pi^+ \pi^-$,	(5 ± 8) × 10 ⁻⁴	642
$K^*(892)^- \rightarrow K_S^0 \pi^-$,		
no ρ^0		
$K^*(892)^- \rho^0 \pi^+$,	(1.6 ± 0.6) × 10 ⁻³	230
$K^*(892)^- \rightarrow K_S^0 \pi^-$		
$K_S^0 2\pi^+ 2\pi^-$ nonresonant	< 1.2 × 10 ⁻³ CL=90%	768
$K^- 3\pi^+ 2\pi^-$	(2.2 ± 0.6) × 10 ⁻⁴	713

Fractions of many of the following modes with resonances have already appeared above as submodes of particular charged-particle modes. (Modes for which there are only upper limits and $\bar{K}^*(892)\rho$ submodes only appear below.)

$K_S^0 \eta$	(4.78 ± 0.30) × 10 ⁻³	772
$K_S^0 \omega$	(1.11 ± 0.06) %	670
$K_S^0 \eta'(958)$	(9.4 ± 0.5) × 10 ⁻³	565
$K^- a_1(1260)^+$	(7.8 ± 1.1) %	327
$K^- a_2(1320)^+$	< 2 × 10 ⁻³ CL=90%	198
$\bar{K}^*(892)^0 \pi^+ \pi^-$ total	(2.4 ± 0.5) %	685
$\bar{K}^*(892)^0 \pi^+ \pi^-$ 3-body	(1.48 ± 0.34) %	685
$\bar{K}^*(892)^0 \rho^0$	(1.57 ± 0.34) %	417

$\bar{K}^*(892)^0 \rho^0$ transverse	(1.7 ± 0.6) %		417
$\bar{K}^*(892)^0 \rho^0$ S-wave	(3.0 ± 0.6) %		417
$\bar{K}^*(892)^0 \rho^0$ S-wave long.	< 3	$\times 10^{-3}$	CL=90% 417
$\bar{K}^*(892)^0 \rho^0$ P-wave	< 3	$\times 10^{-3}$	CL=90% 417
$\bar{K}^*(892)^0 \rho^0$ D-wave	(2.1 ± 0.6) %		417
$K_1(1270)^- \pi^+$	[o] (1.6 ± 0.8) %		484
$K_1(1400)^- \pi^+$	< 1.2	%	CL=90% 386
$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0$	(1.9 ± 0.9) %		644
$K^- \pi^+ \omega$	(3.0 ± 0.6) %		605
$\bar{K}^*(892)^0 \omega$	(1.1 ± 0.5) %		410
$K^- \pi^+ \eta'(958)$	(7.5 ± 1.9) $\times 10^{-3}$		479
$\bar{K}^*(892)^0 \eta'(958)$	< 1.1	$\times 10^{-3}$	CL=90% 120

Hadronic modes with three K's

$K_S^0 K^+ K^-$	(4.45 ± 0.34) $\times 10^{-3}$		544
$K_S^0 a_0(980)^0, a_0^0 \rightarrow K^+ K^-$	(3.0 ± 0.4) $\times 10^{-3}$		–
$K^- a_0(980)^+, a_0^+ \rightarrow K^+ K_S^0$	(6.0 ± 1.8) $\times 10^{-4}$		–
$K^+ a_0(980)^-, a_0^- \rightarrow K^- K_S^0$	< 1.1	$\times 10^{-4}$	CL=95% –
$K_S^0 f_0(980), f_0 \rightarrow K^+ K^-$	< 9	$\times 10^{-5}$	CL=95% –
$K_S^0 \phi, \phi \rightarrow K^+ K^-$	(2.04 ± 0.16) $\times 10^{-3}$		520
$K_S^0 f_0(1370), f_0 \rightarrow K^+ K^-$	(1.7 ± 1.1) $\times 10^{-4}$		–
$3K_S^0$	(9.1 ± 1.3) $\times 10^{-4}$		539
$K^+ 2K^- \pi^+$	(2.21 ± 0.31) $\times 10^{-4}$		434
$K^+ K^- \bar{K}^*(892)^0,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(4.4 ± 1.7) $\times 10^{-5}$		†
$K^- \pi^+ \phi, \phi \rightarrow K^+ K^-$	(4.0 ± 1.7) $\times 10^{-5}$		422
$\phi \bar{K}^*(892)^0,$ $\phi \rightarrow K^+ K^-,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(1.06 ± 0.20) $\times 10^{-4}$		†
$K^+ 2K^- \pi^+$ nonresonant	(3.3 ± 1.5) $\times 10^{-5}$		434
$2K_S^0 K^\pm \pi^\mp$	(6.0 ± 1.3) $\times 10^{-4}$		427

Pionic modes

$\pi^+ \pi^-$	(1.401 ± 0.027) $\times 10^{-3}$	S=1.1	922
$2\pi^0$	(8.0 ± 0.5) $\times 10^{-4}$		923
$\pi^+ \pi^- \pi^0$	(1.43 ± 0.06) %	S=1.9	907
$\rho^+ \pi^-$	(9.8 ± 0.4) $\times 10^{-3}$		764
$\rho^0 \pi^0$	(3.72 ± 0.22) $\times 10^{-3}$		764
$\rho^- \pi^+$	(4.96 ± 0.24) $\times 10^{-3}$		764
$\rho(1450)^+ \pi^-, \rho(1450)^+ \rightarrow$ $\pi^+ \pi^0$	(1.6 ± 2.0) $\times 10^{-5}$		–
$\rho(1450)^0 \pi^0, \rho(1450)^0 \rightarrow$ $\pi^+ \pi^-$	(4.3 ± 1.9) $\times 10^{-5}$		–
$\rho(1450)^- \pi^+, \rho(1450)^- \rightarrow$ $\pi^- \pi^0$	(2.6 ± 0.4) $\times 10^{-4}$		–

$\rho(1700)^+ \pi^- , \rho(1700)^+ \rightarrow \pi^+ \pi^0$	$(5.9 \pm 1.4) \times 10^{-4}$	—
$\rho(1700)^0 \pi^0 , \rho(1700)^0 \rightarrow \pi^+ \pi^-$	$(7.2 \pm 1.7) \times 10^{-4}$	—
$\rho(1700)^- \pi^+ , \rho(1700)^- \rightarrow \pi^- \pi^0$	$(4.6 \pm 1.1) \times 10^{-4}$	—
$f_0(980) \pi^0 , f_0(980) \rightarrow \pi^+ \pi^-$	$(3.6 \pm 0.8) \times 10^{-5}$	—
$f_0(500) \pi^0 , f_0(500) \rightarrow \pi^+ \pi^-$	$(1.18 \pm 0.21) \times 10^{-4}$	—
$f_0(1370) \pi^0 , f_0(1370) \rightarrow \pi^+ \pi^-$	$(5.3 \pm 2.0) \times 10^{-5}$	—
$f_0(1500) \pi^0 , f_0(1500) \rightarrow \pi^+ \pi^-$	$(5.6 \pm 1.5) \times 10^{-5}$	—
$f_0(1710) \pi^0 , f_0(1710) \rightarrow \pi^+ \pi^-$	$(4.4 \pm 1.5) \times 10^{-5}$	—
$f_2(1270) \pi^0 , f_2(1270) \rightarrow \pi^+ \pi^-$	$(1.89 \pm 0.20) \times 10^{-4}$	—
$\pi^+ \pi^- \pi^0$ nonresonant	$(1.20 \pm 0.35) \times 10^{-4}$	907
$3\pi^0$	$< 3.5 \times 10^{-4}$	CL=90% 908
$2\pi^+ 2\pi^-$	$(7.42 \pm 0.21) \times 10^{-3}$	S=1.1 880
$a_1(1260)^+ \pi^- , a_1^+ \rightarrow 2\pi^+ \pi^-$ total	$(4.45 \pm 0.31) \times 10^{-3}$	—
$a_1(1260)^+ \pi^- , a_1^+ \rightarrow \rho^0 \pi^+$ S-wave	$(3.21 \pm 0.25) \times 10^{-3}$	—
$a_1(1260)^+ \pi^- , a_1^+ \rightarrow \rho^0 \pi^+$ D-wave	$(1.9 \pm 0.5) \times 10^{-4}$	—
$a_1(1260)^+ \pi^- , a_1^+ \rightarrow \sigma \pi^+$	$(6.2 \pm 0.7) \times 10^{-4}$	—
$2\rho^0$ total	$(1.82 \pm 0.13) \times 10^{-3}$	518
$2\rho^0$, parallel helicities	$(8.2 \pm 3.2) \times 10^{-5}$	—
$2\rho^0$, perpendicular helicities	$(4.7 \pm 0.6) \times 10^{-4}$	—
$2\rho^0$, longitudinal helicities	$(1.25 \pm 0.10) \times 10^{-3}$	—
Resonant $(\pi^+ \pi^-) \pi^+ \pi^-$	$(1.48 \pm 0.12) \times 10^{-3}$	—
3-body total		
$\sigma \pi^+ \pi^-$	$(6.1 \pm 0.9) \times 10^{-4}$	—
$f_0(980) \pi^+ \pi^- , f_0 \rightarrow \pi^+ \pi^-$	$(1.8 \pm 0.5) \times 10^{-4}$	—
$f_2(1270) \pi^+ \pi^- , f_2 \rightarrow \pi^+ \pi^-$	$(3.6 \pm 0.6) \times 10^{-4}$	—
$\pi^+ \pi^- 2\pi^0$	$(10.0 \pm 0.9) \times 10^{-3}$	882
$\eta \pi^0$	[q] $(6.8 \pm 0.7) \times 10^{-4}$	846
$\omega \pi^0$	[q] $< 2.6 \times 10^{-4}$	CL=90% 761
$2\pi^+ 2\pi^- \pi^0$	$(4.1 \pm 0.5) \times 10^{-3}$	844
$\eta \pi^+ \pi^-$	[q] $(1.09 \pm 0.16) \times 10^{-3}$	827

$\omega\pi^+\pi^-$	[q]	$(1.6 \pm 0.5) \times 10^{-3}$	738
$3\pi^+3\pi^-$		$(4.2 \pm 1.2) \times 10^{-4}$	795
$\eta'(958)\pi^0$		$(8.9 \pm 1.4) \times 10^{-4}$	678
$\eta'(958)\pi^+\pi^-$		$(4.5 \pm 1.7) \times 10^{-4}$	650
2η		$(1.67 \pm 0.20) \times 10^{-3}$	755
$\eta\eta'(958)$		$(1.05 \pm 0.26) \times 10^{-3}$	537

Hadronic modes with a $K\bar{K}$ pair

K^+K^-		$(3.96 \pm 0.08) \times 10^{-3}$	S=1.4	791
$2K_S^0$		$(1.7 \pm 0.4) \times 10^{-4}$	S=2.5	789
$K_S^0K^-\pi^+$		$(3.3 \pm 0.5) \times 10^{-3}$	S=1.1	739
$\bar{K}^*(892)^0K_S^0,$ $\bar{K}^*(892)^0 \rightarrow K^-\pi^+$		$< 5 \times 10^{-4}$	CL=90%	608
$K_S^0K^+\pi^-$		$(2.6 \pm 0.5) \times 10^{-3}$		739
$K^*(892)^0K_S^0,$ $K^*(892)^0 \rightarrow K^+\pi^-$		$< 2.8 \times 10^{-4}$	CL=90%	608
$K^+K^-\pi^0$		$(3.28 \pm 0.14) \times 10^{-3}$		743
$K^*(892)^+K^-,$ $K^*(892)^+ \rightarrow K^+\pi^0$		$(1.46 \pm 0.07) \times 10^{-3}$		—
$K^*(892)^-K^+,$ $K^*(892)^- \rightarrow K^-\pi^0$		$(5.2 \pm 0.4) \times 10^{-4}$		—
$(K^+\pi^0)_{S-wave}K^-$		$(2.34 \pm 0.17) \times 10^{-3}$		743
$(K^-\pi^0)_{S-wave}K^+$		$(1.3 \pm 0.4) \times 10^{-4}$		743
$f_0(980)\pi^0, f_0 \rightarrow K^+K^-$		$(3.4 \pm 0.6) \times 10^{-4}$		—
$\phi\pi^0, \phi \rightarrow K^+K^-$		$(6.4 \pm 0.4) \times 10^{-4}$		—
$2K_S^0\pi^0$		$< 5.9 \times 10^{-4}$		740
$K^+K^-\pi^+\pi^-$	[r]	$(2.43 \pm 0.12) \times 10^{-3}$		677
$\phi\pi^+\pi^-\text{3-body}, \phi \rightarrow$ K^+K^-		$(2.4 \pm 2.4) \times 10^{-5}$		614
$\phi\rho^0, \phi \rightarrow K^+K^-$		$(7.0 \pm 0.6) \times 10^{-4}$		250
$K^+K^-\rho^0\text{3-body}$		$(5 \pm 7) \times 10^{-5}$		302
$f_0(980)\pi^+\pi^-, f_0 \rightarrow K^+K^-$		$(3.6 \pm 0.9) \times 10^{-4}$		—
$K^*(892)^0K^\mp\pi^\pm\text{3-body},$ $K^{*0} \rightarrow K^\pm\pi^\mp$	[s]	$(2.7 \pm 0.6) \times 10^{-4}$		531
$K^*(892)^0\bar{K}^*(892)^0, K^{*0} \rightarrow$ $K^\pm\pi^\mp$		$(7 \pm 5) \times 10^{-5}$		272
$K_1(1270)^\pm K^\mp,$ $K_1(1270)^\pm \rightarrow K^\pm\pi^+\pi^-$		$(8.0 \pm 1.8) \times 10^{-4}$		—
$K_1(1400)^\pm K^\mp,$ $K_1(1400)^\pm \rightarrow K^\pm\pi^+\pi^-$		$(5.3 \pm 1.2) \times 10^{-4}$		—
$2K_S^0\pi^+\pi^-$		$(1.23 \pm 0.23) \times 10^{-3}$		673
$K_S^0K^-\pi^+\pi^-$		$< 1.4 \times 10^{-4}$	CL=90%	595
$K^+K^-\pi^+\pi^-\pi^0$		$(3.1 \pm 2.0) \times 10^{-3}$		600

Other $K\bar{K}X$ modes. They include all decay modes of the ϕ , η , and ω .

$\phi\eta$		$(1.4 \pm 0.5) \times 10^{-4}$		489
$\phi\omega$		$< 2.1 \times 10^{-3}$	CL=90%	238

Radiative modes

$\rho^0\gamma$		$< 2.4 \times 10^{-4}$	CL=90%	771
$\omega\gamma$		$< 2.4 \times 10^{-4}$	CL=90%	768
$\phi\gamma$		$(2.70 \pm 0.35) \times 10^{-5}$		654
$\bar{K}^*(892)^0\gamma$		$(3.27 \pm 0.34) \times 10^{-4}$		719

Doubly Cabibbo suppressed (DC) modes or $\Delta C = 2$ forbidden via mixing (C2M) modes

$K^+\ell^-\bar{\nu}_\ell$ via \bar{D}^0		$< 2.2 \times 10^{-5}$	CL=90%	—
K^+ or $K^*(892)^+$ $e^-\bar{\nu}_e$ via \bar{D}^0		$< 6 \times 10^{-5}$	CL=90%	—
$K^+\pi^-$	DC	$(1.47 \pm 0.07) \times 10^{-4}$	S=2.8	861
$K^+\pi^-$ via DCS		$(1.31 \pm 0.08) \times 10^{-4}$		—
$K^+\pi^-$ via \bar{D}^0		$< 1.6 \times 10^{-5}$	CL=95%	861
$K_S^0\pi^+\pi^-$ in $D^0 \rightarrow \bar{D}^0$		$< 1.8 \times 10^{-4}$	CL=95%	—
$K^*(892)^+\pi^-$, $K^*(892)^+ \rightarrow K_S^0\pi^+$	DC	$(1.13 \pm 0.60 \mp 0.34) \times 10^{-4}$		711
$K_0^*(1430)^+\pi^-$, $K_0^*(1430)^+ \rightarrow K_S^0\pi^+$	DC	$< 1.4 \times 10^{-5}$		—
$K_2^*(1430)^+\pi^-$, $K_2^*(1430)^+ \rightarrow K_S^0\pi^+$	DC	$< 3.4 \times 10^{-5}$		—
$K^+\pi^-\pi^0$	DC	$(3.04 \pm 0.17) \times 10^{-4}$		844
$K^+\pi^-\pi^0$ via \bar{D}^0		$(7.3 \pm 0.5) \times 10^{-4}$		—
$K^+\pi^+2\pi^-$	DC	$(2.61 \pm 0.21 \mp 0.19) \times 10^{-4}$		813
$K^+\pi^+2\pi^-$ via \bar{D}^0		$< 4 \times 10^{-4}$	CL=90%	812
μ^- anything via \bar{D}^0		$< 4 \times 10^{-4}$	CL=90%	—

**$\Delta C = 1$ weak neutral current (C1) modes,
Lepton Family number (LF) violating modes,
Lepton (L) or Baryon (B) number violating modes**

$\gamma\gamma$	C1	$< 2.6 \times 10^{-5}$	CL=90%	932
e^+e^-	C1	$< 7.9 \times 10^{-8}$	CL=90%	932
$\mu^+\mu^-$	C1	$< 1.4 \times 10^{-7}$	CL=90%	926
$\pi^0e^+e^-$	C1	$< 4.5 \times 10^{-5}$	CL=90%	928
$\pi^0\mu^+\mu^-$	C1	$< 1.8 \times 10^{-4}$	CL=90%	915
ηe^+e^-	C1	$< 1.1 \times 10^{-4}$	CL=90%	852
$\eta\mu^+\mu^-$	C1	$< 5.3 \times 10^{-4}$	CL=90%	838
$\pi^+\pi^-e^+e^-$	C1	$< 3.73 \times 10^{-4}$	CL=90%	922
$\rho^0e^+e^-$	C1	$< 1.0 \times 10^{-4}$	CL=90%	771
$\pi^+\pi^-\mu^+\mu^-$	C1	$< 3.0 \times 10^{-5}$	CL=90%	894

$\rho^0 \mu^+ \mu^-$	<i>CI</i>	< 2.2	$\times 10^{-5}$	CL=90%	754
$\omega e^+ e^-$	<i>CI</i>	< 1.8	$\times 10^{-4}$	CL=90%	768
$\omega \mu^+ \mu^-$	<i>CI</i>	< 8.3	$\times 10^{-4}$	CL=90%	751
$K^- K^+ e^+ e^-$	<i>CI</i>	< 3.15	$\times 10^{-4}$	CL=90%	791
$\phi e^+ e^-$	<i>CI</i>	< 5.2	$\times 10^{-5}$	CL=90%	654
$K^- K^+ \mu^+ \mu^-$	<i>CI</i>	< 3.3	$\times 10^{-5}$	CL=90%	710
$\phi \mu^+ \mu^-$	<i>CI</i>	< 3.1	$\times 10^{-5}$	CL=90%	631
$\overline{K}^0 e^+ e^-$		[<i>h</i>] < 1.1	$\times 10^{-4}$	CL=90%	866
$\overline{K}^0 \mu^+ \mu^-$		[<i>h</i>] < 2.6	$\times 10^{-4}$	CL=90%	852
$K^- \pi^+ e^+ e^-$	<i>CI</i>	< 3.85	$\times 10^{-4}$	CL=90%	861
$\overline{K}^*(892)^0 e^+ e^-$		[<i>h</i>] < 4.7	$\times 10^{-5}$	CL=90%	719
$K^- \pi^+ \mu^+ \mu^-$	<i>CI</i>	< 3.59	$\times 10^{-4}$	CL=90%	829
$\overline{K}^*(892)^0 \mu^+ \mu^-$		[<i>h</i>] < 2.4	$\times 10^{-5}$	CL=90%	700
$\pi^+ \pi^- \pi^0 \mu^+ \mu^-$	<i>CI</i>	< 8.1	$\times 10^{-4}$	CL=90%	863
$\mu^\pm e^\mp$	<i>LF</i>	[<i>t</i>] < 2.6	$\times 10^{-7}$	CL=90%	929
$\pi^0 e^\pm \mu^\mp$	<i>LF</i>	[<i>t</i>] < 8.6	$\times 10^{-5}$	CL=90%	924
$\eta e^\pm \mu^\mp$	<i>LF</i>	[<i>t</i>] < 1.0	$\times 10^{-4}$	CL=90%	848
$\pi^+ \pi^- e^\pm \mu^\mp$	<i>LF</i>	[<i>t</i>] < 1.5	$\times 10^{-5}$	CL=90%	911
$\rho^0 e^\pm \mu^\mp$	<i>LF</i>	[<i>t</i>] < 4.9	$\times 10^{-5}$	CL=90%	767
$\omega e^\pm \mu^\mp$	<i>LF</i>	[<i>t</i>] < 1.2	$\times 10^{-4}$	CL=90%	764
$K^- K^+ e^\pm \mu^\mp$	<i>LF</i>	[<i>t</i>] < 1.8	$\times 10^{-4}$	CL=90%	754
$\phi e^\pm \mu^\mp$	<i>LF</i>	[<i>t</i>] < 3.4	$\times 10^{-5}$	CL=90%	648
$\overline{K}^0 e^\pm \mu^\mp$	<i>LF</i>	[<i>t</i>] < 1.0	$\times 10^{-4}$	CL=90%	863
$K^- \pi^+ e^\pm \mu^\mp$	<i>LF</i>	[<i>t</i>] < 5.53	$\times 10^{-4}$	CL=90%	848
$\overline{K}^*(892)^0 e^\pm \mu^\mp$	<i>LF</i>	[<i>t</i>] < 8.3	$\times 10^{-5}$	CL=90%	714
$2\pi^- 2e^+ + \text{c.c.}$	<i>L</i>	< 1.12	$\times 10^{-4}$	CL=90%	922
$2\pi^- 2\mu^+ + \text{c.c.}$	<i>L</i>	< 2.9	$\times 10^{-5}$	CL=90%	894
$K^- \pi^- 2e^+ + \text{c.c.}$	<i>L</i>	< 2.06	$\times 10^{-4}$	CL=90%	861
$K^- \pi^- 2\mu^+ + \text{c.c.}$	<i>L</i>	< 3.9	$\times 10^{-4}$	CL=90%	829
$2K^- 2e^+ + \text{c.c.}$	<i>L</i>	< 1.52	$\times 10^{-4}$	CL=90%	791
$2K^- 2\mu^+ + \text{c.c.}$	<i>L</i>	< 9.4	$\times 10^{-5}$	CL=90%	710
$\pi^- \pi^- e^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 7.9	$\times 10^{-5}$	CL=90%	911
$K^- \pi^- e^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 2.18	$\times 10^{-4}$	CL=90%	848
$2K^- e^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 5.7	$\times 10^{-5}$	CL=90%	754
$p e^-$	<i>L,B</i>	[<i>u</i>] < 1.0	$\times 10^{-5}$	CL=90%	696
$\overline{p} e^+$	<i>L,B</i>	[<i>v</i>] < 1.1	$\times 10^{-5}$	CL=90%	696

$D^*(2007)^0$

$$I(J^P) = \frac{1}{2}(1^-)$$

I, J, P need confirmation.

Mass $m = 2006.98 \pm 0.15$ MeV

$m_{D^{*0}} - m_{D^0} = 142.12 \pm 0.07$ MeV

Full width $\Gamma < 2.1$ MeV, CL = 90%

$\bar{D}^*(2007)^0$ modes are charge conjugates of modes below.

$D^*(2007)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 \pi^0$	(61.9±2.9) %	43
$D^0 \gamma$	(38.1±2.9) %	137

$D^*(2010)^\pm$

$$I(J^P) = \frac{1}{2}(1^-)$$

I, J, P need confirmation.

Mass $m = 2010.28 \pm 0.13$ MeV

$m_{D^*(2010)^+} - m_{D^+} = 140.66 \pm 0.10$ MeV (S = 1.1)

$m_{D^*(2010)^+} - m_{D^0} = 145.421 \pm 0.010$ MeV (S = 1.1)

Full width $\Gamma = 96 \pm 22$ keV

$D^*(2010)^-$ modes are charge conjugates of the modes below.

$D^*(2010)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 \pi^+$	(67.7±0.5) %	39
$D^+ \pi^0$	(30.7±0.5) %	38
$D^+ \gamma$	(1.6±0.4) %	136

$D_0^*(2400)^0$

$$I(J^P) = \frac{1}{2}(0^+)$$

Mass $m = 2318 \pm 29$ MeV (S = 1.7)

Full width $\Gamma = 267 \pm 40$ MeV

$D_0^*(2400)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^+ \pi^-$	seen	385

$D_1(2420)^0$

$$I(J^P) = \frac{1}{2}(1^+)$$

I needs confirmation.

Mass $m = 2421.3 \pm 0.6$ MeV (S = 1.2)

$m_{D_1^0} - m_{D^{*+}} = 411.0 \pm 0.6$ (S = 1.2)

Full width $\Gamma = 27.1 \pm 2.7$ MeV (S = 2.4)

$\bar{D}_1(2420)^0$ modes are charge conjugates of modes below.

$D_1(2420)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^*(2010)^+ \pi^-$	seen	354
$D^0 \pi^+ \pi^-$	seen	425
$D^+ \pi^-$	not seen	473
$D^{*0} \pi^+ \pi^-$	not seen	279

$D_2^*(2460)^0$

$$I(J^P) = \frac{1}{2}(2^+)$$

$J^P = 2^+$ assignment strongly favored.

$$\text{Mass } m = 2462.6 \pm 0.7 \text{ MeV} \quad (S = 1.3)$$

$$m_{D_2^{*0}} - m_{D^+} = 593.0 \pm 0.7 \text{ MeV} \quad (S = 1.3)$$

$$m_{D_2^{*0}} - m_{D^{*+}} = 452.3 \pm 0.7 \text{ MeV} \quad (S = 1.3)$$

$$\text{Full width } \Gamma = 49.0 \pm 1.4 \text{ MeV} \quad (S = 1.7)$$

$\bar{D}_2^*(2460)^0$ modes are charge conjugates of modes below.

$D_2^*(2460)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^+ \pi^-$	seen	507
$D^*(2010)^+ \pi^-$	seen	391
$D^0 \pi^+ \pi^-$	not seen	463
$D^{*0} \pi^+ \pi^-$	not seen	326

$D_2^*(2460)^\pm$

$$I(J^P) = \frac{1}{2}(2^+)$$

$J^P = 2^+$ assignment strongly favored.

$$\text{Mass } m = 2464.4 \pm 1.9 \text{ MeV} \quad (S = 1.9)$$

$$m_{D_2^*(2460)^\pm} - m_{D_2^*(2460)^0} = 2.4 \pm 1.7 \text{ MeV}$$

$$\text{Full width } \Gamma = 37 \pm 6 \text{ MeV} \quad (S = 1.4)$$

$D_2^*(2460)^-$ modes are charge conjugates of modes below.

$D_2^*(2460)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 \pi^+$	seen	512
$D^{*0} \pi^+$	seen	395
$D^+ \pi^+ \pi^-$	not seen	461
$D^{*+} \pi^+ \pi^-$	not seen	325

NOTES

- [a] This result applies to $Z^0 \rightarrow c\bar{c}$ decays only. Here ℓ^+ is an average (not a sum) of e^+ and μ^+ decays.
- [b] See the Particle Listings for the (complicated) definition of this quantity.
- [c] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers in the Particle Listings.
- [d] These subfractions of the $K^- 2\pi^+$ mode are uncertain: see the Particle Listings.
- [e] Submodes of the $D^+ \rightarrow K^- 2\pi^+ \pi^0$ and $K_S^0 2\pi^+ \pi^-$ modes were studied by ANJOS 92C and COFFMAN 92B, but with at most 142 events for the first mode and 229 for the second – not enough for precise results. With nothing new for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.
- [f] The unseen decay modes of the resonances are included.
- [g] This is *not* a test for the $\Delta C=1$ weak neutral current, but leads to the $\pi^+ \ell^+ \ell^-$ final state.
- [h] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.
- [i] In the 2010 *Review*, the values for these quantities were given using a measure of the asymmetry that was inconsistent with the usual definition.
- [j] This value is obtained by subtracting the branching fractions for 2-, 4- and 6-prongs from unity.
- [k] This is the sum of our $K^- 2\pi^+ \pi^-$, $K^- 2\pi^+ \pi^- \pi^0$, $K^0 2\pi^+ 2\pi^-$, $K^+ 2K^- \pi^+$, $2\pi^+ 2\pi^-$, $2\pi^+ 2\pi^- \pi^0$, $K^+ K^- \pi^+ \pi^-$, and $K^+ K^- \pi^+ \pi^- \pi^0$, branching fractions.
- [l] This is the sum of our $K^- 3\pi^+ 2\pi^-$ and $3\pi^+ 3\pi^-$ branching fractions.
- [m] The branching fractions for the $K^- e^+ \nu_e$, $K^*(892)^- e^+ \nu_e$, $\pi^- e^+ \nu_e$, and $\rho^- e^+ \nu_e$ modes add up to 6.19 ± 0.17 %.
- [n] This is a doubly Cabibbo-suppressed mode.
- [o] The two experiments measuring this fraction are in serious disagreement. See the Particle Listings.
- [p] Submodes of the $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ mode with a K^* and/or ρ were studied by COFFMAN 92B, but with only 140 events. With nothing new for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.
- [q] This branching fraction includes all the decay modes of the resonance in the final state.

- [r] The experiments on the division of this charge mode amongst its sub-modes disagree, and the submode branching fractions here add up to considerably more than the charged-mode fraction.
- [s] However, these upper limits are in serious disagreement with values obtained in another experiment.
- [t] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [u] This limit is for either D^0 or \bar{D}^0 to $p e^-$.
- [v] This limit is for either D^0 or \bar{D}^0 to $\bar{p} e^+$.