



$$I(J^P) = 0(?^?)$$

J^P is natural, width and decay modes consistent with 1^- .

$D_s^{*\pm}$ MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

2112.3 ± 0.5 OUR FIT Error includes scale factor of 1.1.

2106.6 ± 2.1 ± 2.7 ¹BLAYLOCK 87 MRK3 $e^+e^- \rightarrow D_s^\pm \gamma X$

¹ Assuming D_s^\pm mass = 1968.7 ± 0.9 MeV.

$m_{D_s^{*\pm}} - m_{D_s^\pm}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

143.8 ± 0.4 OUR FIT

143.9 ± 0.4 OUR AVERAGE

143.76 ± 0.39 ± 0.40

GRONBERG 95 CLE2 e^+e^-

144.22 ± 0.47 ± 0.37

BROWN 94 CLE2 e^+e^-

142.5 ± 0.8 ± 1.5

²ALBRECHT 88 ARG $e^+e^- \rightarrow D_s^\pm \gamma X$

139.5 ± 8.3 ± 9.7

60 AIHARA 84D TPC $e^+e^- \rightarrow$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

143.0 ± 18.0

8 ASRATYAN 85 HLBC FNAL 15-ft, ν -²H

110 ± 46

BRANDELIK 79 DASP $e^+e^- \rightarrow D_s^\pm \gamma X$

² Result includes data of ALBRECHT 84B.

$D_s^{*\pm}$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
-------------	-----	-------------	------	---------

< 1.9

90 GRONBERG 95 CLE2 e^+e^-

< 4.5

90 ALBRECHT 88 ARG $E_{cm}^{ee} = 10.2$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.9

90 BROWN 94 CLE2 e^+e^-

< 22

90 BLAYLOCK 87 MRK3 $e^+e^- \rightarrow D_s^\pm \gamma X$

$\Gamma(D_s^+ \pi^0) / \Gamma(D_s^+ \gamma)$				Γ_2 / Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.062 ± 0.008 OUR FIT				
0.062 ± 0.008 OUR AVERAGE				
0.062 ± 0.005 ± 0.006	AUBERT, BE	05G BABR	10.6 e ⁺ e ⁻ → hadrons	
0.062 ^{+0.020} _{-0.018} ± 0.022	GRONBERG	95 CLE2	e ⁺ e ⁻	

³ Derived from the ratio $\Gamma(D_s^+ \pi^0) / \Gamma(D_s^+ \gamma)$ assuming that the branching fractions of $D_s^{*+} \rightarrow D_s^+ \pi^0$ and $D_s^{*+} \rightarrow D_s^+ \gamma$ decays sum to 100%.

$D_s^{*\pm}$ REFERENCES

AUBERT, BE	05G	PR D72 091101	B. Aubert <i>et al.</i>	(BABAR Collab.)
GRONBERG	95	PRL 75 3232	J. Gronberg <i>et al.</i>	(CLEO Collab.)
BROWN	94	PR D50 1884	D. Brown <i>et al.</i>	(CLEO Collab.)
ASRATYAN	91	PL B257 525	A.E. Asratyan <i>et al.</i>	(ITEP, BELG, SACL+)
ALBRECHT	88	PL B207 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BLAYLOCK	87	PRL 58 2171	G.T. Blaylock <i>et al.</i>	(Mark III Collab.)
ASRATYAN	85	PL 156B 441	A.E. Asratyan <i>et al.</i>	(ITEP, SERP)
AIHARA	84D	PRL 53 2465	H. Aihara <i>et al.</i>	(TPC Collab.)
ALBRECHT	84B	PL 146B 111	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BRANDELIK	79	PL 80B 412	R. Brandelik <i>et al.</i>	(DASP Collab.)