

# QUARKS

The  $u$ -,  $d$ -, and  $s$ -quark masses are estimates of so-called “current-quark masses,” in a mass-independent subtraction scheme such as  $\overline{MS}$  at a scale  $\mu \approx 2$  GeV. The  $c$ - and  $b$ -quark masses are the “running” masses in the  $\overline{MS}$  scheme. For the  $b$ -quark we also quote the 1S mass. These can be different from the heavy quark masses obtained in potential models.

**u**

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$$m_u = 2.3^{+0.7}_{-0.5} \text{ MeV} \quad \text{Charge} = \frac{2}{3} e \quad I_z = +\frac{1}{2}$$

$$m_u/m_d = 0.38\text{--}0.58$$

**d**

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$$m_d = 4.8^{+0.5}_{-0.3} \text{ MeV} \quad \text{Charge} = -\frac{1}{3} e \quad I_z = -\frac{1}{2}$$

$$m_s/m_d = 17\text{--}22$$

$$\bar{m} = (m_u + m_d)/2 = 3.5^{+0.7}_{-0.2} \text{ MeV}$$

**s**

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$m_s = 95 \pm 5 \text{ MeV} \quad \text{Charge} = -\frac{1}{3} e \quad \text{Strangeness} = -1$$

$$m_s / ((m_u + m_d)/2) = 27.5 \pm 1.0$$

**c**

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$m_c = 1.275 \pm 0.025 \text{ GeV} \quad \text{Charge} = \frac{2}{3} e \quad \text{Charm} = +1$$

**b**

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$\text{Charge} = -\frac{1}{3} e \quad \text{Bottom} = -1$$

$$m_b(\overline{MS}) = 4.18 \pm 0.03 \text{ GeV}$$

$$m_b(1S) = 4.66 \pm 0.03 \text{ GeV}$$

**t**

$$I(J^P) = 0(\frac{1}{2}^+)$$

Charge =  $\frac{2}{3} e$       Top = +1

Mass (direct measurements)  $m = 173.21 \pm 0.51 \pm 0.71 \text{ GeV} [a,b]$   
 Mass ( $\overline{MS}$  from cross-section measurements)  $m = 160_{-4}^{+5} \text{ GeV} [a]$   
 Mass (Pole from cross-section measurements)  $m = 176.7_{-3.4}^{+4.0} \text{ GeV}$   
 $m_t - m_{\bar{t}} = -0.2 \pm 0.5 \text{ GeV} \quad (S = 1.1)$   
 Full width  $\Gamma = 2.0 \pm 0.5 \text{ GeV}$   
 $\Gamma(Wb)/\Gamma(Wq(q = b, s, d)) = 0.91 \pm 0.04$

**t-quark EW Couplings**

$F_0 = 0.690 \pm 0.030$   
 $F_- = 0.314 \pm 0.025$   
 $F_+ = 0.008 \pm 0.016$   
 $F_{V+A} < 0.29, \text{ CL} = 95\%$

<b>t DECAY MODES</b>	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level	$\frac{p}{\text{MeV}/c}$
$Wq(q = b, s, d)$			—
$Wb$			—
$\ell\nu_\ell$ anything	[c,d] (9.4±2.4) %		—
$\gamma q(q=u,c)$	[e] < 5.9	$\times 10^{-3}$	95%
<b><math>\Delta T = 1</math> weak neutral current (T1) modes</b>			
$Zq(q=u,c)$	T1 [f] < 2.1	$\times 10^{-3}$	95%

**b' (4<sup>th</sup> Generation) Quark, Searches for**

Mass  $m > 190 \text{ GeV}, \text{ CL} = 95\%$  ( $p\bar{p}$ , quasi-stable  $b'$ )  
 Mass  $m > 400 \text{ GeV}, \text{ CL} = 95\%$  ( $pp$ , neutral-current decays)  
 Mass  $m > 675 \text{ GeV}, \text{ CL} = 95\%$  ( $pp$ , charged-current decays)  
 Mass  $m > 46.0 \text{ GeV}, \text{ CL} = 95\%$  ( $e^+e^-$ , all decays)

**t' (4<sup>th</sup> Generation) Quark, Searches for**

Mass  $m > 782 \text{ GeV}, \text{ CL} = 95\%$  ( $pp$ , neutral-current decays)  
 Mass  $m > 700 \text{ GeV}, \text{ CL} = 95\%$  ( $pp$ , charged-current decays)

**Free Quark Searches**

All searches since 1977 have had negative results.

## NOTES

- [a] A discussion of the definition of the top quark mass in these measurements can be found in the review “The Top Quark.”
- [b] Based on published top mass measurements using data from Tevatron Run-I and Run-II and LHC at  $\sqrt{s} = 7$  TeV. Including the most recent unpublished results from Tevatron Run-II, the Tevatron Electroweak Working Group reports a top mass of  $173.2 \pm 0.9$  GeV. See the note “The Top Quark’ in the Quark Particle Listings of this *Review*.
- [c]  $\ell$  means  $e$  or  $\mu$  decay mode, not the sum over them.
- [d] Assumes lepton universality and  $W$ -decay acceptance.
- [e] This limit is for  $\Gamma(t \rightarrow \gamma q)/\Gamma(t \rightarrow W b)$ .
- [f] This limit is for  $\Gamma(t \rightarrow Z q)/\Gamma(t \rightarrow W b)$ .