

$N(1700) \ 3/2^-$ $I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$ Status: ***

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, *Physics Letters* **111B** 1 (1982). Some further obsolete results published before 1984 were last included in our 2006 edition, *Journal of Physics* **G33** 1 (2006).

The various partial-wave analyses do not agree very well.

The latest GWU analysis (ARNDT 06) finds no evidence for this resonance.

$N(1700)$ BREIT-WIGNER MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1650 to 1750 (≈ 1700) OUR ESTIMATE			
1790 \pm 40	ANISOVICH	12A DPWA	Multichannel
1675 \pm 25	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$
1731 \pm 15	HOEHLER	79 IPWA	$\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1665 \pm 3	SHRESTHA	12A DPWA	Multichannel
1817 \pm 22	BATINIC	10 DPWA	$\pi N \rightarrow N\pi, N\eta$
1740 \pm 20	THOMA	08 DPWA	Multichannel
1736 \pm 33	VRANA	00 DPWA	Multichannel
1737 \pm 44	MANLEY	92 IPWA	$\pi N \rightarrow \pi N \ \& \ N\pi\pi$
1650	SAXON	80 DPWA	$\pi^- p \rightarrow \Lambda K^0$
1690 to 1710	BAKER	78 DPWA	$\pi^- p \rightarrow \Lambda K^0$
1719	BARBOUR	78 DPWA	$\gamma N \rightarrow \pi N$
1670 \pm 10	¹ BAKER	77 IPWA	$\pi^- p \rightarrow \Lambda K^0$
1690	¹ BAKER	77 DPWA	$\pi^- p \rightarrow \Lambda K^0$
1660	² LONGACRE	77 IPWA	$\pi N \rightarrow N\pi\pi$
1710	³ LONGACRE	75 IPWA	$\pi N \rightarrow N\pi\pi$

$N(1700)$ BREIT-WIGNER WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
100 to 250 (≈ 150) OUR ESTIMATE			
390 \pm 140	ANISOVICH	12A DPWA	Multichannel
90 \pm 40	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$
110 \pm 30	HOEHLER	79 IPWA	$\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
56 \pm 8	SHRESTHA	12A DPWA	Multichannel
134 \pm 37	BATINIC	10 DPWA	$\pi N \rightarrow N\pi, N\eta$
180 \pm 30	THOMA	08 DPWA	Multichannel
175 \pm 133	VRANA	00 DPWA	Multichannel
250 \pm 220	MANLEY	92 IPWA	$\pi N \rightarrow \pi N \ \& \ N\pi\pi$
70	SAXON	80 DPWA	$\pi^- p \rightarrow \Lambda K^0$

70 to 100	BAKER	78	DPWA	$\pi^- p \rightarrow \Lambda K^0$
126	BARBOUR	78	DPWA	$\gamma N \rightarrow \pi N$
90 ± 25	¹ BAKER	77	IPWA	$\pi^- p \rightarrow \Lambda K^0$
100	¹ BAKER	77	DPWA	$\pi^- p \rightarrow \Lambda K^0$
600	² LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
300	³ LONGACRE	75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1700) POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1650 to 1750 (≈ 1700) OUR ESTIMATE			
1757 ± 4 ± 1	⁴ SVARC	14	MLS $\pi N \rightarrow \pi N$
1770 ± 40	ANISOVICH	12A	DPWA Multichannel
1700	⁵ HOEHLER	93	SPED $\pi N \rightarrow \pi N$
1660 ± 30	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1662	SHRESTHA	12A	DPWA Multichannel
1806 ± 23	BATINIC	10	DPWA $\pi N \rightarrow N\pi, N\eta$
1710 ± 15	THOMA	08	DPWA Multichannel
1704	VRANA	00	DPWA Multichannel
not seen	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
1710 or 1678	⁶ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
1616 or 1613	² LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

−2×IMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
100 to 300 OUR ESTIMATE			
136 ± 7 ± 4	⁴ SVARC	14	MLS $\pi N \rightarrow \pi N$
420 ± 180	ANISOVICH	12A	DPWA Multichannel
120	⁵ HOEHLER	93	SPED $\pi N \rightarrow \pi N$
90 ± 40	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
55	SHRESTHA	12A	DPWA Multichannel
129 ± 33	BATINIC	10	DPWA $\pi N \rightarrow N\pi, N\eta$
155 ± 25	THOMA	08	DPWA Multichannel
156	VRANA	00	DPWA Multichannel
not seen	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
607 or 567	⁶ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
577 or 575	² LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

N(1700) ELASTIC POLE RESIDUE

MODULUS |r|

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
5 to 50 OUR ESTIMATE			
7 ± 1 ± 1	⁴ SVARC	14	MLS $\pi N \rightarrow \pi N$
50 ± 40	ANISOVICH	12A	DPWA Multichannel
5	HOEHLER	93	SPED $\pi N \rightarrow \pi N$
6 ± 3	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
7	BATINIC	10	DPWA $\pi N \rightarrow N\pi, N\eta$

PHASE θ

<u>VALUE ($^{\circ}$)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
–120 to 20 OUR ESTIMATE			
–113 \pm 4 \pm 2	⁴ SVARC	14	MLS $\pi N \rightarrow \pi N$
–100 \pm 40	ANISOVICH	12A	DPWA Multichannel
0 \pm 50	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
– 34	BATINIC	10	DPWA $\pi N \rightarrow N\pi, N\eta$

N(1700) INELASTIC POLE RESIDUE

The “normalized residue” is the residue divided by $\Gamma_{pole}/2$.

Normalized residue in $N\pi \rightarrow N(1700) \rightarrow \Delta\pi$, S-wave

<u>MODULUS (%)</u>	<u>PHASE ($^{\circ}$)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
34\pm21	–60 \pm 40	ANISOVICH	12A	DPWA Multichannel

Normalized residue in $N\pi \rightarrow N(1700) \rightarrow \Delta\pi$, D-wave

<u>MODULUS (%)</u>	<u>PHASE ($^{\circ}$)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8\pm6	90 \pm 35	ANISOVICH	12A	DPWA Multichannel

N(1700) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	(12 \pm 5) %
Γ_2 $N\eta$	(0.0 \pm 1.0) %
Γ_3 ΛK	< 3 %
Γ_4 ΣK	
Γ_5 $N\pi\pi$	85–95 %
Γ_6 $\Delta\pi$	
Γ_7 $\Delta(1232)\pi$, S-wave	10–90 %
Γ_8 $\Delta(1232)\pi$, D-wave	< 20 %
Γ_9 $N\rho$	< 35 %
Γ_{10} $N\rho$, S=1/2, D-wave	
Γ_{11} $N\rho$, S=3/2, S-wave	(7.0 \pm 1.0) %
Γ_{12} $N\rho$, S=3/2, D-wave	
Γ_{13} $N(\pi\pi)_{S\text{-wave}}^{I=0}$	
Γ_{14} $p\gamma$	0.01–0.05 %
Γ_{15} $p\gamma$, helicity=1/2	0.0–0.024 %
Γ_{16} $p\gamma$, helicity=3/2	0.002–0.026 %
Γ_{17} $n\gamma$	0.01–0.13 %
Γ_{18} $n\gamma$, helicity=1/2	0.0–0.09 %
Γ_{19} $n\gamma$, helicity=3/2	0.01–0.05 %

N(1700) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$ Γ_1/Γ
VALUE (%) DOCUMENT ID TECN COMMENT

12 ±5 OUR ESTIMATE

12 ±5	ANISOVICH	12A	DPWA	Multichannel
11 ±5	CUTKOSKY	80	IPWA	$\pi N \rightarrow \pi N$
8 ±3	HOEHLER	79	IPWA	$\pi N \rightarrow \pi N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2.8±0.5	SHRESTHA	12A	DPWA	Multichannel
9 ±6	BATINIC	10	DPWA	$\pi N \rightarrow N\pi, N\eta$
8 +8 -4	THOMA	08	DPWA	Multichannel
4 ±2	VRANA	00	DPWA	Multichannel
1 ±2	MANLEY	92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$

$\Gamma(N\eta)/\Gamma_{\text{total}}$ Γ_2/Γ
VALUE (%) DOCUMENT ID TECN COMMENT

0±1

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
14±5	BATINIC	10	DPWA	$\pi N \rightarrow N\pi, N\eta$
10±5	THOMA	08	DPWA	Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1700) \rightarrow \Lambda K$ $(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
VALUE DOCUMENT ID TECN COMMENT

-0.06 to +0.04 OUR ESTIMATE

-0.012	BELL	83	DPWA	$\pi^- p \rightarrow \Lambda K^0$
-0.012	SAXON	80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
-0.04	⁷ BAKER	78	DPWA	See SAXON 80
-0.03 ±0.004	¹ BAKER	77	IPWA	$\pi^- p \rightarrow \Lambda K^0$
-0.03	¹ BAKER	77	DPWA	$\pi^- p \rightarrow \Lambda K^0$
+0.026±0.019	DEVENISH	74B		Fixed- <i>t</i> dispersion rel.

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1700) \rightarrow \Sigma K$ $(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
VALUE DOCUMENT ID TECN COMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

not seen	LIVANOS	80	DPWA	$\pi p \rightarrow \Sigma K$
<0.017	⁸ DEANS	75	DPWA	$\pi N \rightarrow \Sigma K$

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1700) \rightarrow \Delta(1232)\pi, S\text{-wave}$ $(\Gamma_1\Gamma_7)^{1/2}/\Gamma$
VALUE DOCUMENT ID TECN COMMENT

0.00	² LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
-0.16	³ LONGACRE	75	IPWA	$\pi N \rightarrow N\pi\pi$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

+0.02±0.03 MANLEY 92 IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$

$\Gamma(\Delta(1232)\pi, S\text{-wave})/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (%) DOCUMENT ID TECN COMMENT

10 to 90 OUR ESTIMATE

72±23 ANISOVICH 12A DPWA Multichannel
 11±1 VRANA 00 DPWA Multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

31±9 SHRESTHA 12A DPWA Multichannel
 10±5 THOMA 08 DPWA Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1700) \rightarrow \Delta(1232)\pi, D\text{-wave}$ $(\Gamma_1\Gamma_8)^{1/2}/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

−0.12 ² LONGACRE 77 IPWA $\pi N \rightarrow N\pi\pi$
 +0.14 ³ LONGACRE 75 IPWA $\pi N \rightarrow N\pi\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

+0.10±0.09 MANLEY 92 IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$

$\Gamma(\Delta(1232)\pi, D\text{-wave})/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (%) DOCUMENT ID TECN COMMENT

<20 OUR ESTIMATE

<10 ANISOVICH 12A DPWA Multichannel
 79±56 VRANA 00 DPWA Multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

3±2 SHRESTHA 12A DPWA Multichannel
 20±11 THOMA 08 DPWA Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1700) \rightarrow N\rho, S=3/2, S\text{-wave}$ $(\Gamma_1\Gamma_{11})^{1/2}/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

±0.01 to ±0.13 OUR ESTIMATE

−0.07 ² LONGACRE 77 IPWA $\pi N \rightarrow N\pi\pi$
 +0.07 ³ LONGACRE 75 IPWA $\pi N \rightarrow N\pi\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

−0.04±0.06 MANLEY 92 IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$

$\Gamma(N\rho, S=3/2, S\text{-wave})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (%) DOCUMENT ID TECN COMMENT

7±1 VRANA 00 DPWA Multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

38±6 SHRESTHA 12A DPWA Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1700) \rightarrow N(\pi\pi)_{S\text{-wave}}^{J=0}$ $(\Gamma_1\Gamma_{13})^{1/2}/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

±0.02 to ±0.28 OUR ESTIMATE

0.00 ² LONGACRE 77 IPWA $\pi N \rightarrow N\pi\pi$
 +0.2 ³ LONGACRE 75 IPWA $\pi N \rightarrow N\pi\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

+0.02±0.02 MANLEY 92 IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$

$\Gamma(N(\pi\pi)_{S\text{-wave}}^{I=0})/\Gamma_{\text{total}}$					Γ_{13}/Γ
VALUE (%)	DOCUMENT ID	TECN	COMMENT		
0 ± 1	VRANA	00	DPWA	Multichannel	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
24 ± 6	SHRESTHA	12A	DPWA	Multichannel	
18 ± 12	THOMA	08	DPWA	Multichannel	

$N(1700)$ PHOTON DECAY AMPLITUDES

Papers on γN amplitudes predating 1981 may be found in our 2006 edition, *Journal of Physics* **G33** 1 (2006).

$N(1700) \rightarrow p\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE ($\text{GeV}^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
0.015 ± 0.025 OUR ESTIMATE			
0.041 ± 0.017	ANISOVICH	12A	DPWA Multichannel
-0.016 ± 0.014	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
-0.002 ± 0.013	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.021 ± 0.005	SHRESTHA	12A	DPWA Multichannel
-0.033 ± 0.021	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$

$N(1700) \rightarrow p\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE ($\text{GeV}^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
-0.015 ± 0.025 OUR ESTIMATE			
-0.034 ± 0.013	ANISOVICH	12A	DPWA Multichannel
-0.009 ± 0.012	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
0.029 ± 0.014	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.050 ± 0.009	SHRESTHA	12A	DPWA Multichannel
-0.014 ± 0.025	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$

$N(1700) \rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE ($\text{GeV}^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
0.020 ± 0.015 OUR ESTIMATE			
0.025 ± 0.010	ANISOVICH	13B	DPWA Multichannel
0.006 ± 0.024	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.002 ± 0.013	FUJII	81	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.049 ± 0.008	SHRESTHA	12A	DPWA Multichannel
$+0.050 \pm 0.042$	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$

$N(1700) \rightarrow n\gamma$, helicity-3/2 amplitude $A_{3/2}$

<u>VALUE (GeV^{-1/2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
−0.030±0.020 OUR ESTIMATE			
−0.032±0.018	ANISOVICH	13B	DPWA Multichannel
−0.033±0.017	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
0.018±0.018	FUJII	81	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
−0.092±0.014	SHRESTHA	12A	DPWA Multichannel
+0.035±0.030	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$

 $N(1700) \quad \gamma p \rightarrow \Lambda K^+$ AMPLITUDES **$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $p\gamma \rightarrow N(1700) \rightarrow \Lambda K^+$ (E_{2-} amplitude)**

<u>VALUE (units 10⁻³)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •		
4.09	TANABE	89 DPWA

 $(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $p\gamma \rightarrow N(1700) \rightarrow \Lambda K^+$ (M_{2-} amplitude)

<u>VALUE (units 10⁻³)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •		
−7.09	TANABE	89 DPWA

 $p\gamma \rightarrow N(1700) \rightarrow \Lambda K^+$ phase angle θ (E_{2-} amplitude)

<u>VALUE (degrees)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •		
−35.9	TANABE	89 DPWA

 $N(1700)$ FOOTNOTES

- ¹ The two BAKER 77 entries are from an IPWA using the Barrelet-zero method and from a conventional energy-dependent analysis.
- ² LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- ³ From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- ⁴ Fit to the amplitudes of HOEHLER 79.
- ⁵ See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- ⁶ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- ⁷ The overall phase of BAKER 78 couplings has been changed to agree with previous conventions.
- ⁸ The range given is from the four best solutions.

N(1700) REFERENCESFor early references, see *Physics Letters* **111B** 1 (1982).

SVARC	14	PR C89 045205	A. Svarc <i>et al.</i>	
ANISOVICH	13B	EPJ A49 67	A.V. Anisovich <i>et al.</i>	
ANISOVICH	12A	EPJ A48 15	A.V. Anisovich <i>et al.</i>	(BONN, PNPI)
SHRESTHA	12A	PR C86 055203	M. Shrestha, D.M. Manley	(KSU)
BATINIC	10	PR C82 038203	M. Batinic <i>et al.</i>	(ZAGR)
THOMA	08	PL B659 87	U. Thoma <i>et al.</i>	(CB-ELSA Collab.)
ARNDT	06	PR C74 045205	R.A. Arndt <i>et al.</i>	(GWU)
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
VRANA	00	PRPL 328 181	T.P. Vrana, S.A. Dytman,, T.-S.H. Lee	(PITT+)
HOEHLER	93	π N Newsletter 9 1	G. Hohler	(KARL)
MANLEY	92	PR D45 4002	D.M. Manley, E.M. Saleski	(KSA) IJP
Also		PR D30 904	D.M. Manley <i>et al.</i>	(VPI)
ARNDT	91	PR D43 2131	R.A. Arndt <i>et al.</i>	(VPI, TELE) IJP
TANABE	89	PR C39 741	H. Tanabe, M. Kohno, C. Bennhold	(MANZ)
Also		NC 102A 193	M. Kohno, H. Tanabe, C. Bennhold	(MANZ)
BELL	83	NP B222 389	K.W. Bell <i>et al.</i>	(RL) IJP
CRAWFORD	83	NP B211 1	R.L. Crawford, W.T. Morton	(GLAS)
PDG	82	PL 111B 1	M. Roos <i>et al.</i>	(HELSE, CIT, CERN)
AWAJI	81	Bonn Conf. 352	N. Awaji, R. Kajikawa	(NAGO)
Also		NP B197 365	K. Fujii <i>et al.</i>	(NAGO)
FUJII	81	NP B187 53	K. Fujii <i>et al.</i>	(NAGO, OSAK)
CUTKOSKY	80	Toronto Conf. 19	R.E. Cutkosky <i>et al.</i>	(CMU, LBL) IJP
Also		PR D20 2839	R.E. Cutkosky <i>et al.</i>	(CMU, LBL) IJP
LIVANOS	80	Toronto Conf. 35	P. Livanos <i>et al.</i>	(SACL) IJP
SAXON	80	NP B162 522	D.H. Saxon <i>et al.</i>	(RHEL, BRIS) IJP
HOEHLER	79	PDAT 12-1	G. Hohler <i>et al.</i>	(KARLT) IJP
Also		Toronto Conf. 3	R. Koch	(KARLT) IJP
BAKER	78	NP B141 29	R.D. Baker <i>et al.</i>	(RL, CAVE) IJP
BARBOUR	78	NP B141 253	I.M. Barbour, R.L. Crawford, N.H. Parsons	(GLAS)
LONGACRE	78	PR D17 1795	R.S. Longacre <i>et al.</i>	(LBL, SLAC)
BAKER	77	NP B126 365	R.D. Baker <i>et al.</i>	(RHEL) IJP
LONGACRE	77	NP B122 493	R.S. Longacre, J. Dolbeau	(SACL) IJP
Also		NP B108 365	J. Dolbeau <i>et al.</i>	(SACL) IJP
DEANS	75	NP B96 90	S.R. Deans <i>et al.</i>	(SFLA, ALAH) IJP
LONGACRE	75	PL 55B 415	R.S. Longacre <i>et al.</i>	(LBL, SLAC) IJP
DEVENISH	74B	NP B81 330	R.C.E. Devenish, C.D. Froggatt, B.R. Martin	(DESY+)