

# $\rho(1700)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

A REVIEW GOES HERE – Check our WWW List of Reviews

---

## $\rho(1700)$ MASS

### $\eta\rho^0$ AND $\pi^+\pi^-$ MODES

| VALUE (MeV)                 | DOCUMENT ID |
|-----------------------------|-------------|
| <b>1720±20 OUR ESTIMATE</b> |             |

### $\eta\rho^0$ MODE

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|---------|
|-------------|-------------|------|---------|

The data in this block is included in the average printed for a previous datablock.

• • • We do not use the following data for averages, fits, limits, etc. • • •

|         |                    |    |      |   |
|---------|--------------------|----|------|---|
| 1740±20 | ANTONELLI          | 88 | DM2  | $e^+e^- \rightarrow \eta\pi^+\pi^-$         |
| 1701±15 | <sup>1</sup> FUKUI | 88 | SPEC | $8.95 \pi^- p \rightarrow \eta\pi^+\pi^- n$ |

<sup>1</sup> Assuming  $\rho^+ f_0(1370)$  decay mode interferes with  $a_1(1260)^+ \pi$  background. From a two Breit-Wigner fit.

### $\pi\pi$ MODE

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------|
|-------------|------|-------------|------|---------|

The data in this block is included in the average printed for a previous datablock.

• • • We do not use the following data for averages, fits, limits, etc. • • •

|   |       |                           |      |   |
|---|-------|---------------------------|------|---|
| 1780 ±20 $\begin{smallmatrix} +15 \\ -20 \end{smallmatrix}$ | 63.5k | <sup>2</sup> ABRAMOWICZ12 | ZEUS | $ep \rightarrow e\pi^+\pi^- p$                  |
| 1861 ±17  |       | <sup>3</sup> LEES         | 12G  | BABR $e^+e^- \rightarrow \pi^+\pi^-\gamma$      |
| 1728 ±17 $\pm 89$   | 5.4M  | <sup>4,5</sup> FUJIKAWA   | 08   | BELL $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$    |
| 1780 $\begin{smallmatrix} +37 \\ -29 \end{smallmatrix}$     |       | <sup>6</sup> ABELE        | 97   | CBAR $\bar{p}n \rightarrow \pi^-\pi^0\pi^0$     |
| 1719 ±15  |       | <sup>6</sup> BERTIN       | 97C  | OBLX $0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$ |
| 1730 ±30  |       | CLEGG                     | 94   | RVUE $e^+e^- \rightarrow \pi^+\pi^-$            |
| 1768 ±21  |       | BISELLO                   | 89   | DM2 $e^+e^- \rightarrow \pi^+\pi^-$             |
| 1745.7±91.9   |       | DUBNICKA                  | 89   | RVUE $e^+e^- \rightarrow \pi^+\pi^-$            |
| 1546 ±26  |       | GESHKEN...                | 89   | RVUE  |
| 1650  |       | <sup>7</sup> ERKAL        | 85   | RVUE $20-70 \gamma p \rightarrow \gamma\pi$     |
| 1550 ±70  |       | ABE                       | 84B  | HYBR $20 \gamma p \rightarrow \pi^+\pi^- p$     |
| 1590 ±20  |       | <sup>8</sup> ASTON        | 80   | OMEG $20-70 \gamma p \rightarrow p2\pi$         |
| 1600 ±10  |       | <sup>9</sup> ATIYA        | 79B  | SPEC $50 \gamma C \rightarrow C2\pi$            |
| 1598 $\begin{smallmatrix} +24 \\ -22 \end{smallmatrix}$     |       | BECKER                    | 79   | ASPK $17 \pi^- p$ polarized                     |
| 1659 ±25  |       | <sup>7</sup> LANG         | 79   | RVUE  |
| 1575  |       | <sup>7</sup> MARTIN       | 78C  | RVUE $17 \pi^- p \rightarrow \pi^+\pi^- n$      |
| 1610 ±30  |       | <sup>7</sup> FROGGATT     | 77   | RVUE $17 \pi^- p \rightarrow \pi^+\pi^- n$      |
| 1590 ±20  |       | <sup>10</sup> HYAMS       | 73   | ASPK $17 \pi^- p \rightarrow \pi^+\pi^- n$      |

- <sup>2</sup> Using the KUHN 90 parametrization of the pion form factor, neglecting  $\rho$ - $\omega$  interference.  
<sup>3</sup> Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the  $\rho(1450)$ ,  $\rho(1700)$ , and  $\rho(2150)$  resonances as free parameters of the fit.  
<sup>4</sup>  $|F_\pi(0)|^2$  fixed to 1.  
<sup>5</sup> From the GOUNARIS 68 parametrization of the pion form factor.  
<sup>6</sup> T-matrix pole.  
<sup>7</sup> From phase shift analysis of HYAMS 73 data.  
<sup>8</sup> Simple relativistic Breit-Wigner fit with constant width.  
<sup>9</sup> An additional 40 MeV uncertainty in both the mass and width is present due to the choice of the background shape.  
<sup>10</sup> Included in BECKER 79 analysis.

### $\pi\omega$ MODE

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------|
|-------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|               |      |                       |     |      |   |
|---------------|------|-----------------------|-----|------|---|
| $1708 \pm 41$ | 7815 | <sup>11</sup> ACHASOV | 13  | SND  | 1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$ |
| 1550 to 1620  |      | <sup>12</sup> ACHASOV | 00i | SND  | $e^+e^- \rightarrow \pi^0\pi^0\gamma$           |
| 1580 to 1710  |      | <sup>13</sup> ACHASOV | 00i | SND  | $e^+e^- \rightarrow \pi^0\pi^0\gamma$           |
| $1710 \pm 90$ |      | ACHASOV               | 97  | RVUE | $e^+e^- \rightarrow \omega\pi^0$                |

- <sup>11</sup> From a phenomenological model based on vector meson dominance with the interfering  $\rho(1450)$  and  $\rho(1700)$  and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.  
<sup>12</sup> Taking into account both  $\rho(1450)$  and  $\rho(1700)$  contributions. Using the data of ACHASOV 00i on  $e^+e^- \rightarrow \omega\pi^0$  and of EDWARDS 00A on  $\tau^- \rightarrow \omega\pi^- \nu_\tau$ .  $\rho(1450)$  mass and width fixed at 1400 MeV and 500 MeV respectively.  
<sup>13</sup> Taking into account the  $\rho(1700)$  contribution only. Using the data of ACHASOV 00i on  $e^+e^- \rightarrow \omega\pi^0$  and of EDWARDS 00A on  $\tau^- \rightarrow \omega\pi^- \nu_\tau$ .

### $K\bar{K}$ MODE

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | CHG | COMMENT |
|-------------|------|-------------|------|-----|---------|
|-------------|------|-------------|------|-----|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                   |      |                     |     |      |  |
|-------------------|------|---------------------|-----|------|--|
| $1740.8 \pm 22.2$ | 27k  | <sup>14</sup> ABELE | 99D | CBAR | $\pm$ 0.0 $\bar{p}p \rightarrow K^+K^-\pi^0$ |
| $1582 \pm 36$     | 1600 | CLELAND             | 82B | SPEC | $\pm$ 50 $\pi p \rightarrow K_S^0 K^\pm p$   |

- <sup>14</sup> K-matrix pole. Isospin not determined, could be  $\omega(1650)$  or  $\phi(1680)$ .

### $2(\pi^+\pi^-)$ MODE

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------|
|-------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                    |     |                         |     |      |                                       |
|--------------------|-----|-------------------------|-----|------|---------------------------------------|
| $1851^{+27}_{-24}$ |     | ACHASOV                 | 97  | RVUE | $e^+e^- \rightarrow 2(\pi^+\pi^-)$    |
| $1570 \pm 20$      |     | <sup>15</sup> CORDIER   | 82  | DM1  | $e^+e^- \rightarrow 2(\pi^+\pi^-)$    |
| $1520 \pm 30$      |     | <sup>16</sup> ASTON     | 81E | OMEG | 20–70 $\gamma p \rightarrow p4\pi$    |
| $1654 \pm 25$      |     | <sup>17</sup> DIBIANCA  | 81  | DBC  | $\pi^+ d \rightarrow pp2(\pi^+\pi^-)$ |
| $1666 \pm 39$      |     | <sup>15</sup> BACCI     | 80  | FRAG | $e^+e^- \rightarrow 2(\pi^+\pi^-)$    |
| 1780               | 34  | KILLIAN                 | 80  | SPEC | 11 $e^- p \rightarrow 2(\pi^+\pi^-)$  |
| 1500               |     | <sup>18</sup> ATIYA     | 79B | SPEC | 50 $\gamma C \rightarrow C4\pi^\pm$   |
| $1570 \pm 60$      | 65  | <sup>19</sup> ALEXANDER | 75  | HBC  | 7.5 $\gamma p \rightarrow p4\pi$      |
| $1550 \pm 60$      |     | <sup>16</sup> CONVERSI  | 74  | OSPK | $e^+e^- \rightarrow 2(\pi^+\pi^-)$    |
| $1550 \pm 50$      | 160 | SCHACHT                 | 74  | STRC | 5.5–9 $\gamma p \rightarrow p4\pi$    |
| $1450 \pm 100$     | 340 | SCHACHT                 | 74  | STRC | 9–18 $\gamma p \rightarrow p4\pi$     |
| $1430 \pm 50$      | 400 | BINGHAM                 | 72B | HBC  | 9.3 $\gamma p \rightarrow p4\pi$      |

<sup>15</sup> Simple relativistic Breit-Wigner fit with model dependent width.

<sup>16</sup> Simple relativistic Breit-Wigner fit with constant width.

<sup>17</sup> One peak fit result.

<sup>18</sup> Parameters roughly estimated, not from a fit.

<sup>19</sup> Skew mass distribution compensated by Ross-Stodolsky factor.

### $\pi^+\pi^-\pi^0\pi^0$ MODE

| VALUE (MeV)   | DOCUMENT ID | TECN | COMMENT               |
|---|-------------|------|-----------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |      |                       |
| 1660 ± 30   | ATKINSON    | 85B  | OMEG 20-70 $\gamma p$ |

### $3(\pi^+\pi^-)$ AND $2(\pi^+\pi^-\pi^0)$ MODES

| VALUE (MeV)   | DOCUMENT ID            | TECN | COMMENT   |
|---|------------------------|------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                        |      |   |
| 1730 ± 34   | <sup>20</sup> FRABETTI | 04   | E687 $\gamma p \rightarrow 3\pi^+ 3\pi^- p$               |
| 1783 ± 15   | CLEGG                  | 90   | RVUE $e^+e^- \rightarrow 3(\pi^+\pi^-)2(\pi^+\pi^-\pi^0)$ |
| <sup>20</sup> From a fit with two resonances with the JACOB 72 continuum.     |                        |      |   |

### $\rho(1700)$ WIDTH

### $\eta\rho^0$ AND $\pi^+\pi^-$ MODES

| VALUE (MeV)                   | DOCUMENT ID |
|-------------------------------|-------------|
| <b>250 ± 100 OUR ESTIMATE</b> |             |

### $\eta\rho^0$ MODE

| VALUE (MeV)   | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|---------|
| The data in this block is included in the average printed for a previous datablock. |             |      |         |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|          |                     |    |  |
|----------|---------------------|----|--|
| 150 ± 30 | ANTONELLI           | 88 | DM2 $e^+e^- \rightarrow \eta\pi^+\pi^-$          |
| 282 ± 44 | <sup>21</sup> FUKUI | 88 | SPEC 8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$ |

<sup>21</sup> Assuming  $\rho^+ f_0(1370)$  decay mode interferes with  $a_1(1260)^+ \pi$  background. From a two Breit-Wigner fit.

### $\pi\pi$ MODE

| VALUE (MeV)   | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---------|
| The data in this block is included in the average printed for a previous datablock. |      |             |      |         |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                         |       |                            |          |  |
|-------------------------|-------|----------------------------|----------|--|
| 310 ± 30 $^{+25}_{-35}$ | 63.5k | <sup>22</sup> ABRAMOWICZ12 | ZEUS     | $ep \rightarrow e\pi^+\pi^- p$             |
| 316 ± 26                |       | <sup>23</sup> LEES         | 12G BABR | $e^+e^- \rightarrow \pi^+\pi^-\gamma$      |
| 164 ± 21 $^{+89}_{-26}$ | 5.4M  | <sup>24,25</sup> FUJIKAWA  | 08 BELL  | $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$    |
| 275 ± 45                |       | <sup>26</sup> ABELE        | 97 CBAR  | $\bar{p}n \rightarrow \pi^-\pi^0\pi^0$     |
| 310 ± 40                |       | <sup>26</sup> BERTIN       | 97C OBLX | $0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$ |
| 400 ± 100               |       | CLEGG                      | 94 RVUE  | $e^+e^- \rightarrow \pi^+\pi^-$            |
| 224 ± 22                |       | BISELLO                    | 89 DM2   | $e^+e^- \rightarrow \pi^+\pi^-$            |
| 242.5 ± 163.0           |       | DUBNICKA                   | 89 RVUE  | $e^+e^- \rightarrow \pi^+\pi^-$            |
| 620 ± 60                |       | GESHKEN...                 | 89 RVUE  |  |

|           |  |             |          |   |
|-----------|--|-------------|----------|---|
| <315      |  | 27 ERKAL    | 85 RVUE  | 20–70 $\gamma p \rightarrow \gamma \pi$ |
| 280 + 30  |  | ABE         | 84B HYBR | 20 $\gamma p \rightarrow \pi^+ \pi^- p$ |
| 230 ± 80  |  | 28 ASTON    | 80 OMEG  | 20–70 $\gamma p \rightarrow p 2\pi$     |
| 283 ± 14  |  | 29 ATIYA    | 79B SPEC | 50 $\gamma C \rightarrow C 2\pi$        |
| 175 + 98  |  | BECKER      | 79 ASPK  | 17 $\pi^- p$ polarized                  |
| 175 - 53  |  |             |          |   |
| 232 ± 34  |  | 27 LANG     | 79 RVUE  |   |
| 340       |  | 27 MARTIN   | 78C RVUE | 17 $\pi^- p \rightarrow \pi^+ \pi^- n$  |
| 300 ± 100 |  | 27 FROGGATT | 77 RVUE  | 17 $\pi^- p \rightarrow \pi^+ \pi^- n$  |
| 180 ± 50  |  | 30 HYAMS    | 73 ASPK  | 17 $\pi^- p \rightarrow \pi^+ \pi^- n$  |

<sup>22</sup> Using the KUHN 90 parametrization of the pion form factor, neglecting  $\rho$ - $\omega$  interference.

<sup>23</sup> Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the  $\rho(1450)$ ,  $\rho(1700)$ , and  $\rho(2150)$  resonances as free parameters of the fit.

<sup>24</sup>  $|F_\pi(0)|^2$  fixed to 1.

<sup>25</sup> From the GOUNARIS 68 parametrization of the pion form factor.

<sup>26</sup> T-matrix pole.

<sup>27</sup> From phase shift analysis of HYAMS 73 data.

<sup>28</sup> Simple relativistic Breit-Wigner fit with constant width.

<sup>29</sup> An additional 40 MeV uncertainty in both the mass and width is present due to the choice of the background shape.

<sup>30</sup> Included in BECKER 79 analysis.

## $K\bar{K}$ MODE

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | CHG | COMMENT |
|-------------|------|-------------|------|-----|---------|
|-------------|------|-------------|------|-----|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|              |      |                     |          |   |   |
|--------------|------|---------------------|----------|---|---|
| 187.2 ± 26.7 | 27k  | <sup>31</sup> ABELE | 99D CBAR | ± | 0.0 $\bar{p} p \rightarrow K^+ K^- \pi^0$ |
| 265 ± 120    | 1600 | CLELAND             | 82B SPEC | ± | 50 $\pi p \rightarrow K_S^0 K^\pm p$      |

<sup>31</sup> K-matrix pole. Isospin not determined, could be  $\omega(1650)$  or  $\phi(1680)$ .

## $2(\pi^+ \pi^-)$ MODE

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------|
|-------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|           |     |                         |          |  |
|-----------|-----|-------------------------|----------|--|
| 510 ± 40  |     | <sup>32</sup> CORDIER   | 82 DM1   | $e^+ e^- \rightarrow 2(\pi^+ \pi^-)$     |
| 400 ± 50  |     | <sup>33</sup> ASTON     | 81E OMEG | 20–70 $\gamma p \rightarrow p 4\pi$      |
| 400 ± 146 |     | <sup>34</sup> DIBIANCA  | 81 DBC   | $\pi^+ d \rightarrow p p 2(\pi^+ \pi^-)$ |
| 700 ± 160 |     | <sup>32</sup> BACCI     | 80 FRAG  | $e^+ e^- \rightarrow 2(\pi^+ \pi^-)$     |
| 100       | 34  | KILLIAN                 | 80 SPEC  | 11 $e^- p \rightarrow 2(\pi^+ \pi^-)$    |
| 600       |     | <sup>35</sup> ATIYA     | 79B SPEC | 50 $\gamma C \rightarrow C 4\pi^\pm$     |
| 340 ± 160 | 65  | <sup>36</sup> ALEXANDER | 75 HBC   | 7.5 $\gamma p \rightarrow p 4\pi$        |
| 360 ± 100 |     | <sup>33</sup> CONVERSI  | 74 OSPK  | $e^+ e^- \rightarrow 2(\pi^+ \pi^-)$     |
| 400 ± 120 | 160 | <sup>37</sup> SCHACHT   | 74 STRC  | 5.5–9 $\gamma p \rightarrow p 4\pi$      |
| 850 ± 200 | 340 | <sup>37</sup> SCHACHT   | 74 STRC  | 9–18 $\gamma p \rightarrow p 4\pi$       |
| 650 ± 100 | 400 | BINGHAM                 | 72B HBC  | 9.3 $\gamma p \rightarrow p 4\pi$        |

<sup>32</sup> Simple relativistic Breit-Wigner fit with model-dependent width.

<sup>33</sup> Simple relativistic Breit-Wigner fit with constant width.

<sup>34</sup> One peak fit result.

<sup>35</sup> Parameters roughly estimated, not from a fit.

<sup>36</sup> Skew mass distribution compensated by Ross-Stodolsky factor.

<sup>37</sup> Width errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

### $\pi^+\pi^-\pi^0\pi^0$ MODE

| VALUE (MeV)   | DOCUMENT ID | TECN | COMMENT               |
|---|-------------|------|-----------------------|
| ••• We do not use the following data for averages, fits, limits, etc. ••• |             |      |                       |
| $300 \pm 50$  | ATKINSON    | 85B  | OMEG $20-70 \gamma p$ |

### $\omega\pi^0$ MODE

| VALUE (MeV)   | DOCUMENT ID           | TECN | COMMENT                                   |
|---|-----------------------|------|---|
| ••• We do not use the following data for averages, fits, limits, etc. ••• |                       |      |   |
| 350 to 580  | <sup>38</sup> ACHASOV | 00i  | SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$ |
| 490 to 1040   | <sup>39</sup> ACHASOV | 00i  | SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$ |

<sup>38</sup>Taking into account both  $\rho(1450)$  and  $\rho(1700)$  contributions. Using the data of ACHASOV 00i on  $e^+e^- \rightarrow \omega\pi^0$  and of EDWARDS 00A on  $\tau^- \rightarrow \omega\pi^-\nu_\tau$ .  $\rho(1450)$  mass and width fixed at 1400 MeV and 500 MeV respectively.

<sup>39</sup>Taking into account the  $\rho(1700)$  contribution only. Using the data of ACHASOV 00i on  $e^+e^- \rightarrow \omega\pi^0$  and of EDWARDS 00A on  $\tau^- \rightarrow \omega\pi^-\nu_\tau$ .

### $3(\pi^+\pi^-)$ AND $2(\pi^+\pi^-\pi^0)$ MODES

| VALUE (MeV)   | DOCUMENT ID            | TECN | COMMENT   |
|---|------------------------|------|---|
| ••• We do not use the following data for averages, fits, limits, etc. ••• |                        |      |   |
| $315 \pm 100$   | <sup>40</sup> FRABETTI | 04   | E687 $\gamma p \rightarrow 3\pi^+3\pi^-\rho$              |
| $285 \pm 20$  | CLEGG                  | 90   | RVUE $e^+e^- \rightarrow 3(\pi^+\pi^-)2(\pi^+\pi^-\pi^0)$ |

<sup>40</sup>From a fit with two resonances with the JACOB 72 continuum.

### $\rho(1700)$ DECAY MODES

| Mode  | Fraction ( $\Gamma_i/\Gamma$ ) |
|---|--------------------------------|
| $\Gamma_1$ $4\pi$                             |                                |
| $\Gamma_2$ $2(\pi^+\pi^-)$                    | large                          |
| $\Gamma_3$ $\rho\pi\pi$                       | dominant                       |
| $\Gamma_4$ $\rho^0\pi^+\pi^-$                 | large                          |
| $\Gamma_5$ $\rho^0\pi^0\pi^0$                 |                                |
| $\Gamma_6$ $\rho^\pm\pi^\mp\pi^0$             | large                          |
| $\Gamma_7$ $a_1(1260)\pi$                     | seen                           |
| $\Gamma_8$ $h_1(1170)\pi$                     | seen                           |
| $\Gamma_9$ $\pi(1300)\pi$                     | seen                           |
| $\Gamma_{10}$ $\rho\rho$                      | seen                           |
| $\Gamma_{11}$ $\pi^+\pi^-$                    | seen                           |
| $\Gamma_{12}$ $\pi\pi$                        | seen                           |
| $\Gamma_{13}$ $K\bar{K}^*(892) + \text{c.c.}$ | seen                           |
| $\Gamma_{14}$ $\eta\rho$                      | seen                           |
| $\Gamma_{15}$ $a_2(1320)\pi$                  | not seen                       |
| $\Gamma_{16}$ $K\bar{K}$                      | seen                           |
| $\Gamma_{17}$ $e^+e^-$                        | seen                           |
| $\Gamma_{18}$ $\pi^0\omega$                   | seen                           |

### $\rho(1700) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into  $e^+e^-$  and with the total width is obtained from the cross-section into channel<sub>l</sub> in  $e^+e^-$  annihilation.

#### $\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_2\Gamma_{17}/\Gamma$

| <u>VALUE (keV)</u>  | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                     |
|---|--------------------|-------------|------------------------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                    |             |                                    |
| 2.6 ± 0.2   | DEL COURT          | 81B DM1     | $e^+e^- \rightarrow 2(\pi^+\pi^-)$ |
| 2.83 ± 0.42   | BACCI              | 80 FRAG     | $e^+e^- \rightarrow 2(\pi^+\pi^-)$ |

#### $\Gamma(\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{11}\Gamma_{17}/\Gamma$

| <u>VALUE (keV)</u>  | <u>DOCUMENT ID</u>    | <u>TECN</u> | <u>COMMENT</u>                           |
|---|-----------------------|-------------|--|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                       |             |  |
| 0.13  | <sup>41</sup> DIEKMAN | 88 RVUE     | $e^+e^- \rightarrow \pi^+\pi^-$          |
| 0.029 <sup>+0.016</sup> <sub>-0.012</sub>                                     | KURDADZE              | 83 OLYA     | 0.64–1.4 $e^+e^- \rightarrow \pi^+\pi^-$ |

<sup>41</sup> Using total width = 220 MeV.

#### $\Gamma(K\bar{K}^*(892) + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{13}\Gamma_{17}/\Gamma$

| <u>VALUE (keV)</u>  | <u>DOCUMENT ID</u>  | <u>TECN</u> | <u>COMMENT</u> |
|---|---------------------|-------------|----------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                     |             |                |
| 0.305 ± 0.071   | <sup>42</sup> BIZOT | 80 DM1      | $e^+e^-$       |

<sup>42</sup> Model dependent.

#### $\Gamma(\eta\rho) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{14}\Gamma_{17}/\Gamma$

| <u>VALUE (eV)</u>   | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                      |
|---|--------------------|-------------|-------------------------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                    |             |                                     |
| 7 ± 3   | ANTONELLI          | 88 DM2      | $e^+e^- \rightarrow \eta\pi^+\pi^-$ |

#### $\Gamma(K\bar{K}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{16}\Gamma_{17}/\Gamma$

| <u>VALUE (keV)</u>  | <u>DOCUMENT ID</u>  | <u>TECN</u> | <u>COMMENT</u> |
|---|---------------------|-------------|----------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                     |             |                |
| 0.035 ± 0.029   | <sup>43</sup> BIZOT | 80 DM1      | $e^+e^-$       |

<sup>43</sup> Model dependent.

#### $\Gamma(\rho\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_3\Gamma_{17}/\Gamma$

| <u>VALUE (keV)</u>  | <u>DOCUMENT ID</u>  | <u>TECN</u> | <u>COMMENT</u> |
|---|---------------------|-------------|----------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                     |             |                |
| 3.510 ± 0.090   | <sup>44</sup> BIZOT | 80 DM1      | $e^+e^-$       |

<sup>44</sup> Model dependent.

$\rho(1700) \Gamma(i)/\Gamma(\text{total}) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(\pi^0 \omega)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \qquad \Gamma_{18}/\Gamma \times \Gamma_{17}/\Gamma$

| VALUE (units $10^{-6}$ )  | EVTS | DOCUMENT ID           | TECN   | COMMENT  |
|---|------|-----------------------|--------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • •   |      |                       |        |  |
| $1.7 \pm 0.4$   | 7815 | <sup>45</sup> ACHASOV | 13 SND | $1.05\text{--}2.00 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$ |
| <sup>45</sup> From a phenomenological model based on vector meson dominance with the interfering $\rho(1450)$ and $\rho(1700)$ and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated. |      |                       |        |  |

$\rho(1700)$  BRANCHING RATIOS

$\Gamma(\rho \pi \pi)/\Gamma(4\pi) \qquad \Gamma_3/\Gamma_1$

| VALUE   | DOCUMENT ID         | TECN     | COMMENT                          |
|---|---------------------|----------|----------------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                     |          |                                  |
| $0.28 \pm 0.06$   | <sup>46</sup> ABELE | 01B CBAR | $0.0 \bar{p} n \rightarrow 5\pi$ |
| <sup>46</sup> $\omega \pi$ not included.                                      |                     |          |                                  |

$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma(2(\pi^+ \pi^-)) \qquad \Gamma_4/\Gamma_2$

| VALUE   | EVTS | DOCUMENT ID           | TECN    | COMMENT   |
|---|------|-----------------------|---------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |                       |         |   |
| $\sim 1.0$  |      | DELCOURT              | 81B DM1 | $e^+ e^- \rightarrow 2(\pi^+ \pi^-)$            |
| $0.7 \pm 0.1$   | 500  | SCHACHT               | 74 STRC | $5.5\text{--}18 \gamma p \rightarrow \rho 4\pi$ |
| 0.80  |      | <sup>47</sup> BINGHAM | 72B HBC | $9.3 \gamma p \rightarrow \rho 4\pi$            |
| <sup>47</sup> The $\pi \pi$ system is in <i>S</i> -wave.                      |      |                       |         |   |

$\Gamma(\rho^0 \pi^0 \pi^0)/\Gamma(\rho^\pm \pi^\mp \pi^0) \qquad \Gamma_5/\Gamma_6$

| VALUE   | DOCUMENT ID | TECN      | CHG | COMMENT  |
|---|-------------|-----------|-----|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |           |     |  |
| $< 0.10$  | ATKINSON    | 85B OMEG  |     | $20\text{--}70 \gamma p$                       |
| $< 0.15$  | ATKINSON    | 82 OMEG 0 |     | $20\text{--}70 \gamma p \rightarrow \rho 4\pi$ |

$\Gamma(a_1(1260)\pi)/\Gamma(4\pi) \qquad \Gamma_7/\Gamma_1$

| VALUE   | DOCUMENT ID         | TECN     | COMMENT                          |
|---|---------------------|----------|----------------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                     |          |                                  |
| $0.16 \pm 0.05$   | <sup>48</sup> ABELE | 01B CBAR | $0.0 \bar{p} n \rightarrow 5\pi$ |
| <sup>48</sup> $\omega \pi$ not included.                                      |                     |          |                                  |

$\Gamma(h_1(1170)\pi)/\Gamma(4\pi) \qquad \Gamma_8/\Gamma_1$

| VALUE   | DOCUMENT ID         | TECN     | COMMENT                          |
|---|---------------------|----------|----------------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                     |          |                                  |
| $0.17 \pm 0.06$   | <sup>49</sup> ABELE | 01B CBAR | $0.0 \bar{p} n \rightarrow 5\pi$ |
| <sup>49</sup> $\omega \pi$ not included.                                      |                     |          |                                  |



$\Gamma(K\bar{K}^*(892)+c.c.)/\Gamma(2(\pi^+\pi^-))$   $\Gamma_{13}/\Gamma_2$

| VALUE   | CL% | DOCUMENT ID                | TECN | COMMENT                          |
|---|-----|----------------------------|------|----------------------------------|
| 0.15±0.03   |     | <sup>60</sup> DELCOURT 81B | DM1  | $e^+e^- \rightarrow \bar{K}K\pi$ |
| <sup>60</sup> Assuming $\rho(1700)$ and $\omega$ radial excitations to be degenerate in mass. |     |                            |      |                                  |

$\Gamma(\eta\rho)/\Gamma_{total}$   $\Gamma_{14}/\Gamma$

| VALUE | CL% | DOCUMENT ID    | TECN | COMMENT                             |
|-------|-----|----------------|------|-------------------------------------|
| <0.04 |     | AKHMETSHIN 00D | CMD2 | $e^+e^- \rightarrow \eta\pi^+\pi^-$ |
| <0.02 | 58  | DONNACHIE 87B  | RVUE |                                     |
|       |     | ATKINSON 86B   | OMEG | 20–70 $\gamma p$                    |

$\Gamma(\eta\rho)/\Gamma(2(\pi^+\pi^-))$   $\Gamma_{14}/\Gamma_2$

| VALUE       | DOCUMENT ID | TECN | COMMENT                           |
|-------------|-------------|------|-----------------------------------|
| 0.123±0.027 | DELCOURT 82 | DM1  | $e^+e^- \rightarrow \pi^+\pi^-MM$ |
| ~ 0.1       | ASTON 80    | OMEG | 20–70 $\gamma p$                  |

$\Gamma(\pi^+\pi^- \text{ neutrals})/\Gamma(2(\pi^+\pi^-))$   $(\Gamma_5+\Gamma_6+0.714\Gamma_{14})/\Gamma_2$

| VALUE   | DOCUMENT ID             | TECN | COMMENT        |
|---|-------------------------|------|----------------|
| 2.6±0.4   | <sup>61</sup> BALLAM 74 | HBC  | 9.3 $\gamma p$ |
| <sup>61</sup> Upper limit. Background not subtracted. |                         |      |                |

$\Gamma(a_2(1320)\pi)/\Gamma_{total}$   $\Gamma_{15}/\Gamma$

| VALUE    | DOCUMENT ID | TECN | COMMENT                                   |
|----------|-------------|------|---|
| not seen | AMELIN 00   | VES  | 37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$ |

$\Gamma(K\bar{K})/\Gamma(2(\pi^+\pi^-))$   $\Gamma_{16}/\Gamma_2$

| VALUE   | CL% | DOCUMENT ID                | TECN | CHG | COMMENT                       |
|---|-----|----------------------------|------|-----|-------------------------------|
| 0.015±0.010   |     | <sup>62</sup> DELCOURT 81B | DM1  |     | $e^+e^- \rightarrow \bar{K}K$ |
| <0.04   | 95  | BINGHAM 72B                | HBC  | 0   | 9.3 $\gamma p$                |
| <sup>62</sup> Assuming $\rho(1700)$ and $\omega$ radial excitations to be degenerate in mass. |     |                            |      |     |                               |

$\Gamma(K\bar{K})/\Gamma(K\bar{K}^*(892)+c.c.)$   $\Gamma_{16}/\Gamma_{13}$

| VALUE       | DOCUMENT ID | TECN | COMMENT                             |
|-------------|-------------|------|-------------------------------------|
| 0.052±0.026 | BUON 82     | DM1  | $e^+e^- \rightarrow \text{hadrons}$ |

$\Gamma(\pi^0\omega)/\Gamma_{total}$   $\Gamma_{18}/\Gamma$

| VALUE    | EVTS | DOCUMENT ID    | TECN | COMMENT                               |
|----------|------|----------------|------|---------------------------------------|
| seen     | 1.6k | ACHASOV 12     | SND  | $e^+e^- \rightarrow \pi^0\pi^0\gamma$ |
| not seen | 2382 | AKHMETSHIN 03B | CMD2 | $e^+e^- \rightarrow \pi^0\pi^0\gamma$ |
| seen     |      | ACHASOV 97     | RVUE | $e^+e^- \rightarrow \omega\pi^0$      |

$\rho(1700)$  REFERENCES

|            |     |  |  |                                 |
|------------|-----|--|--|---------------------------------|
| ACHASOV    | 13  | PR D88 054013                                  | M.N. Achasov <i>et al.</i>                   | (SND Collab.)                   |
| ABRAMOWICZ | 12  | EPJ C72 1869                                   | H. Abramowicz <i>et al.</i>                  | (ZEUS Collab.)                  |
| ACHASOV    | 12  | JETPL 94 734                                   | M.N. Achasov <i>et al.</i>                   |                                 |
| LEES       | 12G | Translated from ZETFP 94 796.<br>PR D86 032013 | J.P. Lees <i>et al.</i>                      | (BABAR Collab.)                 |
| FUJIKAWA   | 08  | PR D78 072006                                  | M. Fujikawa <i>et al.</i>                    | (BELLE Collab.)                 |
| COAN       | 04  | PRL 92 232001                                  | T.E. Coan <i>et al.</i>                      | (CLEO Collab.)                  |
| FRABETTI   | 04  | PL B578 290                                    | P.L. Frabetti <i>et al.</i>                  | (FNAL E687 Collab.)             |
| AKHMETSHIN | 03B | PL B562 173                                    | R.R. Akhmetshin <i>et al.</i>                | (Novosibirsk CMD-2 Collab.)     |
| ABELE      | 01B | EPJ C21 261                                    | A. Abele <i>et al.</i>                       | (Crystal Barrel Collab.)        |
| ACHASOV    | 00I | PL B486 29                                     | M.N. Achasov <i>et al.</i>                   | (Novosibirsk SND Collab.)       |
| AKHMETSHIN | 00D | PL B489 125                                    | R.R. Akhmetshin <i>et al.</i>                | (Novosibirsk CMD-2 Collab.)     |
| AMELIN     | 00  | NP A668 83                                     | D. Amelin <i>et al.</i>                      | (VES Collab.)                   |
| EDWARDS    | 00A | PR D61 072003                                  | K.W. Edwards <i>et al.</i>                   | (CLEO Collab.)                  |
| ABELE      | 99D | PL B468 178                                    | A. Abele <i>et al.</i>                       | (Crystal Barrel Collab.)        |
| ABELE      | 97  | PL B391 191                                    | A. Abele <i>et al.</i>                       | (Crystal Barrel Collab.)        |
| ACHASOV    | 97  | PR D55 2663                                    | N.N. Achasov <i>et al.</i>                   | (NOVM)                          |
| BERTIN     | 97C | PL B408 476                                    | A. Bertin <i>et al.</i>                      | (OBELIX Collab.)                |
| CLEGG      | 94  | ZPHY C62 455                                   | A.B. Clegg, A. Donnachie                     | (LANC, MCHS)                    |
| CLEGG      | 90  | ZPHY C45 677                                   | A.B. Clegg, A. Donnachie                     | (LANC, MCHS)                    |
| KUHN       | 90  | ZPHY C48 445                                   | J.H. Kuhn <i>et al.</i>                      | (MPIM)                          |
| BISELLO    | 89  | PL B220 321                                    | D. Bisello <i>et al.</i>                     | (DM2 Collab.)                   |
| DUBNICKA   | 89  | JP G15 1349                                    | S. Dubnicka <i>et al.</i>                    | (JINR, SLOV)                    |
| GESHKEN... | 89  | ZPHY C45 351                                   | B.V. Geshkenbein                             | (ITEP)                          |
| ANTONELLI  | 88  | PL B212 133                                    | A. Antonelli <i>et al.</i>                   | (DM2 Collab.)                   |
| DIEKMANN   | 88  | PRPL 159 99                                    | B. Diekmann                                  | (BONN)                          |
| FUKUI      | 88  | PL B202 441                                    | S. Fukui <i>et al.</i>                       | (SUGI, NAGO, KEK, KYOT+)        |
| DONNACHIE  | 87B | ZPHY C34 257                                   | A. Donnachie, A.B. Clegg                     | (MCHS, LANC)                    |
| ATKINSON   | 86B | ZPHY C30 531                                   | M. Atkinson <i>et al.</i>                    | (BONN, CERN, GLAS+)             |
| ATKINSON   | 85B | ZPHY C26 499                                   | M. Atkinson <i>et al.</i>                    | (BONN, CERN, GLAS+)             |
| ERKAL      | 85  | ZPHY C29 485                                   | C. Erkal, M.G. Olsson                        | (WISC)                          |
| ABE        | 84B | PRL 53 751                                     | K. Abe <i>et al.</i>                         | (SLAC HFP Collab.)              |
| KURDADZE   | 83  | JETPL 37 733                                   | L.M. Kurdadze <i>et al.</i>                  | (NOVO)                          |
| ATKINSON   | 82  | Translated from ZETFP 37 613.<br>PL 108B 55    | M. Atkinson <i>et al.</i>                    | (BONN, CERN, GLAS+)             |
| BUON       | 82  | PL 118B 221                                    | J. Buon <i>et al.</i>                        | (LALO, MONP)                    |
| CLELAND    | 82B | NP B208 228                                    | W.E. Cleland <i>et al.</i>                   | (DURH, GEVA, LAUS+)             |
| CORDIER    | 82  | PL 109B 129                                    | A. Cordier <i>et al.</i>                     | (LALO)                          |
| DELCOURT   | 82  | PL 113B 93                                     | B. Delcourt <i>et al.</i>                    | (LALO)                          |
| ASTON      | 81E | NP B189 15                                     | D. Aston                                     | (BONN, CERN, EPOL, GLAS, LANC+) |
| DELCOURT   | 81B | Bonn Conf. 205                                 | B. Delcourt                                  | (ORSAY)                         |
| Also       |     | PL 109B 129                                    | A. Cordier <i>et al.</i>                     | (LALO)                          |
| DIBIANCA   | 81  | PR D23 595                                     | F.A. di Bianca <i>et al.</i>                 | (CASE, CMU)                     |
| ASTON      | 80  | PL 92B 215                                     | D. Aston                                     | (BONN, CERN, EPOL, GLAS, LANC+) |
| BACCI      | 80  | PL 95B 139                                     | C. Bacci <i>et al.</i>                       | (ROMA, FRAS)                    |
| BIZOT      | 80  | Madison Conf. 546                              | J.C. Bizot <i>et al.</i>                     | (LALO, MONP)                    |
| KILLIAN    | 80  | PR D21 3005                                    | T.J. Killian <i>et al.</i>                   | (CORN)                          |
| ATIYA      | 79B | PRL 43 1691                                    | M.S. Atiya <i>et al.</i>                     | (COLU, ILL, FNAL)               |
| BECKER     | 79  | NP B151 46                                     | H. Becker <i>et al.</i>                      | (MPIM, CERN, ZEEM, CRAC)        |
| LANG       | 79  | PR D19 956                                     | C.B. Lang, A. Mas-Parareda                   | (GRAZ)                          |
| MARTIN     | 78C | ANP 114 1                                      | A.D. Martin, M.R. Pennington                 | (CERN)                          |
| COSTA...   | 77B | PL 71B 345                                     | B. Costa de Beauregard, B. Pire, T.N. Truong | (EPOL)                          |
| FROGGATT   | 77  | NP B129 89                                     | C.D. Froggatt, J.L. Petersen                 | (GLAS, NORD)                    |
| ALEXANDER  | 75  | PL 57B 487                                     | G. Alexander <i>et al.</i>                   | (TELA)                          |
| BALLAM     | 74  | NP B76 375                                     | J. Ballam <i>et al.</i>                      | (SLAC, LBL, MPIM)               |
| CONVERSI   | 74  | PL 52B 493                                     | M. Conversi <i>et al.</i>                    | (ROMA, FRAS)                    |
| SCHACHT    | 74  | NP B81 205                                     | P. Schacht <i>et al.</i>                     | (MPIM)                          |
| DAVIER     | 73  | NP B58 31                                      | M. Davier <i>et al.</i>                      | (SLAC)                          |
| EISENBERG  | 73  | PL 43B 149                                     | Y. Eisenberg <i>et al.</i>                   | (REHO)                          |
| HYAMS      | 73  | NP B64 134                                     | B.D. Hyams <i>et al.</i>                     | (CERN, MPIM)                    |
| BINGHAM    | 72B | PL 41B 635                                     | H.H. Bingham <i>et al.</i>                   | (LBL, UCB, SLAC) IGJP           |
| JACOB      | 72  | PR D5 1847                                     | M. Jacob, R. Slansky                         |                                 |
| GOUNARIS   | 68  | PRL 21 244                                     | G.J. Gounaris, J.J. Sakurai                  |                                 |