

# CHARMED BARYONS ( $C = +1$ )

$$\Lambda_c^+ = udc, \quad \Sigma_c^{++} = uuc, \quad \Sigma_c^+ = udc, \quad \Sigma_c^0 = ddc,$$

$$\Xi_c^+ = usc, \quad \Xi_c^0 = dsc, \quad \Omega_c^0 = ssc$$

$\Lambda_c^+$

$$I(J^P) = 0(\frac{1}{2}^+)$$

$J$  is not well measured;  $\frac{1}{2}$  is the quark-model prediction.

$$\text{Mass } m = 2286.46 \pm 0.14 \text{ MeV}$$

$$\text{Mean life } \tau = (200 \pm 6) \times 10^{-15} \text{ s} \quad (S = 1.6)$$

$$c\tau = 59.9 \text{ } \mu\text{m}$$

### Decay asymmetry parameters

$$\Lambda\pi^+ \quad \alpha = -0.91 \pm 0.15$$

$$\Sigma^+\pi^0 \quad \alpha = -0.45 \pm 0.32$$

$$\Lambda\ell^+\nu_\ell \quad \alpha = -0.86 \pm 0.04$$

$$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha}) \text{ in } \Lambda_c^+ \rightarrow \Lambda\pi^+, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda}\pi^- = -0.07 \pm 0.31$$

$$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha}) \text{ in } \Lambda_c^+ \rightarrow \Lambda e^+\nu_e, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda}e^-\bar{\nu}_e = 0.00 \pm 0.04$$

### $\Lambda_c^+$ DECAY MODES

Scale factor/  $\rho$   
Confidence level (MeV/c)

#### Hadronic modes with a $p$ : $S = -1$ final states

$p\bar{K}^0$	( 3.21 ± 0.30 ) %	873
$pK^-\pi^+$	( 6.84 <sup>+</sup> <sub>-</sub> 0.32 / 0.40 ) %	823
$p\bar{K}^*(892)^0$	[a] ( 2.13 ± 0.30 ) %	685
$\Delta(1232)^{++}K^-$	( 1.18 ± 0.27 ) %	710
$\Lambda(1520)\pi^+$	[a] ( 2.4 ± 0.6 ) %	627
$pK^-\pi^+$ nonresonant	( 3.8 ± 0.4 ) %	823
$p\bar{K}^0\pi^0$	( 4.5 ± 0.6 ) %	823
$p\bar{K}^0\eta$	( 1.7 ± 0.4 ) %	568
$p\bar{K}^0\pi^+\pi^-$	( 3.5 ± 0.4 ) %	754
$pK^-\pi^+\pi^0$	( 4.6 ± 0.8 ) %	759
$pK^*(892)^-\pi^+$	[a] ( 1.5 ± 0.5 ) %	580
$p(K^-\pi^+)_{\text{nonresonant}}\pi^0$	( 5.0 ± 0.9 ) %	759
$\Delta(1232)\bar{K}^*(892)$	seen	419
$pK^-\pi^+\pi^+\pi^-$	( 1.5 ± 1.0 ) × 10 <sup>-3</sup>	671
$pK^-\pi^+\pi^0\pi^0$	( 1.1 ± 0.5 ) %	678

**Hadronic modes with a  $p$ :  $S = 0$  final states**

$p\pi^+\pi^-$	$(4.7 \pm 2.5) \times 10^{-3}$	927
$pf_0(980)$	[a] $(3.8 \pm 2.5) \times 10^{-3}$	614
$p\pi^+\pi^+\pi^-\pi^-$	$(2.5 \pm 1.6) \times 10^{-3}$	852
$pK^+K^-$	$(1.1 \pm 0.4) \times 10^{-3}$	616
$p\phi$	[a] $(1.12 \pm 0.23) \times 10^{-3}$	590
$pK^+K^- \text{ non-}\phi$	$(4.8 \pm 1.9) \times 10^{-4}$	616

**Hadronic modes with a hyperon:  $S = -1$  final states**

$\Lambda\pi^+$	$(1.46 \pm 0.13) \%$	864
$\Lambda\pi^+\pi^0$	$(5.0 \pm 1.3) \%$	844
$\Lambda\rho^+$	$< 6 \%$	CL=95% 636
$\Lambda\pi^+\pi^+\pi^-$	$(3.59 \pm 0.28) \%$	807
$\Sigma(1385)^+\pi^+\pi^-, \Sigma^{*+} \rightarrow$	$(1.0 \pm 0.5) \%$	688
$\Lambda\pi^+$		
$\Sigma(1385)^-\pi^+\pi^+, \Sigma^{*-} \rightarrow$	$(7.5 \pm 1.4) \times 10^{-3}$	688
$\Lambda\pi^0$		
$\Lambda\pi^+\rho^0$	$(1.4 \pm 0.6) \%$	524
$\Sigma(1385)^+\rho^0, \Sigma^{*+} \rightarrow \Lambda\pi^+$	$(5 \pm 4) \times 10^{-3}$	363
$\Lambda\pi^+\pi^+\pi^- \text{ nonresonant}$	$< 1.1 \%$	CL=90% 807
$\Lambda\pi^+\pi^+\pi^-\pi^0 \text{ total}$	$(2.5 \pm 0.9) \%$	757
$\Lambda\pi^+\eta$	[a] $(2.4 \pm 0.5) \%$	691
$\Sigma(1385)^+\eta$	[a] $(1.16 \pm 0.35) \%$	570
$\Lambda\pi^+\omega$	[a] $(1.6 \pm 0.6) \%$	517
$\Lambda\pi^+\pi^+\pi^-\pi^0, \text{ no } \eta \text{ or } \omega$	$< 9 \times 10^{-3}$	CL=90% 757
$\Lambda K^+\bar{K}^0$	$(6.4 \pm 1.3) \times 10^{-3}$	S=1.6 443
$\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Lambda\bar{K}^0$	$(1.8 \pm 0.6) \times 10^{-3}$	286
$\Sigma^0\pi^+$	$(1.43 \pm 0.14) \%$	825
$\Sigma^+\pi^0$	$(1.37 \pm 0.30) \%$	827
$\Sigma^+\eta$	$(7.5 \pm 2.5) \times 10^{-3}$	713
$\Sigma^+\pi^+\pi^-$	$(4.9 \pm 0.5) \%$	804
$\Sigma^+\rho^0$	$< 1.8 \%$	CL=95% 575
$\Sigma^-\pi^+\pi^+$	$(2.3 \pm 0.4) \%$	799
$\Sigma^0\pi^+\pi^0$	$(2.5 \pm 0.9) \%$	803
$\Sigma^0\pi^+\pi^+\pi^-$	$(1.13 \pm 0.31) \%$	763
$\Sigma^+\pi^+\pi^-\pi^0$	—	767
$\Sigma^+\omega$	[a] $(3.7 \pm 1.0) \%$	569
$\Sigma^+K^+K^-$	$(3.8 \pm 0.6) \times 10^{-3}$	349
$\Sigma^+\phi$	[a] $(4.3 \pm 0.7) \times 10^{-3}$	295
$\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow$	$(1.11 \pm 0.29) \times 10^{-3}$	286
$\Sigma^+K^-$		
$\Sigma^+K^+K^- \text{ nonresonant}$	$< 9 \times 10^{-4}$	CL=90% 349
$\Xi^0 K^+$	$(5.3 \pm 1.3) \times 10^{-3}$	653
$\Xi^- K^+\pi^+$	$(7.0 \pm 0.8) \times 10^{-3}$	S=1.1 565
$\Xi(1530)^0 K^+$	[a] $(3.5 \pm 1.0) \times 10^{-3}$	473

**Hadronic modes with a hyperon:  $S = 0$  final states**

$\Lambda K^+$	$( 6.9 \pm 1.4 ) \times 10^{-4}$		781
$\Lambda K^+ \pi^+ \pi^-$	$< 6 \times 10^{-4}$	CL=90%	637
$\Sigma^0 K^+$	$( 5.7 \pm 1.0 ) \times 10^{-4}$		735
$\Sigma^0 K^+ \pi^+ \pi^-$	$< 2.9 \times 10^{-4}$	CL=90%	574
$\Sigma^+ K^+ \pi^-$	$( 2.3 \pm 0.7 ) \times 10^{-3}$		670
$\Sigma^+ K^*(892)^0$	[a] $( 3.8 \pm 1.2 ) \times 10^{-3}$		470
$\Sigma^- K^+ \pi^+$	$< 1.3 \times 10^{-3}$	CL=90%	664

**Doubly Cabibbo-suppressed modes**

$p K^+ \pi^-$	$< 3.1 \times 10^{-4}$	CL=90%	823
---------------	------------------------	--------	-----

**Semileptonic modes**

$\Lambda \ell^+ \nu_\ell$	[b] $( 2.8 \pm 0.4 ) \%$	871
$\Lambda e^+ \nu_e$	$( 2.9 \pm 0.5 ) \%$	871
$\Lambda \mu^+ \nu_\mu$	$( 2.7 \pm 0.6 ) \%$	867

**Inclusive modes**

$e^+$ anything	$( 4.5 \pm 1.7 ) \%$	—
$p e^+$ anything	$( 1.8 \pm 0.9 ) \%$	—
$p$ anything	$( 50 \pm 16 ) \%$	—
$p$ anything (no $\Lambda$ )	$( 12 \pm 19 ) \%$	—
$n$ anything	$( 50 \pm 16 ) \%$	—
$n$ anything (no $\Lambda$ )	$( 29 \pm 17 ) \%$	—
$\Lambda$ anything	$( 35 \pm 11 ) \%$	S=1.4 —
$\Sigma^\pm$ anything	[c] $( 10 \pm 5 ) \%$	—
3prongs	$( 24 \pm 8 ) \%$	—

**$\Delta C = 1$  weak neutral current ( $CI$ ) modes, or  
Lepton Family number ( $LF$ ), or Lepton number ( $L$ ), or  
Baryon number ( $B$ ) violating modes**

$p e^+ e^-$	$CI$	$< 5.5 \times 10^{-6}$	CL=90%	951
$p \mu^+ \mu^-$	$CI$	$< 4.4 \times 10^{-5}$	CL=90%	937
$p e^+ \mu^-$	$LF$	$< 9.9 \times 10^{-6}$	CL=90%	947
$p e^- \mu^+$	$LF$	$< 1.9 \times 10^{-5}$	CL=90%	947
$\bar{p} 2e^+$	$L, B$	$< 2.7 \times 10^{-6}$	CL=90%	951
$\bar{p} 2\mu^+$	$L, B$	$< 9.4 \times 10^{-6}$	CL=90%	937
$\bar{p} e^+ \mu^+$	$L, B$	$< 1.6 \times 10^{-5}$	CL=90%	947
$\Sigma^- \mu^+ \mu^+$	$L$	$< 7.0 \times 10^{-4}$	CL=90%	812

**$\Lambda_c(2595)^+$** 

$$I(J^P) = 0(\frac{1}{2}^-)$$

The spin-parity follows from the fact that  $\Sigma_c(2455)\pi$  decays, with little available phase space, are dominant. This assumes that  $J^P = 1/2^+$  for the  $\Sigma_c(2455)$ .

$$\text{Mass } m = 2592.25 \pm 0.28 \text{ MeV}$$

$$m - m_{\Lambda_c^+} = 305.79 \pm 0.24 \text{ MeV}$$

$$\text{Full width } \Gamma = 2.6 \pm 0.6 \text{ MeV}$$

$\Lambda_c^+ \pi \pi$  and its submode  $\Sigma_c(2455)\pi$  — the latter just barely — are the only strong decays allowed to an excited  $\Lambda_c^+$  having this mass; and the submode seems to dominate.

<b><math>\Lambda_c(2595)^+</math> DECAY MODES</b>	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	[d] $\approx 67\%$	117
$\Sigma_c(2455)^{++} \pi^-$	$24 \pm 7\%$	†
$\Sigma_c(2455)^0 \pi^+$	$24 \pm 7\%$	†
$\Lambda_c^+ \pi^+ \pi^-$ 3-body	$18 \pm 10\%$	117
$\Lambda_c^+ \pi^0$	[e] not seen	258
$\Lambda_c^+ \gamma$	not seen	288

 **$\Lambda_c(2625)^+$** 

$$I(J^P) = 0(\frac{3}{2}^-)$$

$J^P$  has not been measured;  $\frac{3}{2}^-$  is the quark-model prediction.

$$\text{Mass } m = 2628.11 \pm 0.19 \text{ MeV} \quad (S = 1.1)$$

$$m - m_{\Lambda_c^+} = 341.65 \pm 0.13 \text{ MeV} \quad (S = 1.1)$$

$$\text{Full width } \Gamma < 0.97 \text{ MeV, CL} = 90\%$$

$\Lambda_c^+ \pi \pi$  and its submode  $\Sigma(2455)\pi$  are the only strong decays allowed to an excited  $\Lambda_c^+$  having this mass.

<b><math>\Lambda_c(2625)^+</math> DECAY MODES</b>	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level	$p$ (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	[d] $\approx 67\%$		184
$\Sigma_c(2455)^{++} \pi^-$	$< 5$	90%	102
$\Sigma_c(2455)^0 \pi^+$	$< 5$	90%	102
$\Lambda_c^+ \pi^+ \pi^-$ 3-body	large		184
$\Lambda_c^+ \pi^0$	[e] not seen		293
$\Lambda_c^+ \gamma$	not seen		319

**$\Lambda_c(2880)^+$**

$I(J^P) = 0(\frac{5}{2}^+)$

There is some good evidence that indeed  $J^P = 5/2^+$

Mass  $m = 2881.53 \pm 0.35$  MeV

$m - m_{\Lambda_c^+} = 595.1 \pm 0.4$  MeV

Full width  $\Gamma = 5.8 \pm 1.1$  MeV

$\Lambda_c(2880)^+$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	seen	471
$\Sigma_c(2455)^{0,++} \pi^\pm$	seen	376
$\Sigma_c(2520)^{0,++} \pi^\pm$	seen	317
$pD^0$	seen	316

**$\Lambda_c(2940)^+$**

$I(J^P) = 0(?^?)$

Mass  $m = 2939.3^{+1.4}_{-1.5}$  MeV

Full width  $\Gamma = 17^{+8}_{-6}$  MeV

$\Lambda_c(2940)^+$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$pD^0$	seen	420
$\Sigma_c(2455)^{0,++} \pi^\pm$	seen	-

**$\Sigma_c(2455)$**

$I(J^P) = 1(\frac{1}{2}^+)$

$\Sigma_c(2455)^{++}$  mass  $m = 2453.97 \pm 0.14$  MeV

$\Sigma_c(2455)^+$  mass  $m = 2452.9 \pm 0.4$  MeV

$\Sigma_c(2455)^0$  mass  $m = 2453.75 \pm 0.14$  MeV

$m_{\Sigma_c^{++}} - m_{\Lambda_c^+} = 167.510 \pm 0.017$  MeV

$m_{\Sigma_c^+} - m_{\Lambda_c^+} = 166.4 \pm 0.4$  MeV

$m_{\Sigma_c^0} - m_{\Lambda_c^+} = 167.290 \pm 0.017$  MeV

$m_{\Sigma_c^{++}} - m_{\Sigma_c^0} = 0.220 \pm 0.013$  MeV

$m_{\Sigma_c^+} - m_{\Sigma_c^0} = -0.9 \pm 0.4$  MeV

$\Sigma_c(2455)^{++}$  full width  $\Gamma = 1.89^{+0.09}_{-0.18}$  MeV (S = 1.1)

$\Sigma_c(2455)^+$  full width  $\Gamma < 4.6$  MeV, CL = 90%

$\Sigma_c(2455)^0$  full width  $\Gamma = 1.83^{+0.11}_{-0.19}$  MeV (S = 1.2)

$\Lambda_c^+ \pi$  is the only strong decay allowed to a  $\Sigma_c$  having this mass.

$\Sigma_c(2455)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \pi$	$\approx 100\%$	94

### $\Sigma_c(2520)$

$$I(J^P) = 1(\frac{3}{2}^+)$$

$J^P$  has not been measured;  $\frac{3}{2}^+$  is the quark-model prediction.

$$\Sigma_c(2520)^{++} \text{ mass } m = 2518.41_{-0.19}^{+0.21} \text{ MeV} \quad (S = 1.1)$$

$$\Sigma_c(2520)^+ \text{ mass } m = 2517.5 \pm 2.3 \text{ MeV}$$

$$\Sigma_c(2520)^0 \text{ mass } m = 2518.48 \pm 0.20 \text{ MeV} \quad (S = 1.1)$$

$$m_{\Sigma_c(2520)^{++}} - m_{\Lambda_c^+} = 231.95_{-0.12}^{+0.17} \text{ MeV} \quad (S = 1.3)$$

$$m_{\Sigma_c(2520)^+} - m_{\Lambda_c^+} = 231.0 \pm 2.3 \text{ MeV}$$

$$m_{\Sigma_c(2520)^0} - m_{\Lambda_c^+} = 232.02_{-0.14}^{+0.15} \text{ MeV} \quad (S = 1.3)$$

$$m_{\Sigma_c(2520)^{++}} - m_{\Sigma_c(2520)^0} = 0.01 \pm 0.15 \text{ MeV}$$

$$\Sigma_c(2520)^{++} \text{ full width } \Gamma = 14.78_{-0.40}^{+0.30} \text{ MeV}$$

$$\Sigma_c(2520)^+ \text{ full width } \Gamma < 17 \text{ MeV, CL} = 90\%$$

$$\Sigma_c(2520)^0 \text{ full width } \Gamma = 15.3_{-0.5}^{+0.4} \text{ MeV}$$

$\Lambda_c^+ \pi$  is the only strong decay allowed to a  $\Sigma_c$  having this mass.

$\Sigma_c(2520)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \pi$	$\approx 100\%$	179

### $\Sigma_c(2800)$

$$I(J^P) = 1(?^?)$$

$$\Sigma_c(2800)^{++} \text{ mass } m = 2801_{-6}^{+4} \text{ MeV}$$

$$\Sigma_c(2800)^+ \text{ mass } m = 2792_{-5}^{+14} \text{ MeV}$$

$$\Sigma_c(2800)^0 \text{ mass } m = 2806_{-7}^{+5} \text{ MeV} \quad (S = 1.3)$$

$$m_{\Sigma_c(2800)^{++}} - m_{\Lambda_c^+} = 514_{-6}^{+4} \text{ MeV}$$

$$m_{\Sigma_c(2800)^+} - m_{\Lambda_c^+} = 505_{-5}^{+14} \text{ MeV}$$


$$m_{\Sigma_c(2800)^0} - m_{\Lambda_c^+} = 519_{-7}^{+5} \text{ MeV} \quad (S = 1.3)$$

$$\Sigma_c(2800)^{++} \text{ full width } \Gamma = 75_{-17}^{+22} \text{ MeV}$$

$$\Sigma_c(2800)^+ \text{ full width } \Gamma = 62_{-40}^{+60} \text{ MeV}$$

$$\Sigma_c(2800)^0 \text{ full width } \Gamma = 72_{-15}^{+22} \text{ MeV}$$

$\Sigma_c(2800)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \pi$	seen	443



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

$$\text{Mass } m = 2467.93^{+0.28}_{-0.40} \text{ MeV}$$

$$\text{Mean life } \tau = (442 \pm 26) \times 10^{-15} \text{ s} \quad (S = 1.3)$$

$$c\tau = 132 \mu\text{m}$$

$\Xi_c^+$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level	$p$ (MeV/c)
-----------------------	--------------------------------	------------------	-------------

**No absolute branching fractions have been measured.**  
**The following are branching *ratios* relative to  $\Xi^- 2\pi^+$ .**

**Cabibbo-favored ( $S = -2$ ) decays — relative to  $\Xi^- 2\pi^+$**

$p 2K_S^0$	$0.087 \pm 0.021$		767
$\Lambda \bar{K}^0 \pi^+$	—		852
$\Sigma(1385)^+ \bar{K}^0$	[a] 1.0 $\pm 0.5$		746
$\Lambda K^- 2\pi^+$	$0.323 \pm 0.033$		787
$\Lambda \bar{K}^*(892)^0 \pi^+$	[a] $< 0.16$	90%	608
$\Sigma(1385)^+ K^- \pi^+$	[a] $< 0.23$	90%	678
$\Sigma^+ K^- \pi^+$	$0.94 \pm 0.10$		811
$\Sigma^+ \bar{K}^*(892)^0$	[a] $0.81 \pm 0.15$		658
$\Sigma^0 K^- 2\pi^+$	$0.27 \pm 0.12$		735
$\Xi^0 \pi^+$	$0.55 \pm 0.16$		877
$\Xi^- 2\pi^+$	<b>DEFINED AS 1</b>		851
$\Xi(1530)^0 \pi^+$	[a] $< 0.10$	90%	750
$\Xi^0 \pi^+ \pi^0$	$2.3 \pm 0.7$		856
$\Xi^0 \pi^- 2\pi^+$	$1.7 \pm 0.5$		818
$\Xi^0 e^+ \nu_e$	$2.3 \begin{smallmatrix} +0.7 \\ -0.8 \end{smallmatrix}$		884
$\Omega^- K^+ \pi^+$	$0.07 \pm 0.04$		399

**Cabibbo-suppressed decays — relative to  $\Xi^- 2\pi^+$**

$p K^- \pi^+$	$0.21 \pm 0.04$		944
$p \bar{K}^*(892)^0$	[a] $0.116 \pm 0.030$		828
$\Sigma^+ \pi^+ \pi^-$	$0.48 \pm 0.20$		922

$\Sigma^- 2\pi^+$	$0.18 \pm 0.09$		918
$\Sigma^+ K^+ K^-$	$0.15 \pm 0.06$		580
$\Sigma^+ \phi$	[a] $< 0.11$	90%	549
$\Xi(1690)^0 K^+, \Xi^0 \rightarrow \Sigma^+ K^-$	$< 0.05$	90%	501



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

$$\text{Mass } m = 2470.85^{+0.28}_{-0.40} \text{ MeV}$$

$$m_{\Xi_c^0} - m_{\Xi_c^+} = 2.93 \pm 0.24 \text{ MeV}$$

$$\text{Mean life } \tau = (112^{+13}_{-10}) \times 10^{-15} \text{ s}$$

$$c\tau = 33.6 \text{ } \mu\text{m}$$

### Decay asymmetry parameters

$$\Xi^- \pi^+ \quad \alpha = -0.6 \pm 0.4$$

No absolute branching fractions have been measured. Several measurements of ratios of fractions may be found in the Listings that follow.

$\Xi_c^0$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
-----------------------	--------------------------------	-------------

**No absolute branching fractions have been measured.  
The following are branching *ratios* relative to  $\Xi^- \pi^+$ .**

### Cabibbo-favored ( $S = -2$ ) decays — relative to $\Xi^- \pi^+$

$p K^- K^- \pi^+$	$0.34 \pm 0.04$	676
$p K^- \bar{K}^*(892)^0$	$0.21 \pm 0.05$	413
$p K^- K^- \pi^+$ (no $\bar{K}^{*0}$ )	$0.21 \pm 0.04$	676
$\Lambda K_S^0$	$0.210 \pm 0.028$	906
$\Lambda K^- \pi^+$	$1.07 \pm 0.14$	856
$\Lambda \bar{K}^0 \pi^+ \pi^-$	seen	787
$\Lambda K^- \pi^+ \pi^+ \pi^-$	seen	703
$\Xi^- \pi^+$	<b>DEFINED AS 1</b>	875
$\Xi^- \pi^+ \pi^+ \pi^-$	$3.3 \pm 1.4$	816
$\Omega^- K^+$	$0.297 \pm 0.024$	522
$\Xi^- e^+ \nu_e$	$3.1 \pm 1.1$	882
$\Xi^- \ell^+$ anything	$1.0 \pm 0.5$	—



**Cabibbo-suppressed decays — relative to  $\Xi^- \pi^+$**

$\Xi^- K^+$	$0.028 \pm 0.006$	790
$\Lambda K^+ K^-$ (no $\phi$ )	$0.029 \pm 0.007$	648
$\Lambda \phi$	$0.034 \pm 0.007$	621

$$\Xi_c^{'+}$$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2575.7 \pm 3.0$  MeV

$$m_{\Xi_c^{'+}} - m_{\Xi_c^+} = 107.8 \pm 3.0 \text{ MeV}$$

The  $\Xi_c^{'+} - \Xi_c^+$  mass difference is too small for any strong decay to occur.

$\Xi_c^{'+}$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Xi_c^+ \gamma$	seen	106

$$\Xi_c^{'0}$$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2577.9 \pm 2.9$  MeV

$$m_{\Xi_c^{'0}} - m_{\Xi_c^0} = 107.0 \pm 2.9 \text{ MeV}$$

The  $\Xi_c^{'0} - \Xi_c^0$  mass difference is too small for any strong decay to occur.

$\Xi_c^{'0}$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Xi_c^0 \gamma$	seen	105

$$\Xi_c(2645)$$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

$J^P$  has not been measured;  $\frac{3}{2}^+$  is the quark-model prediction.

$\Xi_c(2645)^+$  mass  $m = 2645.9 \pm 0.5$  MeV ( $S = 1.1$ )

$\Xi_c(2645)^0$  mass  $m = 2645.9 \pm 0.5$  MeV

$$m_{\Xi_c(2645)^+} - m_{\Xi_c^0} = 175.0 \pm 0.6 \text{ MeV} \quad (S = 1.1)$$

$$m_{\Xi_c(2645)^0} - m_{\Xi_c^+} = 178.0 \pm 0.6 \text{ MeV}$$

$$m_{\Xi_c(2645)^+} - m_{\Xi_c(2645)^0} = 0.0 \pm 0.5 \text{ MeV}$$

$\Xi_c(2645)^+$  full width  $\Gamma = 2.6 \pm 0.4$  MeV

$\Xi_c(2645)^0$  full width  $\Gamma < 5.5$  MeV, CL = 90%

$\Xi_c \pi$  is the only strong decay allowed to a  $\Xi_c$  resonance having this mass.

$\Xi_c(2645)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Xi_c^0 \pi^+$	seen	102
$\Xi_c^+ \pi^-$	seen	107

### $\Xi_c(2790)$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

$J^P$  has not been measured;  $\frac{1}{2}^-$  is the quark-model prediction.

$$\begin{aligned} \Xi_c(2790)^+ \text{ mass} &= 2789.1 \pm 3.2 \text{ MeV} \\ \Xi_c(2790)^0 \text{ mass} &= 2791.9 \pm 3.3 \text{ MeV} \\ m_{\Xi_c(2790)^+} - m_{\Xi_c^0} &= 318.2 \pm 3.2 \text{ MeV} \\ m_{\Xi_c(2790)^0} - m_{\Xi_c^+} &= 324.0 \pm 3.3 \text{ MeV} \\ \Xi_c(2790)^+ \text{ width} &< 15 \text{ MeV, CL} = 90\% \\ \Xi_c(2790)^0 \text{ width} &< 12 \text{ MeV, CL} = 90\% \end{aligned}$$

$\Xi_c(2790)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Xi_c^+ \pi^-$	seen	159

### $\Xi_c(2815)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

$J^P$  has not been measured;  $\frac{3}{2}^-$  is the quark-model prediction.

$$\begin{aligned} \Xi_c(2815)^+ \text{ mass } m &= 2816.6 \pm 0.9 \text{ MeV} \\ \Xi_c(2815)^0 \text{ mass } m &= 2819.6 \pm 1.2 \text{ MeV} \\ m_{\Xi_c(2815)^+} - m_{\Xi_c^+} &= 348.7 \pm 0.9 \text{ MeV} \\ m_{\Xi_c(2815)^0} - m_{\Xi_c^0} &= 348.8 \pm 1.2 \text{ MeV} \\ m_{\Xi_c(2815)^+} - m_{\Xi_c(2815)^0} &= -3.0 \pm 1.3 \text{ MeV} \\ \Xi_c(2815)^+ \text{ full width } \Gamma &< 3.5 \text{ MeV, CL} = 90\% \\ \Xi_c(2815)^0 \text{ full width } \Gamma &< 6.5 \text{ MeV, CL} = 90\% \end{aligned}$$

The  $\Xi_c \pi \pi$  modes are consistent with being entirely via  $\Xi_c(2645) \pi$ .

$\Xi_c(2815)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Xi_c^+ \pi^+ \pi^-$	seen	196
$\Xi_c^0 \pi^+ \pi^-$	seen	191

**$\Xi_c(2980)$** 

$$I(J^P) = \frac{1}{2}(??)$$

$$\begin{aligned} \Xi_c(2980)^+ m &= 2970.7 \pm 2.2 \text{ MeV} \quad (S = 1.5) \\ \Xi_c(2980)^0 m &= 2968.0 \pm 2.6 \text{ MeV} \quad (S = 1.2) \\ \Xi_c(2980)^+ \text{ width } \Gamma &= 17.9 \pm 3.5 \text{ MeV} \\ \Xi_c(2980)^0 \text{ width } \Gamma &= 20 \pm 7 \text{ MeV} \quad (S = 1.3) \end{aligned}$$

$\Xi_c(2980)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \bar{K} \pi$	seen	231
$\Sigma_c(2455) \bar{K}$	seen	134
$\Lambda_c^+ \bar{K}$	not seen	414
$\Xi_c 2\pi$	seen	385
$\Xi_c(2645) \pi$	seen	277

 **$\Xi_c(3055)$** 

$$I(J^P) = ?(??)$$

$$\begin{aligned} \text{Mass } m &= 3055.1 \pm 1.7 \text{ MeV} \quad (S = 1.5) \\ \text{Full width } \Gamma &= 11 \pm 4 \text{ MeV} \end{aligned}$$

 **$\Xi_c(3080)$** 

$$I(J^P) = \frac{1}{2}(??)$$

$$\begin{aligned} \Xi_c(3080)^+ m &= 3076.94 \pm 0.28 \text{ MeV} \\ \Xi_c(3080)^0 m &= 3079.9 \pm 1.4 \text{ MeV} \quad (S = 1.3) \\ \Xi_c(3080)^+ \text{ width } \Gamma &= 4.3 \pm 1.5 \text{ MeV} \quad (S = 1.3) \\ \Xi_c(3080)^0 \text{ width } \Gamma &= 5.6 \pm 2.2 \text{ MeV} \end{aligned}$$

$\Xi_c(3080)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \bar{K} \pi$	seen	415
$\Sigma_c(2455) \bar{K}$	seen	342
$\Sigma_c(2455) \bar{K} + \Sigma_c(2520) \bar{K}$	seen	–
$\Lambda_c^+ \bar{K}$	not seen	536
$\Lambda_c^+ \bar{K} \pi^+ \pi^-$	not seen	143

 **$\Omega_c^0$** 

$$I(J^P) = 0(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

$$\begin{aligned} \text{Mass } m &= 2695.2 \pm 1.7 \text{ MeV} \quad (S = 1.3) \\ \text{Mean life } \tau &= (69 \pm 12) \times 10^{-15} \text{ s} \\ c\tau &= 21 \text{ } \mu\text{m} \end{aligned}$$

No absolute branching fractions have been measured.

$\Omega_c^0$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Sigma^+ K^- K^- \pi^+$	seen	689
$\Xi^0 K^- \pi^+$	seen	901
$\Xi^- K^- \pi^+ \pi^+$	seen	830
$\Omega^- e^+ \nu_e$	seen	829
$\Omega^- \pi^+$	seen	821
$\Omega^- \pi^+ \pi^0$	seen	797
$\Omega^- \pi^- \pi^+ \pi^+$	seen	753

**$\Omega_c(2770)^0$**

$$I(J^P) = 0(\frac{3}{2}^+)$$

$J^P$  has not been measured;  $\frac{3}{2}^+$  is the quark-model prediction.

$$\text{Mass } m = 2765.9 \pm 2.0 \text{ MeV} \quad (S = 1.2)$$

$$m_{\Omega_c(2770)^0} - m_{\Omega_c^0} = 70.7^{+0.8}_{-0.9} \text{ MeV}$$

The  $\Omega_c(2770)^0 - \Omega_c^0$  mass difference is too small for any strong decay to occur.

$\Omega_c(2770)^0$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Omega_c^0 \gamma$	presumably 100%	70

### NOTES

- [a] This branching fraction includes all the decay modes of the final-state resonance.
- [b] An  $\ell$  indicates an  $e$  or a  $\mu$  mode, not a sum over these modes.
- [c] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [d] Assuming isospin conservation, so that the other third is  $\Lambda_c^+ \pi^0 \pi^0$ .
- [e] A test that the isospin is indeed 0, so that the particle is indeed a  $\Lambda_c^+$ .