

X(4100)[±]

$$I^G(J^{PC}) = 1^-(?^{??})$$

OMITTED FROM SUMMARY TABLE

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

Reported by AAIJ 18AN in the $\eta_c(1S)\pi^-$ invariant mass distribution in $B^0 \rightarrow \eta_c(1S)K^+\pi^-$ decays with a significance of 3.4σ . $J^P = 0^+$ or 1^- assignment consistent with data.

X(4100)[±] MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4096 ± 20⁺¹⁸₋₂₂	AAIJ	18AN LHCb	$B^0 \rightarrow \eta_c(1S)K^+\pi^-$

X(4100)[±] WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
152 ± 58⁺⁶⁰₋₃₅	AAIJ	18AN LHCb	$B^0 \rightarrow \eta_c(1S)K^+\pi^-$

X(4100)[±] DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta_c(1S)\pi^-$	seen
Γ_2 $\pi^\pm\psi(3770)$	not seen

X(4100)[±] BRANCHING RATIOS

$\Gamma(\eta_c(1S)\pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	18AN LHCb	$B^0 \rightarrow \eta_c(1S)K^+\pi^-$
¹ AAIJ 18AN quotes a fit fraction for $B^0 \rightarrow X(4100)^- K^+ \rightarrow \eta_c(1S)\pi^- K^+$ of $(3.3 \pm 1.1^{+1.2}_{-1.1})\%$ from an amplitude analysis.			

$\Gamma(\pi^\pm\psi(3770))/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ ABLIKIM	19AR BES3	$e^+e^- \rightarrow \pi^+\pi^- D\bar{D}$
¹ From a measurement of $\sigma(e^+e^- \rightarrow \pi^+\pi^- D\bar{D})$ between $\sqrt{s} = 4.08$ and 4.6 GeV.			

X(4100)[±] REFERENCES

ABLIKIM	19AR PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	18AN EPJ C78 1019	R. Aaij <i>et al.</i>	(LHCb Collab.)