

CHARMED MESONS ($C = \pm 1$)

$$D^+ = c\bar{d}, D^0 = c\bar{u}, \bar{D}^0 = \bar{c}u, D^- = \bar{c}d, \quad \text{similarly for } D^{*'}\text{'s}$$

D^\pm

$$I(J^P) = \frac{1}{2}(0^-)$$

$$\text{Mass } m = 1869.66 \pm 0.05 \text{ MeV}$$

$$\text{Mean life } \tau = (1033 \pm 5) \times 10^{-15} \text{ s}$$

$$c\tau = 309.8 \text{ } \mu\text{m}$$

c-quark decays

$$\Gamma(c \rightarrow \ell^+ \text{ anything}) / \Gamma(c \rightarrow \text{ anything}) = 0.096 \pm 0.004 \text{ [a]}$$

$$\Gamma(c \rightarrow D^*(2010)^+ \text{ anything}) / \Gamma(c \rightarrow \text{ anything}) = 0.255 \pm 0.017$$

CP-violation decay-rate asymmetries

$$A_{CP}(\mu^\pm \nu) = (8 \pm 8)\%$$

$$A_{CP}(K_L^0 e^\pm \nu) = (-0.6 \pm 1.6)\%$$

$$A_{CP}(K_S^0 \pi^\pm) = (-0.41 \pm 0.09)\%$$

$$A_{CP}(K_L^0 K^\pm) \text{ in } D^\pm \rightarrow K_L^0 K^\pm = (-4.2 \pm 3.4) \times 10^{-2}$$

$$A_{CP}(K^\mp 2\pi^\pm) = (-0.18 \pm 0.16)\%$$

$$A_{CP}(K^\mp \pi^\pm \pi^\pm \pi^0) = (-0.3 \pm 0.7)\%$$

$$A_{CP}(K_S^0 \pi^\pm \pi^0) = (-0.1 \pm 0.7)\%$$

$$A_{CP}(K_S^0 \pi^\pm \eta) \text{ in } D^\pm \rightarrow K_S^0 \pi^\pm \eta = (-0.9 \pm 3.1) \times 10^{-2}$$

$$A_{CP}(K_S^0 \pi^\pm \pi^+ \pi^-) = (0.0 \pm 1.2)\%$$

$$A_{CP}(K^\pm \pi^+ \pi^- \pi^0) \text{ in } D^\pm \rightarrow K^\pm \pi^+ \pi^- \pi^0 = -0.04 \pm 0.06$$

$$A_{CP}(\pi^\pm \pi^0) = (0.4 \pm 1.3)\% \quad (S = 1.7)$$

$$A_{CP}(\pi^\pm \eta) = (0.3 \pm 0.8)\% \quad (S = 1.2)$$

$$A_{CP}(\pi^\pm \pi^0 \eta) \text{ in } D^\pm \rightarrow \pi^\pm \pi^0 \eta = (-6 \pm 7) \times 10^{-2}$$

$$A_{CP}(\pi^\pm \eta \eta) \text{ in } D^\pm \rightarrow \pi^\pm \eta \eta = (8 \pm 9) \times 10^{-2}$$

$$A_{CP}(\pi^\pm \eta'(958)) = (-0.6 \pm 0.7)\%$$

$$A_{CP}(\bar{K}^0 / K^0 K^\pm) = (0.11 \pm 0.17)\%$$

$$A_{CP}(K_S^0 K^\pm) = (-0.01 \pm 0.07)\%$$

$$A_{CP}(K_S^0 K^\pm \pi^0) \text{ in } D^\pm \rightarrow K_S^0 K^\pm \pi^0 = (1 \pm 4) \times 10^{-2}$$

$$A_{CP}(K_L^0 K^\pm \pi^0) \text{ in } D^\pm \rightarrow K_L^0 K^\pm \pi^0 = (-1 \pm 4) \times 10^{-2}$$

$$A_{CP}(K^+ K^- \pi^\pm) = (0.37 \pm 0.29)\%$$

$$A_{CP}(K^\pm K^{*0}) = (-0.3 \pm 0.4)\%$$

$$A_{CP}(\phi \pi^\pm) = (0.01 \pm 0.09)\% \quad (S = 1.8)$$

$$A_{CP}(K^\pm K_0^*(1430)^0) = (8_{-6}^{+7})\%$$

$$A_{CP}(K^\pm K_2^*(1430)^0) = (43_{-26}^{+20})\%$$

$$A_{CP}(K^\pm K_0^*(700)) = (-12_{-13}^{+18})\%$$

$$\begin{aligned}
 A_{CP}(a_0(1450)^0 \pi^\pm) &= (-19^{+14}_{-16})\% \\
 A_{CP}(\phi(1680) \pi^\pm) &= (-9 \pm 26)\% \\
 A_{CP}(\pi^\pm 2\pi^0) \text{ in } D^\pm \rightarrow \pi^\pm 2\pi^0 &= (5.6 \pm 2.7)\% \\
 A_{CP}(\pi^+ \pi^- \pi^\pm) &= (0.5 \pm 2.0)\% \\
 A_{CP}(2\pi^\pm \pi^\mp \pi^0) \text{ in } D^\pm \rightarrow 2\pi^\pm \pi^\mp \pi^0 &= (0.3 \pm 2.0)\% \\
 A_{CP}(2\pi^\pm \pi^\mp 2\pi^0) \text{ in } D^\pm \rightarrow 2\pi^\pm \pi^\mp 2\pi^0 &= (-4 \pm 4)\% \\
 A_{CP}(\pi^+ \pi^- \pi^\pm \eta) \text{ in } D^\pm \rightarrow \pi^+ \pi^- \pi^\pm \eta &= (3 \pm 5) \times 10^{-2} \\
 A_{CP}(K_S^0 K^\pm \pi^+ \pi^-) &= (-4 \pm 7)\% \\
 A_{CP}(K^\pm \pi^0) &= (-3 \pm 5)\% \\
 A_{CP}(K^\pm \eta) \text{ in } D^\pm \rightarrow K^\pm \eta &= (-6 \pm 11) \times 10^{-2}
 \end{aligned}$$

χ^2 tests of CP-violation (CPV)

$$\begin{aligned}
 \text{Local CPV in } D^\pm \rightarrow \pi^+ \pi^- \pi^\pm &= 78.1\% \\
 \text{Local CPV in } D^\pm \rightarrow K^+ K^- \pi^\pm &= 31\%
 \end{aligned}$$

CP violating asymmetries of P-odd (T-odd) moments

$$A_T(K_S^0 K^\pm \pi^+ \pi^-) = (-12 \pm 11) \times 10^{-3} [b]$$

D^+ form factors

$$\begin{aligned}
 f_+(0) |V_{cs}| \text{ in } \bar{K}^0 \ell^+ \nu_\ell &= 0.719 \pm 0.011 \quad (S = 1.6) \\
 r_1 \equiv a_1/a_0 \text{ in } \bar{K}^0 \ell^+ \nu_\ell &= -2.13 \pm 0.14 \\
 r_2 \equiv a_2/a_0 \text{ in } \bar{K}^0 \ell^+ \nu_\ell &= -3 \pm 12 \quad (S = 1.5) \\
 f_+(0) |V_{cd}| \text{ in } \pi^0 \ell^+ \nu_\ell &= 0.1407 \pm 0.0025 \\
 r_1 \equiv a_1/a_0 \text{ in } \pi^0 \ell^+ \nu_\ell &= -2.00 \pm 0.13 \\
 r_2 \equiv a_2/a_0 \text{ in } \pi^0 \ell^+ \nu_\ell &= -4 \pm 5 \\
 f_+(0) |V_{cd}| \text{ in } D^+ \rightarrow \eta \ell^+ \nu_\ell \ (\ell = e \text{ or } \nu) &= (8.4 \pm 0.4) \times 10^{-2} \\
 r_1 \equiv a_1/a_0 \text{ in } D^+ \rightarrow \eta e^+ \nu_e &= -5.3 \pm 2.7 \quad (S = 1.9) \\
 r_\nu \equiv V(0)/A_1(0) \text{ in } D^+ \rightarrow \omega e^+ \nu_e &= 1.24 \pm 0.11 \\
 r_2 \equiv A_2(0)/A_1(0) \text{ in } D^+ \rightarrow \omega e^+ \nu_e &= 1.06 \pm 0.16 \\
 r_\nu \equiv V(0)/A_1(0) \text{ in } D^+, D^0 \rightarrow \rho e^+ \nu_e &= 1.64 \pm 0.10 \quad (S = 1.2) \\
 r_2 \equiv A_2(0)/A_1(0) \text{ in } D^+, D^0 \rightarrow \rho e^+ \nu_e &= 0.84 \pm 0.06 \\
 r_\nu \equiv V(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 1.49 \pm 0.05 \quad (S = 2.1) \\
 r_2 \equiv A_2(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 0.802 \pm 0.021 \\
 r_3 \equiv A_3(0)/A_1(0) \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 0.0 \pm 0.4 \\
 \Gamma_L/\Gamma_T \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 1.13 \pm 0.08 \\
 \Gamma_+/\Gamma_- \text{ in } \bar{K}^*(892)^0 \ell^+ \nu_\ell &= 0.22 \pm 0.06 \quad (S = 1.6)
 \end{aligned}$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as \bar{K}^0 modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D⁺ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Inclusive modes			
e^+ semileptonic	(16.07 ± 0.30) %		—
μ^+ anything	(17.6 ± 3.2) %		—
K^- anything	(25.7 ± 1.4) %		—
\bar{K}^0 anything + K^0 anything	(61 ± 5) %		—
K^+ anything	(5.9 ± 0.8) %		—
$K^*(892)^-$ anything	(6 ± 5) %		—
$\bar{K}^*(892)^0$ anything	(23 ± 5) %		—
$K^*(892)^0$ anything	< 6.6	% CL=90%	—
η anything	(6.3 ± 0.7) %		—
η' anything	(1.04 ± 0.18) %		—
ϕ anything	(1.12 ± 0.04) %		—
Leptonic and semileptonic modes			
$e^+ \nu_e$	< 8.8	$\times 10^{-6}$ CL=90%	935
$\gamma e^+ \nu_e$	< 3.0	$\times 10^{-5}$ CL=90%	935
$\mu^+ \nu_\mu$	(3.74 ± 0.17)	$\times 10^{-4}$	932
$\tau^+ \nu_\tau$	(1.20 ± 0.27)	$\times 10^{-3}$	90
$\bar{K}^0 e^+ \nu_e$	(8.72 ± 0.09) %		869
$\bar{K}^0 \mu^+ \nu_\mu$	(8.76 ± 0.19) %		865
$K^- \pi^+ e^+ \nu_e$	(4.02 ± 0.18) %	S=3.2	864
$\bar{K}^*(892)^0 e^+ \nu_e, \bar{K}^*(892)^0 \rightarrow$ $K^- \pi^+$	(3.77 ± 0.17) %		722
$(K^- \pi^+) [0.8-1.0]\text{GeV} e^+ \nu_e$	(3.39 ± 0.09) %		864
$(K^- \pi^+)_{S\text{-wave}} e^+ \nu_e$	(2.28 ± 0.11)	$\times 10^{-3}$	—
$\bar{K}^*(1410)^0 e^+ \nu_e,$ $\bar{K}^*(1410)^0 \rightarrow K^- \pi^+$	< 6	$\times 10^{-3}$ CL=90%	—
$\bar{K}_2^*(1430)^0 e^+ \nu_e,$ $\bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$	< 5	$\times 10^{-4}$ CL=90%	—
$K^- \pi^+ e^+ \nu_e$ nonresonant	< 7	$\times 10^{-3}$ CL=90%	864
$\bar{K}^*(892)^0 e^+ \nu_e$	(5.40 ± 0.10) %	S=1.1	722
$K^- \pi^+ \mu^+ \nu_\mu$	(3.65 ± 0.34) %		851
$\bar{K}^*(892)^0 \mu^+ \nu_\mu,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(3.52 ± 0.10) %		717
$K^- \pi^+ \mu^+ \nu_\mu$ nonresonant	(1.9 ± 0.5)	$\times 10^{-3}$	851
$\bar{K}^*(892)^0 \mu^+ \nu_\mu$	(5.27 ± 0.15) %		717

$K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	< 1.5	$\times 10^{-3}$	CL=90%	825
$\bar{K}_1(1270)^0 e^+ \nu_e, \bar{K}_1^0 \rightarrow$	(1.06 ± 0.15)	$\times 10^{-3}$		–
$K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	< 2.3	$\times 10^{-4}$	CL=90%	380
$\bar{K}_0^*(1430)^0 \mu^+ \nu_\mu$	< 1.5	$\times 10^{-3}$	CL=90%	105
$\bar{K}^*(1680)^0 \mu^+ \nu_\mu$	(3.72 ± 0.17)	$\times 10^{-3}$	S=2.0	930
$\pi^0 e^+ \nu_e$	(3.50 ± 0.15)	$\times 10^{-3}$		927
$\pi^0 \mu^+ \nu_\mu$	(1.11 ± 0.07)	$\times 10^{-3}$		855
$\eta e^+ \nu_e$	(1.04 ± 0.11)	$\times 10^{-3}$		851
$\eta \mu^+ \nu_\mu$	(2.49 ± 0.11)	$\times 10^{-3}$	S=1.2	924
$\pi^- \pi^+ e^+ \nu_e$	(6.4 ± 0.6)	$\times 10^{-4}$		–
$f_0(500)^0 e^+ \nu_e, f_0(500)^0 \rightarrow$				
$\pi^+ \pi^-$				
$\rho^0 e^+ \nu_e$	(1.90 ± 0.10)	$\times 10^{-3}$	S=1.2	774
$\rho^0 \mu^+ \nu_\mu$	(2.4 ± 0.4)	$\times 10^{-3}$		770
$\omega e^+ \nu_e$	(1.69 ± 0.11)	$\times 10^{-3}$		771
$\omega \mu^+ \nu_\mu$	(1.77 ± 0.21)	$\times 10^{-3}$		767
$\eta'(958) e^+ \nu_e$	(2.0 ± 0.4)	$\times 10^{-4}$		690
$a(980)^0 e^+ \nu_e, a(980)^0 \rightarrow \eta \pi^0$	$(1.7 \begin{smallmatrix} + 0.8 \\ - 0.7 \end{smallmatrix})$	$\times 10^{-4}$		–
$b_1(1235)^0 e^+ \nu_e, b_1^0 \rightarrow \omega \pi^0$	< 1.75	$\times 10^{-4}$	CL=90%	–
$\phi e^+ \nu_e$	< 1.3	$\times 10^{-5}$	CL=90%	657
$D^0 e^+ \nu_e$	< 1.0	$\times 10^{-4}$	CL=90%	5
Hadronic modes with a \bar{K} or $\bar{K}K\bar{K}$				
$K_S^0 \pi^+$	(1.562 ± 0.031)	%	S=1.7	863
$K_L^0 \pi^+$	(1.46 ± 0.05)	%		863
$K^- 2\pi^+$	[c] (9.38 ± 0.16)	%	S=1.6	846
$(K^- \pi^+)_{S\text{-wave}} \pi^+$	(7.52 ± 0.17)	%		846
$\bar{K}_0^*(1430)^0 \pi^+,$	[d] (1.25 ± 0.06)	%		382
$\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$				
$\bar{K}^*(892)^0 \pi^+,$	(1.04 ± 0.12)	%		714
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$				
$\bar{K}^*(1410)^0 \pi^+, \bar{K}^{*0} \rightarrow$	not seen			381
$K^- \pi^+$				
$\bar{K}_2^*(1430)^0 \pi^+,$	[d] (2.3 ± 0.7)	$\times 10^{-4}$		371
$\bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$				
$\bar{K}^*(1680)^0 \pi^+,$	[d] (2.2 ± 1.1)	$\times 10^{-4}$		58
$\bar{K}^*(1680)^0 \rightarrow K^- \pi^+$				
$K^- (2\pi^+)_{I=2}$	(1.45 ± 0.26)	%		–
$K_S^0 \pi^+ \pi^0$	[c] (7.36 ± 0.21)	%		845
$K_S^0 \rho^+$	$(6.14 \begin{smallmatrix} + 0.60 \\ - 0.35 \end{smallmatrix})$	%		677
$K_S^0 \rho(1450)^+, \rho^+ \rightarrow \pi^+ \pi^0$	$(1.5 \begin{smallmatrix} + 1.2 \\ - 1.4 \end{smallmatrix})$	$\times 10^{-3}$		–
$\bar{K}^*(892)^0 \pi^+,$	(2.64 ± 0.32)	$\times 10^{-3}$		714
$\bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$				

$\bar{K}_0^*(1430)^0 \pi^+, \bar{K}_0^{*0} \rightarrow K_S^0 \pi^0$	$(2.7 \pm 0.9) \times 10^{-3}$	—
$\bar{K}_0^*(1680)^0 \pi^+, \bar{K}_0^{*0} \rightarrow K_S^0 \pi^0$	$(10 \pm_{-10}^7) \times 10^{-4}$	—
$\bar{\kappa}^0 \pi^+, \bar{\kappa}^0 \rightarrow K_S^0 \pi^0$	$(6 \pm_{-4}^5) \times 10^{-3}$	—
$K_S^0 \pi^+ \pi^0$ nonresonant	$(3 \pm 4) \times 10^{-3}$	845
$K_S^0 \pi^+ \pi^0$ nonresonant and $\bar{\kappa}^0 \pi^+$	$(1.37 \pm_{-0.40}^{0.21}) \%$	—
$(K_S^0 \pi^0)_{S\text{-wave}} \pi^+$	$(1.27 \pm_{-0.33}^{0.27}) \%$	845
$K_S^0 \pi^+ \omega$	$(7.1 \pm 0.5) \times 10^{-3}$	606
$K_S^0 \pi^+ \eta$	$(1.31 \pm 0.05) \%$	722
$K_S^0 \pi^+ \eta'(958)$	$(1.90 \pm 0.21) \times 10^{-3}$	481
$K^- 2\pi^+ \pi^0$	[e] $(6.25 \pm 0.18) \%$	817
$K_S^0 2\pi^+ \pi^-$	[e] $(3.10 \pm 0.09) \%$	814
$K_S^0 \pi^+ 2\pi^0$	$(2.90 \pm 0.11) \%$	817
$K^- 2\pi^+ \eta$	$(1.35 \pm 0.12) \times 10^{-3}$	657
$K_S^0 \pi^+ \pi^0 \eta$	$(1.22 \pm 0.25) \times 10^{-3}$	657
$K^- 3\pi^+ \pi^-$	[c] $(5.7 \pm 0.5) \times 10^{-3}$	S=1.1 772
$\bar{K}^*(892)^0 2\pi^+ \pi^-, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(1.2 \pm 0.4) \times 10^{-3}$	645
$\bar{K}^*(892)^0 \rho^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(2.3 \pm 0.4) \times 10^{-3}$	239
$\bar{K}^*(892)^0 a_1(1260)^+$	[f] $(9.3 \pm 1.9) \times 10^{-3}$	†
$K^- \rho^0 2\pi^+$	$(1.72 \pm 0.28) \times 10^{-3}$	524
$K^- 3\pi^+ \pi^-$ nonresonant	$(4.0 \pm 2.9) \times 10^{-4}$	772
$K_S^0 2\pi^+ \pi^- \pi^0$	$(1.53 \pm 0.08) \%$	773
$K_S^0 \pi^+ 3\pi^0$	$(5.5 \pm 0.5) \times 10^{-3}$	776
$K^- 2\pi^+ 2\pi^0$	$(4.95 \pm 0.32) \times 10^{-3}$	776
$K^+ 2K_S^0$	$(2.54 \pm 0.13) \times 10^{-3}$	545
$K^+ K^- K_S^0 \pi^+$	$(2.4 \pm 0.5) \times 10^{-4}$	436
Pionic modes		
$\pi^+ \pi^0$	$(1.247 \pm 0.033) \times 10^{-3}$	925
$2\pi^+ \pi^-$	$(3.27 \pm 0.09) \times 10^{-3}$	909
$\rho^0 \pi^+$	$(8.3 \pm 1.4) \times 10^{-4}$	767
$\pi^+ (\pi^+ \pi^-)_{S\text{-wave}}$	$(1.83 \pm 0.14) \times 10^{-3}$	909
$\sigma \pi^+, \sigma \rightarrow \pi^+ \pi^-$	$(1.38 \pm 0.10) \times 10^{-3}$	—
$f_0(980) \pi^+, f_0(980) \rightarrow \pi^+ \pi^-$	$(1.57 \pm 0.32) \times 10^{-4}$	669
$f_0(1370) \pi^+, f_0(1370) \rightarrow \pi^+ \pi^-$	$(8 \pm 4) \times 10^{-5}$	—

$f_2(1270)\pi^+$,	$(5.0 \pm 0.8) \times 10^{-4}$	485
$f_2(1270) \rightarrow \pi^+\pi^-$		
$\rho(1450)^0\pi^+$,	$< 8 \times 10^{-5}$	338
$\rho(1450)^0 \rightarrow \pi^+\pi^-$		
$f_0(1500)\pi^+$,	$(1.1 \pm 0.4) \times 10^{-4}$	—
$f_0(1500) \rightarrow \pi^+\pi^-$		
$f_0(1710)\pi^+$,	$< 5 \times 10^{-5}$	—
$f_0(1710) \rightarrow \pi^+\pi^-$		
$f_0(1790)\pi^+$,	$< 7 \times 10^{-5}$	—
$f_0(1790) \rightarrow \pi^+\pi^-$		
$(\pi^+\pi^+)_{S\text{-wave}}\pi^-$	$< 1.2 \times 10^{-4}$	909
$2\pi^+\pi^-$ nonresonant	$< 1.1 \times 10^{-4}$	909
$\pi^+2\pi^0$	$(4.61 \pm 0.15) \times 10^{-3}$	910
$2\pi^+\pi^-\pi^0$	$(1.165 \pm 0.030) \%$	883
$\pi^+3\pi^0$	$(4.17 \pm 0.26) \times 10^{-3}$	885
$\pi^+4\pi^0$	$(1.9 \pm 0.4) \times 10^{-3}$	851
$2\pi^+\pi^-2\pi^0$	$(1.07 \pm 0.05) \%$	848
$3\pi^+2\pi^-$	$(1.66 \pm 0.16) \times 10^{-3}$	S=1.1 845
$2\pi^+\pi^-3\pi^0$	$(3.42 \pm 0.35) \times 10^{-3}$	803
$3\pi^+2\pi^-\pi^0$	$(2.34 \pm 0.27) \times 10^{-3}$	799
$\eta\pi^+$	$(3.77 \pm 0.09) \times 10^{-3}$	848
$\eta\pi^+\pi^0$	$(2.05 \pm 0.35) \times 10^{-3}$	S=2.2 831
$\eta2\pi^+\pi^-$	$(3.41 \pm 0.20) \times 10^{-3}$	798
$\eta\pi^+2\pi^0$	$(3.20 \pm 0.33) \times 10^{-3}$	801
$\eta\pi^+3\pi^0$	$(2.9 \pm 0.5) \times 10^{-3}$	759
$\eta2\pi^+\pi^-\pi^0$	$(3.88 \pm 0.34) \times 10^{-3}$	755
$\eta\eta\pi^+$	$(2.96 \pm 0.26) \times 10^{-3}$	700
$\omega\pi^+$	$(2.8 \pm 0.6) \times 10^{-4}$	764
$\omega\pi^+\pi^0$	$(3.9 \pm 0.9) \times 10^{-3}$	742
$\eta'(958)\pi^+$	$(4.97 \pm 0.19) \times 10^{-3}$	681
$\eta'(958)\pi^+\pi^0$	$(1.6 \pm 0.5) \times 10^{-3}$	654

Hadronic modes with a $K\bar{K}$ pair

$K_S^0 K^+$	$(3.04 \pm 0.09) \times 10^{-3}$	S=2.2 793
$K_L^0 K^+$	$(3.21 \pm 0.16) \times 10^{-3}$	793
$K_S^0 K^+\pi^0$	$(5.07 \pm 0.30) \times 10^{-3}$	744
$K^*(892)^+ K_S^0, K^{*+} \rightarrow$ $K^+\pi^0$	$(2.89 \pm 0.30) \times 10^{-3}$	612
$\bar{K}^*(892)^0 K^+, \bar{K}^{*0} \rightarrow$ $K_S^0\pi^0$	$(5.2 \pm 1.4) \times 10^{-4}$	613
$K_L^0 K^+\pi^0$	$(5.24 \pm 0.31) \times 10^{-3}$	744
$K^+ K^- \pi^+$	[c] $(9.68 \pm 0.18) \times 10^{-3}$	744
$K^+ \bar{K}^*(892)^0,$ $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	$(2.49 \pm_{-0.13}^{+0.08}) \times 10^{-3}$	613

$K^+ \bar{K}_0^*(1430)^0,$ $\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$	$(1.82 \pm 0.35) \times 10^{-3}$	—
$K^+ \bar{K}_2^*(1430)^0, \bar{K}_2^* \rightarrow$ $K^- \pi^+$	$(1.6 \pm_{-0.8}^{1.2}) \times 10^{-4}$	—
$K^+ \bar{K}_0^*(700), \bar{K}_0^* \rightarrow K^- \pi^+$	$(6.8 \pm_{-2.1}^{3.5}) \times 10^{-4}$	—
$a_0(1450)^0 \pi^+, a_0^0 \rightarrow$ $K^+ K^-$	$(4.5 \pm_{-1.8}^{7.0}) \times 10^{-4}$	—
$\phi(1680) \pi^+, \phi \rightarrow K^+ K^-$	$(4.9 \pm_{-1.9}^{4.0}) \times 10^{-5}$	—
$\phi \pi^+, \phi \rightarrow K^+ K^-$	$(2.69 \pm_{-0.08}^{0.07}) \times 10^{-3}$	647
$\phi \pi^+$	$(5.70 \pm 0.14) \times 10^{-3}$	647
$K^+ K^- \pi^+ \pi^0$	$(6.62 \pm 0.32) \times 10^{-3}$	682
$K_S^0 K_S^0 \pi^+$	$(2.70 \pm 0.13) \times 10^{-3}$	741
$K_S^0 K_S^0 \pi^+ \pi^0$	$(1.34 \pm 0.21) \times 10^{-3}$	679
$K_S^0 K^+ \eta$	$(1.8 \pm 0.5) \times 10^{-4}$	516
$K^+ K_S^0 \pi^+ \pi^-$	$(1.89 \pm 0.13) \times 10^{-3}$	678
$K_S^0 K^+ \pi^0 \pi^0$	$(5.8 \pm 1.3) \times 10^{-4}$	683
$K_S^0 K^- 2\pi^+$	$(2.27 \pm 0.13) \times 10^{-3}$	678
$K^+ K^- 2\pi^+ \pi^-$	$(2.3 \pm 1.2) \times 10^{-4}$	601
A few poorly measured branching fractions:		
$\phi \pi^+ \pi^0$	$(2.3 \pm 1.0) \%$	619
$\phi \rho^+$	$< 1.5 \%$ CL=90%	260
$K^+ K^- \pi^+ \pi^0$ non- ϕ	$(1.5 \pm_{-0.6}^{0.7}) \%$	682

Doubly Cabibbo-suppressed modes

$K^+ \pi^0$	$(2.08 \pm 0.21) \times 10^{-4}$	S=1.4	864
$K^+ \eta$	$(1.25 \pm 0.16) \times 10^{-4}$	S=1.1	776
$K^+ \eta'(958)$	$(1.85 \pm 0.20) \times 10^{-4}$		571
$K^+ 2\pi^0$	$(2.1 \pm 0.4) \times 10^{-4}$		847
$K^*(892)^+ \pi^0$	$(3.4 \pm 1.4) \times 10^{-4}$		714
$K^+ \pi^+ \pi^-$	$(4.91 \pm 0.09) \times 10^{-4}$		846
$K^+ \rho^0$	$(1.9 \pm 0.5) \times 10^{-4}$		679
$K^+ \eta \pi^0$	$(2.1 \pm 0.5) \times 10^{-4}$		726
$K^*(892)^+ \eta$	$(4.4 \pm_{-1.5}^{1.8}) \times 10^{-4}$		586
$K^*(892)^0 \pi^+, K^*(892)^0 \rightarrow$ $K^+ \pi^-$	$(2.3 \pm 0.4) \times 10^{-4}$		714
$K^+ f_0(980), f_0(980) \rightarrow$ $\pi^+ \pi^-$	$(4.4 \pm 2.6) \times 10^{-5}$		—
$K_2^*(1430)^0 \pi^+, K_2^*(1430)^0 \rightarrow$ $K^+ \pi^-$	$(3.9 \pm 2.7) \times 10^{-5}$		—
$K^+ \pi^+ \pi^-$ nonresonant	not seen		846

$K^+ \pi^+ \pi^- \pi^0$		$(1.21 \pm 0.09) \times 10^{-3}$	817
$K^+ \pi^+ \pi^- \pi^0$ nonresonant		$(1.10 \pm 0.07) \times 10^{-3}$	817
$K^+ \omega$		$(5.7 \begin{smallmatrix} + 2.5 \\ - 2.1 \end{smallmatrix}) \times 10^{-5}$	675
$2K^+ K^-$		$(6.14 \pm 0.11) \times 10^{-5}$	550
$\phi(1020)^0 K^+$		$< 2.1 \times 10^{-5}$ CL=90%	–
$K^+ \phi(1020), \phi \rightarrow K^+ K^-$		$(4.4 \pm 0.6) \times 10^{-6}$	–
$K^+ (K^+ K^-)_{S-wave}$		$(5.77 \pm 0.12) \times 10^{-5}$	550

**$\Delta C = 1$ weak neutral current (C1) modes, or Lepton Family number (LF) ,
or Lepton number (L), or Baryon number (B) violating modes**

$\pi^+ e^+ e^-$	C1	$< 1.1 \times 10^{-6}$ CL=90%	930
$\pi^+ \pi^0 e^+ e^-$		$< 1.4 \times 10^{-5}$ CL=90%	925
$\pi^+ \phi, \phi \rightarrow e^+ e^-$	[g]	$(1.7 \begin{smallmatrix} + 1.4 \\ - 0.9 \end{smallmatrix}) \times 10^{-6}$	–
$\pi^+ \mu^+ \mu^-$	C1	$< 6.7 \times 10^{-8}$ CL=90%	918
$\pi^+ \phi, \phi \rightarrow \mu^+ \mu^-$	[g]	$(1.8 \pm 0.8) \times 10^{-6}$	–
$\rho^+ \mu^+ \mu^-$	C1	$< 5.6 \times 10^{-4}$ CL=90%	757
$K^+ e^+ e^-$	[h]	$< 8.5 \times 10^{-7}$ CL=90%	870
$K^+ \pi^0 e^+ e^-$		$< 1.5 \times 10^{-5}$ CL=90%	864
$K_S^0 \pi^+ e^+ e^-$		$< 2.6 \times 10^{-5}$ CL=90%	–
$K_S^0 K^+ e^+ e^-$		$< 1.1 \times 10^{-5}$ CL=90%	792
$K^+ \mu^+ \mu^-$	[h]	$< 5.4 \times 10^{-8}$ CL=90%	856
$\pi^+ e^+ \mu^-$	LF	$< 2.1 \times 10^{-7}$ CL=90%	927
$\pi^+ e^- \mu^+$	LF	$< 2.2 \times 10^{-7}$ CL=90%	927
$K^+ e^+ \mu^-$	LF	$< 7.5 \times 10^{-8}$ CL=90%	866
$K^+ e^- \mu^+$	LF	$< 1.0 \times 10^{-7}$ CL=90%	866
$\pi^- 2e^+$	L	$< 5.3 \times 10^{-7}$ CL=90%	930
$\pi^- 2\mu^+$	L	$< 1.4 \times 10^{-8}$ CL=90%	918
$\pi^- e^+ \mu^+$	L	$< 1.3 \times 10^{-7}$ CL=90%	927
$\rho^- 2\mu^+$	L	$< 5.6 \times 10^{-4}$ CL=90%	757
$K^- 2e^+$	L	$< 9 \times 10^{-7}$ CL=90%	870
$K_S^0 \pi^- 2e^+$		$< 3.3 \times 10^{-6}$ CL=90%	863
$K^- \pi^0 2e^+$		$< 8.5 \times 10^{-6}$ CL=90%	864
$K^- 2\mu^+$	L	$< 1.0 \times 10^{-5}$ CL=90%	856
$K^- e^+ \mu^+$	L	$< 1.9 \times 10^{-6}$ CL=90%	866
$K^*(892)^- 2\mu^+$	L	$< 8.5 \times 10^{-4}$ CL=90%	703
Λe^+	L,B	$< 1.1 \times 10^{-6}$ CL=90%	602
$\bar{\Lambda} e^+$	L,B	$< 6.5 \times 10^{-7}$ CL=90%	602
$\Sigma^0 e^+$	L,B	$< 1.7 \times 10^{-6}$ CL=90%	554
$\bar{\Sigma}^0 e^+$	L,B	$< 1.3 \times 10^{-6}$ CL=90%	554
$\bar{n} e^+$		$< 1.43 \times 10^{-5}$ CL=90%	699
$n e^+$		$< 2.91 \times 10^{-5}$ CL=90%	699



$$I(J^P) = \frac{1}{2}(0^-)$$

$$\text{Mass } m = 1864.84 \pm 0.05 \text{ MeV}$$

$$m_{D^\pm} - m_{D^0} = 4.822 \pm 0.015 \text{ MeV}$$

$$\text{Mean life } \tau = (410.3 \pm 1.0) \times 10^{-15} \text{ s}$$

$$c\tau = 123.01 \text{ } \mu\text{m}$$

Mixing and related parameters

$$|m_{D_1^0} - m_{D_2^0}| = (0.997 \pm 0.116) \times 10^{10} \hbar \text{ s}^{-1}$$

$$(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma = 2y = (1.394 \pm 0.056) \times 10^{-2}$$

$$|q/p| = 0.995 \pm 0.016$$

$$A_\Gamma = (0.089 \pm 0.113) \times 10^{-3}$$

$$\phi_S^{K_S^0 \pi \pi} = 0.02_{-0.05}^{+0.04}$$

$$K^+ \pi^- \text{ relative strong phase: } \cos \delta = 0.990 \pm 0.025$$

$$K^- \pi^+ \pi^0 \text{ coherence factor } R_{K \pi \pi^0} = 0.792 \pm 0.033$$

$$K^- \pi^+ \pi^0 \text{ average relative strong phase } \delta^{K \pi \pi^0} = (198 \pm 10)^\circ$$

$$K^- \pi^- 2\pi^+ \text{ coherence factor } R_{K 3\pi} = 0.52_{-0.09}^{+0.10}$$

$$K^- \pi^- 2\pi^+ \text{ average relative strong phase } \delta^{K 3\pi} = (149_{-16}^{+26})^\circ \quad (S = 1.4)$$

$$D^0 \rightarrow K^- \pi^- 2\pi^+, R_{K 3\pi} (y \cos \delta^{K 3\pi} - x \sin \delta^{K 3\pi}) = (-3.0 \pm 0.7) \times 10^{-3} \text{ TeV}^{-1}$$

$$K_S^0 K^+ \pi^- \text{ coherence factor } R_{K_S^0 K \pi} = 0.70 \pm 0.08$$

$$K_S^0 K^+ \pi^- \text{ average relative strong phase } \delta^{K_S^0 K \pi} = (0 \pm 16)^\circ$$

$$K^* K \text{ coherence factor } R_{K^* K} = 0.94 \pm 0.12$$

$$K^* K \text{ average relative strong phase } \delta^{K^* K} = (-17 \pm 18)^\circ$$

CP-even fractions (labeled by the D^0 decay)

$$\text{CP-even fraction in } D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0 \text{ decays} = (23.8 \pm 1.7)\%$$

$$\text{CP-even fraction in } D^0 \rightarrow \pi^+ \pi^- \pi^0 \text{ decays} = (97.3 \pm 1.7)\%$$

$$\text{CP-even fraction in } D^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^- \text{ decays} = (74.6 \pm 1.6)\% \quad (S = 1.2)$$

$$\text{CP-even fraction in } D^0 \rightarrow \pi^+ \pi^- 2\pi^0 \text{ decays} = 0.68 \pm 0.08$$

$$\text{CP-even fraction in } D^0 \rightarrow 2\pi^+ 2\pi^- \pi^0 \text{ decays} = 0.44 \pm 0.10$$

$$\text{CP-even fraction in } D^0 \rightarrow \pi^+ \pi^- 3\pi^0 \text{ decays} = 0.52_{-0.27}^{+0.34}$$

$$\text{CP-even fraction in } D^0 \rightarrow 2\pi^+ 2\pi^- 2\pi^0 \text{ decays} = 0.79 \pm 0.26$$

$$\text{CP-even fraction in } D^0 \rightarrow K^+ K^- \pi^0 \text{ decays} = (73 \pm 6)\%$$

$$\text{CP-even fraction in } D^0 \rightarrow K^+ K^- \pi^+ \pi^- \text{ decays} = (75 \pm 4)\%$$

CP-violation decay-rate asymmetries (labeled by the D^0 decay)

$$A_{CP}(K^+ K^-) = (-0.07 \pm 0.11)\%$$

$$\begin{aligned}
 A_{CP}(2K_S^0) &= (-1.9 \pm 1.1)\% \quad (S = 1.1) \\
 A_{CP}(\pi^+\pi^-) &= (0.13 \pm 0.14)\% \\
 A_{CP}(\pi^0\pi^0) &= (0.0 \pm 0.6)\% \\
 A_{CP}(\rho\gamma) &= (6 \pm 15) \times 10^{-2} \\
 A_{CP}(\phi\gamma) &= (-9 \pm 7) \times 10^{-2} \\
 A_{CP}(\overline{K}^*(892)^0\gamma) &= (-0.3 \pm 2.0) \times 10^{-2} \\
 A_{CP}(\pi^+\pi^-\pi^0) &= (0.4 \pm 0.4)\% \\
 A_{CP}(\eta\pi^+\pi^-) \text{ in } D^0, \overline{D}^0 \rightarrow \eta\pi^+\pi^- &= (0.9 \pm 1.3) \times 10^{-2} \\
 A_{CP}(\rho(770)^+\pi^- \rightarrow \pi^+\pi^-\pi^0) &= (1.2 \pm 0.9)\% [i] \\
 A_{CP}(\rho(770)^0\pi^0 \rightarrow \pi^+\pi^-\pi^0) &= (-3.1 \pm 3.0)\% [i] \\
 A_{CP}(\rho(770)^-\pi^+ \rightarrow \pi^+\pi^-\pi^0) &= (-1.0 \pm 1.7)\% [i] \\
 A_{CP}(\rho(1450)^+\pi^- \rightarrow \pi^+\pi^-\pi^0) &= (0 \pm 70)\% [i] \\
 A_{CP}(\rho(1450)^0\pi^0 \rightarrow \pi^+\pi^-\pi^0) &= (-20 \pm 40)\% [i] \\
 A_{CP}(\rho(1450)^-\pi^+ \rightarrow \pi^+\pi^-\pi^0) &= (6 \pm 9)\% [i] \\
 A_{CP}(\rho(1700)^+\pi^- \rightarrow \pi^+\pi^-\pi^0) &= (-5 \pm 14)\% [i] \\
 A_{CP}(\rho(1700)^0\pi^0 \rightarrow \pi^+\pi^-\pi^0) &= (13 \pm 9)\% [i] \\
 A_{CP}(\rho(1700)^-\pi^+ \rightarrow \pi^+\pi^-\pi^0) &= (8 \pm 11)\% [i] \\
 A_{CP}(f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0) &= (0 \pm 35)\% [i] \\
 A_{CP}(f_0(1370)\pi^0 \rightarrow \pi^+\pi^-\pi^0) &= (25 \pm 18)\% [i] \\
 A_{CP}(f_0(1500)\pi^0 \rightarrow \pi^+\pi^-\pi^0) &= (0 \pm 18)\% [i] \\
 A_{CP}(f_0(1710)\pi^0 \rightarrow \pi^+\pi^-\pi^0) &= (0 \pm 24)\% [i] \\
 A_{CP}(f_2(1270)\pi^0 \rightarrow \pi^+\pi^-\pi^0) &= (-4 \pm 6)\% [i] \\
 A_{CP}(\sigma(400)\pi^0 \rightarrow \pi^+\pi^-\pi^0) &= (6 \pm 8)\% [i] \\
 A_{CP}(\text{nonresonant } \pi^+\pi^-\pi^0) &= (-13 \pm 23)\% [i] \\
 A_{CP}(\pi^+\pi^-2\pi^0) \text{ in } D^0, \overline{D}^0 \rightarrow \pi^+\pi^-2\pi^0 &= (-2.5 \pm 2.0)\% \\
 A_{CP}(a_1(1260)^+\pi^- \rightarrow 2\pi^+2\pi^-) &= (5 \pm 6)\% \\
 A_{CP}(a_1(1260)^-\pi^+ \rightarrow 2\pi^+2\pi^-) &= (14 \pm 18)\% \\
 A_{CP}(\pi(1300)^+\pi^- \rightarrow 2\pi^+2\pi^-) &= (-2 \pm 15)\% \\
 A_{CP}(\pi(1300)^-\pi^+ \rightarrow 2\pi^+2\pi^-) &= (-6 \pm 30)\% \\
 A_{CP}(a_1(1640)^+\pi^- \rightarrow 2\pi^+2\pi^-) &= (9 \pm 26)\% \\
 A_{CP}(\pi_2(1670)^+\pi^- \rightarrow 2\pi^+2\pi^-) &= (7 \pm 18)\% \\
 A_{CP}(\sigma f_0(1370) \rightarrow 2\pi^+2\pi^-) &= (-15 \pm 19)\% \\
 A_{CP}(\sigma\rho(770)^0 \rightarrow 2\pi^+2\pi^-) &= (3 \pm 27)\% \\
 A_{CP}(2\rho(770)^0 \rightarrow 2\pi^+2\pi^-) &= (-6 \pm 6)\% \\
 A_{CP}(2f_2(1270) \rightarrow 2\pi^+2\pi^-) &= (-28 \pm 24)\% \\
 A_{CP}(\pi^+\pi^-\pi^0\eta) \text{ in } D^0, \overline{D}^0 \rightarrow \pi^+\pi^-\pi^0\eta &= (-6 \pm 6) \times 10^{-2} \\
 A_{CP}(K^+K^-\pi^0) &= (-1.0 \pm 1.7)\% \\
 A_{CP}(K^*(892)^+K^- \rightarrow K^+K^-\pi^0) &= (-0.9 \pm 1.3)\% [i] \\
 A_{CP}(K^*(1410)^+K^- \rightarrow K^+K^-\pi^0) &= (-21 \pm 24)\% [i] \\
 A_{CP}((K^+\pi^0)_{S\text{-wave}}K^- \rightarrow K^+K^-\pi^0) &= (7 \pm 15)\% [i] \\
 A_{CP}(\phi(1020)\pi^0 \rightarrow K^+K^-\pi^0) &= (1.1 \pm 2.2)\% [i] \\
 A_{CP}(f_0(980)\pi^0 \rightarrow K^+K^-\pi^0) &= (-3 \pm 19)\% [i]
 \end{aligned}$$

$$\begin{aligned}
 A_{CP}(a_0(980)^0 \pi^0 \rightarrow K^+ K^- \pi^0) &= (-5 \pm 16)\% [i] \\
 A_{CP}(f_2'(1525) \pi^0 \rightarrow K^+ K^- \pi^0) &= (0 \pm 160)\% [i] \\
 A_{CP}(K^*(892)^- K^+ \rightarrow K^+ K^- \pi^0) &= (-5 \pm 4)\% [i] \\
 A_{CP}(K^*(1410)^- K^+ \rightarrow K^+ K^- \pi^0) &= (-17 \pm 29)\% [i] \\
 A_{CP}((K^- \pi^0)_{S\text{-wave}} K^+ \rightarrow K^+ K^- \pi^0) &= (-10 \pm 40)\% [i] \\
 A_{CP}(K^+ K^- \eta) \text{ in } D^0, \bar{D}^0 \rightarrow K^+ K^- \eta &= (-1.4 \pm 3.5) \times 10^{-2} \\
 A_{CP}(\phi(1020) \eta \rightarrow K^+ K^- \eta) \text{ in } D^0, \bar{D}^0 \rightarrow \phi(1020) \eta &= (-2 \pm 4) \times 10^{-2} \\
 A_{CP}(K_S^0 \pi^0) &= (-0.20 \pm 0.17)\% \\
 A_{CP}(K_S^0 \eta) &= (0.5 \pm 0.5)\% \\
 A_{CP}(K_S^0 \eta') &= (1.0 \pm 0.7)\% \\
 A_{CP}(K_S^0 \phi) &= (-3 \pm 9)\% \\
 A_{CP}(K^- \pi^+) &= (0.2 \pm 0.5)\% \\
 A_{CP}(K^+ \pi^-) &= (-0.9 \pm 1.4)\% \\
 A_{CP}(D_{CP}(\pm 1) \rightarrow K^\mp \pi^\pm) &= (13.1 \pm 1.0)\% \\
 A_{CP}(K^- \pi^+ \pi^0) &= (0.1 \pm 0.5)\% \\
 A_{CP}(K^+ \pi^- \pi^0) &= (0 \pm 5)\% \\
 A_{CP}(K_S^0 \pi^+ \pi^-) &= (-0.1 \pm 0.8)\% \\
 A_{CP}(K^\mp \pi^\pm \eta) \text{ in } D^0, \bar{D}^0 \rightarrow K^\mp \pi^\pm \eta &= (-1.9 \pm 1.6) \times 10^{-2} \\
 A_{CP}(K_S^0 \pi^0 \eta) \text{ in } D^0, \bar{D}^0 \rightarrow K_S^0 \pi^0 \eta &= (-3.9 \pm 3.3) \times 10^{-2} \\
 A_{CP}(K^\mp \pi^\pm \pi^0 \eta) \text{ in } D^0, \bar{D}^0 \rightarrow K^\mp \pi^\pm \pi^0 \eta &= (-8 \pm 5) \times 10^{-2} \\
 A_{CP}(K^*(892)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &= (0.4 \pm 0.5)\% \\
 A_{CP}(K^*(892)^+ \pi^- \rightarrow K_S^0 \pi^+ \pi^-) &= (1 \pm 6)\% \\
 A_{CP}(\bar{K}^0 \rho^0 \rightarrow K_S^0 \pi^+ \pi^-) &= (-0.1 \pm 0.5)\% \\
 A_{CP}(\bar{K}^0 \omega \rightarrow K_S^0 \pi^+ \pi^-) &= (-13 \pm 7)\% \\
 A_{CP}(\bar{K}^0 f_0(980) \rightarrow K_S^0 \pi^+ \pi^-) &= (-0.4 \pm 2.7)\% \\
 A_{CP}(\bar{K}^0 f_2(1270) \rightarrow K_S^0 \pi^+ \pi^-) &= (-4 \pm 5)\% \\
 A_{CP}(\bar{K}^0 f_0(1370) \rightarrow K_S^0 \pi^+ \pi^-) &= (-1 \pm 9)\% \\
 A_{CP}(\bar{K}^0 \rho^0(1450) \rightarrow K_S^0 \pi^+ \pi^-) &= (-4 \pm 10)\% \\
 A_{CP}(\bar{K}^0 f_0(600) \rightarrow K_S^0 \pi^+ \pi^-) &= (-3 \pm 5)\% \\
 A_{CP}(K^*(1410)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &= (-2 \pm 9)\% \\
 A_{CP}(K_0^*(1430)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &= (4 \pm 4)\% \\
 A_{CP}(K_0^*(1430)^+ \pi^- \rightarrow K_S^0 \pi^+ \pi^-) &= (12 \pm 15)\% \\
 A_{CP}(K_2^*(1430)^- \pi^+ \rightarrow K_S^0 \pi^+ \pi^-) &= (3 \pm 6)\% \\
 A_{CP}(K_2^*(1430)^+ \pi^- \rightarrow K_S^0 \pi^+ \pi^-) &= (-10 \pm 32)\% \\
 A_{CP}(K^- \pi^+ \pi^+ \pi^-) &= (0.2 \pm 0.5)\% \\
 A_{CP}(K^+ \pi^- \pi^+ \pi^-) &= (-2 \pm 4)\% \\
 A_{CP}(K^+ K^- \pi^+ \pi^-) &= (1.3 \pm 1.7)\% \\
 A_{CP}(K_1^*(1270)^+ K^- \rightarrow K^+ K^- \pi^+ \pi^-) &= (-2.3 \pm 1.7)\% \\
 A_{CP}(K_1^*(1270)^+ K^- \rightarrow K^{*0} \pi^+ K^-) &= (-1 \pm 10)\% \\
 A_{CP}(K_1^*(1270)^- K^+ \rightarrow \bar{K}^{*0} \pi^- K^+) &= (-10 \pm 32)\% \\
 A_{CP}(K_1^*(1270)^- K^+ \rightarrow K^+ K^- \pi^+ \pi^-) &= (1.7 \pm 3.5)\%
 \end{aligned}$$

$$\begin{aligned}
 A_{CP}(K_1^*(1270)^+ K^- \rightarrow \rho^0 K^+ K^-) &= (-7 \pm 17)\% \\
 A_{CP}(K_1^*(1270)^- K^+ \rightarrow \rho^0 K^- K^+) &= (10 \pm 13)\% \\
 A_{CP}(K_1(1400)^+ K^- \rightarrow K^+ K^- \pi^+ \pi^-) &= (-4.4 \pm 2.1)\% \\
 A_{CP}(K^*(1410)^+ K^- \rightarrow K^{*0} \pi^+ K^-) &= (-20 \pm 17)\% \\
 A_{CP}(K^*(1410)^- K^+ \rightarrow \bar{K}^{*0} \pi^- K^+) &= (-1 \pm 14)\% \\
 A_{CP}(K^*(1680)^+ K^- \rightarrow K^+ K^- \pi^+ \pi^-) &= (-17 \pm 29)\% \\
 A_{CP}(K^{*0} \bar{K}^{*0}) \text{ in } D^0, \bar{D}^0 \rightarrow K^{*0} \bar{K}^{*0} &= (-5 \pm 14)\% \\
 A_{CP}(K^{*0} \bar{K}^{*0} \text{ S-wave}) &= (-3.9 \pm 2.2)\% \\
 A_{CP}(\phi \rho^0) \text{ in } D^0, \bar{D}^0 \rightarrow \phi \rho^0 &= (1 \pm 9)\% \\
 A_{CP}(\phi \rho^0 \text{ S-wave}) &= (-3 \pm 5)\% \\
 A_{CP}(\phi \rho^0 \text{ D-wave}) &= (-37 \pm 19)\% \\
 A_{CP}(\phi(\pi^+ \pi^-)_{\text{S-wave}}) &= (6 \pm 6)\% \\
 A_{CP}(K^*(892)^0 (K^- \pi^+)_{\text{S-wave}}) &= (-10 \pm 40)\% \\
 A_{CP}(K^+ K^- \pi^+ \pi^- \text{ non-resonant}) &= (8 \pm 20)\% \\
 A_{CP}((K^- \pi^+)_{\text{P-wave}} (K^+ \pi^-)_{\text{S-wave}}) &= (3 \pm 11)\% \\
 A_{CP}(K^+ K^- \mu^+ \mu^-) \text{ in } D^0, \bar{D}^0 \rightarrow K^+ K^- \mu^+ \mu^- &= (-2 \pm 6)\% \\
 A_{CP}(\pi^+ \pi^- \mu^+ \mu^-) \text{ in } D^0, \bar{D}^0 \rightarrow \pi^+ \pi^- \mu^+ \mu^- &= (2.9 \pm 2.1)\%
 \end{aligned}$$

CP-violation asymmetry difference

$$\Delta A_{CP} = A_{CP}(K^+ K^-) - A_{CP}(\pi^+ \pi^-) = (-0.154 \pm 0.029)\%$$

χ^2 tests of CP-violation (CPV) p-values

$$\begin{aligned}
 \text{Local CPV in } D^0, \bar{D}^0 \rightarrow \pi^+ \pi^- \pi^0 &= 4.9\% \\
 \text{Local CPV in } D^0, \bar{D}^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^- &= (0.6 \pm 0.2)\% \\
 \text{Local CPV in } D^0, \bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^- &= 96\% \\
 \text{Local CPV in } D^0, \bar{D}^0 \rightarrow K^+ K^- \pi^0 &= 16.6\% \\
 \text{Local CPV in } D^0, \bar{D}^0 \rightarrow K^+ K^- \pi^+ \pi^- &= 9.1\%
 \end{aligned}$$

T-violation decay-rate asymmetry

$$\begin{aligned}
 A_T(K^+ K^- \pi^+ \pi^-) &= (2.9 \pm 2.2) \times 10^{-3} [b] \\
 A_{T\text{viol}}(K_S \pi^+ \pi^- \pi^0) \text{ in } D^0, \bar{D}^0 \rightarrow K_S \pi^+ \pi^- \pi^0 &= (-0.3^{+1.4}_{-1.6}) \times 10^{-3}
 \end{aligned}$$

CPT-violation decay-rate asymmetry

$$A_{CPT}(K^\mp \pi^\pm) = 0.008 \pm 0.008$$

Form factors

$$\begin{aligned}
 r_V \equiv V(0)/A_1(0) \text{ in } D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell &= 1.46 \pm 0.07 \\
 r_2 \equiv A_2(0)/A_1(0) \text{ in } D^0 \rightarrow K^*(892)^- \ell^+ \nu_\ell &= 0.68 \pm 0.06 \\
 f_+(0) \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell &= 0.736 \pm 0.004 \\
 f_+(0) |V_{cs}| \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell &= 0.7166 \pm 0.0030 \\
 r_1 \equiv a_1/a_0 \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell &= -2.40 \pm 0.16 \\
 r_2 \equiv a_2/a_0 \text{ in } D^0 \rightarrow K^- \ell^+ \nu_\ell &= 5 \pm 4 \\
 f_+(0) \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell &= 0.637 \pm 0.009 \\
 f_+(0) |V_{cd}| \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell &= 0.1436 \pm 0.0026 \quad (S = 1.5) \\
 r_1 \equiv a_1/a_0 \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell &= -1.97 \pm 0.28 \quad (S = 1.4) \\
 r_2 \equiv a_1/a_0 \text{ in } D^0 \rightarrow \pi^- \ell^+ \nu_\ell &= -0.2 \pm 2.2 \quad (S = 1.7)
 \end{aligned}$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as \bar{K}^0 modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D⁰ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Topological modes			
0-prongs	[j] (15 ± 6) %		—
2-prongs	(71 ± 6) %		—
4-prongs	[k] (14.6 ± 0.5) %		—
6-prongs	[l] (6.5 ± 1.3) × 10 ⁻⁴		—
Inclusive modes			
e ⁺ anything	[n] (6.49 ± 0.11) %		—
μ ⁺ anything	(6.8 ± 0.6) %		—
K ⁻ anything	(54.7 ± 2.8) %	S=1.3	—
\bar{K}^0 anything + K ⁰ anything	(47 ± 4) %		—
K ⁺ anything	(3.4 ± 0.4) %		—
K [*] (892) ⁻ anything	(15 ± 9) %		—
\bar{K}^* (892) ⁰ anything	(9 ± 4) %		—
K [*] (892) ⁺ anything	< 3.6 %	CL=90%	—
K [*] (892) ⁰ anything	(2.8 ± 1.3) %		—
η anything	(9.5 ± 0.9) %		—
η' anything	(2.48 ± 0.27) %		—
φ anything	(1.08 ± 0.04) %		—
invisibles	< 9.4 × 10 ⁻⁵	CL=90%	—
Semileptonic modes			
K ⁻ e ⁺ ν _e	(3.549 ± 0.026) %	S=1.2	867
K ⁻ μ ⁺ ν _μ	(3.41 ± 0.04) %		864
K [*] (892) ⁻ e ⁺ ν _e	(2.15 ± 0.16) %		719
K [*] (892) ⁻ μ ⁺ ν _μ	(1.89 ± 0.24) %		714
K ⁻ π ⁰ e ⁺ ν _e	(1.6 ^{+1.3} / _{-0.5}) %		861
\bar{K}^0 π ⁻ e ⁺ ν _e	(1.44 ± 0.04) %		860
(\bar{K}^0 π ⁻) S-wave e ⁺ ν _e	(7.9 ± 1.7) × 10 ⁻⁴		860
K ⁻ π ⁺ π ⁻ e ⁺ ν _e	(2.8 ^{+1.4} / _{-1.1}) × 10 ⁻⁴		843
K ₁ (1270) ⁻ e ⁺ ν _e	(1.01 ± 0.18) × 10 ⁻³		511
K ⁻ π ⁺ π ⁻ μ ⁺ ν _μ	< 1.3 × 10 ⁻³	CL=90%	821
(\bar{K}^* (892)π) ⁻ μ ⁺ ν _μ	< 1.5 × 10 ⁻³	CL=90%	692
π ⁻ e ⁺ ν _e	(2.91 ± 0.04) × 10 ⁻³		927
π ⁻ μ ⁺ ν _μ	(2.67 ± 0.12) × 10 ⁻³	S=1.3	924
π ⁻ π ⁰ e ⁺ ν _e	(1.45 ± 0.07) × 10 ⁻³		922

$\rho^- e^+ \nu_e$	$(1.50 \pm 0.12) \times 10^{-3}$	S=1.9	771
$\rho^- \mu^+ \nu_\mu$	$(1.35 \pm 0.13) \times 10^{-3}$		767
$a(980)^- e^+ \nu_e, a^- \rightarrow \eta \pi^-$	$(1.33 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.34 \\ 0.30 \end{smallmatrix}) \times 10^{-4}$		—
$b_1(1235)^- e^+ \nu_e, b_1^- \rightarrow \omega \pi^-$	$< 1.12 \times 10^{-4}$	CL=90%	—

Hadronic modes with one \bar{K}

$K^- \pi^+$	$(3.947 \pm 0.030) \%$	S=1.2	861
$K_S^0 \pi^0$	$(1.240 \pm 0.022) \%$		860
$K_L^0 \pi^0$	$(9.76 \pm 0.32) \times 10^{-3}$		860
$K_L^0 \eta$	$(4.34 \pm 0.16) \times 10^{-3}$		772
$K_L^0 \eta'$	$(8.12 \pm 0.35) \times 10^{-3}$	S=1.3	565
$K_L^0 \omega$	$(1.16 \pm 0.04) \%$		670
$K_S^0 \pi^+ \pi^-$	[c] $(2.80 \pm 0.18) \%$	S=1.1	842
$K_S^0 \rho^0$	$(6.3 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.6 \\ 0.8 \end{smallmatrix}) \times 10^{-3}$		674
$K_S^0 \omega, \omega \rightarrow \pi^+ \pi^-$	$(2.0 \pm 0.6) \times 10^{-4}$		670
$K_S^0 (\pi^+ \pi^-)_{S-wave}$	$(3.3 \pm 0.8) \times 10^{-3}$		842
$K_S^0 f_0(980), f_0 \rightarrow \pi^+ \pi^-$	$(1.20 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.40 \\ 0.23 \end{smallmatrix}) \times 10^{-3}$		549
$K_S^0 f_0(1370), f_0 \rightarrow \pi^+ \pi^-$	$(2.8 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.9 \\ 1.3 \end{smallmatrix}) \times 10^{-3}$		†
$K_S^0 f_2(1270), f_2 \rightarrow \pi^+ \pi^-$	$(9 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 10 \\ 6 \end{smallmatrix}) \times 10^{-5}$		262
$K^*(892)^- \pi^+, K^{*-} \rightarrow K_S^0 \pi^-$	$(1.64 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.14 \\ 0.17 \end{smallmatrix}) \%$		711
$K_0^*(1430)^- \pi^+, K_0^{*-} \rightarrow K_S^0 \pi^-$	$(2.67 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.40 \\ 0.33 \end{smallmatrix}) \times 10^{-3}$		378
$K_2^*(1430)^- \pi^+, K_2^{*-} \rightarrow K_S^0 \pi^-$	$(3.4 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 1.9 \\ 1.0 \end{smallmatrix}) \times 10^{-4}$		367
$K^*(1680)^- \pi^+, K^{*-} \rightarrow K_S^0 \pi^-$	$(4.4 \pm 3.5) \times 10^{-4}$		46
$K^*(892)^+ \pi^-, K^{*+} \rightarrow K_S^0 \pi^+$	[o] $(1.13 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.60 \\ 0.34 \end{smallmatrix}) \times 10^{-4}$		711
$K_0^*(1430)^+ \pi^-, K_0^{*+} \rightarrow K_S^0 \pi^+$	[o] $< 1.4 \times 10^{-5}$	CL=95%	—
$K_2^*(1430)^+ \pi^-, K_2^{*+} \rightarrow K_S^0 \pi^+$	[o] $< 3.4 \times 10^{-5}$	CL=95%	—
$K_S^0 \pi^+ \pi^-$ nonresonant	$(2.5 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 6.0 \\ 1.6 \end{smallmatrix}) \times 10^{-4}$		842
$K^- \pi^+ \pi^0$	[c] $(14.4 \pm 0.6) \%$	S=2.2	844
$K^- \rho^+$	$(11.2 \pm 0.7) \%$		675
$K^- \rho(1700)^+, \rho^+ \rightarrow \pi^+ \pi^0$	$(8.2 \pm 1.8) \times 10^{-3}$		†

$K^*(892)^- \pi^+, K^*(892)^- \rightarrow$	$(2.31 \begin{smallmatrix} + 0.40 \\ - 0.20 \end{smallmatrix}) \%$		711
$\bar{K}^*(892)^0 \pi^0, \bar{K}^*(892)^0 \rightarrow$	$(1.95 \pm 0.25) \%$		711
$K_0^*(1430)^- \pi^+, K_0^{*-} \rightarrow$	$(4.8 \pm 2.2) \times 10^{-3}$		378
$\bar{K}_0^*(1430)^0 \pi^0, \bar{K}_0^{*0} \rightarrow$	$(5.9 \begin{smallmatrix} + 5.0 \\ - 1.6 \end{smallmatrix}) \times 10^{-3}$		379
$K^*(1680)^- \pi^+, K^{*-} \rightarrow$	$(1.9 \pm 0.7) \times 10^{-3}$		46
$K^- \pi^+ \pi^0$ nonresonant	$(1.15 \begin{smallmatrix} + 0.60 \\ - 0.20 \end{smallmatrix}) \%$		844
$K_S^0 2\pi^0$	$(9.1 \pm 1.1) \times 10^{-3}$	S=2.2	843
$K_L^0 \pi^0 \pi^0$	$(1.26 \pm 0.06) \%$		843
$K_S^0 (2\pi^0)_{S-wave}$	$(2.6 \pm 0.7) \times 10^{-3}$		—
$\bar{K}^*(892)^0 \pi^0, \bar{K}^{*0} \rightarrow K_S^0 \pi^0$	$(8.1 \pm 0.7) \times 10^{-3}$		711
$\bar{K}^*(1430)^0 \pi^0, \bar{K}^{*0} \rightarrow$	$(4 \pm 23) \times 10^{-5}$		—
$\bar{K}^*(1680)^0 \pi^0, \bar{K}^{*0} \rightarrow$	$(1.0 \pm 0.4) \times 10^{-3}$		—
$K_S^0 f_2(1270), f_2 \rightarrow 2\pi^0$	$(2.3 \pm 1.1) \times 10^{-4}$		—
$2K_S^0, \text{one } K_S^0 \rightarrow 2\pi^0$	$(3.2 \pm 1.1) \times 10^{-4}$		—
$K_S^0 3\pi^0$	$(7.6 \pm 0.4) \times 10^{-3}$		815
$K^- 2\pi^+ \pi^-$	[c] $(8.22 \pm 0.14) \%$	S=1.1	813
$K^- \pi^+ \rho^0$ total	$(6.87 \pm 0.31) \%$		609
$K^- \pi^+ \rho^0$ 3-body	$(6.1 \pm 1.6) \times 10^{-3}$		609
$\bar{K}^*(892)^0 \rho^0, \bar{K}^{*0} \rightarrow$	$(1.01 \pm 0.05) \%$		416
$\bar{K}^*(892)^0 \rho^0$ transverse,	$(1.2 \pm 0.4) \%$		417
$\bar{K}^{*0} \rightarrow K^- \pi^+$			
$K^- a_1(1260)^+, a_1^+ \rightarrow$	$(4.32 \pm 0.32) \%$		327
$\rho^0 \pi^+$			
$K_1(1270)^- \pi^+, K_1^- \rightarrow$	$(3.9 \pm 0.4) \times 10^{-3}$		—
$K^- \pi^+ \pi^-$ total			
$K_1(1270)^- \pi^+, K_1^- \rightarrow$	$(6.6 \pm 2.3) \times 10^{-4}$		484
$\bar{K}^*(892)^0 \pi^-, \bar{K}^{*0} \rightarrow$			
$K^- \pi^+$			
$K^- 2\pi^+ \pi^-$ nonresonant	$(1.81 \pm 0.07) \%$		813
$K_S^0 \pi^+ \pi^- \pi^0$	[p] $(5.2 \pm 0.6) \%$		813
$K_S^0 \eta, \eta \rightarrow \pi^+ \pi^- \pi^0$	$(1.17 \pm 0.03) \times 10^{-3}$		772
$K_S^0 \omega, \omega \rightarrow \pi^+ \pi^- \pi^0$	$(9.9 \pm 0.6) \times 10^{-3}$		670
$K^- \pi^+ 2\pi^0$	$(8.86 \pm 0.23) \%$		815
$K^- \pi^+ 3\pi^0$	$(9.5 \pm 0.4) \times 10^{-3}$		774
$K^- \pi^+ \pi^- 2\pi^0$	$(1.27 \pm 0.06) \%$		773

$K^- 2\pi^+ \pi^- \pi^0$	(4.3 ± 0.4) %		771
$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0, \bar{K}^{*0} \rightarrow$	(1.3 ± 0.6) %		643
$\bar{K}^*(892)^0 \omega, \bar{K}^{*0} \rightarrow$	(6.5 ± 3.0) × 10 ⁻³		410
$K^- \pi^+ \omega$	(3.39 ± 0.10) %		605
$\bar{K}^*(892)^0 \omega$	(1.1 ± 0.5) %		410
$K_S^0 \pi^0 \omega$	(8.5 ± 0.6) × 10 ⁻³		605
$K_S^0 \eta \pi^0$	(1.01 ± 0.05) %		721
$K_S^0 a_0(980), a_0 \rightarrow \eta \pi^0$	(1.20 ± 0.28) %		–
$\bar{K}^*(892)^0 \eta, \bar{K}^{*0} \rightarrow K_S^0 \pi^0$	(2.9 ± 0.7) × 10 ⁻³		–
$K^- \pi^+ \eta$	(1.88 ± 0.05) %	S=1.4	721
$K^*(892)^0 \eta, K^{*0} \rightarrow K^- \pi^+$	(8.9 ± 0.8) × 10 ⁻³		–
$a_0(980)^+ K^-, a_0^+ \rightarrow \eta \pi^+$	(7.4 ± 0.9) × 10 ⁻³		–
$K_2^*(1980)^- \pi^+, K_2^{*-} \rightarrow$	(2.2 ± 1.7) × 10 ⁻⁴		–
$K^- \eta$			
$K^- \pi^+ \pi^0 \eta$	(4.49 ± 0.27) × 10 ⁻³		656
$K_S^0 \pi^+ \pi^- \eta$	(2.80 ± 0.21) × 10 ⁻³		651
$K_S^0 2\pi^0 \eta$	(1.76 ± 0.26) × 10 ⁻³		656
$K_S^0 2\pi^+ 2\pi^-$	(2.66 ± 0.30) × 10 ⁻³		768
$K_S^0 \rho^0 \pi^+ \pi^-, \text{no } K^*(892)^-$	(1.1 ± 0.7) × 10 ⁻³		–
$K^*(892)^- 2\pi^+ \pi^-,$	(5 ± 7) × 10 ⁻⁴		642
$K^*(892)^- \rightarrow K_S^0 \pi^-,$			
$\text{no } \rho^0$			
$K^*(892)^- \rho^0 \pi^+,$	(1.6 ± 0.6) × 10 ⁻³		230
$K^*(892)^- \rightarrow K_S^0 \pi^-$			
$K_S^0 2\pi^+ 2\pi^- \text{ nonresonant}$	< 1.2 × 10 ⁻³	CL=90%	768
$K^- 3\pi^+ 2\pi^-$	(2.2 ± 0.6) × 10 ⁻⁴		713

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes. These nine modes below are all corrected for unseen decays of the resonances.

$K_S^0 \eta$	(5.09 ± 0.13) × 10 ⁻³		772
$K_S^0 \omega$	(1.11 ± 0.06) %		670
$K_S^0 \eta'(958)$	(9.49 ± 0.32) × 10 ⁻³		565
$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0$	(1.9 ± 0.9) %		643
$\bar{K}^*(892)^0 \eta$	(1.41 ± 0.12) %		583
$K^- \pi^+ \eta'(958)$	(6.43 ± 0.34) × 10 ⁻³		479
$K_S^0 \eta'(958) \pi^0$	(2.52 ± 0.27) × 10 ⁻³		479
$\bar{K}^*(892)^0 \eta'(958)$	< 1.0 × 10 ⁻³	CL=90%	119

Hadronic modes with three K's

$K_S^0 K^+ K^-$	$(4.42 \pm 0.32) \times 10^{-3}$		544
$K_S^0 a_0(980)^0, a_0^0 \rightarrow K^+ K^-$	$(2.9 \pm 0.4) \times 10^{-3}$		—
$K^- a_0(980)^+, a_0^+ \rightarrow K^+ K_S^0$	$(5.9 \pm 1.8) \times 10^{-4}$		—
$K^+ a_0(980)^-, a_0^- \rightarrow K^- K_S^0$	< 1.1	$\times 10^{-4}$	CL=95% —
$K_S^0 f_0(980), f_0 \rightarrow K^+ K^-$	< 9	$\times 10^{-5}$	CL=95% —
$K_S^0 \phi, \phi \rightarrow K^+ K^-$	$(2.03 \pm 0.15) \times 10^{-3}$		520
$K_L^0 \phi$	$(4.14 \pm 0.23) \times 10^{-3}$		521
$K_S^0 f_0(1370), f_0 \rightarrow K^+ K^-$	$(1.7 \pm 1.1) \times 10^{-4}$		—
$3K_S^0$	$(7.5 \pm 0.7) \times 10^{-4}$		S=1.4 539
$K^+ 2K^- \pi^+$	$(2.25 \pm 0.32) \times 10^{-4}$		434
$K^+ K^- \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+$	$(4.5 \pm 1.8) \times 10^{-5}$		†
$K^- \pi^+ \phi, \phi \rightarrow K^+ K^-$	$(4.0 \pm 1.7) \times 10^{-5}$		422
$\phi \bar{K}^*(892)^0, \phi \rightarrow K^+ K^-, \bar{K}^{*0} \rightarrow K^- \pi^+$	$(1.08 \pm 0.21) \times 10^{-4}$		†
$K^+ 2K^- \pi^+$ nonresonant	$(3.4 \pm 1.5) \times 10^{-5}$		434
$2K_S^0 K^\pm \pi^\mp$	$(5.9 \pm 1.3) \times 10^{-4}$		427

Pionic modes

$\pi^+ \pi^-$	$(1.454 \pm 0.024) \times 10^{-3}$	S=1.4	922
$2\pi^0$	$(8.26 \pm 0.25) \times 10^{-4}$		923
$\pi^+ \pi^- \pi^0$	$(1.49 \pm 0.07) \%$	S=2.3	907
$\rho^+ \pi^-$	$(1.01 \pm 0.05) \%$		764
$\rho^0 \pi^0$	$(3.86 \pm 0.24) \times 10^{-3}$		764
$\rho^- \pi^+$	$(5.15 \pm 0.26) \times 10^{-3}$		764
$\rho(1450)^+ \pi^-, \rho^+ \rightarrow \pi^+ \pi^0$	$(1.6 \pm 2.1) \times 10^{-5}$		—
$\rho(1450)^0 \pi^0, \rho^0 \rightarrow \pi^+ \pi^-$	$(4.5 \pm 2.0) \times 10^{-5}$		—
$\rho(1450)^- \pi^+, \rho^- \rightarrow \pi^- \pi^0$	$(2.7 \pm 0.4) \times 10^{-4}$		—
$\rho(1700)^+ \pi^-, \rho^+ \rightarrow \pi^+ \pi^0$	$(6.1 \pm 1.5) \times 10^{-4}$		—
$\rho(1700)^0 \pi^0, \rho^0 \rightarrow \pi^+ \pi^-$	$(7.4 \pm 1.8) \times 10^{-4}$		—
$\rho(1700)^- \pi^+, \rho^- \rightarrow \pi^- \pi^0$	$(4.8 \pm 1.1) \times 10^{-4}$		—
$f_0(980) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	$(3.7 \pm 0.9) \times 10^{-5}$		—
$f_0(500) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	$(1.22 \pm 0.22) \times 10^{-4}$		—
$f_0(1370) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	$(5.5 \pm 2.1) \times 10^{-5}$		—
$f_0(1500) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	$(5.8 \pm 1.6) \times 10^{-5}$		—
$f_0(1710) \pi^0, f_0 \rightarrow \pi^+ \pi^-$	$(4.6 \pm 1.6) \times 10^{-5}$		—
$f_2(1270) \pi^0, f_2 \rightarrow \pi^+ \pi^-$	$(1.97 \pm 0.21) \times 10^{-4}$		—
$\pi^+ \pi^- \pi^0$ nonresonant	$(1.3 \pm 0.4) \times 10^{-4}$		907
$3\pi^0$	$(2.0 \pm 0.5) \times 10^{-4}$		908
$2\pi^+ 2\pi^-$	$(7.56 \pm 0.20) \times 10^{-3}$		880

$a_1(1260)^+ \pi^-$, $a_1^+ \rightarrow$	$(4.53 \pm 0.31) \times 10^{-3}$	—
$2\pi^+ \pi^-$ total		
$a_1(1260)^+ \pi^-$, $a_1^+ \rightarrow$	$(3.13 \pm 0.21) \times 10^{-3}$	—
$\rho^0 \pi^+$ S-wave		
$a_1(1260)^+ \pi^-$, $a_1^+ \rightarrow$	$(1.9 \pm 0.5) \times 10^{-4}$	—
$\rho^0 \pi^+$ D-wave		
$a_1(1260)^+ \pi^-$, $a_1^+ \rightarrow$	$(6.4 \pm 0.7) \times 10^{-4}$	—
$\sigma \pi^+$		
$a_1(1260)^- \pi^+$, $a_1^- \rightarrow$	$(2.3 \pm 0.9) \times 10^{-4}$	—
$\rho^0 \pi^-$ S-wave		
$a_1(1260)^- \pi^+$, $a_1^- \rightarrow \sigma \pi^-$	$(6.0 \pm 3.4) \times 10^{-5}$	—
$\pi(1300)^+ \pi^-$, $\pi(1300)^+ \rightarrow$	$(5.1 \pm 2.7) \times 10^{-4}$	—
$\sigma \pi^+$		
$\pi(1300)^- \pi^+$, $\pi(1300)^- \rightarrow$	$(2.3 \pm 2.2) \times 10^{-4}$	—
$\sigma \pi^-$		
$a_1(1640)^+ \pi^-$, $a_1^+ \rightarrow$	$(3.2 \pm 1.6) \times 10^{-4}$	—
$\rho^0 \pi^+$ D-wave		
$a_1(1640)^+ \pi^-$, $a_1^+ \rightarrow \sigma \pi^+$	$(1.8 \pm 1.4) \times 10^{-4}$	—
$\pi_2(1670)^+ \pi^-$, $\pi_2^+ \rightarrow$	$(2.0 \pm 0.9) \times 10^{-4}$	—
$f_2(1270)^0 \pi^+$, $f_2^0 \rightarrow$		
$\pi^+ \pi^-$		
$\pi_2(1670)^+ \pi^-$, $\pi_2^+ \rightarrow \sigma \pi^+$	$(2.6 \pm 1.0) \times 10^{-4}$	—
$2\rho^0$ total	$(1.85 \pm 0.13) \times 10^{-3}$	518
$2\rho^0$, parallel helicities	$(8.3 \pm 3.2) \times 10^{-5}$	—
$2\rho^0$, perpendicular helicities	$(4.8 \pm 0.6) \times 10^{-4}$	—
$2\rho^0$, longitudinal helicities	$(1.27 \pm 0.10) \times 10^{-3}$	—
$2\rho(770)^0$, S-wave	$(1.8 \pm 1.3) \times 10^{-4}$	—
$2\rho(770)^0$, P-wave	$(5.3 \pm 1.3) \times 10^{-4}$	—
$2\rho(770)^0$, D-wave	$(6.2 \pm 3.0) \times 10^{-4}$	—
Resonant $(\pi^+ \pi^-) \pi^+ \pi^-$	$(1.51 \pm 0.12) \times 10^{-3}$	—
3-body total		
$\sigma \pi^+ \pi^-$	$(6.2 \pm 0.9) \times 10^{-4}$	—
$\sigma \rho(770)^0$	$(5.0 \pm 2.5) \times 10^{-4}$	—
$f_0(980) \pi^+ \pi^-$, $f_0 \rightarrow$	$(1.8 \pm 0.5) \times 10^{-4}$	—
$\pi^+ \pi^-$		
$f_2(1270) \pi^+ \pi^-$, $f_2 \rightarrow$	$(3.7 \pm 0.6) \times 10^{-4}$	—
$\pi^+ \pi^-$		
$2f_2(1270)$, $f_2 \rightarrow \pi^+ \pi^-$	$(1.6 \pm 1.8) \times 10^{-4}$	—
$f_0(1370) \sigma$, $f_0 \rightarrow$	$(1.6 \pm 0.5) \times 10^{-3}$	—
$\pi^+ \pi^- 2\pi^0$	$(1.002 \pm 0.031) \%$	882
$4\pi^0$	$(7.6 \pm 1.1) \times 10^{-4}$	883
$\eta \pi^0$	[q] $(6.3 \pm 0.6) \times 10^{-4}$	S=1.1 846
$\omega \pi^0$	[q] $(1.17 \pm 0.35) \times 10^{-4}$	761

$\omega\eta$	$(1.98 \pm 0.18) \times 10^{-3}$	S=1.1	648
$2\pi^+ 2\pi^- \pi^0$	$(3.46 \pm 0.21) \times 10^{-3}$		844
$\pi^+ \pi^- 3\pi^0$	$(1.53 \pm 0.21) \times 10^{-3}$		847
$2\pi^+ 2\pi^- 2\pi^0$	$(4.8 \pm 0.4) \times 10^{-3}$		798
$\eta\pi^+ \pi^-$	[q] $(1.16 \pm 0.07) \times 10^{-3}$		827
$\omega\pi^+ \pi^-$	[q] $(1.33 \pm 0.20) \times 10^{-3}$		738
$\omega\pi^0 \pi^0$	$< 1.10 \times 10^{-3}$	CL=90%	740
$\eta 2\pi^0$	$(3.8 \pm 1.3) \times 10^{-4}$		829
$\pi^+ \pi^- \pi^0 \eta$	$(3.23 \pm 0.22) \times 10^{-3}$		797
$\eta 3\pi^0$	$(2.36 \pm 0.28) \times 10^{-3}$		799
$\eta 2\pi^+ 2\pi^-$	$(6.0 \pm 1.2) \times 10^{-4}$		751
$3\pi^+ 3\pi^-$	$(4.3 \pm 1.2) \times 10^{-4}$		795
$\eta'(958)\pi^0$	$(9.2 \pm 1.0) \times 10^{-4}$		678
$\eta'(958)\pi^+ \pi^-$	$(4.5 \pm 1.7) \times 10^{-4}$		650
2η	$(2.11 \pm 0.19) \times 10^{-3}$	S=2.2	754
$2\eta\pi^0$	$(7.3 \pm 2.2) \times 10^{-4}$		699
$2\eta\pi^+ \pi^-$	$(8.5 \pm 1.4) \times 10^{-4}$		623
3η	$< 1.3 \times 10^{-4}$	CL=90%	421
$\eta\eta'(958)$	$(1.01 \pm 0.19) \times 10^{-3}$		537

Hadronic modes with a $K\bar{K}$ pair

$K^+ K^-$	$(4.08 \pm 0.06) \times 10^{-3}$	S=1.6	791
$2K_S^0$	$(1.41 \pm 0.05) \times 10^{-4}$	S=1.1	789
$K_S^0 K^- \pi^+$	$(3.3 \pm 0.5) \times 10^{-3}$	S=1.1	739
$\bar{K}^*(892)^0 K_S^0, \bar{K}^{*0} \rightarrow$ $K^- \pi^+$	$(8.2 \pm 1.6) \times 10^{-5}$		608
$K^*(892)^+ K^-, K^{*+} \rightarrow$ $K_S^0 \pi^+$	$(1.89 \pm 0.30) \times 10^{-3}$		—
$\bar{K}^*(1410)^0 K_S^0, \bar{K}^{*0} \rightarrow$ $K^- \pi^+$	$(1.3 \pm 1.9) \times 10^{-4}$		—
$K^*(1410)^+ K^-, K^{*+} \rightarrow$ $K_S^0 \pi^+$	$(3.2 \pm 1.9) \times 10^{-4}$		—
$(K^- \pi^+)_{S\text{-wave}} K_S^0$	$(6.0 \pm 2.9) \times 10^{-4}$		739
$(K_S^0 \pi^+)_{S\text{-wave}} K^-$	$(3.9 \pm 1.0) \times 10^{-4}$		739
$a_0(980)^- \pi^+, a_0^- \rightarrow K_S^0 K^-$	$(1.3 \pm 1.4) \times 10^{-4}$		—
$a_0(1450)^- \pi^+, a_0^- \rightarrow$ $K_S^0 K^-$	$(2.5 \pm 2.0) \times 10^{-5}$		—
$a_2(1320)^- \pi^+, a_2^- \rightarrow$ $K_S^0 K^-$	$(5 \pm 5) \times 10^{-6}$		—
$\rho(1450)^- \pi^+, \rho^- \rightarrow K_S^0 K^-$	$(4.6 \pm 2.5) \times 10^{-5}$		—
$K_S^0 K^+ \pi^-$	$(2.17 \pm 0.34) \times 10^{-3}$	S=1.1	739
$K^*(892)^0 K_S^0, K^{*0} \rightarrow$ $K^+ \pi^-$	$(1.12 \pm 0.21) \times 10^{-4}$		608

$K^*(892)^- K^+, K^{*-} \rightarrow K_S^0 \pi^-$	$(6.2 \pm 1.0) \times 10^{-4}$	—
$K^*(1410)^0 K_S^0, K^{*0} \rightarrow K^+ \pi^+$	$(5 \pm 8) \times 10^{-5}$	—
$K^*(1410)^- K^+, K^{*-} \rightarrow K_S^0 \pi^-$	$(2.6 \pm 2.0) \times 10^{-4}$	—
$(K^+ \pi^-)_{S-wave} K_S^0$	$(3.7 \pm 1.9) \times 10^{-4}$	739
$(K_S^0 \pi^-)_{S-wave} K^+$	$(1.4 \pm 0.6) \times 10^{-4}$	739
$a_0(980)^+ \pi^-, a_0^+ \rightarrow K_S^0 K^+$	$(6 \pm 4) \times 10^{-4}$	—
$a_0(1450)^+ \pi^-, a_0^+ \rightarrow K_S^0 K^+$	$(3.2 \pm 2.5) \times 10^{-5}$	—
$\rho(1700)^+ \pi^-, \rho^+ \rightarrow K_S^0 K^+$	$(1.1 \pm 0.6) \times 10^{-5}$	—
$K^+ K^- \pi^0$	$(3.42 \pm 0.15) \times 10^{-3}$	743
$K^*(892)^+ K^-, K^*(892)^+ \rightarrow K^+ \pi^0$	$(1.52 \pm 0.08) \times 10^{-3}$	—
$K^*(892)^- K^+, K^*(892)^- \rightarrow K^- \pi^0$	$(5.4 \pm 0.4) \times 10^{-4}$	—
$(K^+ \pi^0)_{S-wave} K^-$	$(2.43 \pm 0.18) \times 10^{-3}$	743
$(K^- \pi^0)_{S-wave} K^+$	$(1.3 \pm 0.5) \times 10^{-4}$	743
$f_0(980) \pi^0, f_0 \rightarrow K^+ K^-$	$(3.6 \pm 0.6) \times 10^{-4}$	—
$\phi \pi^0, \phi \rightarrow K^+ K^-$	$(6.6 \pm 0.4) \times 10^{-4}$	—
$2K_S^0 \pi^0$	$< 1.45 \times 10^{-4}$	CL=90% 740
$K^+ K^- \eta$	$(5.9 \pm 1.9) \times 10^{-5}$	514
$\phi(1020) \eta$	$(1.84 \pm 0.12) \times 10^{-4}$	489
$K^+ K^- \eta$ nonresonant	$(9.9 \pm 0.9) \times 10^{-5}$	514
$2K_S^0 \eta$	$(1.3 \pm 0.6) \times 10^{-4}$	508
$K^+ K^- \pi^0 \pi^0$	$(6.9 \pm 0.8) \times 10^{-4}$	681
$K^+ K^- \pi^+ \pi^-$	$(2.47 \pm 0.11) \times 10^{-3}$	677
$\phi(\pi^+ \pi^-)_{S-wave}, \phi \rightarrow K^+ K^-$	$(10 \pm 5) \times 10^{-5}$	614
$(\phi \rho^0)_{S-wave}, \phi \rightarrow K^+ K^-$	$(6.9 \pm 0.6) \times 10^{-4}$	250
$(\phi \rho^0)_{P-wave}, \phi \rightarrow K^+ K^-$	$(4.0 \pm 1.9) \times 10^{-5}$	—
$(\phi \rho^0)_{D-wave}, \phi \rightarrow K^+ K^-$	$(4.2 \pm 1.4) \times 10^{-5}$	—
$(K^*(892)^0 \bar{K}^*(892)^0)_{S-wave}, K^{*0} \rightarrow K^\pm \pi^\mp$	$(2.24 \pm 0.13) \times 10^{-4}$	—
$(K^*(892)^0 \bar{K}^*(892)^0)_{P-wave}, K^* \rightarrow K^\pm \pi^\mp$	$(1.20 \pm 0.08) \times 10^{-4}$	—
$(K^*(892)^0 \bar{K}^*(892)^0)_{D-wave}, K^* \rightarrow K^\pm \pi^\mp$	$(4.7 \pm 0.4) \times 10^{-5}$	—
$K^*(892)^0 (K^- \pi^+)_{S-wave}$	$(1.4 \pm 0.6) \times 10^{-4}$	—
3-body, $K^{*0} \rightarrow K^+ \pi^-$		
$K_1(1270)^+ K^-, K_1^+ \rightarrow K^{*0} \pi^+$	$(1.4 \pm 0.9) \times 10^{-4}$	—

$K_1(1270)^+ K^-$, $K_1^+ \rightarrow K^*(1430)^0 \pi^+$, $K^{*0} \rightarrow K^+ \pi^-$	$(1.5 \pm 0.5) \times 10^{-4}$	—
$K_1(1270)^+ K^-$, $K_1^+ \rightarrow \rho^0 K^+$	$(2.2 \pm 0.6) \times 10^{-4}$	—
$K_1(1270)^+ K^-$, $K_1^+ \rightarrow \omega(782) K^+$, $\omega \rightarrow \pi^+ \pi^-$	$(1.5 \pm 1.2) \times 10^{-5}$	—
$K_1(1270)^- K^+$, $K_1^- \rightarrow \rho^0 K^-$	$(1.3 \pm 0.4) \times 10^{-4}$	—
$K_1(1400)^+ K^-$, $K_1^+ \rightarrow K^*(892)^0 \pi^+$, $K^{*0} \rightarrow K^+ \pi^-$	$(4.6 \pm 0.4) \times 10^{-4}$	—
$K^*(1410)^- K^+$, $K^{*-} \rightarrow \bar{K}^{*0} \pi^-$	$(7.0 \pm 1.1) \times 10^{-5}$	—
$K_1(1680)^+ K^-$, $K_1^+ \rightarrow K^{*0} \pi^+$, $K^{*0} \rightarrow K^+ \pi^-$	$(8.9 \pm 3.2) \times 10^{-5}$	—
$K^+ K^- \pi^+ \pi^-$ non-resonant	$(2.7 \pm 0.6) \times 10^{-4}$	—
$2K_S^0 \pi^+ \pi^-$	$(5.3 \pm 0.9) \times 10^{-4}$	673
$K_S^0 K^- \pi^+ \pi^0$	$(1.32 \pm 0.16) \times 10^{-3}$	677
$K_S^0 K^+ \pi^- \pi^0$	$(6.5 \pm 0.7) \times 10^{-4}$	677
$K_S^0 K^- 2\pi^+ \pi^-$	$< 1.4 \times 10^{-4}$	CL=90% 595
$K^+ K^- \pi^+ \pi^- \pi^0$	$(3.1 \pm 2.0) \times 10^{-3}$	600

Other $K\bar{K}X$ modes. They include all decay modes of the ϕ , η , and ω .

$\phi \pi^0$	$(1.17 \pm 0.04) \times 10^{-3}$	645
$\phi \eta$	$(1.8 \pm 0.5) \times 10^{-4}$	489
$\phi \omega$	$(6.5 \pm 1.0) \times 10^{-4}$	238

Radiative modes

$\rho^0 \gamma$	$(1.82 \pm 0.32) \times 10^{-5}$	771
$\omega \gamma$	$< 2.4 \times 10^{-4}$	CL=90% 768
$\phi \gamma$	$(2.81 \pm 0.19) \times 10^{-5}$	654
$\bar{K}^*(892)^0 \gamma$	$(4.1 \pm 0.7) \times 10^{-4}$	719

Doubly Cabibbo suppressed (DC) modes or $\Delta C = 2$ forbidden via mixing (C2M) modes

$K^+ \ell^- \bar{\nu}_\ell$ via \bar{D}^0	$[r] < 2.2 \times 10^{-5}$	CL=90% —
K^+ or $K^*(892)^+ e^- \bar{\nu}_e$ via \bar{D}^0	$< 6 \times 10^{-5}$	CL=90% —
$K^+ \pi^-$ DC	$(1.50 \pm 0.07) \times 10^{-4}$	S=3.0 861
$K^+ \pi^-$ via DCS	$(1.363 \pm 0.025) \times 10^{-4}$	—
$K^+ \pi^-$ via \bar{D}^0	$< 1.6 \times 10^{-5}$	CL=95% 861
$K_S^0 \pi^+ \pi^-$ in $D^0 \rightarrow \bar{D}^0$	$< 1.8 \times 10^{-4}$	CL=95% —
$K^*(892)^+ \pi^-$, $K^{*+} \rightarrow K_S^0 \pi^+$ DC	$(1.13 \pm_{-0.34}^{+0.60}) \times 10^{-4}$	711

$K_0^*(1430)^+ \pi^-$, $K_0^{*+} \rightarrow DC$	< 1.4	$\times 10^{-5}$		—
$K_S^0 \pi^+$				
$K_2^*(1430)^+ \pi^-$, $K_2^{*+} \rightarrow DC$	< 3.4	$\times 10^{-5}$		—
$K_S^0 \pi^+$				
$K^+ \pi^- \pi^0$	DC	$(3.06 \pm 0.16) \times 10^{-4}$	$S=1.4$	844
$K^+ \pi^- \pi^0$ via \bar{D}^0		$(7.6 \begin{smallmatrix} + 0.5 \\ - 0.6 \end{smallmatrix}) \times 10^{-4}$		—
$K^+ \pi^- 2\pi^0$		< 3.6	$\times 10^{-4}$	$CL=90\%$ 815
$K^+ \pi^+ 2\pi^-$ via DCS		$(2.49 \pm 0.07) \times 10^{-4}$		—
$K^+ \pi^+ 2\pi^-$	DC	$(2.65 \pm 0.06) \times 10^{-4}$		813
$K^+ \pi^+ 2\pi^-$ via \bar{D}^0		$(7.9 \pm 3.0) \times 10^{-6}$		812
μ^- anything via \bar{D}^0		< 4	$\times 10^{-4}$	$CL=90\%$ —

**$\Delta C = 1$ weak neutral current (C1) modes,
Lepton Family number (LF) violating modes,
Lepton (L) or Baryon (B) number violating modes**

$\gamma\gamma$	$C1$	< 8.5	$\times 10^{-7}$	$CL=90\%$	932
$e^+ e^-$	$C1$	< 7.9	$\times 10^{-8}$	$CL=90\%$	932
$\mu^+ \mu^-$	$C1$	< 6.2	$\times 10^{-9}$	$CL=90\%$	926
$\pi^0 e^+ e^-$	$C1$	< 4	$\times 10^{-6}$	$CL=90\%$	928
$\pi^0 \mu^+ \mu^-$	$C1$	< 1.8	$\times 10^{-4}$	$CL=90\%$	915
$\pi^0 \nu \bar{\nu}$		< 2.1	$\times 10^{-4}$	$CL=90\%$	928
$\eta e^+ e^-$	$C1$	< 3	$\times 10^{-6}$	$CL=90\%$	852
$\eta \mu^+ \mu^-$	$C1$	< 5.3	$\times 10^{-4}$	$CL=90\%$	838
$\pi^+ \pi^- e^+ e^-$	$C1$	< 7	$\times 10^{-6}$	$CL=90\%$	922
$\rho^0 e^+ e^-$	$C1$	< 1.0	$\times 10^{-4}$	$CL=90\%$	771
$\pi^+ \pi^- \mu^+ \mu^-$	$C1$	$(9.6 \pm 1.2) \times 10^{-7}$			894
$\pi^+ \pi^- \mu^+ \mu^-$ (non-res)		< 5.5	$\times 10^{-7}$	$CL=90\%$	—
$\rho^0 \mu^+ \mu^-$	$C1$	< 2.2	$\times 10^{-5}$	$CL=90\%$	754
$\omega e^+ e^-$	$C1$	< 6	$\times 10^{-6}$	$CL=90\%$	768
$\omega \mu^+ \mu^-$	$C1$	< 8.3	$\times 10^{-4}$	$CL=90\%$	751
$K^- K^+ e^+ e^-$	$C1$	< 1.1	$\times 10^{-5}$	$CL=90\%$	791
$\phi e^+ e^-$	$C1$	< 5.2	$\times 10^{-5}$	$CL=90\%$	654
$K^- K^+ \mu^+ \mu^-$	$C1$	$(1.54 \pm 0.32) \times 10^{-7}$			710
$K^- K^+ \mu^+ \mu^-$ (non-res)		< 3.3	$\times 10^{-5}$	$CL=90\%$	—
$\phi \mu^+ \mu^-$	$C1$	< 3.1	$\times 10^{-5}$	$CL=90\%$	631
$\bar{K}^0 e^+ e^-$		$[h] < 2.4$	$\times 10^{-5}$	$CL=90\%$	866
$\bar{K}^0 \mu^+ \mu^-$		$[h] < 2.6$	$\times 10^{-4}$	$CL=90\%$	852
$K^- \pi^+ e^+ e^-$, $675 < m_{ee} < 875$ MeV		$(4.0 \pm 0.5) \times 10^{-6}$			—
$K^- \pi^+ e^+ e^-$, $1.005 < m_{ee} < 1.035$ GeV		< 5	$\times 10^{-7}$	$CL=90\%$	—
$\bar{K}^*(892)^0 e^+ e^-$		$[h] < 4.7$	$\times 10^{-5}$	$CL=90\%$	719
$K^- \pi^+ \mu^+ \mu^-$	$C1$	< 3.59	$\times 10^{-4}$	$CL=90\%$	829

$K^- \pi^+ \mu^+ \mu^-$, $675 < m_{\mu\mu} < 875$ MeV			$(4.2 \pm 0.4) \times 10^{-6}$	—
$\bar{K}^*(892)^0 \mu^+ \mu^-$		$[h] < 2.4$	$\times 10^{-5}$	CL=90% 700
$\pi^+ \pi^- \pi^0 \mu^+ \mu^-$	CI	< 8.1	$\times 10^{-4}$	CL=90% 863
$\mu^\pm e^\mp$	LF	$[s] < 1.3$	$\times 10^{-8}$	CL=90% 929
$\pi^0 e^\pm \mu^\mp$	LF	$[s] < 8.0$	$\times 10^{-7}$	CL=90% 924
$\eta e^\pm \mu^\mp$	LF	$[s] < 2.25$	$\times 10^{-6}$	CL=90% 848
$\pi^+ \pi^- e^\pm \mu^\mp$	LF	$[s] < 1.71$	$\times 10^{-6}$	CL=90% 911
$\rho^0 e^\pm \mu^\mp$	LF	$[s] < 5.0$	$\times 10^{-7}$	CL=90% 767
$\omega e^\pm \mu^\mp$	LF	$[s] < 1.71$	$\times 10^{-6}$	CL=90% 764
$K^- K^+ e^\pm \mu^\mp$	LF	$[s] < 1.00$	$\times 10^{-6}$	CL=90% 754
$\phi e^\pm \mu^\mp$	LF	$[s] < 5.1$	$\times 10^{-7}$	CL=90% 648
$\bar{K}^0 e^\pm \mu^\mp$	LF	$[s] < 1.74$	$\times 10^{-6}$	CL=90% 863
$K^- \pi^+ e^\pm \mu^\mp$	LF	$[s] < 1.90$	$\times 10^{-6}$	CL=90% 848
$\bar{K}^*(892)^0 e^\pm \mu^\mp$	LF	$[s] < 1.25$	$\times 10^{-6}$	CL=90% 714
$2\pi^- 2e^+$	L	< 9.1	$\times 10^{-7}$	CL=90% 922
$2\pi^- 2\mu^+$	L	< 1.52	$\times 10^{-6}$	CL=90% 894
$K^- \pi^- 2e^+$	L	< 5.0	$\times 10^{-7}$	CL=90% 861
$K^- \pi^- 2\mu^+$	L	< 5.3	$\times 10^{-7}$	CL=90% 829
$2K^- 2e^+$	L	< 3.4	$\times 10^{-7}$	CL=90% 791
$2K^- 2\mu^+$	L	< 1.0	$\times 10^{-7}$	CL=90% 710
$\pi^- \pi^- e^+ \mu^+$	L	< 3.06	$\times 10^{-6}$	CL=90% 911
$K^- \pi^- e^+ \mu^+$	L	< 2.10	$\times 10^{-6}$	CL=90% 848
$2K^- e^+ \mu^+$	L	< 5.8	$\times 10^{-7}$	CL=90% 754
$p e^-$	L,B	< 2.2	$\times 10^{-6}$	CL=90% 696
$\bar{p} e^+$	L,B	< 1.2	$\times 10^{-6}$	CL=90% 696

$D^*(2007)^0$

$$I(J^P) = \frac{1}{2}(1^-)$$

I, J, P need confirmation.

Mass $m = 2006.85 \pm 0.05$ MeV ($S = 1.1$)

$m_{D^{*0}} - m_{D^0} = 142.014 \pm 0.030$ MeV ($S = 1.5$)

Full width $\Gamma < 2.1$ MeV, CL = 90%

$\bar{D}^*(2007)^0$ modes are charge conjugates of modes below.

$D^*(2007)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 \pi^0$	$(64.7 \pm 0.9) \%$	43
$D^0 \gamma$	$(35.3 \pm 0.9) \%$	137
$D^0 e^+ e^-$	$(3.91 \pm 0.33) \times 10^{-3}$	137

$D^*(2010)^\pm$

$I(J^P) = \frac{1}{2}(1^-)$
 I, J, P need confirmation.

Mass $m = 2010.26 \pm 0.05$ MeV

$m_{D^*(2010)^+} - m_{D^+} = 140.603 \pm 0.015$ MeV

$m_{D^*(2010)^+} - m_{D^0} = 145.4258 \pm 0.0017$ MeV

Full width $\Gamma = 83.4 \pm 1.8$ keV

$D^*(2010)^-$ modes are charge conjugates of the modes below.

$D^*(2010)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 \pi^+$	(67.7±0.5) %	39
$D^+ \pi^0$	(30.7±0.5) %	38
$D^+ \gamma$	(1.6±0.4) %	136

$D_0^*(2300)$

$I(J^P) = \frac{1}{2}(0^+)$

was $D_0^*(2400)$

Mass $m = 2343 \pm 10$ MeV ($S = 1.5$)

Full width $\Gamma = 229 \pm 16$ MeV

$D_0^*(2300)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D \pi^\pm$	seen	411

$D_1(2420)$

$I(J^P) = \frac{1}{2}(1^+)$

Mass $m = 2422.1 \pm 0.6$ MeV ($S = 1.7$)

$m_{D_1(2420)^0} - m_{D^{*+}} = 411.8 \pm 0.6$ MeV ($S = 1.7$)

$m_{D_1(2420)^\pm} - m_{D_1(2420)^0} = 4 \pm 4$ MeV

Full width $\Gamma = 31.3 \pm 1.9$ MeV ($S = 2.8$)

$\bar{D}_1(2420)$ modes are charge conjugates of modes below.

$D_1(2420)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^*(2007)^0 \pi$	seen	359

$D_1(2430)^0$

$$I(J^P) = \frac{1}{2}(1^+)$$

Mass $m = 2412 \pm 9$ MeV

Full width $\Gamma = 314 \pm 29$ MeV

$D_1(2430)^0$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^*(2010)^+ \pi^-$	seen	345

$D_2^*(2460)$

$$I(J^P) = \frac{1}{2}(2^+)$$

Mass $m = 2461.1^{+0.7}_{-0.8}$ MeV ($S = 6.2$)

$m_{D_2^*(2460)^0} - m_{D^+} = 591.5^{+0.7}_{-0.8}$ MeV ($S = 5.9$)

$m_{D_2^*(2460)^0} - m_{D^{*+}} = 450.9^{+0.7}_{-0.8}$ MeV ($S = 5.9$)

$m_{D_2^*(2460)^\pm} - m_{D_2^*(2460)^0} = 2.4 \pm 1.7$ MeV

Full width $\Gamma = 47.3 \pm 0.8$ MeV ($S = 1.5$)

$\bar{D}_2^*(2460)$ modes are charge conjugates of modes below.

$D_2^*(2460)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D \pi^-$	seen	509
$D^*(2010) \pi^-$	seen	389

$D_3^*(2750)$

$$I(J^P) = \frac{1}{2}(3^-)$$

Mass $m = 2763.1 \pm 3.2$ MeV ($S = 2.1$)

Full width $\Gamma = 66 \pm 5$ MeV

$D_3^*(2750)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D \pi$	seen	743
$D^+ \pi^-$	seen	739
$D^0 \pi^\pm$	seen	743
$D^* \pi$	seen	639
$D^{*+} \pi^-$	seen	639

NOTES

- [a] This result applies to $Z^0 \rightarrow c\bar{c}$ decays only. Here ℓ^+ is an average (not a sum) of e^+ and μ^+ decays.
- [b] See the Particle Listings for the (complicated) definition of this quantity.
- [c] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers in the Particle Listings.
- [d] These subfractions of the $K^- 2\pi^+$ mode are uncertain: see the Particle Listings.
- [e] See the listings under " $D \rightarrow K\pi\pi\pi$ partial wave analyses" and our 2008 Review (Physics Letters **B667** 1 (2008)) for measurements of submodes of this mode.
- [f] The unseen decay modes of the resonances are included.
- [g] This is *not* a test for the $\Delta C=1$ weak neutral current, but leads to the $\pi^+ \ell^+ \ell^-$ final state.
- [h] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.
- [i] In the 2010 *Review*, the values for these quantities were given using a measure of the asymmetry that was inconsistent with the usual definition.
- [j] This value is obtained by subtracting the branching fractions for 2-, 4- and 6-prongs from unity.
- [k] This is the sum of our $K^- 2\pi^+ \pi^-$, $K^- 2\pi^+ \pi^- \pi^0$, $\bar{K}^0 2\pi^+ 2\pi^-$, $K^+ 2K^- \pi^+$, $2\pi^+ 2\pi^-$, $2\pi^+ 2\pi^- \pi^0$, $K^+ K^- \pi^+ \pi^-$, and $K^+ K^- \pi^+ \pi^- \pi^0$, branching fractions.
- [l] This is the sum of our $K^- 3\pi^+ 2\pi^-$ and $3\pi^+ 3\pi^-$ branching fractions.
- [n] The branching fractions for the $K^- e^+ \nu_e$, $K^*(892)^- e^+ \nu_e$, $\pi^- e^+ \nu_e$, and $\rho^- e^+ \nu_e$ modes add up to 6.17 ± 0.17 %.
- [o] This is a doubly Cabibbo-suppressed mode.
- [p] Submodes of the $D^0 \rightarrow K_S^0 \pi^+ \pi^- \pi^0$ mode with a K^* and/or ρ were studied by COFFMAN 92B, but with only 140 events. With nothing new for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.
- [q] This branching fraction includes all the decay modes of the resonance in the final state.
- [r] This limit assumes the average of $B(D^0 \rightarrow K^- e^+ \nu_e)$ and $B(D^0 \rightarrow K^- \mu^+ \nu_\mu)$ for the $B(D^0 \rightarrow K^- \ell^+ \nu_\ell)$ value.
- [s] The value is for the sum of the charge states or particle/antiparticle states indicated.