

$N(1680) F_{15}$ $I(J^P) = \frac{1}{2}(\frac{5}{2}^+)$ Status: ***

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

 $N(1680)$ BREIT-WIGNER MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1675 to 1690 (≈ 1680) OUR ESTIMATE			
1684 \pm 4	MANLEY	92	IPWA $\pi N \rightarrow \pi N & N\pi\pi$
1680 \pm 10	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
1684 \pm 3	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1679 \pm 5	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
1678	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
1674 \pm 12	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
1682	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
1680	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
1660	77	IPWA $\pi N \rightarrow N\pi\pi$	
1685	KNASEL	75	DPWA $\pi^- p \rightarrow \Lambda K^0$
1670	75	IPWA $\pi N \rightarrow N\pi\pi$	

 $N(1680)$ BREIT-WIGNER WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
120 to 140 (≈ 130) OUR ESTIMATE			
139 \pm 8	MANLEY	92	IPWA $\pi N \rightarrow \pi N & N\pi\pi$
120 \pm 10	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
128 \pm 8	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
124 \pm 4	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
126	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
126 \pm 20	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$
121	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
119	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
150	77	IPWA $\pi N \rightarrow N\pi\pi$	
155	KNASEL	75	DPWA $\pi^- p \rightarrow \Lambda K^0$
130	75	IPWA $\pi N \rightarrow N\pi\pi$	

 $N(1680)$ POLE POSITION

REAL PART	DOCUMENT ID	TECN	COMMENT
1665 to 1675 (≈ 1670) OUR ESTIMATE			
1670	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
1673	93	ARGD $\pi N \rightarrow \pi N$	
1667 \pm 5	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1670	ARNDT	91	DPWA	$\pi N \rightarrow \pi N$	Soln SM90
1668 or 1674	⁴ LONGACRE	78	IPWA	$\pi N \rightarrow N\pi\pi$	
1656 or 1653	¹ LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$	

-2×IMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
105 to 135 (≈ 120) OUR ESTIMATE			
120	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
135	³ HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
110 ± 10	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
116	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
132 or 137	⁴ LONGACRE	78	IPWA $\pi N \rightarrow N\pi\pi$
145 or 143	¹ LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

N(1680) ELASTIC POLE RESIDUE

MODULUS |r|

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
40	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
44	HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
34 ± 2	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
37	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
+ 1	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
- 17	HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
-25 ± 5	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
- 14	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90

N(1680) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 N\pi$	60–70 %
$\Gamma_2 N\eta$	
$\Gamma_3 \Lambda K$	
$\Gamma_4 \Sigma K$	
$\Gamma_5 N\pi\pi$	30–40 %
$\Gamma_6 \Delta\pi$	5–15 %
$\Gamma_7 \Delta(1232)\pi, P\text{-wave}$	6–14 %
$\Gamma_8 \Delta(1232)\pi, F\text{-wave}$	<2 %
$\Gamma_9 N\rho$	3–15 %

Γ_{10}	$N\rho, S=1/2, F\text{-wave}$	
Γ_{11}	$N\rho, S=3/2, P\text{-wave}$	<12 %
Γ_{12}	$N\rho, S=3/2, F\text{-wave}$	1–5 %
Γ_{13}	$N(\pi\pi)_{S\text{-wave}}^{I=0}$	5–20 %
Γ_{14}	$p\gamma$	0.21–0.32 %
Γ_{15}	$p\gamma, \text{ helicity}=1/2$	0.001–0.011 %
Γ_{16}	$p\gamma, \text{ helicity}=3/2$	0.20–0.32 %
Γ_{17}	$n\gamma$	0.021–0.046 %
Γ_{18}	$n\gamma, \text{ helicity}=1/2$	0.004–0.029 %
Γ_{19}	$n\gamma, \text{ helicity}=3/2$	0.01–0.024 %

$N(1680)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.6 to 0.7 OUR ESTIMATE				
0.70±0.03	MANLEY	92	IPWA $\pi N \rightarrow \pi N & N\pi\pi$	
0.62±0.05	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$	
0.65±0.02	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.68	ARNDT	95	DPWA $\pi N \rightarrow N\pi$	
0.69±0.04	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$	

$(\Gamma_f;\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1680) \rightarrow N\eta$

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	BAKER	79	DPWA $\pi^- p \rightarrow n\eta$	

$\Gamma(N\eta)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.01 ± 0.004	BATINIC	95	DPWA $\pi N \rightarrow N\pi, N\eta$	
0.0005 or 0.001	5 CARRERAS	70	MPWA t pole + resonance	
0.0004	5 BOTKE	69	MPWA t pole + resonance	
0.003 ± 0.002	5 DEANS	69	MPWA t pole + resonance	

$\Gamma(N\eta)/\Gamma(N\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.027	HEUSCH	66	RVUE π^0, η photoproduction	

$(\Gamma_f;\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1680) \rightarrow \Lambda K$

Coupling to ΛK not required in the analyses of BAKER 77, SAXON 80, or BELL 83.

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.01	KNASEL	75	DPWA $\pi^- p \rightarrow \Lambda K^0$	
-0.009 ± 0.009	DEVENISH	74B	Fixed- t dispersion rel.	

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1680) \rightarrow \Sigma K$	$(\Gamma_1 \Gamma_4)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.001	6 DEANS	75 DPWA	$\pi N \rightarrow \Sigma K$

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1680) \rightarrow \Delta(1232)\pi$, P-wave	$(\Gamma_1 \Gamma_7)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.31 to -0.21 OUR ESTIMATE			
-0.26 ± 0.04	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
-0.27	1,7 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
-0.25	2 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.38	8 NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1680) \rightarrow \Delta(1232)\pi$, F-wave	$(\Gamma_1 \Gamma_8)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
+0.03 to +0.11 OUR ESTIMATE			
+0.07 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
+0.07	1,7 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.08	2 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.05	8 NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1680) \rightarrow N\rho, S=3/2$, P-wave	$(\Gamma_1 \Gamma_{11})^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.30 to -0.10 OUR ESTIMATE			
-0.20 ± 0.05	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
-0.23	1,7 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
-0.30	2 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.34	8 NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1680) \rightarrow N\rho, S=3/2$, F-wave	$(\Gamma_1 \Gamma_{12})^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.18 to -0.10 OUR ESTIMATE			
-0.13 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
-0.15	1,7 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_f; \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1680) \rightarrow N(\pi\pi)_{S\text{-wave}}^{I=0}$		$(\Gamma_1 \Gamma_{13})^{1/2} / \Gamma$	
VALUE	DOCUMENT ID	TECN	COMMENT
+0.25 to +0.35 OUR ESTIMATE			
+0.29 ± 0.04	MANLEY	92	IPWA $\pi N \rightarrow \pi N & N\pi\pi$
+0.31	^{1,7} LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
+0.30	² LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.42	⁸ NOVOSELLER	78	IPWA $\pi N \rightarrow N\pi\pi$

 $N(1680)$ PHOTON DECAY AMPLITUDES **$N(1680) \rightarrow p\gamma$, helicity-1/2 amplitude $A_{1/2}$**

VALUE (GeV $^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
-0.015 ± 0.006 OUR ESTIMATE			
-0.010 ± 0.004	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
-0.017 ± 0.018	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
-0.009 ± 0.006	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.028 ± 0.003	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
-0.026 ± 0.003	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
-0.018 ± 0.014	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.006 ± 0.002	LI	93	IPWA $\gamma N \rightarrow \pi N$
-0.005 ± 0.015	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
-0.009 ± 0.002	FELLER	76	DPWA $\gamma N \rightarrow \pi N$

 $N(1680) \rightarrow p\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE (GeV $^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
+0.133 ± 0.012 OUR ESTIMATE			
0.145 ± 0.005	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
0.132 ± 0.010	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
0.115 ± 0.008	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
0.115 ± 0.003	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
0.122 ± 0.003	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
0.141 ± 0.014	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.154 ± 0.002	LI	93	IPWA $\gamma N \rightarrow \pi N$
+0.138 ± 0.021	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
+0.121 ± 0.010	FELLER	76	DPWA $\gamma N \rightarrow \pi N$

 $N(1680) \rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV $^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
+0.029 ± 0.010 OUR ESTIMATE			
0.030 ± 0.005	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
0.017 ± 0.014	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
0.032 ± 0.003	FUJII	81	DPWA $\gamma N \rightarrow \pi N$
0.026 ± 0.005	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
0.028 ± 0.014	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
0.044 ± 0.012	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
0.025 ± 0.010	TAKEDA	80	DPWA $\gamma N \rightarrow \pi N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.022±0.002	LI	93	IPWA	$\gamma N \rightarrow \pi N$
+0.037±0.010	BARBOUR	78	DPWA	$\gamma N \rightarrow \pi N$

N(1680) → $n\gamma$, helicity-3/2 amplitude A_{3/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.033±0.009 OUR ESTIMATE			
-0.040±0.015	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
-0.033±0.013	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.023±0.005	FUJII	81	DPWA $\gamma N \rightarrow \pi N$
-0.024±0.009	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
-0.029±0.017	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
-0.033±0.015	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
-0.035±0.012	TAKEDA	80	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.048±0.002	LI	93	IPWA $\gamma N \rightarrow \pi N$
-0.038±0.018	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$

N(1680) FOOTNOTES

¹ LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

² From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

³ See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

⁴ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.

⁵ The parametrization used may be double counting.

⁶ The range given is from 3 of 4 best solutions; not present in solution 1. DEANS 75 disagrees with $\pi^+ p \rightarrow \Sigma^+ K^+$ data of WINNIK 77 around 1920 MeV.

⁷ LONGACRE 77 considers this coupling to be well determined.

⁸ A Breit-Wigner fit to the HERNDON 75 IPWA.

N(1680) REFERENCES

For early references, see Physics Letters **111B** 70 (1982). For very early references, see Reviews of Modern Physics **37** 633 (1965).

ARNDT	96	PR C53 430	+Strakovsky, Workman	(VPI)
ARNDT	95	PR C52 2120	+Strakovsky, Workman, Pavan	(VPI, BRCO)
BATINIC	95	PR C51 2310	+Slaus, Svarc, Nefkens	(BOSK, UCLA)
HOEHLER	93	πN Newsletter 9 1		(KARL)
LI	93	PR C47 2759	+Arndt, Roper, Workman	(VPI)
MANLEY	92	PR D45 4002	+Saleski	(KENT) IJP
Also	84	PR D30 904	Manley, Arndt, Goradia, Teplitz	(VPI)
ARNDT	91	PR D43 2131	+Li, Roper, Workman, Ford	(VPI, TELE) IJP
BELL	83	NP B222 389	+Blissett, Broome, Daley, Hart, Lintern+	(RL) IJP
CRAWFORD	83	NP B211 1	+Morton	(GLAS)
PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
AWAJI	81	Bonn Conf. 352	+Kajikawa	(NAGO)
Also	82	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+	(NAGO)
FUJII	81	NP B187 53	+Hayashii, Iwata, Kajikawa+	(NAGO, OSAK)
ARAI	80	Toronto Conf. 93	Arai, Fujii	(INUS)
Also	82	NP B194 251		(INUS)

CRAWFORD	80	Toronto Conf. 107		(GLAS)
CUTKOSKY	80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP
SAXON	80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+	(RHEL, BRIS) IJP
TAKEDA	80	NP B168 17	+Arai, Fujii, Ikeda, Iwasaki+	(TOKY, INUS)
BAKER	79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+	(RHEL) IJP
HOEHLER	79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80	Toronto Conf. 3	Koch	(KARLT) IJP
BARBOUR	78	NP B141 253	+Crawford, Parsons	(GLAS)
LONGACRE	78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+	(LBL, SLAC)
NOVOSELLER	78	NP B137 509	Novoseller	(CIT) IJP
Also	78B	NP B137 445	+Blissett, Bloodworth, Broome, Hart+	(CIT) IJP
BAKER	77	NP B126 365	+Dolbeau	(RHEL) IJP
LONGACRE	77	NP B122 493	Dolbeau, Triantis, Neveu, Cadet	(SACL) IJP
Also	76	NP B108 365	+Toaff, Revel, Goldberg, Berny	(HAIF) I
WINNIK	77	NP B128 66	+Fukushima, Horikawa, Kajikawa+	(NAGO, OSAK) IJP
FELLER	76	NP B104 219	+Mitchell, Montgomery+	(SFLA, ALAH) IJP
DEANS	75	NP B96 90	+Longacre, Miller, Rosenfeld+	(LBL, SLAC)
HERNDON	75	PR D11 3183	+Lindquist, Nelson+	(CHIC, WUSL, OSU, ANL) IJP
KNASEL	75	PR D11 1	Rosenfeld, Lasinski, Smadja+	(LBL, SLAC) IJP
LONGACRE	75	PL 55B 415	+Froggatt, Martin	(DESY, NORD, LOUC)
DEVENISH	74B	NP B81 330	+Donnachie	(DARE, MCHS)
CARRERAS	70	NP B16 35	+Wooten	(UCSB)
BOTKE	69	PR 180 1417	+Prescott, Dashen	(SFLA)
DEANS	69	PR 185 1797		(CIT)
HEUSCH	66	PRL 17 1019		
