


$f_2(2150)$ $I^G(J^{PC}) = 0^+(2^{++})$

OMMITTED FROM SUMMARY TABLE

This entry was previously called T_0 . **$f_2(2150)$ MASS** **$f_2(2150)$ MASS, COMBINED MODES (MeV)**VALUE (MeV)DOCUMENT ID

2138±18 OUR AVERAGE Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.5. See the ideogram below.

 **$f_2(2150)$ MASS, COMBINED MODES (MeV)** **$\eta\eta$ MODE**VALUE (MeV)DOCUMENT IDTECNCOMMENT

The data in this block is included in the average printed for a previous datablock.

2138±23 OUR AVERAGE Error includes scale factor of 1.8. See the ideogram below.

2175±20


PROKOSHKIN 95D GAM4 $300 \pi^- N \rightarrow \pi^- N 2\eta$,
450 $p p \rightarrow p p 2\eta$

2130±35

SINGOVSKI 94 GAM4 450 $p p \rightarrow p p 2\eta$

2104±20

¹ ARMSTRONG 93C E760 $\bar{p} p \rightarrow \pi^0 \eta\eta \rightarrow 6\gamma$ 1 No J^{PC} determination.

 $f_2(2150)$ MASS, $\eta\eta$ MODE (MeV) $\eta\pi\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.				

2135±20±45 ADOMEIT 96 CBAR 0 1.94 $\bar{p}p \rightarrow \eta 3\pi^0$

 $\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~2226	HASAN 94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~2090	² OAKDEN 94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~2120	³ OAKDEN 94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~2170	⁴ MARTIN 80B	RVUE	
~2150	⁴ MARTIN 80C	RVUE	
~2150	⁵ DULUDE 78B	OSPK	1–2 $\bar{p}p \rightarrow \pi^0 \pi^0$

²OAKDEN 94 makes an amplitude analysis of LEAR data on $\bar{p}p \rightarrow \pi\pi$ using a method based on Barrelet zeros. This is solution A. The amplitude analysis of HASAN 94 includes earlier data as well, and assume that the data can be parametrized in terms of towers of nearly degenerate resonances on the leading Regge trajectory. See also KLOET 96 and MARTIN 97 who make related analyses.

³From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$.

⁴ $I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.


⁵ $I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.

S-CHANNEL $\bar{p}p$, $\bar{N}N$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2139 ± 8	⁶ EVANGELISTA 97 SPEC			$0.6\text{--}2.4 \bar{p}p \rightarrow K_S^0 K_S^0$
~ 2190	⁷ CUTTS	78B CNTR		$0.97\text{--}3 \bar{p}p \rightarrow \bar{N}N$
2155 ± 15	^{7,8} COUPLAND	77 CNTR	0	$0.7\text{--}2.4 \bar{p}p \rightarrow \bar{p}p$
2193 ± 2	^{7,9} ALSPECTOR	73 CNTR		$\bar{p}p$ S channel
⁶ Isospin 0 and 2 not separated.				
⁷ Isospins 0 and 1 not separated.				
⁸ From a fit to the total elastic cross section.				
⁹ Referred to as T or T region by ALSPECTOR 73.				

 $f_2(2150)$ WIDTH **$f_2(2150)$ WIDTH, COMBINED MODES (MeV)**

VALUE (MeV)	DOCUMENT ID
194 ± 15 OUR AVERAGE	Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.6. See the ideogram below.

 **$f_2(2150)$ WIDTH, COMBINED MODES (MeV)**

$\eta\eta$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

193±17 OUR AVERAGE Error includes scale factor of 1.9. See the ideogram below.

150±35

PROKOSHKIN 95D GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$,
450 $p\bar{p} \rightarrow p\bar{p} 2\eta$


130±30

SINGOVSKI 94 GAM4 450 $p\bar{p} \rightarrow p\bar{p} 2\eta$

203±10

10 ARMSTRONG 93C E760 $\bar{p}p \rightarrow \pi^0 \eta\eta \rightarrow 6\gamma$

10 No J^{PC} determination.



$f_2(2150)$ WIDTH, $\eta\eta$ MODE (MeV)

 $\eta\pi\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

250±25±45

ADOMEIT 96 CBAR 0 1.94 $\bar{p}p \rightarrow \eta 3\pi^0$

 $\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

250 OUR ESTIMATE

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 226

HASAN 94 RVUE $\bar{p}p \rightarrow \pi\pi$

~ 70

11 OAKDEN 94 RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$

~ 250

12 MARTIN 80B RVUE

~ 250

12 MARTIN 80C RVUE

~ 250

13 DULUDE 78B OSPK 1–2 $\bar{p}p \rightarrow \pi^0 \pi^0$

¹¹ See however KLOE T 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

¹² $I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.

¹³ $I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.

S-CHANNEL $\bar{p}p$, $\bar{N}N$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
56^{+31}_{-16}	¹⁴ EVANGELISTA 97	SPEC		$0.6-2.4 \bar{p}p \rightarrow K_S^0 K_S^0$
135 ± 75	^{15,16} COUPLAND	77	CNTR	$0.7-2.4 \bar{p}p \rightarrow \bar{p}p$
98 ± 8	¹⁶ ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
¹⁴ Isospin 0 and 2 not separated.				
¹⁵ From a fit to the total elastic cross section.				
¹⁶ Isospins 0 and 1 not separated.				

$f_2(2150)$ DECAY MODES

Mode
$\Gamma_1 \pi\pi$
$\Gamma_2 \eta\eta$
$\Gamma_3 \bar{K}K$
$\Gamma_4 f_2(1270)\eta$
$\Gamma_5 a_2(1320)\pi$

$f_2(2150)$ BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma(\eta\eta)$	Γ_3/Γ_2			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.1	95	¹⁷ PROKOSHIN 95D GAM4	300	$\pi^- N \rightarrow \pi^- N 2\eta$, 450 $p\bar{p} \rightarrow p\bar{p} 2\eta$

¹⁷ Using data from ARMSTRONG 89D.

$\Gamma(\pi\pi)/\Gamma(\eta\eta)$	Γ_1/Γ_2			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.33	95	¹⁸ PROKOSHIN 95D GAM4	300	$\pi^- N \rightarrow \pi^- N 2\eta$, 450 $p\bar{p} \rightarrow p\bar{p} 2\eta$

¹⁸ Derived from a $\pi^0 \pi^0 / \eta\eta$ limit.

$\Gamma(f_2(1270)\eta)/\Gamma(a_2(1320)\pi)$	Γ_4/Γ_5		
VALUE	DOCUMENT ID	TECN	COMMENT
0.79 ± 0.11	¹⁹ ADOMEIT	96	CBAR $1.94 \bar{p}p \rightarrow \eta 3\pi^0$
¹⁹ Using $B(a_2(1320) \rightarrow \eta\pi) = 0.145$			

f₂(2150) REFERENCES

EVANGELISTA	97	PR D56 3803	C. Evangelista, Palano, Drijard+	(LEAR Collab.)
MARTIN	97	PR C56 1114	B.R. Martin, Oades	(LOUC, AARH)
ADOMEIT	96	ZPHY C71 227	+Amsler, Armstrong+	(Crystal Barrel Collab.)
KLOET	96	PR D53 6120	+Myhrer	(RUTG, NORD)
PROKOSHKIN	95D	SPD 40 495		(SERP) IGJPC
		Translated from DANS 344 469.		
HASAN	94	PL B334 215	+Bugg	(LOQM)
OAKDEN	94	NPA 574 731	+Pennington	(DURH)
SINGOVSKI	94	NC 107 1911		(SERP)
ARMSTRONG	93C	PL B307 394	+Bettoni+	(FNAL, FERR, GENO, UCI, NWES+)
ARMSTRONG	89D	PL B227 186	+Benayoun	(ATHU, BARI, BIRM, CERN, CDEF)
MARTIN	80B	NP B176 355	+Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	+Pennington	(DURH) JP
CUTTS	78B	PR D17 16	+Good, Grannis, Green, Lee+	(STON, WISC)
DULUDE	78B	PL 79B 335	+Lanou, Massimo, Peaslee+	(BROW, MIT, BARI) JP
COUPLAND	77	PL 71B 460	+Eisenhandler, Gibson, Astbury+	(LOQM, RHEL)
ALSPECTOR	73	PRL 30 511	+Cohen, Cvijanovich+	(RUTG, UPNJ)

OTHER RELATED PAPERS

FIELDS	71	PRL 27 1749	+Cooper, Rhines, Allison	(ANL, OXF)
YOH	71	PRL 26 922	+Barish, Carroll, Loblkowicz+	(CIT, BNL, ROCH)
