

X(1910)

$$I^G(J^{PC}) = 0^{+}(?^{?+})$$

OMITTED FROM SUMMARY TABLE

We list here two different peaks with close masses and widths seen in the mass distributions of $\omega\omega$ and $\eta\eta'$ final states. ALDE 91B argues that they are of different nature.

X(1910) MASS

VALUE (MeV)
1810 to 1920 OUR ESTIMATE

DOCUMENT ID**X(1910) $\omega\omega$ MODE**VALUE (MeV)DOCUMENT IDTECNCOMMENT**1921 ± 8 OUR AVERAGE**

1920 ± 10

¹ BELADIDZE 92B VES 36 $\pi^- p \rightarrow \omega\omega n$

1924 ± 14

¹ ALDE 90 GAM2 38 $\pi^- p \rightarrow \omega\omega n$ $^1 J^{PC} = 2^{++}$.**X(1910) $\eta\eta'$ MODE**VALUE (MeV)DOCUMENT IDTECNCOMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

1911 ± 10

ALDE 91B GAM2 38 $\pi^- p \rightarrow \eta\eta' n$ **X(1910) WIDTH**VALUE (MeV)DOCUMENT ID**90 to 250 OUR ESTIMATE****X(1910) $\omega\omega$ MODE**VALUE (MeV)DOCUMENT IDTECNCOMMENT**90 ± 19 OUR AVERAGE**

90 ± 20

² BELADIDZE 92B VES 36 $\pi^- p \rightarrow \omega\omega n$

91 ± 50

² ALDE 90 GAM2 38 $\pi^- p \rightarrow \omega\omega n$ $^2 J^{PC} = 2^{++}$.**X(1910) $\eta\eta'$ MODE**VALUE (MeV)DOCUMENT IDTECNCOMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

90 ± 35

ALDE 91B GAM2 38 $\pi^- p \rightarrow \eta\eta' n$

X(1910) DECAY MODES

Mode
$\Gamma_1 \quad \pi^0 \pi^0$
$\Gamma_2 \quad K_S^0 K_S^0$
$\Gamma_3 \quad \eta \eta$
$\Gamma_4 \quad \omega \omega$
$\Gamma_5 \quad \eta \eta'$
$\Gamma_6 \quad \eta' \eta'$

X(1910) BRANCHING RATIOS

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen ALDE 89B GAM2 $38 \pi^- p \rightarrow \omega \omega n$

$\Gamma(\pi^0 \pi^0)/\Gamma(\eta \eta')$ Γ_1/Γ_5

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.1 ALDE 89 GAM2 $38 \pi^- p \rightarrow \eta \eta' n$

$\Gamma(\eta \eta)/\Gamma(\eta \eta')$ Γ_3/Γ_5

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.05 90 ALDE 91B GAM2 $38 \pi^- p \rightarrow \eta \eta' n$

$\Gamma(K_S^0 K_S^0)/\Gamma(\eta \eta')$ Γ_2/Γ_5

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.066 90 BALOSHIN 86 SPEC $40 \pi p \rightarrow K_S^0 K_S^0 n$

$\Gamma(\eta' \eta')/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen BELADIDZE 92D VES $37 \pi^- p \rightarrow \eta' \eta' n$

X(1910) REFERENCES

BELADIDZE	92B	ZPHY C54 367	+Bityukov, Borisov+	(VES Collab.)
BELADIDZE	92D	ZPHY C57 13	+Berdnikov+	(VES Collab.)
ALDE	91B	SJNP 54 455	+Binon+ (SERP, BELG, LANL, LAPP, PISA, KEK)	
		Translated from YAF 54 751.		
Also	92	PL B276 375	Alde, Binon+ (BELG, SERP, KEK, LANL, LAPP)	
ALDE	90	PL B241 600	+Binon+ (SERP, BELG, LANL, LAPP, PISA, KEK)	
ALDE	89	PL B216 447	+Binon, Bricman, Donskov+ (SERP, BELG, LANL, LAPP)	
Also	88E	SJNP 48 1035	Alde, Binon, Bricman+ (BELG, SERP, LANL, LAPP)	
		Translated from YAF 48 1724.		
ALDE	89B	PL B216 451	+Binon, Bricman+ (SERP, BELG, LANL, LAPP, TBIL)	
BALOSHIN	86	SJNP 43 959	+Barkov, Bolonkin, Vladimirkii, Grigoriev+ (ITEP)	
		Translated from YAF 43 1487.		

————— **OTHER RELATED PAPERS** —————

LEE 94 PL B323 227 +Chung, Kirk+ (BNL, IND, KYUN, MASD, RICE)
