

**$\tau$  – THIS IS PART 3 OF 4**

To reduce the size of this section's PostScript file, we have divided it into four PostScript files. We present the following index:

## PART 1

Page #	Section name
1	Mass
1	Mean life
2	Magnetic moment anomaly
2	Electric dipole moment
3	Weak dipole moment
4	Decay modes

## PART 2

Page #	Section name
12	Branching fractions
19	Branching ratios
19	Modes with one charged particle

## PART 3

Page #	Section name
29	Modes with one charged particle (cont.)
33	Modes with $K^0$ 's
38	Modes with three charged particle
47	Modes with five charged particle
48	Misc. other allowedmodes

## PART 4

Page #	Section name
55	Misc. other allowedmodes (cont.)
56	Forbidden modes
64	Decay parameters
70	References


$$\Gamma(h^- \geq 2\pi^0 \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{16} / \Gamma$$

$$\Gamma_{16} / \Gamma = (\Gamma_{19} + \Gamma_{20} + \Gamma_{23} + \Gamma_{24} + \Gamma_{26} + 0.157\Gamma_{32} + 0.157\Gamma_{34} + 0.157\Gamma_{36} + 0.157\Gamma_{38} + 0.0246\Gamma_{41} + 0.319\Gamma_{110}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.79 ± 0.16 OUR FIT</b>					Error includes scale factor of 1.2.
<b>10.3 ± 1.1 OUR AVERAGE</b>					Error includes scale factor of 2.9. See the ideogram below.
9.91 ± 0.31 ± 0.27	f&a		ACKERSTAFF	98M OPAL	1991–1995 LEP runs
14.0 ± 1.2 ± 0.6	avg	938	<sup>60</sup> BEHREND	90 CELL	$E_{\text{cm}}^{\text{ee}} = 35 \text{ GeV}$
12.0 ± 1.4 ± 2.5	f&a		<sup>61</sup> BURCHAT	87 MRK2	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
9.89 ± 0.34 ± 0.55			<sup>62</sup> AKERS	94E OPAL	Repl. by ACKERSTAFF 98M
13.9 ± 2.0 $\begin{smallmatrix} +1.9 \\ -2.2 \end{smallmatrix}$			<sup>63</sup> AIHARA	86E TPC	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$

- <sup>60</sup> No independent of BEHREND 90  $\Gamma(h^- 2\pi^0 \nu_\tau (\text{exp. } K^0))$  and  $\Gamma(h^- \geq 3\pi^0 \nu_\tau)$ .
- <sup>61</sup> Error correlated with BURCHAT 87  $\Gamma(\rho^- \nu_e) / \Gamma(\text{total})$  value.
- <sup>62</sup> AKERS 94E not independent of AKERS 94E  $B(h^- \geq 1\pi^0 \nu_\tau)$  and  $B(h^- \pi^0 \nu_\tau)$  measurements.
- <sup>63</sup> AIHARA 86E (TPC) quote  $B(2\pi^0 \pi^- \nu_\tau) + 1.6B(3\pi^0 \pi^- \nu_\tau) + 1.1B(\pi^0 \eta \pi^- \nu_\tau)$ .



$$\Gamma(h^- \geq 2\pi^0 \nu_\tau) / \Gamma_{\text{total}} (\%)$$

$$\frac{\Gamma(h^- 2\pi^0 \nu_\tau)/\Gamma_{\text{total}}}{\Gamma_{17}/\Gamma = (\Gamma_{19} + \Gamma_{20} + 0.157\Gamma_{32} + 0.157\Gamma_{34})/\Gamma} \quad \Gamma_{17}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.39±0.14 OUR FIT</b>	Error includes scale factor of 1.2.			
<b>9.48±0.13±0.10</b>	12k	<sup>64</sup> BUSKULIC	96	ALEP LEP 1991–1993 data
<sup>64</sup> BUSKULIC 96 quote $9.29 \pm 0.13 \pm 0.10$ . We add 0.19 to undo their correction for $\tau^- \rightarrow h^- K^0 \nu_\tau$ .				

$$\frac{\Gamma(h^- 2\pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}}{\Gamma_{18}/\Gamma = (\Gamma_{19} + \Gamma_{20})/\Gamma} \quad \Gamma_{18}/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. f&a marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.23±0.14 OUR FIT</b>	Error includes scale factor of 1.2.			
<b>8.95±0.33 OUR AVERAGE</b>	Error includes scale factor of 1.1.			
$8.88 \pm 0.37 \pm 0.42$	f&a 1060	ACCIARRI	95 L3	1992 LEP run
$8.96 \pm 0.16 \pm 0.44$	avg	<sup>65</sup> PROCARIO	93 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
$10.38 \pm 0.66 \pm 0.82$	f&a 809	<sup>66</sup> DECAMP	92C ALEP	1989–1990 LEP runs
$5.7 \pm 0.5 \pm \begin{smallmatrix} +1.7 \\ -1.0 \end{smallmatrix}$	f&a 133	<sup>67</sup> ANTREASYAN	91 CBAL	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV
$10.0 \pm 1.5 \pm 1.1$	f&a 333	<sup>68</sup> BEHREND	90 CELL	$E_{\text{cm}}^{ee} = 35$ GeV
$8.7 \pm 0.4 \pm 1.1$	f&a 815	<sup>69</sup> BAND	87 MAC	$E_{\text{cm}}^{ee} = 29$ GeV
$6.0 \pm 3.0 \pm 1.8$	f&a	BEHREND	84 CELL	$E_{\text{cm}}^{ee} = 14,22$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.2 \pm 0.6 \pm 1.2$  <sup>70</sup> GAN 87 MRK2  $E_{\text{cm}}^{ee} = 29$  GeV

<sup>65</sup> PROCARIO 93 entry is obtained from  $B(h^- 2\pi^0 \nu_\tau)/B(h^- \pi^0 \nu_\tau)$  using ARTUSO 94 result for  $B(h^- \pi^0 \nu_\tau)$ .

<sup>66</sup> We subtract 0.0015 to account for  $\tau^- \rightarrow K^*(892)^- \nu_\tau$  contribution.

<sup>67</sup> ANTREASYAN 91 subtract 0.001 to account for the  $\tau^- \rightarrow K^*(892)^- \nu_\tau$  contribution.

<sup>68</sup> BEHREND 90 subtract 0.002 to account for the  $\tau^- \rightarrow K^*(892)^- \nu_\tau$  contribution.

<sup>69</sup> BAND 87 assume  $B(\pi^- 3\pi^0 \nu_\tau) = 0.01$  and  $B(\pi^- \pi^0 \eta \nu_\tau) = 0.005$ .

<sup>70</sup> GAN 87 analysis use photon multiplicity distribution.

$$\frac{\Gamma(h^- 2\pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma(h^- \pi^0 \nu_\tau)}{\Gamma_{18}/\Gamma_{12} = (\Gamma_{19} + \Gamma_{20})/(\Gamma_{13} + \Gamma_{15})} \quad \Gamma_{18}/\Gamma_{12}$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.357±0.006 OUR FIT</b>	Error includes scale factor of 1.2.		
<b>0.342±0.006±0.016</b>	<sup>71</sup> PROCARIO	93 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

<sup>71</sup> PROCARIO 93 quote  $0.345 \pm 0.006 \pm 0.016$  after correction for 2 kaon backgrounds assuming  $B(K^{*-} \nu_\tau) = 1.42 \pm 0.18\%$  and  $B(h^- K^0 \pi^0 \nu_\tau) = 0.48 \pm 0.48\%$ . We multiply by  $0.990 \pm 0.010$  to remove these corrections to  $B(h^- \pi^0 \nu_\tau)$ .

$$\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}} \qquad \Gamma_{19}/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		DOCUMENT ID	TECN	COMMENT
-----------	--	-------------	------	---------

**9.15±0.15 OUR FIT** Error includes scale factor of 1.2.

**9.21±0.13±0.11** avg <sup>72</sup>BUSKULIC 96 ALEP LEP 1991–1993 data

<sup>72</sup>Not independent of BUSKULIC 96  $B(h^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$  and  $B(K^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$  values.

$$\Gamma(K^- 2\pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}} \qquad \Gamma_{20}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

**0.080±0.027 OUR FIT**

**0.081±0.027 OUR AVERAGE**

0.08 ±0.02 ±0.02 59 BUSKULIC 96 ALEP LEP 1991–1993 data

0.09 ±0.10 ±0.03 3 <sup>73</sup>BATTLE 94 CLEO  $E_{\text{cm}}^{ee} \approx 10.6$  GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.04 ±0.03 ±0.02 11 BUSKULIC 94E ALEP Repl. by BUSKULIC 96

<sup>73</sup>BATTLE 94 quote  $0.14 \pm 0.10 \pm 0.03$  or  $< 0.3\%$  at 90% CL. We subtract  $(0.05 \pm 0.02)\%$  to account for  $\tau^- \rightarrow K^- (K^0 \rightarrow \pi^0 \pi^0) \nu_\tau$  background.

$$\Gamma(h^- \geq 3\pi^0 \nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{21}/\Gamma$$

$$\Gamma_{21}/\Gamma = (\Gamma_{23} + \Gamma_{24} + \Gamma_{26} + 0.157\Gamma_{36} + 0.157\Gamma_{38} + 0.0246\Gamma_{41} + 0.319\Gamma_{110})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

**1.40±0.11 OUR FIT** Error includes scale factor of 1.1.

**1.8 ±0.6 OUR AVERAGE** Error includes scale factor of 1.1.

1.53±0.40±0.46 f&a 186 DECAMP 92C ALEP 1989–1990 LEP

3.2 ±1.0 ±1.0 f&a BEHREND 90 CELL  $E_{\text{cm}}^{ee} = 35$  GeV

$$\Gamma(h^- 3\pi^0 \nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{22}/\Gamma$$

$$\Gamma_{22}/\Gamma = (\Gamma_{23} + \Gamma_{24} + 0.157\Gamma_{36} + 0.157\Gamma_{38})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

**1.23±0.10 OUR FIT** Error includes scale factor of 1.1.

**1.22±0.10 OUR AVERAGE**

1.24±0.09±0.11 f&a 2.3k <sup>74</sup>BUSKULIC 96 ALEP LEP 1991–1993 data

1.70±0.24±0.38 f&a 293 ACCIARRI 95 L3 1992 LEP run

1.15±0.08±0.13 avg <sup>75</sup>PROCARIO 93 CLEO  $E_{\text{cm}}^{ee} \approx 10.6$  GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0  $\begin{matrix} +1.4 & +1.1 \\ -0.1 & -0.1 \end{matrix}$  <sup>76</sup>GAN 87 MRK2  $E_{\text{cm}}^{ee} = 29$  GeV

<sup>74</sup>BUSKULIC 96 quote  $B(h^- 3\pi^0 \nu_\tau (\text{ex. } K^0)) = 1.17 \pm 0.09 \pm 0.11$ . We add 0.07 to remove their correction for  $K^0$  backgrounds.

<sup>75</sup> PROCARIO 93 entry is obtained from  $B(h^- 3\pi^0 \nu_\tau)/B(h^- \pi^0 \nu_\tau)$  using ARTUSO 94 result for  $B(h^- \pi^0 \nu_\tau)$ .

<sup>76</sup> Highly correlated with GAN 87  $\Gamma(\eta\pi^-\pi^0 \nu_\tau)/\Gamma_{\text{total}}$  value. Authors quote  $B(\pi^\pm 3\pi^0 \nu_\tau) + 0.67B(\pi^\pm \eta\pi^0 \nu_\tau) = 0.047 \pm 0.010 \pm 0.011$ .

$$\frac{\Gamma(h^- 3\pi^0 \nu_\tau)/\Gamma(h^- \pi^0 \nu_\tau)}{\Gamma_{22}/\Gamma_{12}} = (\Gamma_{23} + \Gamma_{24} + 0.157\Gamma_{36} + 0.157\Gamma_{38})/(\Gamma_{13} + \Gamma_{15})$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.048±0.004 OUR FIT</b>			Error includes scale factor of 1.1.
<b>0.044±0.003±0.005</b>	<sup>77</sup> PROCARIO 93	CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

<sup>77</sup> PROCARIO 93 quote  $0.041 \pm 0.003 \pm 0.005$  after correction for 2 kaon backgrounds assuming  $B(K^{*-} \nu_\tau) = 1.42 \pm 0.18\%$  and  $B(h^- K^0 \pi^0 \nu_\tau) = 0.48 \pm 0.48\%$ . We add  $0.003 \pm 0.003$  and multiply the sum by  $0.990 \pm 0.010$  to remove these corrections.

$$\frac{\Gamma(\pi^- 3\pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}}{\Gamma_{23}/\Gamma}$$

VALUE (%)	DOCUMENT ID
<b>1.11±0.14 OUR FIT</b>	

$$\frac{\Gamma(K^- 3\pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}}{\Gamma_{24}/\Gamma}$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>0.043<sup>+0.100</sup><sub>-0.029</sub> OUR FIT</b>			
<b>0.05 ±0.13</b>	<sup>78</sup> BUSKULIC 94E	ALEP	1991-1992 LEP runs

<sup>78</sup> BUSKULIC 94E quote  $B(K^- \geq 0\pi^0 \geq 0K^0 \nu_\tau) - [B(K^- \nu_\tau) + B(K^- \pi^0 \nu_\tau) + B(K^- K^0 \nu_\tau) + B(K^- \pi^0 \pi^0 \nu_\tau) + B(K^- \pi^0 K^0 \nu_\tau)] = 0.05 \pm 0.13\%$  accounting for common systematic errors in BUSKULIC 94E and BUSKULIC 94F measurements of these modes. We assume  $B(K^- \geq 2K^0 \nu_\tau)$  and  $B(K^- \geq 4\pi^0 \nu_\tau)$  are negligible.

$$\frac{\Gamma(h^- 4\pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}}{\Gamma_{25}/\Gamma} = (\Gamma_{26} + 0.319\Gamma_{110})/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.17±0.06 OUR FIT</b>				
<b>0.16±0.06 OUR AVERAGE</b>				

0.16±0.04±0.09	232	<sup>79</sup> BUSKULIC 96	ALEP	LEP 1991–1993 data
----------------	-----	---------------------------	------	--------------------

0.16±0.05±0.05		<sup>80</sup> PROCARIO 93	CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
----------------	--	---------------------------	------	---------------------------------------

<sup>79</sup> BUSKULIC 96 quote result for  $\tau^- \rightarrow h^- \geq 4\pi^0 \nu_\tau$ . We assume  $B(h^- \geq 5\pi^0 \nu_\tau)$  is negligible.

<sup>80</sup> PROCARIO 93 quotes  $B(h^- 4\pi^0 \nu_\tau)/B(h^- \pi^0 \nu_\tau) = 0.006 \pm 0.002 \pm 0.002$ . We multiply by the ARTUSO 94 result for  $B(h^- \pi^0 \nu_\tau)$  to obtain  $B(h^- 4\pi^0 \nu_\tau)$ . PROCARIO 93 assume  $B(h^- \geq 5\pi^0 \nu_\tau)$  is small and do not correct for it.

$$\frac{\Gamma(h^- 4\pi^0 \nu_\tau (\text{ex. } K^0, \eta))/\Gamma_{\text{total}}}{\Gamma_{26}/\Gamma}$$

VALUE (%)	DOCUMENT ID
<b>0.11±0.06 OUR FIT</b>	

$$\Gamma(K^- \geq 0\pi^0 \geq 0K^0 \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{27} / \Gamma$$

$$\Gamma_{27} / \Gamma = (\Gamma_{10} + \Gamma_{15} + \Gamma_{20} + \Gamma_{24} + \Gamma_{34} + \Gamma_{38}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

**1.66 ± 0.10 OUR FIT**

**1.69 ± 0.07 OUR AVERAGE**

1.70 ± 0.05 ± 0.06	avg	1610	<sup>81</sup> BUSKULIC	96 ALEP	LEP 1991–1993 data
1.54 ± 0.24	f&a		ABREU	94K DLPH	LEP 1992 Z data
1.70 ± 0.12 ± 0.19	f&a	202	<sup>82</sup> BATTLE	94 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
1.6 ± 0.4 ± 0.2	f&a	35	AIHARA	87B TPC	$E_{\text{cm}}^{ee} = 29$ GeV
1.71 ± 0.29	f&a	53	MILLS	84 DLCO	$E_{\text{cm}}^{ee} = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.60 ± 0.07 ± 0.12		967	<sup>83</sup> BUSKULIC	94E ALEP	Repl. by BUSKULIC 96
--------------------	--	-----	------------------------	----------	-------------------------

<sup>81</sup> Not independent of BUSKULIC 96  $B(K^- \nu_\tau)$ ,  $B(K^- \pi^0 \nu_\tau)$ ,  $B(K^- 2\pi^0 \nu_\tau)$ ,  $B(K^- K^0 \nu_\tau)$ , and  $B(K^- K^0 \pi^0 \nu_\tau)$  values.

<sup>82</sup> BATTLE 94 quote  $1.60 \pm 0.12 \pm 0.19$ . We add  $0.10 \pm 0.02$  to correct for their rejection of  $K_S^0 \rightarrow \pi^+ \pi^-$  decays.

<sup>83</sup> Not independent of BUSKULIC 94E  $B(K^- \nu_\tau)$ ,  $B(K^- \pi^0 \nu_\tau)$ ,  $B(K^- 2\pi^0 \nu_\tau)$ ,  $B(K^- K^0 \nu_\tau)$ , and  $B(K^- K^0 \pi^0 \nu_\tau)$  values.

$$\Gamma(K^- \geq 1(\pi^0 \text{ or } K^0) \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{28} / \Gamma$$

$$\Gamma_{28} / \Gamma = (\Gamma_{15} + \Gamma_{20} + \Gamma_{24} + \Gamma_{34} + \Gamma_{38}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

**0.95 ± 0.10 OUR FIT**

**0.76 ± 0.23 OUR AVERAGE**

0.69 ± 0.25	avg		<sup>84</sup> ABREU	94K DLPH	LEP 1992 Z data
1.2 ± 0.5 $\begin{smallmatrix} +0.2 \\ -0.4 \end{smallmatrix}$	f&a	9	AIHARA	87B TPC	$E_{\text{cm}}^{ee} = 29$ GeV

<sup>84</sup> Not independent of ABREU 94K  $B(K^- \nu_\tau)$  and  $B(K^- \geq 0 \text{ neutrals } \nu_\tau)$  measurements.

$$\Gamma(K^0(\text{particles})^- \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{29} / \Gamma$$

$$\Gamma_{29} / \Gamma = (\Gamma_{32} + \Gamma_{34} + \Gamma_{36} + \Gamma_{38} + \Gamma_{41}) / \Gamma$$

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

**1.66 ± 0.09 OUR FIT** Error includes scale factor of 1.4.

**1.94 ± 0.13 OUR AVERAGE**

1.94 ± 0.12 ± 0.12		929	<sup>85</sup> BARATE	98E ALEP	1991–1995 LEP runs
1.94 ± 0.18 ± 0.12		141	<sup>86</sup> AKERS	94G OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>85</sup> BARATE 98E measure  $\Gamma(K_S^0(\text{particles})^- \nu_\tau) / \Gamma_{\text{total}} = (0.970 \pm 0.058 \pm 0.062)\%$ . We multiply this by 2 to obtain the listed value.

<sup>86</sup> AKERS 94G measure  $\Gamma(K_S^0(\text{particles})^- \nu_\tau) / \Gamma_{\text{total}} = 0.97 \pm 0.09 \pm 0.06$ .

$$\Gamma(h^-\bar{K}^0 \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{30} / \Gamma$$

$$\Gamma_{30} / \Gamma = (\Gamma_{32} + \Gamma_{34} + \Gamma_{36} + \Gamma_{38} + 0.657\Gamma_{41}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.62 ± 0.09 OUR FIT</b>	Error includes scale factor of 1.4.			
<b>1.3 ± 0.3</b>	44	TSCHIRHART 88	HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

$$\Gamma(h^-\bar{K}^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{31} / \Gamma = (\Gamma_{32} + \Gamma_{34}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.99 ± 0.08 OUR FIT</b>	Error includes scale factor of 1.5.			
<b>0.90 ± 0.07 OUR AVERAGE</b>				
1.01 ± 0.11 ± 0.07	avg 555	<sup>87</sup> BARATE	98E ALEP	1991–1995 LEP runs
0.855 ± 0.036 ± 0.073	f&a 1242	COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$

<sup>87</sup> Not independent of BARATE 98E  $B(\tau^- \rightarrow \pi^- \bar{K}^0 \nu_\tau)$  and  $B(\tau^- \rightarrow K^- K^0 \nu_\tau)$  values.

$$\Gamma(\pi^-\bar{K}^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{32} / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.83 ± 0.08 OUR FIT</b>	Error includes scale factor of 1.4.			
<b>0.78 ± 0.06 OUR AVERAGE</b>				
0.855 ± 0.117 ± 0.066	avg 509	<sup>88</sup> BARATE	98E ALEP	1991–1995 LEP runs
0.79 ± 0.10 ± 0.09	f&a 98	<sup>89</sup> BUSKULIC	96 ALEP	LEP 1991–1993 data
0.704 ± 0.041 ± 0.072	avg	<sup>90</sup> COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$
0.95 ± 0.15 ± 0.06	f&a	<sup>91</sup> ACCIARRI	95F L3	1991–1993 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.88 ± 0.14 ± 0.09	53	BUSKULIC	94F ALEP	Repl. by BUSKULIC 96
--------------------	----	----------	----------	----------------------

<sup>88</sup> BARATE 98E reconstruct  $K^0$ 's using  $K_S^0 \rightarrow \pi^+ \pi^-$  decays. Not independent of BARATE 98E  $B(K^0 \text{ particles}^- \nu_\tau)$  value.

<sup>89</sup> BUSKULIC 96 measure  $K^0$ 's by detecting  $K_L^0$ 's in their hadron calorimeter.

<sup>90</sup> Not independent of COAN 96  $B(h^- K^0 \nu_\tau)$  and  $B(K^- K^0 \nu_\tau)$  measurements.

<sup>91</sup> ACCIARRI 95F do not identify  $\pi^- / K^-$  and assume  $B(K^- K^0 \nu_\tau) = (0.29 \pm 0.12)\%$ .

$$\Gamma(\pi^-\bar{K}^0 (\text{non-} K^*(892)^- ) \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{33} / \Gamma$$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.17</b>	95	ACCIARRI	95F L3	1991–1993 LEP runs

$$\Gamma(K^- K^0 \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{34} / \Gamma$$

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

**0.159 ± 0.024 OUR FIT**
**0.161 ± 0.024 OUR AVERAGE**

0.158 ± 0.042 ± 0.017		46	<sup>92</sup> BARATE	98E ALEP	1991–1995 LEP runs
-----------------------	--	----	----------------------	----------	--------------------

0.26 ± 0.09 ± 0.02		13	<sup>93</sup> BUSKULIC	96 ALEP	LEP 1991–1993 data
--------------------	--	----	------------------------	---------	--------------------

0.151 ± 0.021 ± 0.022		111	COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$
-----------------------	--	-----	------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.29 ± 0.12 ± 0.03		8	BUSKULIC	94F ALEP	Repl. by BUSKULIC 96
--------------------	--	---	----------	----------	-------------------------

<sup>92</sup> BARATE 98E reconstruct  $K^0$ 's using  $K_S^0 \rightarrow \pi^+ \pi^-$  decays.

<sup>93</sup> BUSKULIC 96 measure  $K^0$ 's by detecting  $K_L^0$ 's in their hadron calorimeter.

$$\Gamma(h^- \bar{K}^0 \pi^0 \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{35} / \Gamma = (\Gamma_{36} + \Gamma_{38}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

**0.55 ± 0.05 OUR FIT**
**0.50 ± 0.06 OUR AVERAGE** Error includes scale factor of 1.2.

0.446 ± 0.052 ± 0.046	avg	157	<sup>94</sup> BARATE	98E ALEP	1991–1995 LEP runs
-----------------------	-----	-----	----------------------	----------	--------------------

0.562 ± 0.050 ± 0.048	f&a	264	COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$
-----------------------	-----	-----	------	---------	---

<sup>94</sup> Not independent of BARATE 98E  $B(\tau^- \rightarrow \pi^- \bar{K}^0 \pi^0 \nu_\tau)$  and  $B(\tau^- \rightarrow K^- K^0 \pi^0 \nu_\tau)$  values.

$$\Gamma(\pi^- \bar{K}^0 \pi^0 \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{36} / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

**0.39 ± 0.05 OUR FIT**
**0.36 ± 0.05 OUR AVERAGE**

0.294 ± 0.073 ± 0.037	f&a	142	<sup>95</sup> BARATE	98E ALEP	1991–1995 LEP runs
-----------------------	-----	-----	----------------------	----------	-----------------------

0.32 ± 0.11 ± 0.05	f&a	23	<sup>96</sup> BUSKULIC	96 ALEP	LEP 1991–1993 data
--------------------	-----	----	------------------------	---------	-----------------------

0.417 ± 0.058 ± 0.044	avg		<sup>97</sup> COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$
-----------------------	-----	--	--------------------	---------	---

0.41 ± 0.12 ± 0.03	f&a		<sup>98</sup> ACCIARRI	95F L3	1991–1993 LEP runs
--------------------	-----	--	------------------------	--------	-----------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.33 ± 0.14 ± 0.07		9	BUSKULIC	94F ALEP	Repl. by BUSKULIC 96
--------------------	--	---	----------	----------	-------------------------

<sup>95</sup> BARATE 98E reconstruct  $K^0$ 's using  $K_S^0 \rightarrow \pi^+ \pi^-$  decays.

<sup>96</sup> BUSKULIC 96 measure  $K^0$ 's by detecting  $K_L^0$ 's in their hadron calorimeter.

<sup>97</sup> Not independent of COAN 96  $B(h^- K^0 \pi^0 \nu_\tau)$  and  $B(K^- K^0 \pi^0 \nu_\tau)$  measurements.

<sup>98</sup> ACCIARRI 95F do not identify  $\pi^- / K^-$  and assume  $B(K^- K^0 \pi^0 \nu_\tau) = (0.05 \pm 0.05)\%$ .

$\Gamma(\overline{K}^0 \rho^- \nu_\tau) / \Gamma_{\text{total}}$   $\Gamma_{37} / \Gamma$ 

VALUE (%)		DOCUMENT ID	TECN	COMMENT
<b>0.188 ± 0.054 ± 0.038</b>		99 BARATE	98E ALEP	1991–1995 LEP runs
99 BARATE 98E determine the $\overline{K}^0 \rho^-$ fraction in $\tau^- \rightarrow \pi^- \overline{K}^0 \pi^0 \nu_\tau$ decays to be $(0.64 \pm 0.09 \pm 0.10)$ and multiply their $B(\pi^- \overline{K}^0 \pi^0 \nu_\tau)$ measurement by this fraction to obtain the quoted result.				

 $\Gamma(K^- K^0 \pi^0 \nu_\tau) / \Gamma_{\text{total}}$   $\Gamma_{38} / \Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.151 ± 0.029 OUR FIT</b>				
<b>0.133 ± 0.031 OUR AVERAGE</b>				
0.152 ± 0.076 ± 0.021	15	100 BARATE	98E ALEP	1991–1995 LEP runs
0.10 ± 0.05 ± 0.03	5	101 BUSKULIC	96 ALEP	LEP 1991–1993 data
0.145 ± 0.036 ± 0.020	32	COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.05 ± 0.05 ± 0.01	1	BUSKULIC	94F ALEP	Repl. by BUSKULIC 96
100 BARATE 98E reconstruct $K^0$ 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.				
101 BUSKULIC 96 measure $K^0$ 's by detecting $K_L^0$ 's in their hadron calorimeter.				

 $\Gamma(\pi^- \overline{K}^0 \pi^0 \pi^0 \nu_\tau) / \Gamma_{\text{total}}$   $\Gamma_{39} / \Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.58 ± 0.33 ± 0.14</b>	5	102 BARATE	98E ALEP	1991–1995 LEP runs
102 BARATE 98E reconstruct $K^0$ 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.				


 $\Gamma(K^- K^0 \pi^0 \pi^0 \nu_\tau) / \Gamma_{\text{total}}$   $\Gamma_{40} / \Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.39 × 10<sup>-3</sup></b>	95	BARATE	98E ALEP	1991–1995 LEP runs

 $\Gamma(\pi^- K^0 \overline{K}^0 \nu_\tau) / \Gamma_{\text{total}}$   $\Gamma_{41} / \Gamma$ 

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.121 ± 0.021 OUR FIT</b>				Error includes scale factor of 1.2.
<b>0.116 ± 0.028 OUR AVERAGE</b>				Error includes scale factor of 1.5. See the ideogram below.
0.153 ± 0.030 ± 0.016	f&a 74	103 BARATE	98E ALEP	1991–1995 LEP runs
0.092 ± 0.020 ± 0.012	avg 42	104 COAN	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
0.31 ± 0.12 ± 0.04	f&a	ACCIARRI	95F L3	1991–1993 LEP runs
103 BARATE 98E obtain this value by adding twice their $B(\pi^- K_S^0 K_S^0 \nu_\tau)$ value to their $B(\pi^- K_S^0 K_L^0 \nu_\tau)$ value.				
104 We multiply the COAN 96 measurement $B(h^- K_S^0 K_S^0 \nu_\tau) = (0.023 \pm 0.005 \pm 0.003)\%$ by 4 to obtain the listed value. This factor of 1/4 is uncertain, and might be as large as 1/2, due to Bose-Einstein correlations and the resonant parentage of this state.				



$$\Gamma(\pi^- K^0 \bar{K}^0 \nu_\tau) / \Gamma_{\text{total}} (\%)$$

$$\Gamma(\pi^- K_S^0 K_S^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{42} / \Gamma = \frac{1}{4} \Gamma_{41} / \Gamma$$

Bose-Einstein correlations might make the mixing fraction different than 1/4.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.030±0.005 OUR FIT</b>				Error includes scale factor of 1.2.
<b>0.024±0.005 OUR AVERAGE</b>				
0.026±0.010±0.005	6	BARATE	98E ALEP	1991–1995 LEP runs
0.023±0.005±0.003	42	COAN	96 CLEO	$E_{\text{cm}}^{e\bar{e}} \approx 10.6 \text{ GeV}$

$$\Gamma(\pi^- K_S^0 K_L^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{43} / \Gamma = \frac{1}{2} \Gamma_{41} / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.060±0.010 OUR FIT</b>				Error includes scale factor of 1.2.
<b>0.101±0.023±0.013 avg</b>	68	BARATE	98E ALEP	1991–1995 LEP runs

$$\Gamma(\pi^- K_S^0 K_S^0 \pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{44} / \Gamma$$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.020</b>	95	BARATE	98E ALEP	1991–1995 LEP runs

$$\Gamma(\pi^- K_S^0 K_L^0 \pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{45} / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.031±0.011±0.005</b>	11	BARATE	98E ALEP	1991–1995 LEP runs

$\Gamma(K^- K^0 \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ 
 $\Gamma_{46}/\Gamma = (\Gamma_{34} + \Gamma_{38})/\Gamma$ 

VALUE (%)	DOCUMENT ID
<b>0.31 ± 0.04 OUR FIT</b>	

 $\Gamma(K^0 h^+ h^- h^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ 
 $\Gamma_{47}/\Gamma$ 

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.17</b>	95	TSCHIRHART 88	HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.27	90	BELTRAMI 85	HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

 $\Gamma(K^0 h^+ h^- h^- \nu_\tau)/\Gamma_{\text{total}}$ 
 $\Gamma_{48}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.023 ± 0.019 ± 0.007</b>	6	<sup>105</sup> BARATE 98E	ALEP	1991–1995 LEP runs
<sup>105</sup> BARATE 98E reconstruct $K^0$ 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.				

 $\Gamma(h^- h^- h^+ \geq 0 \text{ neut. } \nu_\tau (\text{"3-prong"}))/\Gamma_{\text{total}}$ 
 $\Gamma_{49}/\Gamma$ 

$$\Gamma_{49}/\Gamma = (0.3431\Gamma_{32} + 0.3431\Gamma_{34} + 0.3431\Gamma_{36} + 0.3431\Gamma_{38} + 0.4508\Gamma_{41} + \Gamma_{57} + \Gamma_{65} + \Gamma_{73} + \Gamma_{74} + \Gamma_{79} + \Gamma_{81} + \Gamma_{84} + \Gamma_{85} + 0.285\Gamma_{110} + 0.9101\Gamma_{125} + 0.9101\Gamma_{126})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>15.18 ± 0.13 OUR FIT</b>				Error includes scale factor of 1.2.
<b>14.8 ± 0.4 OUR AVERAGE</b>				

14.4 ± 0.6 ± 0.3	f&a	ADEVA	91F L3	$E_{\text{cm}}^{ee} = 88.3\text{--}94.3 \text{ GeV}$
15.0 ± 0.4 ± 0.3	f&a	BEHREND	89B CELL	$E_{\text{cm}}^{ee} = 14\text{--}47 \text{ GeV}$
15.1 ± 0.8 ± 0.6	f&a	AIHARA	87B TPC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
13.5 ± 0.3 ± 0.3		ABACHI	89B HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
12.8 ± 1.0 ± 0.7		<sup>106</sup> BURCHAT	87 MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
12.1 ± 0.5 ± 1.2		RUCKSTUHL	86 DLCO	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
12.8 ± 0.5 ± 0.8	1420	SCHMIDKE	86 MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
15.3 ± 1.1 $\begin{smallmatrix} +1.3 \\ -1.6 \end{smallmatrix}$	367	ALTHOFF	85 TASS	$E_{\text{cm}}^{ee} = 34.5 \text{ GeV}$
13.6 ± 0.5 ± 0.8		BARTEL	85F JADE	$E_{\text{cm}}^{ee} = 34.6 \text{ GeV}$
12.2 ± 1.3 ± 3.9		<sup>107</sup> BERGER	85 PLUT	$E_{\text{cm}}^{ee} = 34.6 \text{ GeV}$
13.3 ± 0.3 ± 0.6		FERNANDEZ	85 MAC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
24 ± 6	35	BRANDELIK	80 TASS	$E_{\text{cm}}^{ee} = 30 \text{ GeV}$
32 ± 5	692	<sup>108</sup> BACINO	78B DLCO	$E_{\text{cm}}^{ee} = 3.1\text{--}7.4 \text{ GeV}$
35 ± 11		<sup>108</sup> BRANDELIK	78 DASP	Assumes $V\text{--}A$ decay
18 ± 6.5	33	<sup>108</sup> JAROS	78 MRK1	$E_{\text{cm}}^{ee} > 6 \text{ GeV}$

<sup>106</sup> BURCHAT 87 value is not independent of SCHMIDKE 86 value.

<sup>107</sup> Not independent of BERGER 85  $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma_{\text{total}}$ ,  $\Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{\text{total}}$ ,  $\Gamma(h^- \geq 1 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ , and  $\Gamma(h^- \geq 0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$ , and therefore not used in the fit.

<sup>108</sup> Low energy experiments are not in average or fit because the systematic errors in background subtraction are judged to be large.

$$\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{50} / \Gamma$$

$$\Gamma_{50} / \Gamma = (\Gamma_{57} + \Gamma_{65} + \Gamma_{73} + \Gamma_{74} + \Gamma_{79} + \Gamma_{81} + \Gamma_{84} + \Gamma_{85} + 0.285\Gamma_{110} + 0.9101\Gamma_{125} + 0.9101\Gamma_{126}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
-----------	--	------	-------------	------	---------

**14.60 ± 0.13 OUR FIT** Error includes scale factor of 1.2.

**14.63 ± 0.25 OUR AVERAGE** Error includes scale factor of 1.4. See the ideogram below.

14.96 ± 0.09 ± 0.22	f&a	10.4k	AKERS	95Y OPAL	1991–1994 LEP runs
14.22 ± 0.10 ± 0.37	avg		<sup>109</sup> BALEST	95C CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV
13.3 ± 0.3 ± 0.8	f&a		<sup>110</sup> ALBRECHT	92D ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV
14.35 <sup>+0.40</sup> <sub>-0.45</sub> ± 0.24	f&a		DECAMP	92C ALEP	1989–1990 LEP runs


• • • We do not use the following data for averages, fits, limits, etc. • • •

15.26 ± 0.26 ± 0.22 ACTON 92H OPAL Repl. by AKERS 95Y

<sup>109</sup> Not independent of BALEST 95C  $B(h^- h^- h^+ \nu_\tau)$  and  $B(h^- h^- h^+ \pi^0 \nu_\tau)$  values, and BORTOLETTO 93  $B(h^- h^- h^+ 2\pi^0 \nu_\tau) / B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau)$  value.

<sup>110</sup> This ALBRECHT 92D value is not independent of their  $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) \Gamma(e^- \bar{\nu}_e \nu_\tau) / \Gamma_{\text{total}}^2$  value.

WEIGHTED AVERAGE  
14.63 ± 0.25 (Error scaled by 1.4)



$$\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}} (\%)$$

$$\Gamma(\pi^- \pi^+ \pi^- \geq 0 \text{ neutrals } \nu_\tau) / \Gamma(h^- h^- h^+ \geq \text{Oneut. } \nu_\tau \text{ ("3-prong")}) \quad \Gamma_{51}/\Gamma_{49}$$

$$\Gamma_{51}/\Gamma_{49} = (0.3431\Gamma_{32} + 0.3431\Gamma_{36} + 0.1078\Gamma_{41} + \Gamma_{57} + \Gamma_{65} + \Gamma_{73} + \Gamma_{74} + 0.285\Gamma_{110} + 0.9101\Gamma_{125} + 0.9101\Gamma_{126}) / (0.3431\Gamma_{32} + 0.3431\Gamma_{34} + 0.3431\Gamma_{36} + 0.3431\Gamma_{38} + 0.4508\Gamma_{41} + \Gamma_{57} + \Gamma_{65} + \Gamma_{73} + \Gamma_{74} + \Gamma_{79} + \Gamma_{81} + \Gamma_{84} + \Gamma_{85} + 0.285\Gamma_{110} + 0.9101\Gamma_{125} + 0.9101\Gamma_{126})$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.962 ± 0.005 OUR FIT</b>				Error includes scale factor of 1.1.
<b>0.945 ± 0.019</b>	490	<sup>111</sup> BAUER	94 TPC	$E_{cm}^{ee} = 29 \text{ GeV}$

<sup>111</sup>BAUER 94 quote  $B(\pi^- \pi^+ \pi^- \geq 0 \text{ neutrals } \nu_\tau) = 0.1329 \pm 0.0027$ . We divide by 0.1406, their assumed value for  $B(\text{"3prong"})$ .

$$\Gamma(h^- h^- h^+ \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{52}/\Gamma$$

$$\Gamma_{52}/\Gamma = (0.3431\Gamma_{32} + 0.3431\Gamma_{34} + \Gamma_{57} + \Gamma_{79} + \Gamma_{84} + 0.0221\Gamma_{125}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.96 ± 0.10 OUR FIT</b>				Error includes scale factor of 1.1.

**9.7 ± 0.4 OUR AVERAGE** Error includes scale factor of 3.1. See the ideogram below.

7.6 ± 0.1 ± 0.5	avg 7.5k	<sup>112</sup> ALBRECHT	96E ARG	$E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$
9.92 ± 0.10 ± 0.09	f&a 11.2k	<sup>113</sup> BUSKULIC	96 ALEP	LEP 1991–1993 data
9.49 ± 0.36 ± 0.63	f&a	DECAMP	92C ALEP	1989–1990 LEP runs
8.7 ± 0.7 ± 0.3	f&a 694	<sup>114</sup> BEHREND	90 CELL	$E_{cm}^{ee} = 35 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.0 ± 0.3 ± 0.7	1566	<sup>115</sup> BAND	87 MAC	$E_{cm}^{ee} = 29 \text{ GeV}$
6.7 ± 0.8 ± 0.9		<sup>116</sup> BURCHAT	87 MRK2	$E_{cm}^{ee} = 29 \text{ GeV}$
6.4 ± 0.4 ± 0.9		<sup>117</sup> RUCKSTUHL	86 DLCO	$E_{cm}^{ee} = 29 \text{ GeV}$
7.8 ± 0.5 ± 0.8	890	SCHMIDKE	86 MRK2	$E_{cm}^{ee} = 29 \text{ GeV}$
8.4 ± 0.4 ± 0.7	1255	<sup>117</sup> FERNANDEZ	85 MAC	$E_{cm}^{ee} = 29 \text{ GeV}$
9.7 ± 2.0 ± 1.3		BEHREND	84 CELL	$E_{cm}^{ee} = 14,22 \text{ GeV}$

<sup>112</sup>ALBRECHT 96E not independent of ALBRECHT 93C  $\Gamma(h^- h^- h^+ \nu_\tau \text{ (ex. } K^0)) \times \Gamma(\text{particle}^- \geq 0 \text{ neutrals } \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}^2$  value.


<sup>113</sup>BUSKULIC 96 quote  $B(h^- h^- h^+ \nu_\tau \text{ (ex. } K^0)) = 9.50 \pm 0.10 \pm 0.11$ . We add 0.42 to remove their  $K^0$  correction and reduce the systematic error accordingly.

<sup>114</sup>BEHREND 90 subtract 0.3% to account for the  $\tau^- \rightarrow K^*(892)^- \nu_\tau$  contribution to measured events.

<sup>115</sup>BAND 87 subtract for charged kaon modes; not independent of FERNANDEZ 85 value.

<sup>116</sup>BURCHAT 87 value is not independent of SCHMIDKE 86 value.

<sup>117</sup>Value obtained by multiplying paper's  $R = B(h^- h^- h^+ \nu_\tau) / B(\text{3-prong})$  by  $B(\text{3-prong}) = 0.143$  and subtracting 0.3% for  $K^*(892)$  background.



$$\Gamma(h^- h^- h^+ \nu_\tau) / \Gamma_{\text{total}} (\%)$$

$$\Gamma(h^- h^- h^+ \nu_\tau) / \Gamma(h^- h^- h^+ \geq 0 \text{ neut. } \nu_\tau \text{ ("3-prong")}) \quad \Gamma_{52} / \Gamma_{49}$$

$$\Gamma_{52} / \Gamma_{49} = (0.3431\Gamma_{32} + 0.3431\Gamma_{34} + \Gamma_{57} + \Gamma_{79} + \Gamma_{84} + 0.0221\Gamma_{125}) / (0.3431\Gamma_{32} + 0.3431\Gamma_{34} + 0.3431\Gamma_{36} + 0.3431\Gamma_{38} + 0.4508\Gamma_{41} + \Gamma_{57} + \Gamma_{65} + \Gamma_{73} + \Gamma_{74} + \Gamma_{79} + \Gamma_{81} + \Gamma_{84} + \Gamma_{85} + 0.285\Gamma_{110} + 0.9101\Gamma_{125} + 0.9101\Gamma_{126})$$

This branching fractions is not independent of values for  $\Gamma(h^- h^- h^+ \nu_\tau) / \Gamma_{\text{total}}$  and  $\Gamma(h^- h^- h^+ \geq 0 \text{ neut. } \nu_\tau \text{ ("3-prong")}) / \Gamma_{\text{total}}$ .

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.656±0.006 OUR FIT</b>	Error includes scale factor of 1.1.		
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
0.47 ±0.03 ±0.06	RUCKSTUHL 86	DLCO	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
0.61 ±0.03 ±0.05	FERNANDEZ 85	MAC	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$

$$\Gamma(h^- h^- h^+ \nu_\tau \text{ (ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{53} / \Gamma$$

$$\Gamma_{53} / \Gamma = (\Gamma_{57} + \Gamma_{79} + \Gamma_{84} + 0.0221\Gamma_{125}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.62±0.10 OUR FIT</b>	Error includes scale factor of 1.1.			
<b>9.57±0.11 OUR AVERAGE</b>				
9.50±0.10±0.11	avg 11.2k	<sup>118</sup> BUSKULIC	96 ALEP	LEP 1991–1993 data
9.87±0.10±0.24	avg	<sup>119</sup> AKERS	95Y OPAL	1991–1994 LEP runs
9.51±0.07±0.20	f&a 37.7k	BALEST	95C CLEO	$E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$

<sup>118</sup> Not independent of BUSKULIC 96  $B(h^- h^- h^+ \nu_\tau)$  value.

<sup>119</sup> Not independent of AKERS 95Y  $B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ (ex. } K_S^0 \rightarrow \pi^+ \pi^-))$  and  $B(h^- h^- h^+ \nu_\tau \text{ (ex. } K^0)) / B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ (ex. } K_S^0 \rightarrow \pi^+ \pi^-))$  values.

$$\Gamma(h^- h^- h^+ \nu_\tau (\text{ex. } K^0)) / \Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-))$$

$$\Gamma_{53}/\Gamma_{50} = (\Gamma_{57} + \Gamma_{79} + \Gamma_{84} + 0.0221\Gamma_{125}) / (\Gamma_{57} + \Gamma_{65} + \Gamma_{73} + \Gamma_{74} + \Gamma_{79} + \Gamma_{81} + \Gamma_{84} + \Gamma_{85} + 0.285\Gamma_{110} + 0.9101\Gamma_{125} + 0.9101\Gamma_{126})$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.659 ± 0.006 OUR FIT</b>			Error includes scale factor of 1.1.
<b>0.660 ± 0.004 ± 0.014</b>	AKERS	95Y OPAL	1991–1994 LEP runs

$$\Gamma(h^- h^- h^+ \nu_\tau (\text{ex. } K^0, \omega)) / \Gamma_{\text{total}} \quad \Gamma_{54}/\Gamma = (\Gamma_{57} + \Gamma_{79} + \Gamma_{84}) / \Gamma$$

$$\Gamma_{54}/\Gamma = (\Gamma_{57} + \Gamma_{79} + \Gamma_{84}) / \Gamma$$

VALUE (%)	DOCUMENT ID
<b>9.57 ± 0.10 OUR FIT</b>	
	Error includes scale factor of 1.1.

$$\Gamma(\pi^- \pi^+ \pi^- \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{55}/\Gamma = (0.3431\Gamma_{32} + \Gamma_{57} + 0.0221\Gamma_{125}) / \Gamma$$

VALUE (%)	DOCUMENT ID
<b>9.56 ± 0.11 OUR FIT</b>	
	Error includes scale factor of 1.1.

$$\Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{56}/\Gamma = (0.3431\Gamma_{32} + \Gamma_{57}) / \Gamma$$

VALUE (%)	DOCUMENT ID
<b>9.52 ± 0.11 OUR FIT</b>	
	Error includes scale factor of 1.1.

$$\Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0, \omega)) / \Gamma_{\text{total}} \quad \Gamma_{57}/\Gamma$$

VALUE (%)	DOCUMENT ID
<b>9.23 ± 0.11 OUR FIT</b>	
	Error includes scale factor of 1.1.

$$\Gamma(h^- h^- h^+ \geq 1 \text{ neutrals } \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{58}/\Gamma$$

$$\Gamma_{58}/\Gamma = (0.3431\Gamma_{36} + 0.3431\Gamma_{38} + 0.1077\Gamma_{41} + \Gamma_{65} + \Gamma_{73} + \Gamma_{74} + \Gamma_{81} + \Gamma_{85} + 0.285\Gamma_{110} + 0.888\Gamma_{125} + 0.9101\Gamma_{126}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.18 ± 0.11 OUR FIT</b>				Error includes scale factor of 1.2.
<b>5.2 ± 0.6 OUR AVERAGE</b>				
5.6 ± 0.7 ± 0.3	avg	352	120	BEHREND 90 CELL $E_{\text{cm}}^{\text{ee}} = 35 \text{ GeV}$
4.2 ± 0.5 ± 0.9	f&a	203	121	ALBRECHT 87L ARG $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$
6.2 ± 2.3 ± 1.7	f&a			BEHREND 84 CELL $E_{\text{cm}}^{\text{ee}} = 14, 22 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.1 ± 0.8 ± 0.9		122	BURCHAT	87	MRK2	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
7.6 ± 0.4 ± 0.9		123, 124	RUCKSTUHL	86	DLCO	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
4.7 ± 0.5 ± 0.8		530	125	SCHMIDKE	86	MRK2 $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
5.6 ± 0.4 ± 0.7		124	FERNANDEZ	85	MAC	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$

<sup>120</sup> BEHREND 90 value is not independent of BEHREND 90  $B(3h\nu_\tau \geq 1 \text{ neutrals}) + B(5\text{-prong})$ .

<sup>121</sup> ALBRECHT 87L measure the product of branching ratios  $B(3\pi^\pm \pi^0 \nu_\tau) B((e\bar{\nu} \text{ or } \mu\bar{\nu} \text{ or } \pi \text{ or } K \text{ or } \rho)\nu_\tau) = 0.029$  and use the PDG 86 values for the second branching ratio which sum to  $0.69 \pm 0.03$  to get the quoted value.

<sup>122</sup> BURCHAT 87 value is not independent of SCHMIDKE 86 value.

<sup>123</sup> Contributions from kaons and from  $>1\pi^0$  are subtracted. Not independent of (3-prong +  $0\pi^0$ ) and (3-prong +  $\geq 0\pi^0$ ) values.

<sup>124</sup> Value obtained using paper's  $R = B(h^- h^- h^+ \nu_\tau)/B(3\text{-prong})$  and current  $B(3\text{-prong}) = 0.143$ .

<sup>125</sup> Not independent of SCHMIDKE 86  $h^- h^- h^+ \nu_\tau$  and  $h^- h^- h^+ (\geq 0\pi^0)\nu_\tau$  values.

$$\Gamma(h^- h^- h^+ \geq 1 \text{ neutrals } \nu_\tau \text{ (ex. } K_S^0 \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{59} / \Gamma$$

$$\Gamma_{59} / \Gamma = (\Gamma_{65} + \Gamma_{73} + \Gamma_{74} + \Gamma_{81} + \Gamma_{85} + 0.285\Gamma_{110} + 0.888\Gamma_{125} + 0.9101\Gamma_{126}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.98 ± 0.11 OUR FIT</b>				Error includes scale factor of 1.2.

**5.07 ± 0.24 OUR AVERAGE**

5.09 ± 0.10 ± 0.23	avg	<sup>126</sup> AKERS	95Y OPAL	1991–1994 LEP runs
4.95 ± 0.29 ± 0.65	f&a	570 DECAMP	92C ALEP	1989–1990 LEP runs

<sup>126</sup> Not independent of AKERS 95Y  $B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ (ex. } K_S^0 \rightarrow \pi^+ \pi^-))$  and  $B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ (ex. } K^0)) / B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ (ex. } K_S^0 \rightarrow \pi^+ \pi^-))$  values.

$$\Gamma(h^- h^- h^+ \pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{60} / \Gamma$$

$$\Gamma_{60} / \Gamma = (0.3431\Gamma_{36} + 0.3431\Gamma_{38} + \Gamma_{65} + \Gamma_{81} + \Gamma_{85} + 0.888\Gamma_{125} + 0.0221\Gamma_{126}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.50 ± 0.09 OUR FIT</b>				Error includes scale factor of 1.1.

<b>4.45 ± 0.09 ± 0.07</b>	6.1k	<sup>127</sup> BUSKULIC	96 ALEP	LEP 1991–1993 data
---------------------------	------	-------------------------	---------	--------------------

<sup>127</sup> BUSKULIC 96 quote  $B(h^- h^- h^+ \pi^0 \nu_\tau \text{ (ex. } K^0)) = 4.30 \pm 0.09 \pm 0.09$ . We add 0.15 to remove their  $K^0$  correction and reduce the systematic error accordingly.

$$\Gamma(h^- h^- h^+ \pi^0 \nu_\tau \text{ (ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{61} / \Gamma$$

$$\Gamma_{61} / \Gamma = (\Gamma_{65} + \Gamma_{81} + \Gamma_{85} + 0.888\Gamma_{125} + 0.0221\Gamma_{126}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.31 ± 0.09 OUR FIT</b>				Error includes scale factor of 1.1.

<b>4.23 ± 0.06 ± 0.22</b>	7.2k	BALEST	95c CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$
---------------------------	------	--------	----------	---

$$\Gamma(h^- h^- h^+ \pi^0 \nu_\tau \text{ (ex. } K^0, \omega)) / \Gamma_{\text{total}} \quad \Gamma_{62} / \Gamma = (\Gamma_{65} + \Gamma_{81} + \Gamma_{85}) / \Gamma$$

VALUE (%)	DOCUMENT ID
<b>2.59 ± 0.09 OUR FIT</b>	

$$\Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{63} / \Gamma = (0.3431\Gamma_{36} + \Gamma_{65} + 0.888\Gamma_{125} + 0.0221\Gamma_{126}) / \Gamma$$

VALUE (%)	DOCUMENT ID
<b>4.35 ± 0.10 OUR FIT</b>	

$$\Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau \text{ (ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{64} / \Gamma = (\Gamma_{65} + 0.888\Gamma_{125} + 0.0221\Gamma_{126}) / \Gamma$$

VALUE (%)	DOCUMENT ID
<b>4.22 ± 0.10 OUR FIT</b>	

$$\Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0, \omega)) / \Gamma_{\text{total}} \quad \Gamma_{65} / \Gamma$$

VALUE (%) DOCUMENT ID  
**2.49 ± 0.10 OUR FIT**

$$\Gamma(h^- (\rho\pi)^0 \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau) \quad \Gamma_{66} / \Gamma_{60}$$

$$\Gamma_{66} / \Gamma_{60} = (\Gamma_{68} + \Gamma_{69} + \Gamma_{70}) / \Gamma_{60}$$

VALUE DOCUMENT ID TECN COMMENT  
**0.64 ± 0.07 ± 0.03** 128 ALBRECHT 91D ARG  $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6$  GeV

128 ALBRECHT 91D not independent of their  $\Gamma(h^- \rho^+ h^- \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$ ,  $\Gamma(h^- \rho^- h^+ \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$ , and  $\Gamma(h^- \rho \pi^0 \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$  values.

$$\Gamma((a_1(1260) h)^- \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau) \quad \Gamma_{67} / \Gamma_{60}$$

VALUE CL% DOCUMENT ID TECN COMMENT  
**<0.44** 95 129 ALBRECHT 91D ARG  $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6$  GeV

129 ALBRECHT 91D not independent of their  $\Gamma(h^- \omega \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau (\text{ex. } K^0))$ ,  $\Gamma(h^- \rho \pi^0 \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$ ,  $\Gamma(h^- \rho^+ h^- \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$ , and  $\Gamma(h^- \rho^- h^+ \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$  values.

$$\Gamma(h^- \rho \pi^0 \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau) \quad \Gamma_{68} / \Gamma_{60}$$

VALUE EVTS DOCUMENT ID TECN COMMENT  
**0.30 ± 0.04 ± 0.02** 393 ALBRECHT 91D ARG  $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6$  GeV

$$\Gamma(h^- \rho^+ h^- \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau) \quad \Gamma_{69} / \Gamma_{60}$$

VALUE EVTS DOCUMENT ID TECN COMMENT  
**0.10 ± 0.03 ± 0.04** 142 ALBRECHT 91D ARG  $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6$  GeV

$$\Gamma(h^- \rho^- h^+ \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau) \quad \Gamma_{70} / \Gamma_{60}$$

VALUE EVTS DOCUMENT ID TECN COMMENT  
**0.26 ± 0.05 ± 0.01** 370 ALBRECHT 91D ARG  $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6$  GeV

$$[\Gamma(h^- \rho^+ h^- \nu_\tau) + \Gamma(h^- \rho^- h^+ \nu_\tau)] / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau) \quad (\Gamma_{69} + \Gamma_{70}) / \Gamma_{60}$$

VALUE EVTS DOCUMENT ID TECN COMMENT  
**0.33 ± 0.06 ± 0.01** 475 130 ALBRECHT 91D ARG  $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6$  GeV

130 ALBRECHT 91D not independent of their  $\Gamma(h^- \rho^+ h^- \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$  and  $\Gamma(h^- \rho^- h^+ \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$  values.

$$\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{71} / \Gamma$$

$$\Gamma_{71} / \Gamma = (0.1077\Gamma_{41} + \Gamma_{73} + 0.236\Gamma_{110} + 0.888\Gamma_{126}) / \Gamma$$

VALUE (%) DOCUMENT ID  
**0.54 ± 0.04 OUR FIT**

$$\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{72} / \Gamma$$

$$\Gamma_{72} / \Gamma = (\Gamma_{73} + 0.236\Gamma_{110} + 0.888\Gamma_{126}) / \Gamma$$

VALUE (%) EVTS DOCUMENT ID TECN COMMENT  
**0.53 ± 0.04 OUR FIT**  
**0.50 ± 0.07 ± 0.07** 1.8k BUSKULIC 96 ALEP LEP 1991–1993 data

$$\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma(h^- h^- h^+ \geq 0 \text{ neut. } \nu_\tau (\text{"3-prong"})) \quad \Gamma_{72}/\Gamma_{49}$$

$$\Gamma_{72}/\Gamma_{49} = (\Gamma_{73} + 0.236\Gamma_{110} + 0.888\Gamma_{126}) / (0.3431\Gamma_{32} + 0.3431\Gamma_{34} + 0.3431\Gamma_{36} + 0.3431\Gamma_{38} + 0.4508\Gamma_{41} + \Gamma_{57} + \Gamma_{65} + \Gamma_{73} + \Gamma_{74} + \Gamma_{79} + \Gamma_{81} + \Gamma_{84} + \Gamma_{85} + 0.285\Gamma_{110} + 0.9101\Gamma_{125} + 0.9101\Gamma_{126})$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.0348 ± 0.0028 OUR FIT</b>				
<b>0.034 ± 0.002 ± 0.003</b>	668	BORTOLETTO93	CLEO	$E_{\text{cm}}^{ee} \approx 10.6$ GeV

$$\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau (\text{ex. } K^0, \omega, \eta)) / \Gamma_{\text{total}} \quad \Gamma_{73}/\Gamma$$

VALUE (%)	DOCUMENT ID
<b>0.11 ± 0.04 OUR FIT</b>	

$$\Gamma(h^- h^- h^+ \geq 3\pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{74}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.14<sup>+0.09</sup><sub>-0.07</sub> OUR FIT</b>				Error includes scale factor of 1.5.
<b>0.11 ± 0.04 ± 0.05</b>	440	BUSKULIC	96 ALEP	LEP 1991–1993 data

$$\Gamma(h^- h^- h^+ 3\pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{75}/\Gamma$$

VALUE (units 10 <sup>-4</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.85 ± 0.56 ± 0.51</b>	57	ANDERSON	97 CLEO	$E_{\text{cm}}^{ee} = 10.6$ GeV

$$\Gamma(K^- h^+ h^- \geq 0 \text{ neutrals } \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{76}/\Gamma = (0.3431\Gamma_{34} + 0.3431\Gamma_{38} + \Gamma_{79} + \Gamma_{81} + \Gamma_{84} + \Gamma_{85}) / \Gamma$$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<b>0.54 ± 0.07 OUR FIT</b>				Error includes scale factor of 1.1.
<b>&lt; 0.6</b>	90	AIHARA	84C TPC	$E_{\text{cm}}^{ee} = 29$ GeV

$$\Gamma(K^- \pi^+ \pi^- \geq 0 \text{ neutrals } \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{77}/\Gamma = (0.3431\Gamma_{34} + 0.3431\Gamma_{38} + \Gamma_{79} + \Gamma_{81}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.31 ± 0.06 OUR FIT</b>				Error includes scale factor of 1.1.
<b>0.30 ± 0.07 OUR AVERAGE</b>				Error includes scale factor of 1.2.
0.275 ± 0.064	avg	<sup>131</sup> BARATE	98 ALEP	1991–1995 LEP runs
0.58 <sup>+0.15</sup> <sub>-0.13</sub> ± 0.12	f&a	<sup>132</sup> BAUER	94 TPC	$E_{\text{cm}}^{ee} = 29$ GeV
0.22 <sup>+0.16</sup> <sub>-0.13</sub> ± 0.05	f&a	<sup>133</sup> MILLS	85 DLCO	$E_{\text{cm}}^{ee} = 29$ GeV

<sup>131</sup> Not independent of BARATE 98  $\Gamma(\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau) / \Gamma_{\text{total}}$  and  $\Gamma(\tau^- \rightarrow K^- \pi^+ \pi^- \pi^0 \nu_\tau) / \Gamma_{\text{total}}$  values.

<sup>132</sup> We multiply 0.58% by 0.20, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

<sup>133</sup> Error correlated with MILLS 85 ( $K K \pi \nu$ ) value. We multiply 0.22% by 0.23, the relative systematic error quoted by MILLS 85, to obtain obtain the systematic error.

$$\Gamma(K^- \pi^+ \pi^- \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{78} / \Gamma = (0.3431 \Gamma_{34} + \Gamma_{79}) / \Gamma$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>0.23 ± 0.04 OUR FIT</b>			
<b>0.214 ± 0.037 ± 0.029</b>	BARATE	98	ALEP 1991–1995 LEP runs

$$\Gamma(K^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \qquad \Gamma_{79} / \Gamma$$

VALUE (%)	DOCUMENT ID
<b>0.18 ± 0.05 OUR FIT</b>	

$$\Gamma(K^- \pi^+ \pi^- \pi^0 \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{80} / \Gamma = (0.3431 \Gamma_{38} + \Gamma_{81}) / \Gamma$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>0.08 ± 0.04 OUR FIT</b>			
<b>0.061 ± 0.039 ± 0.018</b>	BARATE	98	ALEP 1991–1995 LEP runs

$$\Gamma(K^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \qquad \Gamma_{81} / \Gamma$$

VALUE	DOCUMENT ID
<b><math>(2.4^{+4.3}_{-1.6}) \times 10^{-4}</math> OUR FIT</b>	

$$\Gamma(K^- \pi^+ K^- \geq 0 \text{ neut. } \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{82} / \Gamma$$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.09</b>	95	BAUER	94	TPC $E_{\text{cm}}^{ee} = 29 \text{ GeV}$

$$\Gamma(K^- K^+ \pi^- \geq 0 \text{ neut. } \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{83} / \Gamma = (\Gamma_{84} + \Gamma_{85}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.23 ± 0.04 OUR FIT</b>				
<b>0.22 ± 0.04 OUR AVERAGE</b>				
0.238 ± 0.042	avg	134 BARATE	98	ALEP 1991–1995 LEP runs
0.15 $^{+0.09}_{-0.07}$ ± 0.03	f&a	4 135 BAUER	94	TPC $E_{\text{cm}}^{ee} = 29 \text{ GeV}$

<sup>134</sup> Not independent of BARATE 98  $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau) / \Gamma_{\text{total}}$  and  $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \pi^0 \nu_\tau) / \Gamma_{\text{total}}$  values.

<sup>135</sup> We multiply 0.15% by 0.20, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

$$\Gamma(K^- K^+ \pi^- \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{84} / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.161 ± 0.026 OUR FIT</b>				
<b>0.165 ± 0.027 OUR AVERAGE</b>				
0.163 ± 0.021 ± 0.017		BARATE	98	ALEP 1991–1995 LEP runs
0.22 $^{+0.17}_{-0.11}$ ± 0.05	9	136 MILLS	85	DLCO $E_{\text{cm}}^{ee} = 29 \text{ GeV}$

<sup>136</sup> Error correlated with MILLS 85 ( $K \pi \pi \pi^0 \nu$ ) value. We multiply 0.22% by 0.23, the relative systematic error quoted by MILLS 85, to obtain the systematic error.

$$\Gamma(K^- K^+ \pi^- \pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{85} / \Gamma$$

VALUE (%)		DOCUMENT ID	TECN	COMMENT
<b>0.069 ± 0.030 OUR FIT</b>				
<b>0.075 ± 0.029 ± 0.015</b>		BARATE	98	ALEP 1991–1995 LEP runs

$$\Gamma(K^- K^+ K^- \geq 0 \text{ neut. } \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{86} / \Gamma$$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.21</b>	95	BAUER	94	TPC $E_{\text{cm}}^{ee} = 29 \text{ GeV}$

$$\Gamma(K^- K^+ K^- \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{87} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 1.9 × 10<sup>-4</sup></b>	90	BARATE	98	ALEP 1991–1995 LEP runs

$$\Gamma(\pi^- K^+ \pi^- \geq 0 \text{ neut. } \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{88} / \Gamma$$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.25</b>	95	BAUER	94	TPC $E_{\text{cm}}^{ee} = 29 \text{ GeV}$

$$\Gamma(e^- e^- e^+ \bar{\nu}_e \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{89} / \Gamma$$

VALUE (units 10 <sup>-5</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.8 ± 1.4 ± 0.4</b>	5	ALAM	96	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$$\Gamma(\mu^- e^- e^+ \bar{\nu}_\mu \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{90} / \Gamma$$

VALUE (units 10 <sup>-5</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 3.6</b>	90	ALAM	96	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$$\Gamma(3h^- 2h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ (ex. } K_S^0 \rightarrow \pi^- \pi^+ \text{) ("5-prong")}) / \Gamma_{\text{total}} \quad \Gamma_{91} / \Gamma$$

$$\Gamma_{91} / \Gamma = (\Gamma_{92} + \Gamma_{93}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.097 ± 0.007 OUR FIT</b>				
<b>0.102 ± 0.011 OUR AVERAGE</b>				
0.097 ± 0.005 ± 0.011	419	GIBAUT	94B	CLEO $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
0.26 ± 0.06 ± 0.05		ACTON	92H	OPAL $E_{\text{cm}}^{ee} = 88.2\text{--}94.2 \text{ GeV}$
0.10 <sup>+0.05</sup> / <sub>-0.04</sub> ± 0.03		DECAMP	92C	ALEP 1989–1990 LEP runs
0.102 ± 0.029	13	BYLSMA	87	HRS $E_{\text{cm}}^{ee} = 29 \text{ GeV}$
0.16 ± 0.08 ± 0.04	4	BURCHAT	85	MRK2 $E_{\text{cm}}^{ee} = 29 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.16 ± 0.13 ± 0.04		BEHREND	89B	CELL $E_{\text{cm}}^{ee} = 14\text{--}47 \text{ GeV}$
0.3 ± 0.1 ± 0.2		BARTEL	85F	JADE $E_{\text{cm}}^{ee} = 34.6 \text{ GeV}$
0.13 ± 0.04	10	BELTRAMI	85	HRS Repl. by BYLSMA 87
1.0 ± 0.4	10	BEHREND	82	CELL Repl. by BEHREND 89B

$$\left[ \Gamma(h^- h^- h^+ \geq 1 \text{ neutrals } \nu_\tau) + \Gamma(3h^- 2h^+ \geq 0 \text{ neutrals } \nu_\tau) \right. \\ \left. (\text{ex. } K_S^0 \rightarrow \pi^- \pi^+) (\text{"5-prong"}) \right] / \Gamma_{\text{total}} \quad (\Gamma_{58} + \Gamma_{91}) / \Gamma \\ (\Gamma_{58} + \Gamma_{91}) / \Gamma = (0.3431\Gamma_{36} + 0.3431\Gamma_{38} + 0.1077\Gamma_{41} + \Gamma_{65} + \Gamma_{73} + \Gamma_{74} + \Gamma_{81} + \Gamma_{85} + \\ \Gamma_{92} + \Gamma_{93} + 0.285\Gamma_{110} + 0.888\Gamma_{125} + 0.9101\Gamma_{126}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

**5.28 ± 0.11 OUR FIT** Error includes scale factor of 1.2.

**5.4 ± 0.5 OUR AVERAGE**

5.05 ± 0.29 ± 0.65      570      DECAMP      92C ALEP      1989–1990 LEP runs

5.8 ± 0.7 ± 0.2      352      137 BEHREND      90 CELL       $E_{\text{cm}}^{ee} = 35 \text{ GeV}$

137 BEHREND 90 not independent of their  $\Gamma(h^- h^- h^+ \geq 1 \text{ neutrals } \nu_\tau) / \Gamma_{\text{total}}$  measurement.

$$\Gamma(3h^- 2h^+ \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{92} / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

**0.075 ± 0.007 OUR FIT**

**0.073 ± 0.008 OUR AVERAGE**

0.080 ± 0.011 ± 0.013      58      BUSKULIC      96 ALEP      LEP 1991–1993 data

0.077 ± 0.005 ± 0.009      295      GIBAUT      94B CLEO       $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

0.064 ± 0.023 ± 0.01      12      ALBRECHT      88B ARG       $E_{\text{cm}}^{ee} = 10 \text{ GeV}$

0.051 ± 0.020      7      BYLSMA      87 HRS       $E_{\text{cm}}^{ee} = 29 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.067 ± 0.030      5      138 BELTRAMI      85 HRS      Repl. by BYLSMA 87

138 The error quoted is statistical only.

$$\Gamma(3h^- 2h^+ \pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{93} / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

**0.022 ± 0.005 OUR FIT**

**0.021 ± 0.005 OUR AVERAGE**

0.018 ± 0.007 ± 0.012      18      BUSKULIC      96 ALEP      LEP 1991–1993 data

0.019 ± 0.004 ± 0.004      31      GIBAUT      94B CLEO       $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

0.051 ± 0.022      6      BYLSMA      87 HRS       $E_{\text{cm}}^{ee} = 29 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.067 ± 0.030      5      139 BELTRAMI      85 HRS      Repl. by BYLSMA 87

139 The error quoted is statistical only.

$$\Gamma(3h^- 2h^+ 2\pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{94} / \Gamma$$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
-----------	-----	-------------	------	---------

**< 0.011**      90      GIBAUT      94B CLEO       $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$$\Gamma((5\pi)^- \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{95} / \Gamma$$

$$\Gamma_{95} / \Gamma = (\Gamma_{26} + \frac{1}{4}\Gamma_{41} + \Gamma_{73} + \Gamma_{92} + 0.236\Gamma_{110} + 0.888\Gamma_{126}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-----------	-------------	------	---------

**0.74 ± 0.07 OUR FIT**

**0.61 ± 0.06 ± 0.08**      avg      140 GIBAUT      94B CLEO       $E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

140 Not independent of GIBAUT 94B B( $3h^- 2h^+ \nu_\tau$ ), PROCARIO 93 B( $h^- 4\pi^0 \nu_\tau$ ), and BORTOLETTO 93 B( $2h^- h^+ 2\pi^0 \nu_\tau$ )/B("3prong") measurements. Result is corrected for  $\eta$  contributions.

$$\Gamma(4h^-3h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ ("7-prong")})/\Gamma_{\text{total}} \quad \Gamma_{96}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.4 \times 10^{-6}$	90	EDWARDS	97B CLEO	$E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<1.8 \times 10^{-5}$	95	ACKERSTAFF	97J OPAL	1990–1995 LEP runs
$<2.9 \times 10^{-4}$	90	BYLSMA	87 HRS	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$

$$\Gamma(K^*(892)^- \geq 0(h^0 \neq K_S^0)\nu_\tau)/\Gamma_{\text{total}} \quad \Gamma_{97}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.94 \pm 0.27 \pm 0.15$	74	AKERS	94G OPAL	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94 \text{ GeV}$

$$\Gamma(K^*(892)^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}} \quad \Gamma_{98}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.33 ± 0.13 OUR AVERAGE</b>				
$1.19 \pm 0.15^{+0.13}_{-0.18}$	104	ALBRECHT	95H ARG	$E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6 \text{ GeV}$
$1.43 \pm 0.11 \pm 0.13$	475	<sup>141</sup> GOLDBERG	90 CLEO	$E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.9 \text{ GeV}$

<sup>141</sup>GOLDBERG 90 estimates that 10% of observed  $K^*(892)$  are accompanied by a  $\pi^0$ .

$$\Gamma(K^*(892)^- \nu_\tau)/\Gamma_{\text{total}} \quad \Gamma_{99}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.28 ± 0.08 OUR AVERAGE</b>				
$1.39 \pm 0.09 \pm 0.10$		<sup>142</sup> BUSKULIC	96 ALEP	LEP 1991–1993 data
$1.11 \pm 0.12$		<sup>143</sup> COAN	96 CLEO	$E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$
$1.42 \pm 0.22 \pm 0.09$		<sup>144</sup> ACCIARRI	95F L3	1991–1993 LEP runs
$1.23 \pm 0.21^{+0.11}_{-0.21}$	54	<sup>145</sup> ALBRECHT	88L ARG	$E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$
$1.9 \pm 0.3 \pm 0.4$	44	<sup>146</sup> TSCHIRHART	88 HRS	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
$1.5 \pm 0.4 \pm 0.4$	15	<sup>147</sup> AIHARA	87C TPC	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$
$1.3 \pm 0.3 \pm 0.3$	31	YELTON	86 MRK2	$E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.45 \pm 0.13 \pm 0.11$	273	<sup>148</sup> BUSKULIC	94F ALEP	Repl. by BUSKULIC 96
$1.7 \pm 0.7$	11	DORFAN	81 MRK2	$E_{\text{cm}}^{\text{ee}} = 4.2\text{--}6.7 \text{ GeV}$

<sup>142</sup>Not independent of BUSKULIC 96  $B(\pi^- \bar{K}^0 \nu_\tau)$  and  $B(K^- \pi^0 \nu_\tau)$  measurements.

<sup>143</sup>Not independent of COAN 96  $B(\pi^- \bar{K}^0 \nu_\tau)$  and BATTLE 94  $B(K^- \pi^0 \nu_\tau)$  measurements.  $K\pi$  final states are consistent with and assumed to originate from  $K^*(892)^-$  production.

<sup>144</sup>This result is obtained from their  $B(\pi^- \bar{K}^0 \nu_\tau)$  assuming all those decays originate in  $K^*(892)^-$  decays.

<sup>145</sup>The authors divide by  $\Gamma_1/\Gamma = 0.865$  to obtain this result.

<sup>146</sup>Not independent of TSCHIRHART 88  $\Gamma(\tau^- \rightarrow h^- \bar{K}^0 \geq 0 \text{ neutrals } \geq 0 K_L^0 \nu_\tau)/\Gamma(\text{total})$ .

<sup>147</sup>Decay  $\pi^-$  identified in this experiment, is assumed in the others.

<sup>148</sup>BUSKULIC 94F obtain this result from BUSKULIC 94F  $B(\bar{K}^0 \pi^- \nu_\tau)$  and BUSKULIC 94E  $B(K^- \pi^0 \nu_\tau)$  assuming all of those decays originate in  $K^*(892)^-$  decays.

$$\Gamma(K^*(892)^-\nu_\tau)/\Gamma(\pi^-\pi^0\nu_\tau) \quad \Gamma_{99}/\Gamma_{13}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.075±0.027</b>		149 ABREU	94K DLPH	LEP 1992 Z data
149 ABREU 94K quote $B(\tau^- \rightarrow K^*(892)^-\nu_\tau)B(K^*(892)^- \rightarrow K^-\pi^0)/B(\tau^- \rightarrow \rho^-\nu_\tau) = 0.025 \pm 0.009$ . We divide by $B(K^*(892)^- \rightarrow K^-\pi^0) = 0.333$ to obtain this result.				

$$\Gamma(K^*(892)^0 K^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}} \quad \Gamma_{100}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.32±0.08±0.12</b>	119	GOLDBERG	90 CLEO	$E_{\text{cm}}^{ee} = 9.4\text{--}10.9$ GeV

$$\Gamma(K^*(892)^0 K^- \nu_\tau)/\Gamma_{\text{total}} \quad \Gamma_{101}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.21 ±0.04 OUR AVERAGE</b>				
0.213±0.048		150 BARATE	98 ALEP	1991–1995 LEP runs
0.20 ±0.05 ±0.04	47	ALBRECHT	95H ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV
150 BARATE 98 measure the $K^-(\rho^0 \rightarrow \pi^+\pi^-)$ fraction in $\tau^- \rightarrow K^-\pi^+\pi^-\nu_\tau$ decays to be $(35 \pm 11)\%$ and derive this result from their measurement of $\Gamma(\tau^- \rightarrow K^-\pi^+\pi^-\nu_\tau)/\Gamma_{\text{total}}$ assuming the intermediate states are all $K^-\rho$ and $K^-K^*(892)^0$ .				

$$\Gamma(\bar{K}^*(892)^0 \pi^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}} \quad \Gamma_{102}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.38±0.11±0.13</b>	105	GOLDBERG	90 CLEO	$E_{\text{cm}}^{ee} = 9.4\text{--}10.9$ GeV

$$\Gamma(\bar{K}^*(892)^0 \pi^- \nu_\tau)/\Gamma_{\text{total}} \quad \Gamma_{103}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.22 ±0.05 OUR AVERAGE</b>				
0.209±0.058		151 BARATE	98 ALEP	1991–1995 LEP runs
0.25 ±0.10 ±0.05	27	ALBRECHT	95H ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV
151 BARATE 98 measure the $K^-K^*(892)^0$ fraction in $\tau^- \rightarrow K^-K^+\pi^-\nu_\tau$ decays to be $(87 \pm 13)\%$ and derive this result from their measurement of $\Gamma(\tau^- \rightarrow K^-K^+\pi^-\nu_\tau)/\Gamma_{\text{total}}$ .				

$$\Gamma((\bar{K}^*(892)\pi)^-\nu_\tau \rightarrow \pi^-\bar{K}^0\pi^0\nu_\tau)/\Gamma_{\text{total}} \quad \Gamma_{104}/\Gamma$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>0.106±0.037±0.032</b>	152 BARATE	98E ALEP	1991–1995 LEP runs
152 BARATE 98E determine the $\bar{K}^0\rho^-$ fraction in $\tau^- \rightarrow \pi^-\bar{K}^0\pi^0\nu_\tau$ decays to be $(0.64 \pm 0.09 \pm 0.10)$ and multiply their $B(\pi^-\bar{K}^0\pi^0\nu_\tau)$ measurement by one minus this fraction to obtain the quoted result.			

$$\Gamma(K_1(1270)^-\nu_\tau)/\Gamma_{\text{total}} \quad \Gamma_{105}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.41<sup>+0.41</sup><sub>-0.35</sub> ±0.10</b>	5	153 BAUER	94 TPC	$E_{\text{cm}}^{ee} = 29$ GeV

153 We multiply 0.41% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

$$\Gamma(K_1(1400)^- \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{106} / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.76^{+0.40}_{-0.33} \pm 0.20$	11	<sup>154</sup> BAUER	94 TPC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

<sup>154</sup> We multiply 0.76% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

$$[\Gamma(K_1(1270)^- \nu_\tau) + \Gamma(K_1(1400)^- \nu_\tau)] / \Gamma_{\text{total}} \qquad (\Gamma_{105} + \Gamma_{106}) / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.17^{+0.41}_{-0.37} \pm 0.29$	16	<sup>155</sup> BAUER	94 TPC	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

<sup>155</sup> We multiply 1.17% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error. Not independent of BAUER 94  $B(K_1(1270)^- \nu_\tau)$  and BAUER 94  $B(K_1(1400)^- \nu_\tau)$  measurements.

$$\Gamma(K_2^*(1430)^- \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{107} / \Gamma$$

VALUE (%)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 0.3$	95		TSCHIRHART 88	HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 0.33$	95	<sup>156</sup>	ACCIARRI	95F L3	1991–1993 LEP runs
$< 0.9$	95	0	DORFAN	81 MRK2	$E_{\text{cm}}^{ee} = 4.2\text{--}6.7 \text{ GeV}$

<sup>156</sup> ACCIARRI 95F quote  $B(\tau^- \rightarrow K^*(1430)^- \rightarrow \pi^- \bar{K}^0 \nu_\tau) < 0.11\%$ . We divide by  $B(K^*(1430)^- \rightarrow \pi^- \bar{K}^0) = 0.33$  to obtain the limit shown.

$$\Gamma(a_0(980)^- \geq 0 \text{ neutrals } \nu_\tau) / \Gamma_{\text{total}} \times B(a_0(980)^- \rightarrow K^0 K^-) \qquad \Gamma_{108} / \Gamma \times B$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.8 \times 10^{-4}$	90	GOLDBERG 90	CLEO	$E_{\text{cm}}^{ee} = 9.4\text{--}10.9 \text{ GeV}$

$$\Gamma(\eta \pi^- \nu_\tau) / \Gamma_{\text{total}} \qquad \Gamma_{109} / \Gamma$$

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 1.4$	95	0	BARTELT	96 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 6.2$	95		BUSKULIC	97C ALEP	1991–1994 LEP runs
$< 3.4$	95		ARTUSO	92 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$
$< 90$	95		ALBRECHT	88M ARG	$E_{\text{cm}}^{ee} \approx 10 \text{ GeV}$
$< 140$	90		BEHREND	88 CELL	$E_{\text{cm}}^{ee} = 14\text{--}46.8 \text{ GeV}$
$< 180$	95		BARINGER	87 CLEO	$E_{\text{cm}}^{ee} = 10.5 \text{ GeV}$
$< 250$	90	0	COFFMAN	87 MRK3	$E_{\text{cm}}^{ee} = 3.77 \text{ GeV}$
510 $\pm 100 \pm 120$		65	DERRICK	87 HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
$< 100$	95		GAN	87B MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

$$\Gamma(\eta\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{110}/\Gamma$$

VALUE (%)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	-----	------	-------------	------	---------

**0.174±0.024 OUR FIT**

**0.173±0.024 OUR AVERAGE**

0.18 ±0.04 ±0.02			BUSKULIC	97C ALEP	1991–1994 LEP
------------------	--	--	----------	----------	---------------

0.17 ±0.02 ±0.02		125	ARTUSO	92 CLEO	$E_{\text{cm}}^{\text{ee}} \approx 10.6$ runs GeV
------------------	--	-----	--------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.10	95		ALBRECHT	88M ARG	$E_{\text{cm}}^{\text{ee}} \approx 10$ GeV
-------	----	--	----------	---------	---

<2.10	95		BARINGER	87 CLEO	$E_{\text{cm}}^{\text{ee}} = 10.5$ GeV
-------	----	--	----------	---------	--

4.20 $\begin{smallmatrix} +0.70 \\ -1.20 \end{smallmatrix}$ ±1.60		157	GAN	87 MRK2	$E_{\text{cm}}^{\text{ee}} = 29$ GeV
---	--	-----	-----	---------	--------------------------------------

<sup>157</sup> Highly correlated with GAN 87  $\Gamma(\pi^- 3\pi^0\nu_\tau)/\Gamma(\text{total})$  value.

$$\Gamma(\eta\pi^-\pi^0\pi^0\nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{111}/\Gamma$$

VALUE (units 10 <sup>-4</sup> )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	------	-------------	------	---------

**1.4±0.6±0.3** 15 BERGFELD 97 CLEO  $E_{\text{cm}}^{\text{ee}} = 10.6$  GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.3	95		ARTUSO	92 CLEO	$E_{\text{cm}}^{\text{ee}} \approx 10.6$ GeV
-------	----	--	--------	---------	--

<120	95		ALBRECHT	88M ARG	$E_{\text{cm}}^{\text{ee}} \approx 10$ GeV
------	----	--	----------	---------	--

$$\Gamma(\eta K^-\nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{112}/\Gamma$$

VALUE (units 10 <sup>-4</sup> )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	------	-------------	------	---------

**2.7±0.6 OUR AVERAGE**

2.9 $\begin{smallmatrix} +1.3 \\ -1.2 \end{smallmatrix}$ ±0.7			BUSKULIC	97C ALEP	1991–1994 LEP runs
---	--	--	----------	----------	--------------------

2.6±0.5±0.5	85		BARTELT	96 CLEO	$E_{\text{cm}}^{\text{ee}} \approx 10.6$ GeV
-------------	----	--	---------	---------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.7	95		ARTUSO	92 CLEO	$E_{\text{cm}}^{\text{ee}} \approx 10.6$ GeV
------	----	--	--------	---------	--

$$\Gamma(\eta\pi^+\pi^-\pi^-\geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{113}/\Gamma$$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
-----------	-----	-------------	------	---------

**<0.3** 90 ABACHI 87B HRS  $E_{\text{cm}}^{\text{ee}} = 29$  GeV

$$\Gamma(\eta\pi^-\pi^+\pi^-\nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{114}/\Gamma$$

VALUE (units 10 <sup>-4</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

**3.4  $\begin{smallmatrix} +0.6 \\ -0.5 \end{smallmatrix}$  ±0.6** 89 BERGFELD 97 CLEO  $E_{\text{cm}}^{\text{ee}} = 10.6$  GeV

$$\Gamma(\eta a_1(1260)^-\nu_\tau \rightarrow \eta\pi^-\rho^0\nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{115}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**<3.9 × 10<sup>-4</sup>** 90 BERGFELD 97 CLEO  $E_{\text{cm}}^{\text{ee}} = 10.6$  GeV

$$\Gamma(\eta\eta\pi^-\nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{116}/\Gamma$$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; 1.1</b>	95	ARTUSO	92 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<83	95	ALBRECHT	88M ARG	$E_{\text{cm}}^{ee} \approx 10 \text{ GeV}$

$$\Gamma(\eta\eta\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{117}/\Gamma$$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; 2.0</b>	95	ARTUSO	92 CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<90	95	ALBRECHT	88M ARG	$E_{\text{cm}}^{ee} \approx 10 \text{ GeV}$

$$\Gamma(\eta'(958)\pi^-\nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{118}/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; <math>7.4 \times 10^{-5}</math></b>	90	BERGFELD	97 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$$\Gamma(\eta'(958)\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{119}/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; <math>8.0 \times 10^{-5}</math></b>	90	BERGFELD	97 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$$\Gamma(\phi\pi^-\nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{120}/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; <math>2.0 \times 10^{-4}</math></b>	90	<sup>158</sup> AVERY	97 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< $3.5 \times 10^{-4}$	90	ALBRECHT	95H ARG	$E_{\text{cm}}^{ee} = 9.4\text{--}10.6 \text{ GeV}$
<sup>158</sup> AVERY 97 limit varies from $(1.2\text{--}2.0) \times 10^{-4}$ depending on decay model assumptions.				

$$\Gamma(\phi K^-\nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{121}/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; <math>6.7 \times 10^{-5}</math></b>	90	<sup>159</sup> AVERY	97 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$
<sup>159</sup> AVERY 97 limit varies from $(5.4\text{--}6.7) \times 10^{-5}$ depending on decay model assumptions.				

$$\Gamma(f_1(1285)\pi^-\nu_\tau)/\Gamma_{\text{total}} \qquad \Gamma_{122}/\Gamma$$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>5.8^{+1.4}_{-1.3} \pm 1.8</math></b>	54	BERGFELD	97 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$$\Gamma(f_1(1285)\pi^-\nu_\tau \rightarrow \eta\pi^-\pi^+\pi^-\nu_\tau)/\Gamma(\eta\pi^-\pi^+\pi^-\nu_\tau) \qquad \Gamma_{123}/\Gamma_{114}$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.55 \pm 0.14</math></b>	BERGFELD	97 CLEO	$E_{\text{cm}}^{ee} = 10.6 \text{ GeV}$

$$\frac{\Gamma(h^- \omega \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}}{\Gamma_{124}/\Gamma = (\Gamma_{125} + \Gamma_{126})/\Gamma} \quad \Gamma_{124}/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.36 ± 0.08 OUR FIT</b>					
<b>1.65 ± 0.3 ± 0.2</b>	avg	1513	ALBRECHT	88M ARG	$E_{\text{cm}}^{ee} \approx 10 \text{ GeV}$

$$\frac{\Gamma(h^- \omega \nu_\tau)/\Gamma_{\text{total}}}{\Gamma_{125}/\Gamma} \quad \Gamma_{125}/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)		EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.93 ± 0.06 OUR FIT</b>					
<b>1.92 ± 0.07 OUR AVERAGE</b>					
1.91 ± 0.07 ± 0.06	f&a	5803	BUSKULIC	97C ALEP	1991-1994 LEP runs
1.95 ± 0.07 ± 0.11	avg	2223	<sup>160</sup> BALEST	95C CLEO	$E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$
1.60 ± 0.27 ± 0.41	f&a	139	BARINGER	87 CLEO	$E_{\text{cm}}^{ee} = 10.5 \text{ GeV}$

<sup>160</sup> Not independent of BALEST 95C  $B(\tau^- \rightarrow h^- \omega \nu_\tau)/B(\tau^- \rightarrow h^- h^- h^+ \pi^0 \nu_\tau)$  value.

$$\frac{[\Gamma(h^- \rho \pi^0 \nu_\tau) + \Gamma(h^- \rho^+ h^- \nu_\tau) + \Gamma(h^- \rho^- h^+ \nu_\tau) + \Gamma(h^- \omega \nu_\tau)]}{\Gamma(h^- h^- h^+ \pi^0 \nu_\tau)} \quad (\Gamma_{68} + \Gamma_{69} + \Gamma_{70} + \Gamma_{125})/\Gamma_{60}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;0.81</b>	95	<sup>161</sup> ALBRECHT	91D ARG	$E_{\text{cm}}^{ee} = 9.4-10.6 \text{ GeV}$

<sup>161</sup> ALBRECHT 91D not independent of their  $\Gamma(h^- \omega \nu_\tau)/\Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$  (ex.  $K^0$ ),  $\Gamma(h^- \rho \pi^0 \nu_\tau)/\Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$ ,  $\Gamma(h^- \rho^+ h^- \nu_\tau)/\Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$ , and  $\Gamma(h^- \rho^- h^+ \nu_\tau)/\Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$  values.