

Z Boson – THIS IS PART 1 OF 2

To reduce the size of this section's PostScript file, we have divided it into two PostScript files. We present the following index:

PART 1

Page #	Section name
1	Mass
17	Width
18	Decay modes
19	Partial widths
21	Branching ratios

PART 2

Page #	Section name
35	Average particle multiplicities in hadronic Z decay
42	Hadronic pole cross section
42	Vector couplings to charged leptons
44	Axial-vector couplings to charged leptons
45	Couplings to neutral leptons
46	Asymmetry parameters
48	Transverse spin correlations
49	Charge asymmetry
55	References



$$J = 1$$

THE Z BOSON

Revised February 1998 by C. Caso (Univ. of Genova) and A. Gurtu (Tata Inst.)

Precision measurements at the Z -boson resonance using electron–positron colliding beams began in 1989 at the SLC and at LEP. During 1989–95, the four CERN experiments have made high-statistics studies of the Z . The availability of longitudinally polarized electron beams at the SLC since 1993 has enabled a precision determination of the effective electroweak mixing angle $\sin^2\bar{\theta}_W$ that is competitive with the CERN results on this parameter.

The Z -boson properties reported in this section may broadly be categorized as:

- The standard ‘lineshape’ parameters of the Z consisting of its mass, M_Z , its total width, Γ_Z , and its partial decay widths, $\Gamma(\text{hadrons})$, and $\Gamma(\ell\bar{\ell})$ where $\ell = e, \mu, \tau, \nu$;
- Z asymmetries in leptonic decays and extraction of Z couplings to charged and neutral leptons;
- The b - and c -quark-related partial widths and charge asymmetries which require special techniques;
- Determination of Z decay modes and the search for modes that violate known conservation laws;
- Average particle multiplicities in hadronic Z decay.

For the lineshape-related Z properties there are no new published LEP results after those included in the 1994 edition of this compilation. The reason for this is the identification in mid 1995 of a new systematic effect which shifts the LEP energy by a few MeV. This is due to a drift of the dipole field in the LEP magnets caused by parasitic currents generated by electrically powered trains in the Geneva area. The LEP Energy Working Group has been studying the implications of this for the Z -lineshape properties which would be obtained after analysis of the high statistics 1993–95 data. The main consequence of this effect is expected to be in the determination of the Z mass.

Details on Z -parameter determination and the study of $Z \rightarrow b\bar{b}, c\bar{c}$ at LEP and SLC are given in this note.

The standard ‘lineshape’ parameters of the Z are determined with increasing precision from an analysis of the production cross sections of these final states in e^+e^- collisions. The $Z \rightarrow \nu\bar{\nu}(\gamma)$ state is identified directly by detecting single photon production and indirectly by subtracting the visible partial widths from the total width. Inclusion in this analysis of the forward-backward asymmetry of charged leptons, $A_{FB}^{(0,\ell)}$, of the τ polarization, $P(\tau)$, and its forward-backward asymmetry, $P(\tau)^{fb}$, enables the separate determination of the effective vector (\bar{g}_V) and axial vector (\bar{g}_A) couplings of the Z to these leptons and the ratio (\bar{g}_V/\bar{g}_A) which is related to the effective electroweak mixing angle $\sin^2\bar{\theta}_W$ (see the “Electroweak Model and Constraints on New Physics” Review).

Determination of the b - and c -quark-related partial widths and charge asymmetries involves tagging the b and c quarks. Traditionally this was done by requiring the presence of a prompt lepton in the event with high momentum and high transverse momentum (with respect to the accompanying jet).

Precision vertex measurement with silicon detectors has enabled one to do impact parameter and lifetime tagging. Neural-network techniques have also been used to classify events as b or non- b on a statistical basis using event-shape variables. Finally, the presence of a charmed meson (D/D^*) has been used to tag heavy quarks.

Z-parameter determination

LEP is run at a few energy points on and around the Z mass constituting an energy ‘scan.’ The shape of the cross-section variation around the Z peak can be described by a Breit-Wigner *ansatz* with an energy-dependent total width [1–3]. The **three** main properties of this distribution, viz., the **position** of the peak, the **width** of the distribution, and the **height** of the peak, determine respectively the values of M_Z , Γ_Z , and $\Gamma(e^+e^-) \times \Gamma(f\bar{f})$, where $\Gamma(e^+e^-)$ and $\Gamma(f\bar{f})$ are the electron and fermion partial widths of the Z . The quantitative determination of these parameters is done by writing analytic expressions for these cross sections in terms of the parameters and fitting the calculated cross sections to the measured ones by varying these parameters, taking properly into account all the errors. Single-photon exchange (σ_γ^0) and γ - Z interference ($\sigma_{\gamma Z}^0$) are included, and the large ($\sim 25\%$) initial-state radiation (ISR) effects are taken into account by convoluting the analytic expressions over a ‘Radiator Function’ [1–4] $H(s, s')$. Thus for the process $e^+e^- \rightarrow f\bar{f}$:

$$\sigma_f(s) = \int H(s, s') \sigma_f^0(s') ds' \quad (1)$$

$$\sigma_f^0(s) = \sigma_Z^0 + \sigma_\gamma^0 + \sigma_{\gamma Z}^0 \quad (2)$$

$$\sigma_Z^0 = \frac{12\pi}{M_Z^2} \frac{\Gamma(e^+e^-)\Gamma(f\bar{f})}{\Gamma_Z^2} \frac{s \Gamma_Z^2}{(s - M_Z^2)^2 + s^2\Gamma_Z^2/M_Z^2} \quad (3)$$

$$\sigma_\gamma^0 = \frac{4\pi\alpha^2(s)}{3s} Q_f^2 N_c^f \quad (4)$$

$$\begin{aligned} \sigma_{\gamma Z}^0 = & -\frac{2\sqrt{2}\alpha(s)}{3} (Q_f G_F N_c^f g_{Ve} g_{Vf}) \\ & \times \frac{(s - M_Z^2) M_Z^2}{(s - M_Z^2)^2 + s^2 \Gamma_Z^2 / M_Z^2} \end{aligned} \quad (5)$$

where Q_f is the charge of the fermion, $N_c^f = 3(1)$ for quark (lepton) and g_{Vf} is the neutral vector coupling of the Z to the fermion-antifermion pair $f\bar{f}$.

Since $\sigma_{\gamma Z}^0$ is expected to be much less than σ_Z^0 , the LEP Collaborations have generally calculated the interference term in the framework of the Standard Model using the best known values of g_V . This fixing of $\sigma_{\gamma Z}^0$ leads to a tighter constraint on M_Z and consequently a smaller error on its fitted value.

Defining

$$A_f = 2 \frac{g_{Vf} \cdot g_{Af}}{(g_{Vf}^2 + g_{Af}^2)} \quad (6)$$

where g_{Af} is the neutral axial-vector coupling of the Z to $f\bar{f}$, the lowest-order expressions for the various lepton-related asymmetries on the Z pole are [5–7] $A_{FB}^{(0,\ell)} = (3/4)A_e A_f$, $P(\tau) = -A_\tau$, $P(\tau)^{fb} = -(3/4)A_e$, $A_{LR} = A_e$. The full analysis takes into account the energy dependence of the asymmetries. Experimentally A_{LR} is defined as $(\sigma_L - \sigma_R)/(\sigma_L + \sigma_R)$ where $\sigma_{L(R)}$ are the $e^+e^- \rightarrow Z$ production cross sections with left- (right)-handed electrons.

In terms of g_A and g_V , the partial decay width of the Z to $f\bar{f}$ can be written as

$$\Gamma(f\bar{f}) = \frac{G_F M_Z^3}{6\sqrt{2}\pi} (g_{Vf}^2 + g_{Af}^2) N_c^f (1 + \delta_{\text{QED}})(1 + \delta_{\text{QCD}}) \quad (7)$$

where $\delta_{\text{QED}} = 3\alpha Q_f^2/4\pi$ accounts for final-state photonic corrections and $\delta_{\text{QCD}} = 0$ for leptons and $\delta_{\text{QCD}} = (\alpha_s/\pi) + 1.409(\alpha_s/\pi)^2 - 12.77(\alpha_s/\pi)^3$ for quarks, α_s being the strong coupling constant at $\mu = M_Z$.

In the above framework, the QED radiative corrections have been explicitly taken into account by convoluting over the ISR and allowing the electromagnetic coupling constant to run [8]: $\alpha(s) = \alpha/(1 - \Delta\alpha)$. On the other hand, weak radiative corrections that depend upon the assumptions of the electroweak theory and on the values of the unknown M_{top} and M_{Higgs} are accounted for by **absorbing them into the couplings**, which are then called the *effective* couplings \bar{g}_V and \bar{g}_A (or alternatively the effective parameters of the \star scheme of Kennedy and Lynn [9]).

S-matrix approach to the Z

While practically all experimental analyses of LEP/SLC data have followed the ‘Breit-Wigner’ approach described above, an alternative S-matrix-based analysis is also possible. The Z , like all unstable particles, is associated with a complex pole in the S matrix. The pole position is process independent and gauge invariant. The mass, \bar{M}_Z , and width, $\bar{\Gamma}_Z$, can be defined in terms of the pole in the energy plane via [10–13]

$$\bar{s} = \bar{M}_Z^2 - i\bar{M}_Z\bar{\Gamma}_Z \quad (8)$$

leading to the relations

$$\begin{aligned} \bar{M}_Z &= M_Z / \sqrt{1 + \Gamma_Z^2/M_Z^2} \\ &\approx M_Z - 34.1 \text{ MeV} \end{aligned} \quad (9)$$

$$\begin{aligned} \bar{\Gamma}_Z &= \Gamma_Z / \sqrt{1 + \Gamma_Z^2/M_Z^2} \\ &\approx \Gamma_Z - 0.9 \text{ MeV} . \end{aligned} \quad (10)$$

Some authors [14] choose to define the Z mass and width via

$$\bar{s} = (\bar{M}_Z - \frac{i}{2}\bar{\Gamma}_Z)^2 \quad (11)$$

which yields $\bar{M}_Z \approx M_Z - 26$ MeV, $\bar{\Gamma}_Z \approx \Gamma_Z - 1.2$ MeV.

The L3 and OPAL Collaborations at LEP (ACCIARRI 97K and ACKERSTAFF 97C) have analyzed their data using the S-matrix approach as defined in Eq. (8), in addition to the conventional one. They observe a downward shift in the Z mass as expected.

Handling the large-angle e^+e^- final state

Unlike other $f\bar{f}$ decay final states of the Z , the e^+e^- final state has a contribution not only from the s -channel but also from the t -channel and s - t interference. The full amplitude is not amenable to fast calculation, which is essential if one has to carry out minimization fits within reasonable computer time. The usual procedure is to calculate the non- s channel part of the cross section separately using the Standard Model programs ALIBABA [15] or TOPAZ0 [16] with the measured value of M_{top} , and the ‘central’ value of M_{Higgs} (300 GeV) and add it to the s -channel cross section calculated as for other channels. This leads to two additional sources of error in the analysis: firstly, the theoretical calculation in ALIBABA itself is known to be accurate to $\sim 0.5\%$, and secondly, there is uncertainty due to the error on M_{top} and the unknown value of M_{Higgs} (60–1000 GeV). These additional errors are propagated into the analysis by including them in the systematic error on the e^+e^- final state.

Errors due to uncertainty in LEP energy determination [17–21]

The systematic errors related to the LEP energy measurement can be classified as:

- The absolute energy scale error;
- Energy-point-to-energy-point errors due to the non-linear response of the magnets to the exciting currents;
- Energy-point-to-energy-point errors due to possible higher-order effects in the relationship between the dipole field and beam energy;
- Energy reproducibility errors due to various unknown uncertainties in temperatures, tidal effects, corrector settings, RF status, *etc.* Since one groups together data taken at ‘nominally same’ energies in different fills, it can be assumed that these errors are uncorrelated and are reduced by $\sqrt{\overline{N}_{\text{fill}}}$ where $\overline{N}_{\text{fill}}$ is the (luminosity weighted) effective number of fills at a particular energy point.

At each energy point the last two errors can be summed into one point-to-point error.

Choice of fit parameters

The LEP Collaborations have chosen the following primary set of parameters for fitting: M_Z , Γ_Z , σ_{hadron}^0 , $R(\text{lepton})$, $A_{FB}^{(0,\ell)}$, where $R(\text{lepton}) = \Gamma(\text{hadrons})/\Gamma(\text{lepton})$, $\sigma_{\text{hadron}}^0 = 12\pi\Gamma(e^+e^-)\Gamma(\text{hadrons})/M_Z^2\Gamma_Z^2$. With a knowledge of these fitted parameters and their covariance matrix, any other parameter can be derived. The main advantage of these parameters is that they form the **least correlated** set of parameters, so that it becomes easy to combine results from the different LEP experiments.

Thus, the most general fit carried out to cross section and asymmetry data determines the **nine parameters**: M_Z , Γ_Z , σ_{hadron}^0 , $R(e)$, $R(\mu)$, $R(\tau)$, $A_{FB}^{(0,e)}$, $A_{FB}^{(0,\mu)}$, $A_{FB}^{(0,\tau)}$. Assumption

of lepton universality leads to a **five-parameter fit** determining M_Z , Γ_Z , σ_{hadron}^0 , $R(\text{lepton})$, $A_{FB}^{(0,\ell)}$. The use of **only** cross-section data leads to six- or four-parameter fits if lepton universality is or is not assumed, *i.e.*, $A_{FB}^{(0,\ell)}$ values are not determined.

In order to determine the best values of the effective vector and axial vector couplings of the charged leptons to the Z , the above mentioned nine- and five-parameter fits are carried out with added constraints from the measured values of A_τ and A_e obtained from τ polarization studies at LEP and the determination of A_{LR} at SLC.

Combining results from the LEP and SLC experiments [22]

Each LEP experiment provides the values of the parameters mentioned above together with the full covariance matrix. The statistical and experimental systematic errors are assumed to be uncorrelated among the four experiments. The sources of **common** systematic errors are i) the LEP energy uncertainties, and ii) the effect of theoretical uncertainty in calculating the small-angle Bhabha cross section for luminosity determination and in estimating the non- s channel contribution to the large-angle Bhabha cross section. Using this information, a full covariance matrix, V , of all the input parameters is constructed and a combined parameter set is obtained by minimizing $\chi^2 = \Delta^T V^{-1} \Delta$, where Δ is the vector of residuals of the combined parameter set to the results of individual experiments.

Non-LEP measurement of a Z parameter, (*e.g.*, $\Gamma(e^+e^-)$ from SLD) is included in the overall fit by calculating its value using the fit parameters and constraining it to the measurement.

Study of $Z \rightarrow b\bar{b}$ and $Z \rightarrow c\bar{c}$

In the sector of c - and b -physics the LEP experiments have measured the ratios of partial widths $R_b = \Gamma(Z \rightarrow b\bar{b})/\Gamma(Z \rightarrow \text{hadrons})$ and $R_c = \Gamma(Z \rightarrow c\bar{c})/\Gamma(Z \rightarrow \text{hadrons})$ and the forward-backward (charge) asymmetries $A_{FB}^{b\bar{b}}$ and $A_{FB}^{c\bar{c}}$. Several of the analyses have also determined other quantities, in particular the semileptonic branching ratios, $B(b \rightarrow \ell)$ and $B(b \rightarrow c \rightarrow \ell^+)$, the average $B^0\bar{B}^0$ mixing parameter $\bar{\chi}$ and the probabilities for a c -quark to fragment into a D^+ , a D_s , a D^{*+} , or a charmed baryon. The latter measurements do not concern properties of the Z boson and hence they are not covered in this section. However, they are correlated with the electroweak parameters, and since the mixture of b hadrons is different from the one at the $\Upsilon(4S)$, their values might differ from those measured at the $\Upsilon(4S)$.

All the above quantities are correlated to each other since:

- Several analyses (for example the lepton fits) determine more than one parameter simultaneously;
- Some of the electroweak parameters depend explicitly on the values of other parameters (for example R_b depends on R_c);
- Common tagging and analysis techniques produce common systematic uncertainties.

The LEP Electroweak Heavy Flavour Working Group has developed [23] a procedure for combining the measurements taking into account known sources of correlation. The combining procedure determines eleven parameters: the four parameters of interest in the electroweak sector, R_b , R_c , $A_{FB}^{b\bar{b}}$, and $A_{FB}^{c\bar{c}}$ and, in addition, $B(b \rightarrow \ell)$, $B(b \rightarrow c \rightarrow \ell^+)$, $\bar{\chi}$, $f(D^+)$, $f(D_s)$, $f(c_{\text{baryon}})$ and $P(c \rightarrow D^{*+}) \times B(D^{*+} \rightarrow \pi^+ D^0)$, to take into account their correlations with the electroweak parameters.

Before the fit both the peak and off-peak asymmetries are translated to $\sqrt{s} = 91.26$ GeV using the predicted dependence from ZFITTER [4].

Summary of the measurements and of the various kinds of analysis

The measurements of R_b and R_c fall into two classes. In the first, named single-tag measurement, a method for selecting b and c events is applied and the number of tagged events is counted. The second technique, named double-tag measurement, is based on the following principle: if the number of events with a single hemisphere tagged is N_t and with both hemispheres tagged is N_{tt} , then given a total number of N_{had} hadronic Z decays one has:

$$\frac{N_t}{2N_{\text{had}}} = \varepsilon_b R_b + \varepsilon_c R_c + \varepsilon_{uds}(1 - R_b - R_c) \quad (12)$$

$$\frac{N_{tt}}{N_{\text{had}}} = C_b \varepsilon_b^2 R_b + C_c \varepsilon_c^2 R_c + C_{uds} \varepsilon_{uds}^2 (1 - R_b - R_c) \quad (13)$$

where ε_b , ε_c , and ε_{uds} are the tagging efficiencies per hemisphere for b , c , and light quark events, and $C_q \neq 1$ accounts for the fact that the tagging efficiencies between the hemispheres may be correlated. In tagging the b one has $\varepsilon_b \gg \varepsilon_c \gg \varepsilon_{uds}$, $C_b \approx 1$. Neglecting the c and uds background and the hemisphere correlations, these equations give:

$$\varepsilon_b = 2N_{tt}/N_t \quad (14)$$

$$R_b = N_t^2 / (4N_{tt}N_{\text{had}}) . \quad (15)$$

The double-tagging method has thus the great advantage that the tagging efficiency is directly derived from the data, reducing the systematic error of the measurement. The backgrounds, dominated by $c\bar{c}$ events, obviously complicate this

simple picture, and their level must still be inferred by other means. The rate of charm background in these analyses depends explicitly on the value of R_c . The correlations in the tagging efficiencies between the hemispheres (due for instance to correlations in momentum between the b hadrons in the two hemispheres) are small but nevertheless lead to further systematic uncertainties.

The measurements in the b - and c -sector can be grouped in the following categories:

- Lepton fits which use hadronic events with one or more leptons in the final state. Each analysis usually gives several electroweak parameters chosen among: R_b , R_c , $A_{FB}^{b\bar{b}}$, $A_{FB}^{c\bar{c}}$, $B(b \rightarrow \ell)$, $B(b \rightarrow c \rightarrow \ell^+)$ and $\bar{\chi}$. The output parameters are then correlated. The dominant sources of systematics are due to lepton identification, to other semileptonic branching ratios and to the modelling of the semileptonic decay;
- Event shape tag for R_b ;
- Lifetime (and lepton) double-tagging measurements of R_b . These are the most precise measurements of R_b and obviously dominate the combined result. The main sources of systematics come from the charm contamination and from estimating the hemisphere b -tagging efficiency correlation. The charm rejection has been improved (and hence the systematic errors reduced) by using either the information of the secondary vertex invariant mass or the information from the energy of all particles at the secondary vertex and their rapidity;

- Measurements of $A_{FB}^{b\bar{b}}$ using lifetime tagged events with a hemisphere charge measurement. Their contribution to the combined result has roughly the same weight as the lepton fits;
- Analyses with $D/D^{*\pm}$ to measure R_c . These measurements make use of several different tagging techniques (inclusive/exclusive double tag, inclusive single/double tag, exclusive double tag, reconstruction of all weakly decaying D states) and no assumptions are made on the energy dependence of charm fragmentation;
- Analyses with $D/D^{*\pm}$ to measure $A_{FB}^{c\bar{c}}$ or simultaneously $A_{FB}^{b\bar{b}}$ and $A_{FB}^{c\bar{c}}$;
- Measurements of A_b and A_c from SLD, using several tagging methods (lepton, D/D^* , and impact parameter). These quantities are directly extracted from a measurement of the left–right forward–backward asymmetry in $c\bar{c}$ and $b\bar{b}$ production using a polarized electron beam.

Averaging procedure

All the measurements are provided by the LEP Collaborations in the form of tables with a detailed breakdown of the systematic errors of each measurement and its dependence on other electroweak parameters.

The averaging proceeds via the following steps:

- Define and propagate a consistent set of external inputs such as branching ratios, hadron lifetimes, fragmentation models *etc.* All the measurements are also consistently checked to ensure that all use a common set of assumptions (for instance since the QCD corrections for the forward–backward asymmetries are strongly dependent on the experimental conditions, the data are corrected before combining);
- Form the full (statistical and systematic) covariance matrix of the measurements. The systematic correlations between different analyses are calculated from the detailed error breakdown in the measurement tables. The correlations relating several measurements made by the same analysis are also used;
- Take into account any explicit dependence of a measurement on the other electroweak parameters. As an example of this dependence we illustrate the case of the double-tag measurement of R_b , where c -quarks constitute the main background. The normalization of the charm contribution is not usually fixed by the data and the measurement of R_b depends on the assumed value of R_c , which can be written as:

$$R_b = R_b^{\text{meas}} + a(R_c) \frac{(R_c - R_c^{\text{used}})}{R_c}, \quad (16)$$

where R_b^{meas} is the result of the analysis which assumed a value of $R_c = R_c^{\text{used}}$ and $a(R_c)$ is the constant which gives the dependence on R_c ;

- Perform a χ^2 minimization with respect to the combined electroweak parameters.

After the fit the average peak asymmetries $A_{FB}^{c\bar{c}}$ and $A_{FB}^{b\bar{b}}$ are corrected for the energy shift and for QED, γ exchange, and γZ interference effects to obtain the corresponding pole asymmetries $A_{FB}^{0,c}$ and $A_{FB}^{0,b}$. A small correction is also applied to both R_b and R_c to account for the contribution of γ exchange.

References

1. R.N. Cahn, Phys. Rev. **D36**, 2666 (1987).
2. F.A. Berends *et al.*, “Z Physics at LEP 1”, CERN Report 89-08 (1989), Vol. 1, eds. G. Altarelli, R. Kleiss, and C. Verzegnassi, p. 89.
3. A. Borrelli *et al.*, Nucl. Phys. **B333**, 357 (1990).
4. D. Bardin *et al.*, Nucl. Phys. **B351**, 1 (1991).
5. M. Consoli *et al.*, “Z Physics at LEP 1”, CERN Report 89-08 (1989), Vol. 1, eds. G. Altarelli, R. Kleiss, and C. Verzegnassi, p. 7.
6. M. Bohm *et al.*, *ibid*, p. 203.
7. S. Jadach *et al.*, *ibid*, p. 235.
8. G. Burgers *et al.*, *ibid*, p. 55.
9. D.C. Kennedy and B.W. Lynn, SLAC-PUB 4039 (1986, revised 1988).
10. R. Stuart, Phys. Lett. **B262**, 113 (1991).
11. A. Sirlin, Phys. Rev. Lett. **67**, 2127 (1991).
12. A. Leike, T. Riemann, and J. Rose, Phys. Lett. **B273**, 513 (1991).
13. See also D. Bardin *et al.*, Phys. Lett. **B206**, 539 (1988).
14. S. Willenbrock and G. Valencia, Phys. Lett. **B259**, 373 (1991).
15. W. Beenakker, F.A. Berends, and S.C. van der Marck, Nucl. Phys. **B349**, 323 (1991).
16. K. Miyabayashi *et al.* (TOPAZ Collaboration) Phys. Lett. **B347**, 171 (1995).

17. R. Assmann *et al.* (Working Group on LEP Energy), *Z. Phys.* **C66**, 567 (1995).
 18. L. Arnaudon *et al.* (Working Group on LEP Energy and LEP Collaborations), *Phys. Lett.* **B307**, 187 (1993).
 19. L. Arnaudon *et al.* (Working Group on LEP Energy), CERN-PPE/92-125 (1992).
 20. L. Arnaudon *et al.*, *Phys. Lett.* **B284**, 431 (1992).
 21. R. Baily *et al.*, 'LEP Energy Calibration' CERN-SL 90-95.
 22. The LEP Collaborations: ALEPH, DELPHI, L3, OPAL, the LEP Electroweak Working Group, and the SLD Heavy Flavour Group:
CERN-PPE/97-154 (1997); CERN-PPE/96-183 (1996);
CERN-PPE/95-172 (1995); CERN-PPE/94-187 (1994);
CERN-PPE/93-157 (1993).
 23. The LEP Experiments: ALEPH, DELPHI, L3, and OPAL
Nucl. Instrum. Methods **A378**, 101 (1996).
-

Z MASS

The fit is performed using the Z mass and width, the Z hadronic pole cross section, the ratios of hadronic to leptonic partial widths, and the Z pole forward-backward lepton asymmetries. We believe that this set is the most free of correlations. Common systematic errors are taken into account. For more details, see the 'Note on the Z Boson.'

The Z -boson mass listed here corresponds to a Breit-Wigner resonance parameter. The value is 34 MeV greater than the real part of the position of the pole (in the energy-squared plane) in the Z -boson propagator. Also the LEP experiments have generally assumed a fixed value of the $\gamma - Z$ interferences term based on the standard model. Keeping this term as free parameter leads to a somewhat larger error on the fitted Z mass. See ACCIARRI 97K and ACKERSTAFF 97C for a detailed investigation of both these issues.

A new source of LEP energy variation was discovered in mid 1995: an energy change of a few MeV is correlated with the passage of a train on nearby railway tracks. The LEP energy working group is studying the implications of this effect for the high statistics data recorded since 1993. The main consequence of this is expected to be a shift in the overall LEP energy values leading to a corresponding shift in the value of m_Z . The LEP collaborations have consequently deferred publication of their results on Z lineshape and lepton forward-backward asymmetries based on 1993 and later data.

Because of the high current interest, we mention here the following preliminary results, but do not average them or include them in the Listings or Tables.

Combining published and unpublished preliminary LEP results (as of end of February 1998) yields an average Z -boson mass of 91.1867 ± 0.0020 GeV, with a total width of 2.4948 ± 0.0025 GeV.

<u>VALUE (GeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
91.187±0.007 OUR FIT				
91.188±0.007 OUR AVERAGE				
$91.187 \pm 0.007 \pm 0.006$	1.16M	¹ ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV
$91.195 \pm 0.006 \pm 0.007$	1.19M	¹ ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
$91.182 \pm 0.007 \pm 0.006$	1.33M	¹ AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV
$91.187 \pm 0.007 \pm 0.006$	1.27M	¹ BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
91.193 ± 0.010	1.2M	² ACCIARRI	97K L3	$E_{cm}^{ee} = \text{LEP1} + 130-136$ GeV + 161-172 GeV
91.185 ± 0.010		³ ACKERSTAFF	97C OPAL	$E_{cm}^{ee} = \text{LEP1} + 130-136$ GeV + 161 GeV
91.162 ± 0.011	1.2M	⁴ ACCIARRI	96B L3	Repl. by ACCIARRI 97K
91.192 ± 0.011	1.33M	⁵ ALEXANDER	96X OPAL	Repl. by ACKER- STAFF 97C
91.151 ± 0.008		⁶ MIYABAYASHI	95 TOPZ	$E_{cm}^{ee} = 57.8$ GeV
$91.181 \pm 0.007 \pm 0.006$	512k	⁷ ACTON	93D OPAL	Repl. by AKERS 94
91.195 ± 0.009	460k	⁸ ADRIANI	93F L3	Repl. by ACCIARRI 94
91.187 ± 0.009	520k	⁹ BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
$91.74 \pm 0.28 \pm 0.93$	156	¹⁰ ALITTI	92B UA2	$E_{cm}^{pp} = 630$ GeV
$89.2 \begin{smallmatrix} +2.1 \\ -1.8 \end{smallmatrix}$		¹¹ ADACHI	90F RVUE	
$90.9 \pm 0.3 \pm 0.2$	188	¹² ABE	89C CDF	$E_{cm}^{pp} = 1.8$ TeV
91.14 ± 0.12	480	¹³ ABRAMS	89B MRK2	$E_{cm}^{ee} = 89-93$ GeV
$93.1 \pm 1.0 \pm 3.0$	24	^{14,15} ALBAJAR	89 UA1	$E_{cm}^{pp} = 546,630$ GeV

¹ The second error of 6.3 MeV is due to a common LEP energy uncertainty.

² ACCIARRI 97K interpret the s -dependence of the cross sections and lepton forward-backward asymmetries in the framework of the S-matrix formalism with a combined fit to their cross section and asymmetry data at the Z peak (ACCIARRI 94) and their data at 130, 136, 161, and 172 GeV. The authors have corrected the measurement for the 34.1 MeV shift with respect to the Breit-Wigner fits. The error contains a contribution of ± 3 MeV due to the uncertainty on the γZ interference.

³ ACKERSTAFF 97C obtain this using the S-matrix formalism for a combined fit to their cross-section and asymmetry data at the Z peak (AKERS 94) and their data at 130, 136, and 161 GeV. The authors have corrected the measurement for the 34 MeV shift with respect to the Breit-Wigner fits.

⁴ ACCIARRI 96B interpret the s -dependence of the cross sections and lepton forward-backward asymmetries in the framework of the S-matrix ansatz. The 130-136 GeV data constrains the γZ interference terms. As expected, this result is below the mass values obtained with a standard Breit-Wigner parametrization.

⁵ ALEXANDER 96X obtain this using the S-matrix formalism for a combined fit to their cross-section and asymmetry data at the Z peak (AKERS 94) and their data at 130 and 136 GeV. The authors have corrected the measurement for the 34 MeV shift with respect to the Breit-Wigner fits.

⁶ MIYABAYASHI 95 combine their low energy total hadronic cross-section measurement with the ACTON 93D data and perform a fit using an S-matrix formalism. As expected, this result is below the mass values obtained with the standard Breit-Wigner parametrization.

- ⁷ The systematic error in ACTON 93D is from the uncertainty in the LEP energy calibration.
- ⁸ The error in ADRIANI 93F includes 6 MeV due to the uncertainty in LEP energy calibration.
- ⁹ BUSKULIC 93J supersedes DECAMP 92B. The error includes 6 MeV due to the uncertainty in LEP energy calibration.
- ¹⁰ Enters fit through W/Z mass ratio given in the W Particle Listings. The ALITTI 92B systematic error (± 0.93) has two contributions: one (± 0.92) cancels in m_W/m_Z and one (± 0.12) is noncancelling. These were added in quadrature.
- ¹¹ ADACHI 90F use a Breit-Wigner resonance shape fit and combine their results with published data of PEP and PETRA.
- ¹² First error of ABE 89 is combination of statistical and systematic contributions; second is mass scale uncertainty.
- ¹³ ABRAMS 89B uncertainty includes 35 MeV due to the absolute energy measurement.
- ¹⁴ Enters fit through Z - W mass difference given in the W Particle Listings.
- ¹⁵ ALBAJAR 89 result is from a total sample of 33 $Z \rightarrow e^+e^-$ events.

Z WIDTH

<u>VALUE (GeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.490±0.007 OUR FIT				
2.491±0.007 OUR AVERAGE				
2.50 ±0.21 ±0.06		¹⁶ ABREU	96R DLPH	$E_{cm}^{ee} = 91.2$ GeV
2.483±0.011±0.0045	1.16M	¹⁷ ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV
2.494±0.009±0.0045	1.19M	¹⁷ ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
2.483±0.011±0.0045	1.33M	¹⁷ AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV
2.501±0.011±0.0045	1.27M	¹⁷ BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2.494±0.010	1.2M	¹⁸ ACCIARRI	97K L3	$E_{cm}^{ee} = \text{LEP1} + 130-136$ GeV + 161-172 GeV
2.492±0.010	1.2M	¹⁹ ACCIARRI	96B L3	Repl. by ACCIARRI 97K
2.483±0.011±0.004	512k	²⁰ ACTON	93D OPAL	Repl. by AKERS 94
2.490±0.011	460k	²¹ ADRIANI	93F L3	Repl. by ACCIARRI 94
2.501±0.012	520k	²² BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
3.8 ±0.8 ±1.0	188	ABE	89C CDF	$E_{cm}^{p\bar{p}} = 1.8$ TeV
2.42 $\begin{smallmatrix} +0.45 \\ -0.35 \end{smallmatrix}$	480	²³ ABRAMS	89B MRK2	$E_{cm}^{ee} = 89-93$ GeV
2.7 $\begin{smallmatrix} +1.2 \\ -1.0 \end{smallmatrix}$ ±1.3	24	²⁴ ALBAJAR	89 UA1	$E_{cm}^{p\bar{p}} = 546,630$ GeV
2.7 ±2.0 ±1.0	25	²⁵ ANSARI	87 UA2	$E_{cm}^{p\bar{p}} = 546,630$ GeV

- ¹⁶ ABREU 96R obtain this value from a study of the interference between initial and final state radiation in the process $e^+e^- \rightarrow Z \rightarrow \mu^+\mu^-$.
- ¹⁷ The second error of 4.5 MeV is due to a common LEP energy uncertainty.
- ¹⁸ ACCIARRI 97K interpret the s -dependence of the cross sections and lepton forward-backward asymmetries in the framework of the S-matrix formalism with a combined fit to their cross section and asymmetry data at the Z peak (ACCIARRI 94) and their data at 130, 136, 161, and 172 GeV. The authors have corrected the measurement for the 0.9 MeV shift with respect to the Breit-Wigner fits.
- ¹⁹ ACCIARRI 96B interpret the s -dependence of the cross sections and lepton forward-backward asymmetries in the framework of the S-matrix ansatz. The 130-136 GeV data constrains the γZ interference terms. The fitted width is expected to be 0.9 MeV less than that obtained using the standard Breit-Wigner parametrization (see 'Note on the Z Boson').

²⁰The systematic error is from the uncertainty in the LEP energy calibration.

²¹The error in ADRIANI 93F includes 4 MeV due to the uncertainty in LEP energy calibration.

²²The error in BUSKULIC 93J includes 4 MeV due to the uncertainty in LEP energy calibration.

²³ABRAMS 89B uncertainty includes 50 MeV due to the miniSAM background subtraction error.

²⁴ALBAJAR 89 result is from a total sample of 33 $Z \rightarrow e^+e^-$ events.

²⁵Quoted values of ANSARI 87 are from direct fit. Ratio of Z and W production gives either $\Gamma(Z) < (1.09 \pm 0.07) \times \Gamma(W)$, CL = 90% or $\Gamma(Z) = (0.82^{+0.19}_{-0.14} \pm 0.06) \times \Gamma(W)$. Assuming Standard-Model value $\Gamma(W) = 2.65$ GeV then gives $\Gamma(Z) < 2.89 \pm 0.19$ or $= 2.17^{+0.50}_{-0.37} \pm 0.16$.

Z DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 e^+e^-	(3.366 \pm 0.008) %	
Γ_2 $\mu^+\mu^-$	(3.367 \pm 0.013) %	
Γ_3 $\tau^+\tau^-$	(3.360 \pm 0.015) %	
Γ_4 $\ell^+\ell^-$	[a] (3.366 \pm 0.006) %	
Γ_5 invisible	(20.01 \pm 0.16) %	
Γ_6 hadrons	(69.90 \pm 0.15) %	
Γ_7 $(u\bar{u} + c\bar{c})/2$	(10.1 \pm 1.1) %	
Γ_8 $(d\bar{d} + s\bar{s} + b\bar{b})/3$	(16.6 \pm 0.6) %	
Γ_9 $c\bar{c}$	(12.4 \pm 0.6) %	
Γ_{10} $b\bar{b}$	(15.16 \pm 0.09) %	
Γ_{11} ggg	< 1.1	% 95%
Γ_{12} $\pi^0\gamma$	< 5.2	$\times 10^{-5}$ 95%
Γ_{13} $\eta\gamma$	< 5.1	$\times 10^{-5}$ 95%
Γ_{14} $\omega\gamma$	< 6.5	$\times 10^{-4}$ 95%
Γ_{15} $\eta'(958)\gamma$	< 4.2	$\times 10^{-5}$ 95%
Γ_{16} $\gamma\gamma$	< 5.2	$\times 10^{-5}$ 95%
Γ_{17} $\gamma\gamma\gamma$	< 1.0	$\times 10^{-5}$ 95%
Γ_{18} $\pi^\pm W^\mp$	[b] < 7	$\times 10^{-5}$ 95%
Γ_{19} $\rho^\pm W^\mp$	[b] < 8.3	$\times 10^{-5}$ 95%
Γ_{20} $J/\psi(1S)X$	(3.66 \pm 0.23) $\times 10^{-3}$	
Γ_{21} $\psi(2S)X$	(1.60 \pm 0.29) $\times 10^{-3}$	
Γ_{22} $\chi_{c1}(1P)X$	(2.9 \pm 0.7) $\times 10^{-3}$	
Γ_{23} $\chi_{c2}(1P)X$	< 3.2	$\times 10^{-3}$ 90%
Γ_{24} $\Upsilon(1S)X + \Upsilon(2S)X$ $+ \Upsilon(3S)X$	(1.0 \pm 0.5) $\times 10^{-4}$	
Γ_{25} $\Upsilon(1S)X$	< 5.5	$\times 10^{-5}$ 95%
Γ_{26} $\Upsilon(2S)X$	< 1.39	$\times 10^{-4}$ 95%

Γ_{27}	$\gamma(3S)X$		< 9.4	$\times 10^{-5}$	95%	
Γ_{28}	$(D^0/\bar{D}^0)X$		(20.7 ± 2.0)	%		
Γ_{29}	$D^\pm X$		(12.2 ± 1.7)	%		
Γ_{30}	$D^*(2010)^\pm X$	[b]	(11.4 ± 1.3)	%		
Γ_{31}	BX					
Γ_{32}	B^*X					
Γ_{33}	$B_s^0 X$		seen			
Γ_{34}	anomalous $\gamma +$ hadrons	[c]	< 3.2	$\times 10^{-3}$	95%	
Γ_{35}	$e^+ e^- \gamma$	[c]	< 5.2	$\times 10^{-4}$	95%	
Γ_{36}	$\mu^+ \mu^- \gamma$	[c]	< 5.6	$\times 10^{-4}$	95%	
Γ_{37}	$\tau^+ \tau^- \gamma$	[c]	< 7.3	$\times 10^{-4}$	95%	
Γ_{38}	$\ell^+ \ell^- \gamma \gamma$	[d]	< 6.8	$\times 10^{-6}$	95%	
Γ_{39}	$q\bar{q}\gamma\gamma$	[d]	< 5.5	$\times 10^{-6}$	95%	
Γ_{40}	$\nu\bar{\nu}\gamma\gamma$	[d]	< 3.1	$\times 10^{-6}$	95%	
Γ_{41}	$e^\pm \mu^\mp$	LF	[b]	< 1.7	$\times 10^{-6}$	95%
Γ_{42}	$e^\pm \tau^\mp$	LF	[b]	< 9.8	$\times 10^{-6}$	95%
Γ_{43}	$\mu^\pm \tau^\mp$	LF	[b]	< 1.2	$\times 10^{-5}$	95%

[a] ℓ indicates each type of lepton (e , μ , and τ), not sum over them.

[b] The value is for the sum of the charge states of particle/antiparticle states indicated.

[c] See the Particle Listings below for the γ energy range used in this measurement.

[d] For $m_{\gamma\gamma} = (60 \pm 5)$ GeV.

Z PARTIAL WIDTHS

$\Gamma(e^+ e^-)$ Γ_1

For the LEP experiments, this parameter is not directly used in the overall fit but is derived using the fit results; see the 'Note on the Z Boson.'

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
83.82 ± 0.30 OUR FIT				
$82.89 \pm 1.20 \pm 0.89$		²⁶ ABE	95J SLD	$E_{cm}^{ee} = 91.31$ GeV
• • •				We do not use the following data for averages, fits, limits, etc. • • •
83.31 ± 0.54	31.4k	ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV
83.43 ± 0.52	38k	ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
83.63 ± 0.53	42k	AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV
84.61 ± 0.49	45.8k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV

²⁶ ABE 95J obtain this measurement from Bhabha events in a restricted fiducial region to improve systematics. They use the values 91.187 and 2.489 GeV for the Z mass and total decay width to extract this partial width.

$\Gamma(\mu^+ \mu^-)$
 Γ_2

This parameter is not directly used in the overall fit but is derived using the fit results; see the 'Note on the Z Boson.'

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

83.83 ± 0.39 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

84.15 ± 0.77	45.6k	ABREU	94	DLPH	$E_{cm}^{ee} = 88-94$ GeV
83.20 ± 0.79	34k	ACCIARRI	94	L3	$E_{cm}^{ee} = 88-94$ GeV
83.83 ± 0.65	57k	AKERS	94	OPAL	$E_{cm}^{ee} = 88-94$ GeV
83.62 ± 0.75	46.4k	BUSKULIC	94	ALEP	$E_{cm}^{ee} = 88-94$ GeV

 $\Gamma(\tau^+ \tau^-)$
 Γ_3

This parameter is not directly used in the overall fit but is derived using the fit results; see the 'Note on the Z Boson.'

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

83.67 ± 0.44 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

83.55 ± 0.91	25k	ABREU	94	DLPH	$E_{cm}^{ee} = 88-94$ GeV
84.04 ± 0.94	25k	ACCIARRI	94	L3	$E_{cm}^{ee} = 88-94$ GeV
82.90 ± 0.77	47k	AKERS	94	OPAL	$E_{cm}^{ee} = 88-94$ GeV
84.18 ± 0.79	45.1k	BUSKULIC	94	ALEP	$E_{cm}^{ee} = 88-94$ GeV

 $\Gamma(\ell^+ \ell^-)$
 Γ_4

In our fit $\Gamma(\ell^+ \ell^-)$ is defined as the partial Z width for the decay into a pair of massless charged leptons. This parameter is not directly used in the 5-parameter fit assuming lepton universality but is derived using the fit results. See the 'Note on the Z Boson.'

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

83.83 ± 0.27 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

83.56 ± 0.45	102k	ABREU	94	DLPH	$E_{cm}^{ee} = 88-94$ GeV
83.49 ± 0.46	97k	ACCIARRI	94	L3	$E_{cm}^{ee} = 88-94$ GeV
83.55 ± 0.44	146k	AKERS	94	OPAL	$E_{cm}^{ee} = 88-94$ GeV
84.40 ± 0.43	137.3k	BUSKULIC	94	ALEP	$E_{cm}^{ee} = 88-94$ GeV

 $\Gamma(\text{invisible})$
 Γ_5

We use only direct measurements of the invisible partial width to obtain the average value quoted below. The fit value is obtained as a difference between the total and the observed partial widths assuming lepton universality.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

498.3 ± 4.2 OUR FIT
517 ± 22 OUR AVERAGE

539 ± 26 ± 17	410	AKERS	95C	OPAL	$E_{cm}^{ee} = 88-94$ GeV
450 ± 34 ± 34	258	BUSKULIC	93L	ALEP	$E_{cm}^{ee} = 88-94$ GeV
540 ± 80 ± 40	52	ADEVA	92	L3	$E_{cm}^{ee} = 88-94$ GeV
524 ± 40 ± 20	172	²⁷ ADRIANI	92E	L3	$E_{cm}^{ee} = 88-94$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

509.4 ± 7.0	ABREU	94	DLPH	$E_{cm}^{ee} = 88-94$ GeV
496.5 ± 7.9	ACCIARRI	94	L3	$E_{cm}^{ee} = 88-94$ GeV
490.3 ± 7.3	AKERS	94	OPAL	$E_{cm}^{ee} = 88-94$ GeV
501 ± 6	BUSKULIC	94	ALEP	$E_{cm}^{ee} = 88-94$ GeV

²⁷ ADRIANI 92E improves but does not supersede ADEVA 92, obtained with 1990 data only.

$\Gamma(\text{hadrons})$

Γ_6

This parameter is not directly used in the 5-parameter fit assuming lepton universality, but is derived using the fit results. See the 'Note on the Z Boson.'

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1740.7 ± 5.9 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

1723 ± 10	1.05M	ABREU	94	DLPH	$E_{cm}^{ee} = 88-94$ GeV
1748 ± 10	1.09M	ACCIARRI	94	L3	$E_{cm}^{ee} = 88-94$ GeV
1741 ± 10	1.19M	²⁸ AKERS	94	OPAL	$E_{cm}^{ee} = 88-94$ GeV
1746 ± 10	1.27M	BUSKULIC	94	ALEP	$E_{cm}^{ee} = 88-94$ GeV

²⁸ AKERS 94 assumes lepton universality. Without this assumption, it becomes 1742 ± 11 MeV.

Z BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma(e^+e^-)$

Γ_6/Γ_1

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
20.77 ± 0.08 OUR FIT				
20.74 ± 0.18	31.4k	ABREU	94	DLPH $E_{cm}^{ee} = 88-94$ GeV
20.96 ± 0.15	38k	ACCIARRI	94	L3 $E_{cm}^{ee} = 88-94$ GeV
20.83 ± 0.16	42k	AKERS	94	OPAL $E_{cm}^{ee} = 88-94$ GeV
20.59 ± 0.15	45.8k	BUSKULIC	94	ALEP $E_{cm}^{ee} = 88-94$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

20.99 ± 0.25	17k	ACTON	93D	OPAL	Repl. by AKERS 94
20.69 ± 0.21		BUSKULIC	93J	ALEP	Repl. by BUSKULIC 94
$27.0^{+11.7}_{-8.8}$	12	²⁹ ABRAMS	89D	MRK2	$E_{cm}^{ee} = 89-93$ GeV

²⁹ ABRAMS 89D have included both statistical and systematic uncertainties in their quoted errors.

$\Gamma(\text{hadrons})/\Gamma(\mu^+\mu^-)$


Γ_6/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
20.76 ± 0.07 OUR FIT				
20.78 ± 0.09 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
20.54 ± 0.14	45.6k	ABREU	94	DLPH $E_{cm}^{ee} = 88-94$ GeV
21.02 ± 0.16	34k	ACCIARRI	94	L3 $E_{cm}^{ee} = 88-94$ GeV
20.78 ± 0.11	57k	AKERS	94	OPAL $E_{cm}^{ee} = 88-94$ GeV
20.83 ± 0.15	46.4k	BUSKULIC	94	ALEP $E_{cm}^{ee} = 88-94$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

20.65±0.17	23k	ACTON	93D OPAL	Repl. by AKERS 94
20.88±0.20		BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
18.9 $\begin{smallmatrix} +7.1 \\ -5.3 \end{smallmatrix}$	13	³⁰ ABRAMS	89D MRK2	$E_{cm}^{ee} = 89-93$ GeV

³⁰ ABRAMS 89D have included both statistical and systematic uncertainties in their quoted errors.



$$\Gamma(\text{hadrons})/\Gamma(\mu^+\mu^-)$$

$\Gamma(\text{hadrons})/\Gamma(\tau^+\tau^-)$ Γ_6/Γ_3

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
20.80±0.08 OUR FIT				
20.81±0.08 OUR AVERAGE				
20.68±0.18	25k	ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV
20.80±0.20	25k	ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
21.01±0.15	47k	AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV
20.70±0.16	45.1k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

21.22±0.25	18k	ACTON	93D OPAL	Repl. by AKERS 94
20.77±0.23		BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
15.2 $\begin{smallmatrix} +4.8 \\ -3.9 \end{smallmatrix}$	21	³¹ ABRAMS	89D MRK2	$E_{cm}^{ee} = 89-93$ GeV

³¹ ABRAMS 89D have included both statistical and systematic uncertainties in their quoted errors.

$\Gamma(\text{hadrons})/\Gamma(\ell^+\ell^-)$

Γ_6/Γ_4

ℓ indicates each type of lepton (e , μ , and τ), not sum over them.

Our fit result is obtained requiring lepton universality.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------


20.76 ± 0.05 OUR FIT

20.77 ± 0.07 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

20.62 ± 0.10	102k	ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV
20.93 ± 0.10	97k	ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
20.835 ± 0.086	146k	AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV
20.69 ± 0.09	137.3k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

20.88 ± 0.13	58k	ACTON	93D OPAL	Repl. by AKERS 94
21.00 ± 0.15	40k	ADRIANI	93M L3	Repl. by ACCIARRI 94
20.78 ± 0.13		BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
18.9 $^{+3.6}_{-3.2}$	46	ABRAMS	89B MRK2	$E_{cm}^{ee} = 89-93$ GeV



$\Gamma(\text{hadrons})/\Gamma(\ell^+\ell^-)$

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$

Γ_6/Γ

This parameter is not directly used in the overall fit but is derived using the fit results; see the 'Note on the Z Boson.'

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.6990 ± 0.0015 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6983 ± 0.0023	1.14M	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
-----------------	-------	----------	---------	---------------------------

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_1/Γ

This parameter is not directly used in the overall fit but is derived using the fit results; see the 'Note on the Z Boson.'

VALUE EVTS DOCUMENT ID TECN COMMENT

0.03366 ± 0.00008 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.03383 ± 0.00013 45.8k BUSKULIC 94 ALEP $E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_2/Γ

This parameter is not directly used in the overall fit but is derived using the fit results; see the 'Note on the Z Boson.'

VALUE EVTS DOCUMENT ID TECN COMMENT

0.03367 ± 0.00013 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.03344 ± 0.00026 46.4k BUSKULIC 94 ALEP $E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_3/Γ

This parameter is not directly used in the overall fit but is derived using the fit results; see the 'Note on the Z Boson.'

VALUE EVTS DOCUMENT ID TECN COMMENT

0.03360 ± 0.00015 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.03366 ± 0.00028 45.1k BUSKULIC 94 ALEP $E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

$\Gamma(\ell^+\ell^-)/\Gamma_{\text{total}}$ Γ_4/Γ

ℓ indicates each type of lepton (e , μ , and τ), not sum over them.

Our fit result assumes lepton universality.

This parameter is not directly used in the overall fit but is derived using the fit results; see the 'Note on the Z Boson.'

VALUE EVTS DOCUMENT ID TECN COMMENT

0.03366 ± 0.00006 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.03375 ± 0.00009 137.3k BUSKULIC 94 ALEP $E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

$\Gamma(\text{invisible})/\Gamma_{\text{total}}$ Γ_5/Γ

See the data, the note, and the fit result for the partial width, Γ_5 , above.

VALUE DOCUMENT ID

0.2001 ± 0.0016 OUR FIT

$\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$ Γ_2/Γ_1

This parameter is not directly used in the overall fit but is derived using the fit results; see the 'Note on the Z Boson.'

VALUE DOCUMENT ID

1.000 ± 0.005 OUR FIT

$\Gamma(\tau^+\tau^-)/\Gamma(e^+e^-)$ Γ_3/Γ_1

This parameter is not directly used in the overall fit but is derived using the fit results; see the 'Note on the Z Boson.'

VALUE DOCUMENT ID

0.998 ± 0.005 OUR FIT

$\Gamma((u\bar{u} + c\bar{c})/2)/\Gamma(\text{hadrons})$ Γ_7/Γ_6

This quantity is the branching ratio of $Z \rightarrow$ "up-type" quarks to $Z \rightarrow$ hadrons. Except ACKERSTAFF 97T the values of $Z \rightarrow$ "up-type" and $Z \rightarrow$ "down-type" branchings are extracted from measurements of $\Gamma(\text{hadrons})$, and $\Gamma(Z \rightarrow \gamma + \text{jets})$ where γ is a high-energy (>5 GeV) isolated photon. As the experiments use different procedures and slightly different values of M_Z , $\Gamma(\text{hadrons})$ and α_s in their extraction procedures, our average has to be taken with caution.

VALUE	DOCUMENT ID	TECN	COMMENT
0.145 ± 0.015 OUR AVERAGE			
0.160 ± 0.019 ± 0.019	32 ACKERSTAFF 97T	OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.137 ^{+0.038} _{-0.054}	33 ABREU	95X DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.139 ± 0.026	34 ACTON	93F OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.137 ± 0.033	35 ADRIANI	93 L3	$E_{\text{cm}}^{ee} = 91.2$ GeV

³² ACKERSTAFF 97T measure $\Gamma_{u\bar{u}}/(\Gamma_{d\bar{d}} + \Gamma_{u\bar{u}} + \Gamma_{s\bar{s}}) = 0.258 \pm 0.031 \pm 0.032$. To obtain this branching ratio authors use $R_c + R_b = 0.380 \pm 0.010$. This measurement is fully negatively correlated with the measurement of $\Gamma_{d\bar{d},s\bar{s}}/(\Gamma_{d\bar{d}} + \Gamma_{u\bar{u}} + \Gamma_{s\bar{s}})$ given in the next data block.

³³ ABREU 95X use $M_Z = 91.187 \pm 0.009$ GeV, $\Gamma(\text{hadrons}) = 1725 \pm 12$ MeV and $\alpha_s = 0.123 \pm 0.005$. To obtain this branching ratio we divide their value of $C_{2/3} = 0.91^{+0.25}_{-0.36}$ by their value of $(3C_{1/3} + 2C_{2/3}) = 6.66 \pm 0.05$.

³⁴ ACTON 93F use the LEP 92 value of $\Gamma(\text{hadrons}) = 1740 \pm 12$ MeV and $\alpha_s = 0.122^{+0.006}_{-0.005}$.

³⁵ ADRIANI 93 use $M_Z = 91.181 \pm 0.022$ GeV, $\Gamma(\text{hadrons}) = 1742 \pm 19$ MeV and $\alpha_s = 0.125 \pm 0.009$. To obtain this branching ratio we divide their value of $C_{2/3} = 0.92 \pm 0.22$ by their value of $(3C_{1/3} + 2C_{2/3}) = 6.720 \pm 0.076$.

 $\Gamma((d\bar{d} + s\bar{s} + b\bar{b})/3)/\Gamma(\text{hadrons})$ Γ_8/Γ_6

This quantity is the branching ratio of $Z \rightarrow$ "down-type" quarks to $Z \rightarrow$ hadrons. Except ACKERSTAFF 97T the values of $Z \rightarrow$ "up-type" and $Z \rightarrow$ "down-type" branchings are extracted from measurements of $\Gamma(\text{hadrons})$, and $\Gamma(Z \rightarrow \gamma + \text{jets})$ where γ is a high-energy (>5 GeV) isolated photon. As the experiments use different procedures and slightly different values of M_Z , $\Gamma(\text{hadrons})$ and α_s in their extraction procedures, our average has to be taken with caution.

VALUE	DOCUMENT ID	TECN	COMMENT
0.237 ± 0.009 OUR AVERAGE			
0.230 ± 0.010 ± 0.010	36 ACKERSTAFF 97T	OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.243 ^{+0.036} _{-0.026}	37 ABREU	95X DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.241 ± 0.017	38 ACTON	93F OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.243 ± 0.022	39 ADRIANI	93 L3	$E_{\text{cm}}^{ee} = 91.2$ GeV

³⁶ ACKERSTAFF 97T measure $\Gamma_{d\bar{d},s\bar{s}}/(\Gamma_{d\bar{d}} + \Gamma_{u\bar{u}} + \Gamma_{s\bar{s}}) = 0.371 \pm 0.016 \pm 0.016$. To obtain this branching ratio authors use $R_c + R_b = 0.380 \pm 0.010$. This measurement is fully negatively correlated with the measurement of $\Gamma_{u\bar{u}}/(\Gamma_{d\bar{d}} + \Gamma_{u\bar{u}} + \Gamma_{s\bar{s}})$ presented in the previous data block.

³⁷ ABREU 95X use $M_Z = 91.187 \pm 0.009$ GeV, $\Gamma(\text{hadrons}) = 1725 \pm 12$ MeV and $\alpha_s = 0.123 \pm 0.005$. To obtain this branching ratio we divide their value of $C_{1/3} = 1.62^{+0.24}_{-0.17}$ by their value of $(3C_{1/3} + 2C_{2/3}) = 6.66 \pm 0.05$.

³⁸ ACTON 93F use the LEP 92 value of $\Gamma(\text{hadrons}) = 1740 \pm 12 \text{ MeV}$ and $\alpha_s = 0.122^{+0.006}_{-0.005}$.

³⁹ ADRIANI 93 use $M_Z = 91.181 \pm 0.022 \text{ GeV}$, $\Gamma(\text{hadrons}) = 1742 \pm 19 \text{ MeV}$ and $\alpha_s = 0.125 \pm 0.009$. To obtain this branching ratio we divide their value of $C_{1/3} = 1.63 \pm 0.15$ by their value of $(3C_{1/3} + 2C_{2/3}) = 6.720 \pm 0.076$.

$R_c = \Gamma(c\bar{c})/\Gamma(\text{hadrons})$

Γ_9/Γ_6

OUR FIT is obtained by a simultaneous fit to several c - and b -quark measurements as explained in the "Note on the Z boson." As a cross check we have also performed a weighted average of the R_c measurements taking into account the various common systematic errors. Assuming that the smallest common systematic error is fully correlated, we obtain $R_c = 0.171 \pm 0.009$.

Because of the high current interest, we mention the following preliminary results here, but do not average them or include them in the Listings or Tables. Combining published and unpublished preliminary LEP and SLD electroweak results (as of end of February 1998) yields $R_c = 0.1734 \pm 0.0048$. The Standard Model predicts $R_c = 0.1723$ for $m_t = 175 \text{ GeV}$ and $M_H = 300 \text{ GeV}$.

VALUE	DOCUMENT ID	TECN	COMMENT
0.177 ± 0.008 OUR FIT			
0.180 ± 0.011 ± 0.013	⁴⁰ ACKERSTAFF	98E OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
0.167 ± 0.011 ± 0.012	⁴¹ ALEXANDER	96R OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
0.1623 ± 0.0085 ± 0.0209	⁴² ABREU	95D DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
0.165 ± 0.005 ± 0.020	⁴³ BUSKULIC	94G ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
0.187 ± 0.031 ± 0.023	⁴⁴ ABREU	93I DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.142 ± 0.008 ± 0.014	⁴⁵ AKERS	95O OPAL	Repl. by ACKERSTAFF 98E
0.151 ± 0.008 ± 0.041	⁴⁶ ABREU	92O DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$

⁴⁰ ACKERSTAFF 98E use an inclusive/exclusive double tag. In one jet $D^{*\pm}$ mesons are exclusively reconstructed in several decay channels and in the opposite jet a slow pion (opposite charge inclusive $D^{*\pm}$) tag is used. The b content of this sample is measured by the simultaneous detection of a lepton in one jet and an inclusively reconstructed $D^{*\pm}$ meson in the opposite jet. The systematic error includes an uncertainty of ± 0.006 due to the external branching ratios.

⁴¹ ALEXANDER 96R obtain this value via direct charm counting, summing the partial contributions from D^0 , D^+ , D_s^+ , and Λ_c^+ , and assuming that strange-charmed baryons account for the 15% of the Λ_c^+ production. An uncertainty of ± 0.005 due to the uncertainties in the charm hadron branching ratios is included in the overall systematics.

⁴² ABREU 95D perform a maximum likelihood fit to the combined p and p_T distributions of single and dilepton samples. The second error includes an uncertainty of ± 0.0124 due to models and branching ratios.

⁴³ BUSKULIC 94G perform a simultaneous fit to the p and p_T spectra of both single and dilepton events.

⁴⁴ ABREU 93I assume that the D_s and charmed baryons are equally produced at LEP and CLEO (10 GeV) energies.

⁴⁵ AKERS 95O use the presence of a $D^{*\pm}$ to tag $Z \rightarrow c\bar{c}$ with $D^* \rightarrow D^0\pi$ and $D^0 \rightarrow K\pi$. They measure $P_c * \Gamma(c\bar{c})/\Gamma(\text{hadrons})$ to be $(1.006 \pm 0.055 \pm 0.061) \times 10^{-3}$, where P_c is the product branching ratio $B(c \rightarrow D^*)B(D^* \rightarrow D^0\pi)B(D^0 \rightarrow K\pi)$. Assuming that P_c remains unchanged with energy, they use its value $(7.1 \pm 0.5) \times 10^{-3}$ determined

at CESR/PETRA to obtain $\Gamma(c\bar{c})/\Gamma(\text{hadrons})$. The second error of AKERS 950 includes an uncertainty of ± 0.011 from the uncertainty on P_c .

- ⁴⁶ ABREU 920 use the neural network technique to tag heavy flavour events among a sample of 123k selected hadronic events. The systematic error consists of three parts: due to Monte Carlo (MC) parametrization (0.023), choice of MC model (0.033) and detector effects (0.009) added in quadrature.

$R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons})$

Γ_{10}/Γ_6

OUR FIT is obtained by a simultaneous fit to several c - and b -quark measurements as explained in the "Note on the Z boson." As a cross check we have also performed a weighted average of the R_b measurements taking into account the various common systematic errors. We have assumed that the smallest common systematic error is fully correlated. For $R_c = 0.177$ (as given by OUR FIT above), we obtain $R_b = 0.2169 \pm 0.0012$. For an expected Standard Model value of $R_c = 0.1723$, our weighted average gives $R_b = 0.2172 \pm 0.0012$.

Because of the high current interest, we mention the following preliminary results here, but do not average them or include them in the Listings or Tables. Combining published and unpublished preliminary LEP and SLD electroweak results (as of end of February 1998) yields $R_b = 0.2170 \pm 0.0009$. The Standard Model predicts $R_b = 0.2158$ for $m_t = 175$ GeV and $M_H = 300$ GeV.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.2169±0.0012 OUR FIT				
0.2142±0.0034±0.0015		47 ABE	98D SLD	$E_{\text{cm}}^{ee} = 91.2$ GeV
0.2175±0.0014±0.0017		48 ACKERSTAFF	97K OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.2159±0.0009±0.0011		49 BARATE	97F ALEP	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.2216±0.0016±0.0021		50 ABREU	96 DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.2145±0.0089±0.0067		51 ABREU	95D DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.219 ±0.006 ±0.005		52 BUSKULIC	94G ALEP	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.222 ±0.003 ±0.007		53 ADRIANI	93E L3	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.222 ±0.011 ±0.007		54 AKERS	93B OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.251 ±0.049 ±0.030	32	55 JACOBSEN	91 MRK2	$E_{\text{cm}}^{ee} = 91$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.2167±0.0011±0.0013		56 BARATE	97E ALEP	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.229 ±0.011		57 ABE	96E SLD	Repl. by ABE 98D
0.2217±0.0020±0.0033		58 ABREU	95D DLPH	Repl. by ABREU 96
0.2241±0.0063±0.0046		59 ABREU	95J DLPH	Repl. by ABREU 96
0.2171±0.0021±0.0021		60 AKERS	95B OPAL	Repl. by ACKER-STAFF 97K
0.228 ±0.005 ±0.005		61 BUSKULIC	93N ALEP	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.222 $\begin{smallmatrix} +0.033 \\ -0.031 \end{smallmatrix}$ ±0.017		62 ABREU	92 DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.219 ±0.014 ±0.019		63 ABREU	92K DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.232 ±0.005 ±0.017		64 ABREU	92O DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
0.23 $\begin{smallmatrix} +0.10 \\ -0.08 \end{smallmatrix}$ $\begin{smallmatrix} +0.05 \\ -0.04 \end{smallmatrix}$	15	65 KRAL	90 MRK2	$E_{\text{cm}}^{ee} = 89-93$ GeV

- ⁴⁷ ABE 98D use a double tag based on 3D impact parameter with reconstruction of secondary vertices. The charm background is reduced by requiring the invariant mass at the secondary vertex to be above 2 GeV. The systematic error includes an uncertainty of ± 0.0002 due to the uncertainty on R_c .

- 48 ACKERSTAFF 97K use lepton and/or separated decay vertex to tag independently each hemisphere. Comparing the numbers of single- and double-tagged events, they determine the b -tagging efficiency directly from the data.
- 49 BARATE 97F combine the lifetime-mass hemisphere tag (BARATE 97E) with event shape information and lepton tag to identify $Z \rightarrow b\bar{b}$ candidates. They further use c - and uds -selection tags to identify the background. For R_c different from its Standard Model value of 0.172, R_b varies as $-0.019 \times (R_c - 0.172)$.
- 50 ABREU 96 obtain this result combining several analyses (double lifetime tag, mixed tag and multivariate analysis). This value is obtained assuming $R_c = \Gamma(c\bar{c})/\Gamma(\text{hadrons}) = 0.172$. For a value of R_c different from this by an amount ΔR_c the change in the value is given by $-0.087 \cdot \Delta R_c$.
- 51 ABREU 95D perform a maximum likelihood fit to the combined p and p_T distributions of single and dilepton samples. The second error includes an uncertainty of ± 0.0023 due to models and branching ratios.
- 52 BUSKULIC 94G perform a simultaneous fit to the p and p_T spectra of both single and dilepton events.
- 53 ADRIANI 93E use a multidimensional analysis based on a neural network approach.
- 54 AKERS 93B use a simultaneous fit to single and dilepton events (electrons and muons) to tag $Z \rightarrow b\bar{b}$.
- 55 JACOBSEN 91 tagged $b\bar{b}$ events by requiring coincidence of ≥ 3 tracks with significant impact parameters using vertex detector. Systematic error includes lifetime and decay uncertainties (± 0.014).
- 56 BARATE 97E combine a lifetime tag with a mass cut based on the mass difference between c hadrons and b hadrons.
- 57 ABE 96E obtain this value by combining results from three different b -tagging methods (2D impact parameter, 3D impact parameter, and 3D displaced vertex).
- 58 ABREU 95D obtain this result combining several analyses (double-lifetime tag and mixed tags). The second error contains an uncertainty of ± 0.0029 due to the total systematics and an uncertainty of ± 0.0016 due to an 8% variation of $\Gamma(c\bar{c})/\Gamma(\text{hadrons})$ around its Standard Model value (0.171 ± 0.014). Combining with their own lepton analysis, ABREU 95D obtain $0.2210 \pm 0.0033 \pm 0.0003$ (models) ± 0.0014 [$\Gamma(c\bar{c})/\Gamma(\text{hadrons})$].
- 59 ABREU 95J obtain this value with a multivariate analysis based on event shape and particle trajectories near the interaction point. The second error contains an uncertainty of ± 0.0012 due to an 8% variation of $\Gamma(c\bar{c})/\Gamma(\text{hadrons})$ around its Standard Model value (0.171 ± 0.014).
- 60 AKERS 95B select events based on the lepton and/or vertex tag independently in each hemisphere. Comparing the numbers of single- and double-tagged events, they determine the b -tagging efficiency directly from data.
- 61 BUSKULIC 93N use event shape and high p_T lepton discriminators applied to both hemispheres.
- 62 ABREU 92 result is from an indirect technique. They measure the lifetime τ_B , but use a world average of τ_B independent of $\Gamma(b\bar{b})$ and compare to their $\Gamma(b\bar{b})$ dependent lifetime from a hadron sample.
- 63 ABREU 92K use boosted-sphericity technique to tag and enrich the $b\bar{b}$ content with a sample of 50k hadronic events. Most of the systematic error is from hadronization uncertainty.
- 64 ABREU 92O use the neural network technique to tag heavy flavour events among a sample of 123k selected hadronic events. The systematic error consists of three parts: due to Monte Carlo (MC) parametrization (0.010), choice of MC model (0.008), and detector effects (0.011) added in quadrature.
- 65 KRAL 90 used isolated leptons and found $\Gamma(b\bar{b})/\Gamma(\text{total}) = 0.17^{+0.07+0.04}_{-0.06-0.03}$.

$\Gamma(ggg)/\Gamma(\text{hadrons})$
 Γ_{11}/Γ_6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.6 \times 10^{-2}$	95	⁶⁶ ABREU	96S DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV

⁶⁶ This branching ratio is slightly dependent on the jet-finder algorithm. The value we quote is obtained using the JADE algorithm, while using the DURHAM algorithm ABREU 96S obtain an upper limit of 1.5×10^{-2} .

 $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$
 Γ_{12}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-5}$	95	⁶⁷ ACCIARRI	95G L3	$E_{\text{cm}}^{ee} = 88-94$ GeV
$<5.5 \times 10^{-5}$	95	ABREU	94B DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
$<2.1 \times 10^{-4}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{ee} = 88-94$ GeV
$<1.4 \times 10^{-4}$	95	AKRAWY	91F OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.2 \times 10^{-4}$	95	⁶⁸ ADRIANI	92B L3	Repl. by ACCIARRI 95G
-----------------------	----	-----------------------	--------	-----------------------

⁶⁷ This limit is for both decay modes $Z \rightarrow \pi^0\gamma/\gamma\gamma$ which are indistinguishable in ACCIARRI 95G.

⁶⁸ This limit is for both decay modes $Z \rightarrow \pi^0\gamma/\gamma\gamma$ which are indistinguishable in ADRIANI 92B.

 $\Gamma(\eta\gamma)/\Gamma_{\text{total}}$
 Γ_{13}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.6 \times 10^{-5}$	95	ACCIARRI	95G L3	$E_{\text{cm}}^{ee} = 88-94$ GeV
$<8.0 \times 10^{-5}$	95	ABREU	94B DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
$<5.1 \times 10^{-5}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{ee} = 88-94$ GeV
$<2.0 \times 10^{-4}$	95	AKRAWY	91F OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.8 \times 10^{-4}$	95	ADRIANI	92B L3	Repl. by ACCIARRI 95G
-----------------------	----	---------	--------	-----------------------

 $\Gamma(\omega\gamma)/\Gamma_{\text{total}}$
 Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.5 \times 10^{-4}$	95	ABREU	94B DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV

 $\Gamma(\eta'(958)\gamma)/\Gamma_{\text{total}}$
 Γ_{15}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.2 \times 10^{-5}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{ee} = 88-94$ GeV

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$
 Γ_{16}/Γ

This decay would violate the Landau-Yang theorem.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-5}$	95	⁶⁹ ACCIARRI	95G L3	$E_{\text{cm}}^{ee} = 88-94$ GeV
$<5.5 \times 10^{-5}$	95	ABREU	94B DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
$<1.4 \times 10^{-4}$	95	AKRAWY	91F OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.2 \times 10^{-4}$	95	⁷⁰ ADRIANI	92B L3	Repl. by ACCIARRI 95G
-----------------------	----	-----------------------	--------	-----------------------

⁶⁹ This limit is for both decay modes $Z \rightarrow \pi^0\gamma/\gamma\gamma$ which are indistinguishable in ACCIARRI 95G.

⁷⁰ This limit is for both decay modes $Z \rightarrow \pi^0\gamma/\gamma\gamma$ which are indistinguishable in ADRIANI 92B.

$\Gamma(\gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-5}$	95	⁷¹ ACCIARRI	95C L3	$E_{\text{cm}}^{ee} = 88-94$ GeV
$<1.7 \times 10^{-5}$	95	⁷¹ ABREU	94B DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
$<6.6 \times 10^{-5}$	95	AKRAWY	91F OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<3.3 \times 10^{-5}$	95	ADRIANI	92B L3	Repl. by ACCIARRI 95C
-----------------------	----	---------	--------	-----------------------

⁷¹ Limit derived in the context of composite Z model.

 $\Gamma(\pi^{\pm} W^{\mp})/\Gamma_{\text{total}}$ Γ_{18}/Γ

The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7 \times 10^{-5}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{ee} = 88-94$ GeV

 $\Gamma(\rho^{\pm} W^{\mp})/\Gamma_{\text{total}}$ Γ_{19}/Γ

The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.3 \times 10^{-5}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{ee} = 88-94$ GeV

 $\Gamma(J/\psi(1S)X)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.66 ± 0.23 OUR AVERAGE				

$3.40 \pm 0.23 \pm 0.27$	441	⁷² ACCIARRI	97J L3	$E_{\text{cm}}^{ee} = 88-94$ GeV
--------------------------	-----	------------------------	--------	----------------------------------

$3.9 \pm 0.2 \pm 0.3$	511	⁷³ ALEXANDER	96B OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV
-----------------------	-----	-------------------------	----------	----------------------------------

$3.73 \pm 0.39 \pm 0.36$	153	⁷⁴ ABREU	94P DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
--------------------------	-----	---------------------	----------	----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.6 \pm 0.5 \pm 0.4$	121	⁷⁴ ADRIANI	93J L3	Repl. by ACCIARRI 97J
-----------------------	-----	-----------------------	--------	-----------------------

⁷² ACCIARRI 97J combine $\mu^+ \mu^-$ and $e^+ e^-$ $J/\psi(1S)$ decay channels and take into account the common systematic error.

⁷³ ALEXANDER 96B identify $J/\psi(1S)$ from the decays into lepton pairs. $(4.8 \pm 2.4)\%$ of this branching ratio is due to prompt $J/\psi(1S)$ production (ALEXANDER 96N).

⁷⁴ Combining $\mu^+ \mu^-$ and $e^+ e^-$ channels and taking into account the common systematic errors. $(7.7^{+6.3}_{-5.4})\%$ of this branching ratio is due to prompt $J/\psi(1S)$ production.

 $\Gamma(\psi(2S)X)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.60 ± 0.29 OUR AVERAGE				

$1.6 \pm 0.5 \pm 0.3$	39	⁷⁵ ACCIARRI	97J L3	$E_{\text{cm}}^{ee} = 88-94$ GeV
-----------------------	----	------------------------	--------	----------------------------------

$1.6 \pm 0.3 \pm 0.2$	46.9	⁷⁶ ALEXANDER	96B OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV
-----------------------	------	-------------------------	----------	----------------------------------

$1.60 \pm 0.73 \pm 0.33$	5.4	⁷⁷ ABREU	94P DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
--------------------------	-----	---------------------	----------	----------------------------------

⁷⁵ ACCIARRI 97J measure this branching ratio via the decay channel $\psi(2S) \rightarrow \ell^+ \ell^-$ ($\ell = \mu, e$).

⁷⁶ ALEXANDER 96B measure this branching ratio via the decay channel $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$, with $J/\psi \rightarrow \ell^+ \ell^-$.

⁷⁷ ABREU 94P measure this branching ratio via decay channel $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$, with $J/\psi \rightarrow \mu^+ \mu^-$.

$$\Gamma(\chi_{c1}(1P)X)/\Gamma_{\text{total}} \qquad \Gamma_{22}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.9 ± 0.7 OUR AVERAGE

2.7 ± 0.6 ± 0.5	33	⁷⁸ ACCIARRI	97J L3	$E_{\text{cm}}^{ee} = 88-94$ GeV
-----------------	----	------------------------	--------	----------------------------------

5.0 ± 2.1 ^{+1.5} _{-0.9}	6.4	⁷⁹ ABREU	94P DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
---	-----	---------------------	----------	----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.5 ± 2.9 ± 0.6	19	⁷⁹ ADRIANI	93J L3	Repl. by ACCIARRI 97J
-----------------	----	-----------------------	--------	-----------------------

⁷⁸ ACCIARRI 97J measure this branching ratio via the decay channel $\chi_{c1} \rightarrow J/\psi + \gamma$, with $J/\psi \rightarrow \ell^+ \ell^-$ ($\ell = \mu, e$). The $M(\ell^+ \ell^- \gamma) - M(\ell^+ \ell^-)$ mass difference spectrum is fitted with two gaussian shapes for χ_{c1} and χ_{c2} .

⁷⁹ This branching ratio is measured via the decay channel $\chi_{c1} \rightarrow J/\psi + \gamma$, with $J/\psi \rightarrow \mu^+ \mu^-$.

$$\Gamma(\chi_{c2}(1P)X)/\Gamma_{\text{total}} \qquad \Gamma_{23}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 3.2 × 10⁻³	90	⁸⁰ ACCIARRI	97J L3	$E_{\text{cm}}^{ee} = 88-94$ GeV
-----------------------------------	----	------------------------	--------	----------------------------------

⁸⁰ ACCIARRI 97J derive this limit via the decay channel $\chi_{c2} \rightarrow J/\psi + \gamma$, with $J/\psi \rightarrow \ell^+ \ell^-$ ($\ell = \mu, e$). The $M(\ell^+ \ell^- \gamma) - M(\ell^+ \ell^-)$ mass difference spectrum is fitted with two gaussian shapes for χ_{c1} and χ_{c2} .

$$\Gamma(\Upsilon(1S)X + \Upsilon(2S)X + \Upsilon(3S)X)/\Gamma_{\text{total}} \qquad \Gamma_{24}/\Gamma = (\Gamma_{25} + \Gamma_{26} + \Gamma_{27})/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.0 ± 0.4 ± 0.22	6.4	⁸¹ ALEXANDER	96F OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV
-------------------------	-----	-------------------------	----------	----------------------------------

⁸¹ ALEXANDER 96F identify the Υ (which refers to any of the three lowest bound states) through its decay into $e^+ e^-$ and $\mu^+ \mu^-$. The systematic error includes an uncertainty of ± 0.2 due to the production mechanism.

$$\Gamma(\Upsilon(1S)X)/\Gamma_{\text{total}} \qquad \Gamma_{25}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 5.5 × 10⁻⁵	95	⁸² ACCIARRI	97R L3	$E_{\text{cm}}^{ee} = 88-94$ GeV
-----------------------------------	----	------------------------	--------	----------------------------------

⁸² ACCIARRI 97R search for $\Upsilon(1S)$ through its decay into $\ell^+ \ell^-$ ($\ell = e$ or μ).

$$\Gamma(\Upsilon(2S)X)/\Gamma_{\text{total}} \qquad \Gamma_{26}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 13.9 × 10⁻⁵	95	⁸³ ACCIARRI	97R L3	$E_{\text{cm}}^{ee} = 88-94$ GeV
------------------------------------	----	------------------------	--------	----------------------------------

⁸³ ACCIARRI 97R search for $\Upsilon(2S)$ through its decay into $\ell^+ \ell^-$ ($\ell = e$ or μ).

$$\Gamma(\Upsilon(3S)X)/\Gamma_{\text{total}} \qquad \Gamma_{27}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 9.4 × 10⁻⁵	95	⁸⁴ ACCIARRI	97R L3	$E_{\text{cm}}^{ee} = 88-94$ GeV
-----------------------------------	----	------------------------	--------	----------------------------------

⁸⁴ ACCIARRI 97R search for $\Upsilon(3S)$ through its decay into $\ell^+ \ell^-$ ($\ell = e$ or μ).

$$\Gamma((D^0/\bar{D}^0)X)/\Gamma(\text{hadrons}) \qquad \Gamma_{28}/\Gamma_6$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.296 ± 0.019 ± 0.021	369	⁸⁵ ABREU	93I DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
------------------------------	-----	---------------------	----------	----------------------------------

⁸⁵ The (D^0/\bar{D}^0) states in ABREU 93I are detected by the $K\pi$ decay mode. This is a corrected result (see the erratum of ABREU 93I).

$\Gamma(D^\pm X)/\Gamma(\text{hadrons})$ Γ_{29}/Γ_6

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.174±0.016±0.018	539	⁸⁶ ABREU	93I DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

⁸⁶ The D^\pm states in ABREU 93I are detected by the $K\pi\pi$ decay mode. This is a corrected result (see the erratum of ABREU 93I).

 $\Gamma(D^*(2010)^\pm X)/\Gamma(\text{hadrons})$ Γ_{30}/Γ_6

The value is for the sum of the charge states indicated.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.163±0.019 OUR AVERAGE		Error includes scale factor of 1.3.		
0.155±0.010±0.013	358	⁸⁷ ABREU	93I DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.21 ±0.04	362	⁸⁸ DECAMP	91J ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

⁸⁷ $D^*(2010)^\pm$ in ABREU 93I are reconstructed from $D^0\pi^\pm$, with $D^0 \rightarrow K^-\pi^+$. The new CLEO II measurement of $B(D^{*\pm} \rightarrow D^0\pi^\pm) = (68.1 \pm 1.6)\%$ is used. This is a corrected result (see the erratum of ABREU 93I).

⁸⁸ DECAMP 91J report $B(D^*(2010)^+ \rightarrow D^0\pi^+) B(D^0 \rightarrow K^-\pi^+) \Gamma(D^*(2010)^\pm X) / \Gamma(\text{hadrons}) = (5.11 \pm 0.34) \times 10^{-3}$. They obtained the above number assuming $B(D^0 \rightarrow K^-\pi^+) = (3.62 \pm 0.34 \pm 0.44)\%$ and $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (55 \pm 4)\%$. We have rescaled their original result of 0.26 ± 0.05 taking into account the new CLEO II branching ratio $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (68.1 \pm 1.6)\%$.

 $\Gamma(B_s^0 X)/\Gamma(\text{hadrons})$ Γ_{33}/Γ_6

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	⁸⁹ ABREU	92M DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
seen	⁹⁰ ACTON	92N OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
seen	⁹¹ BUSKULIC	92E ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

⁸⁹ ABREU 92M reported value is $\Gamma(B_s^0 X) * B(B_s^0 \rightarrow D_s \mu \nu_\mu X) * B(D_s \rightarrow \phi\pi) / \Gamma(\text{hadrons}) = (18 \pm 8) \times 10^{-5}$.

⁹⁰ ACTON 92N find evidence for B_s^0 production using $D_s\text{-}\ell$ correlations, with $D_s^+ \rightarrow \phi\pi^+$ and $K^*(892)K^+$. Assuming R_b from the Standard Model and averaging over the e and μ channels, authors measure the product branching fraction to be $f(\bar{b} \rightarrow B_s^0) \times B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell X) \times B(D_s^- \rightarrow \phi\pi^-) = (3.9 \pm 1.1 \pm 0.8) \times 10^{-4}$.

⁹¹ BUSKULIC 92E find evidence for B_s^0 production using $D_s\text{-}\ell$ correlations, with $D_s^+ \rightarrow \phi\pi^+$ and $K^*(892)K^+$. Using $B(D_s^+ \rightarrow \phi\pi^+) = (2.7 \pm 0.7)\%$ and summing up the e and μ channels, the weighted average product branching fraction is measured to be $B(\bar{b} \rightarrow B_s^0) \times B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell X) = 0.040 \pm 0.011_{-0.012}^{+0.010}$.

 $\Gamma(B^* X) / [\Gamma(BX) + \Gamma(B^* X)]$ $\Gamma_{32}/(\Gamma_{31} + \Gamma_{32})$

As the experiments assume different values of the b -baryon contribution, our average should be taken with caution. If we assume a common baryon production fraction $f_{\Lambda_b} = (13.2 \pm 4.1)\%$ as given in the 1996 edition of this *Review* OUR AVERAGE becomes 0.77 ± 0.04 .

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.75 ±0.04 OUR AVERAGE				
0.760±0.036±0.083		⁹² ACKERSTAFF	97M OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.771±0.026±0.070		⁹³ BUSKULIC	96D ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.72 ±0.03 ±0.06		⁹⁴ ABREU	95R DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.76 ±0.08 ±0.06	1378	⁹⁵ ACCIARRI	95B L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

⁹² ACKERSTAFF 97M use an inclusive B reconstruction method and assume a $(13.2 \pm 4.1)\%$ b -baryon contribution. The value refers to a b -flavored meson mixture of B_u , B_d , and B_s .

⁹³ BUSKULIC 96D use an inclusive reconstruction of B hadrons and assume a $(12.2 \pm 4.3)\%$ b -baryon contribution. The value refers to a b -flavored mixture of B_u , B_d , and B_s .

⁹⁴ ABREU 95R use an inclusive B -reconstruction method and assume a $(10 \pm 4)\%$ b -baryon contribution. The value refers to a b -flavored meson mixture of B_u , B_d , and B_s .

⁹⁵ ACCIARRI 95B assume a 9.4% b -baryon contribution. The value refers to a b -flavored mixture of B_u , B_d , and B_s .

$\Gamma(\text{anomalous } \gamma + \text{hadrons})/\Gamma_{\text{total}}$

Γ_{34}/Γ

Limits on additional sources of prompt photons beyond expectations for final-state bremsstrahlung.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.2 \times 10^{-3}$	95	⁹⁶ AKRAWY	90J OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

⁹⁶ AKRAWY 90J report $\Gamma(\gamma X) < 8.2$ MeV at 95%CL. They assume a three-body $\gamma q\bar{q}$ distribution and use $E(\gamma) > 10$ GeV.

$\Gamma(e^+ e^- \gamma)/\Gamma_{\text{total}}$

Γ_{35}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-4}$	95	⁹⁷ ACTON	91B OPAL	$E_{\text{cm}}^{ee} = 91.2$ GeV

⁹⁷ ACTON 91B looked for isolated photons with $E > 2\%$ of beam energy (> 0.9 GeV).

$\Gamma(\mu^+ \mu^- \gamma)/\Gamma_{\text{total}}$

Γ_{36}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.6 \times 10^{-4}$	95	⁹⁸ ACTON	91B OPAL	$E_{\text{cm}}^{ee} = 91.2$ GeV

⁹⁸ ACTON 91B looked for isolated photons with $E > 2\%$ of beam energy (> 0.9 GeV).

$\Gamma(\tau^+ \tau^- \gamma)/\Gamma_{\text{total}}$

Γ_{37}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.3 \times 10^{-4}$	95	⁹⁹ ACTON	91B OPAL	$E_{\text{cm}}^{ee} = 91.2$ GeV

⁹⁹ ACTON 91B looked for isolated photons with $E > 2\%$ of beam energy (> 0.9 GeV).

$\Gamma(\ell^+ \ell^- \gamma\gamma)/\Gamma_{\text{total}}$

Γ_{38}/Γ

The value is the sum over $\ell = e, \mu, \tau$.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.8 \times 10^{-6}$	95	¹⁰⁰ ACTON	93E OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

¹⁰⁰ For $m_{\gamma\gamma} = 60 \pm 5$ GeV.

$\Gamma(q\bar{q}\gamma\gamma)/\Gamma_{\text{total}}$

Γ_{39}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.5 \times 10^{-6}$	95	¹⁰¹ ACTON	93E OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

¹⁰¹ For $m_{\gamma\gamma} = 60 \pm 5$ GeV.

$\Gamma(\nu\bar{\nu}\gamma\gamma)/\Gamma_{\text{total}}$

Γ_{40}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.1 \times 10^{-6}$	95	¹⁰² ACTON	93E OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

¹⁰² For $m_{\gamma\gamma} = 60 \pm 5$ GeV.

$$\Gamma(e^\pm \mu^\mp) / \Gamma(e^+ e^-)$$

 Γ_{41} / Γ_1

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.07	90	ALBAJAR	89 UA1	$E_{cm}^{p\bar{p}} = 546,630 \text{ GeV}$

$$\Gamma(e^\pm \mu^\mp) / \Gamma_{\text{total}}$$

 Γ_{41} / Γ

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.5 \times 10^{-6}$	95	ABREU	97C DLPH	$E_{cm}^{ee} = 88-94 \text{ GeV}$
<1.7 $\times 10^{-6}$	95	AKERS	95W OPAL	$E_{cm}^{ee} = 88-94 \text{ GeV}$
$<0.6 \times 10^{-5}$	95	ADRIANI	93i L3	$E_{cm}^{ee} = 88-94 \text{ GeV}$
$<2.6 \times 10^{-5}$	95	DECAMP	92 ALEP	$E_{cm}^{ee} = 88-94 \text{ GeV}$

$$\Gamma(e^\pm \tau^\mp) / \Gamma_{\text{total}}$$

 Γ_{42} / Γ

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-5}$	95	ABREU	97C DLPH	$E_{cm}^{ee} = 88-94 \text{ GeV}$
<9.8 $\times 10^{-6}$	95	AKERS	95W OPAL	$E_{cm}^{ee} = 88-94 \text{ GeV}$
$<1.3 \times 10^{-5}$	95	ADRIANI	93i L3	$E_{cm}^{ee} = 88-94 \text{ GeV}$
$<1.2 \times 10^{-4}$	95	DECAMP	92 ALEP	$E_{cm}^{ee} = 88-94 \text{ GeV}$

$$\Gamma(\mu^\pm \tau^\mp) / \Gamma_{\text{total}}$$

 Γ_{43} / Γ

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.2 $\times 10^{-5}$	95	ABREU	97C DLPH	$E_{cm}^{ee} = 88-94 \text{ GeV}$
$<1.7 \times 10^{-5}$	95	AKERS	95W OPAL	$E_{cm}^{ee} = 88-94 \text{ GeV}$
$<1.9 \times 10^{-5}$	95	ADRIANI	93i L3	$E_{cm}^{ee} = 88-94 \text{ GeV}$
$<1.0 \times 10^{-4}$	95	DECAMP	92 ALEP	$E_{cm}^{ee} = 88-94 \text{ GeV}$

Z Boson – THIS IS PART 2 OF 2

To reduce the size of this section's PostScript file, we have divided it into two PostScript files. We present the following index:

PART 1

Page #	Section name
1	Mass
17	Width
18	Decay modes
19	Partial widths
21	Branching ratios

PART 2

Page #	Section name
35	Average particle multiplicities in hadronic z decay
42	Hadronic pole cross section
42	Vector couplings to charged leptons
44	Axial-vector couplings to charged leptons
45	Couplings to neutral leptons
46	Asymmetry parameters
48	Transverse spin correlations
49	Charge asymmetry
55	References

AVERAGE PARTICLE MULTIPLICITIES IN HADRONIC Z DECAY

Summed over particle and antiparticle, when appropriate.

$\langle N_{\pi^\pm} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
17.05 ± 0.43	AKERS	94P OPAL	$E_{cm}^{ee} = 91.2$ GeV

$\langle N_{\pi^0} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
9.79 ± 0.28 OUR AVERAGE			
9.63 ± 0.13 ± 0.63	BARATE	97J ALEP	$E_{cm}^{ee} = 91.2$ GeV
9.90 ± 0.02 ± 0.33	ACCIARRI	96 L3	$E_{cm}^{ee} = 91.2$ GeV
9.2 ± 0.2 ± 1.0	ADAM	96 DLPH	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
9.18 ± 0.03 ± 0.73	ACCIARRI	94B L3	Repl. by ACCIARRI 96

$\langle N_\eta \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.93 ± 0.01 ± 0.09	ACCIARRI	96 L3	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.91 ± 0.02 ± 0.11	ACCIARRI	94B L3	Repl. by ACCIARRI 96
0.298 ± 0.023 ± 0.021	¹⁰³ BUSKULIC	92D ALEP	$E_{cm}^{ee} = 91.2$ GeV
¹⁰³ BUSKULIC 92D obtain this value for $x > 0.1$.			

$\langle N_{\rho^0} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
1.30 ± 0.12 OUR AVERAGE			
1.45 ± 0.06 ± 0.20	BUSKULIC	96H ALEP	$E_{cm}^{ee} = 91.2$ GeV
1.21 ± 0.04 ± 0.15	ABREU	95L DLPH	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.43 ± 0.12 ± 0.22	ABREU	93 DLPH	Repl. by ABREU 95L

$\langle N_\omega \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
1.11 ± 0.11 OUR AVERAGE			
1.17 ± 0.09 ± 0.15	ACCIARRI	97D L3	$E_{cm}^{ee} = 91.2$ GeV
1.07 ± 0.06 ± 0.13	BUSKULIC	96H ALEP	$E_{cm}^{ee} = 91.2$ GeV

$\langle N_{\eta'} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.25 ± 0.04	¹⁰⁴ ACCIARRI	97D L3	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.068 ± 0.018 ± 0.016	¹⁰⁵ BUSKULIC	92D ALEP	$E_{cm}^{ee} = 91.2$ GeV
¹⁰⁴ ACCIARRI 97D obtain this value averaging over the two decay channels $\eta' \rightarrow \pi^+ \pi^- \eta$ and $\eta' \rightarrow \rho^0 \gamma$.			
¹⁰⁵ BUSKULIC 92D obtain this value for $x > 0.1$.			

$\langle N_{f_0(980)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.098 ± 0.016	¹⁰⁶ ABREU	95L DLPH	$E_{cm}^{ee} = 91.2$ GeV
$0.10 \pm 0.03 \pm 0.019$	¹⁰⁷ ABREU	93 DLPH	Repl. by ABREU 95L

¹⁰⁶ ABREU 95L obtain this value for $0.05 < x < 0.6$.

¹⁰⁷ ABREU 93 obtain this value for $x > 0.05$.

$\langle N_\phi \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.108 ± 0.006 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.


$0.104 \pm 0.003 \pm 0.007$	ABREU	96U DLPH	$E_{cm}^{ee} = 91.2$ GeV
-----------------------------	-------	----------	--------------------------

$0.122 \pm 0.004 \pm 0.008$	BUSKULIC	96H ALEP	$E_{cm}^{ee} = 91.2$ GeV
-----------------------------	----------	----------	--------------------------

$0.100 \pm 0.004 \pm 0.007$	AKERS	95X OPAL	$E_{cm}^{ee} = 91.2$ GeV
-----------------------------	-------	----------	--------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.086 \pm 0.015 \pm 0.010$	ACTON	92O OPAL	Repl. by AKERS 95X
-----------------------------	-------	----------	--------------------



$\langle N_\phi \rangle$

$\langle N_{f_2(1270)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.170 ± 0.043	¹⁰⁸ ABREU	95L DLPH	$E_{cm}^{ee} = 91.2$ GeV
-------------------	----------------------	----------	--------------------------

$0.11 \pm 0.04 \pm 0.03$	¹⁰⁹ ABREU	93 DLPH	Repl. by ABREU 95L
--------------------------	----------------------	---------	--------------------

¹⁰⁸ ABREU 95L obtain this value for $x > 0.05$.

¹⁰⁹ ABREU 93 obtain this value for $x > 0.1$.

$\langle N_{f_2'(1525)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.020 ± 0.005 ± 0.006	ABREU	96C DLPH	$E_{cm}^{ee} = 91.2$ GeV

 $\langle N_{K^\pm} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
2.37 ± 0.11 OUR AVERAGE			
2.26 ± 0.01 ± 0.18	ABREU	95F DLPH	$E_{cm}^{ee} = 91.2$ GeV
2.42 ± 0.13	AKERS	94P OPAL	$E_{cm}^{ee} = 91.2$ GeV

 $\langle N_{K^0} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
2.013 ± 0.023 OUR AVERAGE			
2.024 ± 0.006 ± 0.042	ACCIARRI	97L L3	$E_{cm}^{ee} = 91.2$ GeV
1.962 ± 0.022 ± 0.056	ABREU	95L DLPH	$E_{cm}^{ee} = 91.2$ GeV
1.99 ± 0.01 ± 0.04	AKERS	95U OPAL	$E_{cm}^{ee} = 91.2$ GeV
2.061 ± 0.047	BUSKULIC	94K ALEP	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.04 ± 0.02 ± 0.14	ACCIARRI	94B L3	Repl. by ACCIARRI 97L
2.12 ± 0.05 ± 0.04	ABREU	92G DLPH	Repl. by ABREU 95L

 $\langle N_{K^*(892)^\pm} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.72 ± 0.05 OUR AVERAGE			
0.712 ± 0.031 ± 0.059	ABREU	95L DLPH	$E_{cm}^{ee} = 91.2$ GeV
0.72 ± 0.02 ± 0.08	ACTON	93 OPAL	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.33 ± 0.11 ± 0.24	ABREU	92G DLPH	Repl. by ABREU 95L

 $\langle N_{K^*(892)^0} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.752 ± 0.025 OUR AVERAGE			
0.74 ± 0.02 ± 0.02	ACKERSTAFF	97S OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.77 ± 0.02 ± 0.07	ABREU	96U DLPH	$E_{cm}^{ee} = 91.2$ GeV
0.83 ± 0.01 ± 0.09	BUSKULIC	96H ALEP	$E_{cm}^{ee} = 91.2$ GeV
0.97 ± 0.18 ± 0.31	ABREU	93 DLPH	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.74 ± 0.03 ± 0.03	AKERS	95X OPAL	Repl. by ACKERSTAFF 97S


 $\langle N_{K_2^*(1430)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.079 ± 0.026 ± 0.031	ABREU	96U DLPH	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.19 ± 0.04 ± 0.06	¹¹⁰ AKERS	95X OPAL	$E_{cm}^{ee} = 91.2$ GeV
¹¹⁰ AKERS 95X obtain this value for $x < 0.3$.			

$\langle N_{D^\pm} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.187±0.020 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.		
0.170±0.009±0.014	ALEXANDER	96R OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.251±0.026±0.025	BUSKULIC	94J ALEP	$E_{cm}^{ee} = 91.2$ GeV
0.199±0.019±0.024	¹¹¹ ABREU	93I DLPH	$E_{cm}^{ee} = 91.2$ GeV

¹¹¹ See ABREU 95 (erratum).



$\langle N_{D^\pm} \rangle$

$\langle N_{D^0} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.462±0.026 OUR AVERAGE			
0.465±0.017±0.027	ALEXANDER	96R OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.518±0.052±0.035	BUSKULIC	94J ALEP	$E_{cm}^{ee} = 91.2$ GeV
0.403±0.038±0.044	¹¹² ABREU	93I DLPH	$E_{cm}^{ee} = 91.2$ GeV

¹¹² See ABREU 95 (erratum).

$\langle N_{D_s^\pm} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.131±0.010±0.018	ALEXANDER	96R OPAL	$E_{cm}^{ee} = 91.2$ GeV

$\langle N_{D^*(2010)^\pm} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.183 ± 0.008 OUR AVERAGE

0.1854 ± 0.0041 ± 0.0091	113 ACKERSTAFF	98E OPAL	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$
--------------------------	----------------	----------	---

0.187 ± 0.015 ± 0.013	BUSKULIC	94J ALEP	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$
-----------------------	----------	----------	---

0.171 ± 0.012 ± 0.016	114 ABREU	93I DLPH	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$
-----------------------	-----------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.183 ± 0.009 ± 0.011	115 AKERS	95O OPAL	Repl. by ACKERSTAFF 98E
-----------------------	-----------	----------	-------------------------

113 ACKERSTAFF 98E systematic error includes an uncertainty of ± 0.0069 due to the branching ratios $B(D^{*+} \rightarrow D^0 \pi^+) = 0.683 \pm 0.014$ and $B(D^0 \rightarrow K^- \pi^+) = 0.0383 \pm 0.0012$.

114 See ABREU 95 (erratum).

115 AKERS 95O systematic error includes an uncertainty of ± 0.008 due to the $D^{*\pm}$ and D^0 branching ratios [they use $B(D^* \rightarrow D^0 \pi) = 0.681 \pm 0.016$ and $B(D^0 \rightarrow K \pi) = 0.0401 \pm 0.0014$ to obtain this measurement].

 $\langle N_{D_{s1}(2536)^+} \rangle$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.9^{+0.7}_{-0.6} \pm 0.2$	116 ACKERSTAFF	97W OPAL	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$
-----------------------------	----------------	----------	---

116 ACKERSTAFF 97W obtain this value for $x > 0.6$ and with the assumption that its decay width is saturated by the $D^* K$ final states.

 $\langle N_{B^*} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.28 ± 0.01 ± 0.03	117 ABREU	95R DLPH	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$
---------------------------	-----------	----------	---

117 ABREU 95R quote this value for a flavor-averaged excited state.

 $\langle N_{J/\psi(1S)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.0056 ± 0.0003 ± 0.0004	118 ALEXANDER	96B OPAL	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$
---------------------------------	---------------	----------	---

118 ALEXANDER 96B identify $J/\psi(1S)$ from the decays into lepton pairs.

 $\langle N_{\psi(2S)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.0023 ± 0.0004 ± 0.0003	ALEXANDER	96B OPAL	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$
---------------------------------	-----------	----------	---

 $\langle N_{\rho} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.98 ± 0.09 OUR AVERAGE

1.07 ± 0.01 ± 0.14	ABREU	95F DLPH	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$
--------------------	-------	----------	---

0.92 ± 0.11	AKERS	94P OPAL	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$
-------------	-------	----------	---

$\langle N_{\Delta(1232)^{++}} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.087 ± 0.033 OUR AVERAGE	Error includes scale factor of 2.4.		
0.079 ± 0.009 ± 0.011	ABREU	95W DLPH	$E_{cm}^{ee} = 91.2$ GeV
0.22 ± 0.04 ± 0.04	ALEXANDER	95D OPAL	$E_{cm}^{ee} = 91.2$ GeV

 $\langle N_{\Lambda} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.372 ± 0.007 OUR AVERAGE			
0.364 ± 0.004 ± 0.017	ACCIARRI	97L L3	$E_{cm}^{ee} = 91.2$ GeV
0.374 ± 0.002 ± 0.010	ALEXANDER	97D OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.386 ± 0.016	BUSKULIC	94K ALEP	$E_{cm}^{ee} = 91.2$ GeV
0.357 ± 0.003 ± 0.017	ABREU	93L DLPH	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.37 ± 0.01 ± 0.04	ACCIARRI	94B L3	Repl. by ACCIARRI 97L
0.351 ± 0.019	ACTON	92J OPAL	Repl. by ALEXANDER 97D

 $\langle N_{\Lambda(1520)} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0213 ± 0.0021 ± 0.0019	ALEXANDER	97D OPAL	$E_{cm}^{ee} = 91.2$ GeV

 $\langle N_{\Sigma^+} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.099 ± 0.008 ± 0.013	ALEXANDER	97E OPAL	$E_{cm}^{ee} = 91.2$ GeV

 $\langle N_{\Sigma^-} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.083 ± 0.006 ± 0.009	ALEXANDER	97E OPAL	$E_{cm}^{ee} = 91.2$ GeV

 $\langle N_{\Sigma^+ + \Sigma^-} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.181 ± 0.018 OUR AVERAGE			
0.182 ± 0.010 ± 0.016	¹¹⁹ ALEXANDER	97E OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.170 ± 0.014 ± 0.061	ABREU	95O DLPH	$E_{cm}^{ee} = 91.2$ GeV

¹¹⁹ We have combined the values of $\langle N_{\Sigma^+} \rangle$ and $\langle N_{\Sigma^-} \rangle$ from ALEXANDER 97E adding the statistical and systematic errors of the two final states separately in quadrature. If isospin symmetry is assumed this value becomes $0.174 \pm 0.010 \pm 0.015$.

 $\langle N_{\Sigma^0} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.070 ± 0.011 OUR AVERAGE			
0.071 ± 0.012 ± 0.013	ALEXANDER	97E OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.070 ± 0.010 ± 0.010	ADAM	96B DLPH	$E_{cm}^{ee} = 91.2$ GeV

 $\langle N_{(\Sigma^+ + \Sigma^- + \Sigma^0)/3} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.084 ± 0.005 ± 0.008	ALEXANDER	97E OPAL	$E_{cm}^{ee} = 91.2$ GeV

$\langle N_{\Sigma(1385)^+} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0239 ± 0.0009 ± 0.0012	ALEXANDER	97D OPAL	$E_{cm}^{ee} = 91.2$ GeV

 $\langle N_{\Sigma(1385)^-} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0240 ± 0.0010 ± 0.0014	ALEXANDER	97D OPAL	$E_{cm}^{ee} = 91.2$ GeV

 $\langle N_{\Sigma(1385)^+ + \Sigma(1385)^-} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.046 ± 0.004 OUR AVERAGE	Error includes scale factor of 1.6.		
0.0479 ± 0.0013 ± 0.0026	ALEXANDER	97D OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.0382 ± 0.0028 ± 0.0045	ABREU	95O DLPH	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.0380 ± 0.0062	ACTON	92J OPAL	Repl. by ALEXANDER 97D

 $\langle N_{\Xi^-} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0258 ± 0.0009 OUR AVERAGE			
0.0259 ± 0.0004 ± 0.0009	ALEXANDER	97D OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.0250 ± 0.0009 ± 0.0021	ABREU	95O DLPH	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.020 ± 0.004 ± 0.003	ABREU	92G DLPH	Repl. by ABREU 95O
0.0206 ± 0.0021	ACTON	92J OPAL	Repl. by ALEXANDER 97D

 $\langle N_{\Xi(1530)^0} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0053 ± 0.0013 OUR AVERAGE	Error includes scale factor of 3.2.		
0.0068 ± 0.0005 ± 0.0004	ALEXANDER	97D OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.0041 ± 0.0004 ± 0.0004	ABREU	95O DLPH	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.0063 ± 0.0014	ACTON	92J OPAL	Repl. by ALEXANDER 97D

 $\langle N_{\Omega^-} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.00164 ± 0.00028 OUR AVERAGE			
0.0018 ± 0.0003 ± 0.0002	ALEXANDER	97D OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.0014 ± 0.0002 ± 0.0004	ADAM	96B DLPH	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.0050 ± 0.0015	ACTON	92J OPAL	Repl. by ALEXANDER 97D

 $\langle N_{\Lambda_c^+} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.078 ± 0.012 ± 0.012	ALEXANDER	96R OPAL	$E_{cm}^{ee} = 91.2$ GeV

$\langle N_{\text{charged}} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
21.00±0.13 OUR AVERAGE			
21.05±0.20	AKERS	95Z OPAL	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$
20.91±0.03±0.22	BUSKULIC	95R ALEP	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$
21.40±0.43	ACTON	92B OPAL	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$
20.71±0.04±0.77	ABREU	91H DLPH	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$
20.7 ±0.7	ADEVA	91I L3	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$
20.1 ±1.0 ±0.9	ABRAMS	90 MRK2	$E_{\text{cm}}^{ee} = 91.1 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
20.85±0.02±0.24	DECAMP	91K ALEP	Repl. by BUSKULIC 95R

Z HADRONIC POLE CROSS SECTION

This quantity is defined as

$$\sigma_h^0 = \frac{12\pi}{M_Z^2} \frac{\Gamma(e^+ e^-) \Gamma(\text{hadrons})}{\Gamma_Z^2}$$

It is one of the parameters used in the Z lineshape fit. (See the 'Note on the Z Boson.')

<u>VALUE (nb)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
41.54±0.14 OUR FIT				
41.49±0.10 OUR AVERAGE				
41.23±0.20	1.05M	ABREU	94 DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
41.39±0.26	1.09M	ACCIARRI	94 L3	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
41.70±0.23	1.19M	AKERS	94 OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
41.60±0.16	1.27M	BUSKULIC	94 ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
41.45±0.31	512k	ACTON	93D OPAL	Repl. by AKERS 94
41.34±0.28	460k	ADRIANI	93M L3	Repl. by ACCIARRI 94
41.60±0.27	520k	BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
42 ±4	450	ABRAMS	89B MRK2	$E_{\text{cm}}^{ee} = 89.2\text{--}93.0 \text{ GeV}$

Z VECTOR COUPLINGS TO CHARGED LEPTONS

These quantities are the effective vector couplings of the Z to charged leptons. Their magnitude is derived from a measurement of the Z lineshape and the forward-backward lepton asymmetries as a function of energy around the Z mass. The relative sign among the vector to axial-vector couplings is obtained from a measurement of the Z asymmetry parameters, A_e and A_τ , or ν_e scattering. The fit values quoted below correspond to global nine- or five-parameter fits to lineshape, lepton forward-backward asymmetry, and A_e and A_τ measurements. See "Note on the Z boson" for details.

Within the current data set, the reason for the smallness of g_V^μ compared to g_V^e and g_V^τ is due to the large value of A_e which is heavily weighted by the SLD result. This large value of A_e leads to a large value of g_V^e . Since

g_V^μ is obtained using the relation $A_{FB}^\mu = 0.75 \times A_e \times A_\mu$, a large value of g_V^e leads to a SMALL value of g_V^μ . Concerning the τ , its g_V gets mainly determined directly from A_τ which is obtained from a measurement of the τ polarization (see "Note on the Z boson").

 g_V^e

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.0383 ± 0.0008 OUR FIT				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
-0.0414 ± 0.0020		120 ABE	95J SLD	$E_{cm}^{ee} = 91.31$ GeV
-0.0364 $\begin{smallmatrix} +0.0096 \\ -0.0082 \end{smallmatrix}$	38k	121 ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
-0.036 ± 0.005	45.8k	122 BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
-0.040 $\begin{smallmatrix} +0.013 \\ -0.011 \end{smallmatrix}$		123 ADRIANI	93M L3	Repl. by ACCIARRI 94
-0.034 $\begin{smallmatrix} +0.006 \\ -0.005 \end{smallmatrix}$		121 BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
120 ABE 95J obtain this result combining polarized Bhabha results with the A_{LR} measurement of ABE 94C. The Bhabha results alone give $-0.0507 \pm 0.0096 \pm 0.0020$.				
121 The τ polarization result has been included.				
122 BUSKULIC 94 use the added constraint of τ polarization.				
123 ADRIANI 93M use their measurement of the τ polarization in addition to forward-backward lepton asymmetries.				

 g_V^μ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.0274 ± 0.0047 OUR FIT				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
-0.0402 $\begin{smallmatrix} +0.0153 \\ -0.0211 \end{smallmatrix}$	34k	124 ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
-0.034 ± 0.013	46.4k	125 BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
-0.048 $\begin{smallmatrix} +0.021 \\ -0.033 \end{smallmatrix}$		126 ADRIANI	93M L3	Repl. by ACCIARRI 94
-0.019 $\begin{smallmatrix} +0.018 \\ -0.019 \end{smallmatrix}$		124 BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
124 The τ polarization result has been included.				
125 BUSKULIC 94 use the added constraint of τ polarization.				
126 ADRIANI 93M use their measurement of the τ polarization in addition to forward-backward lepton asymmetries.				

 g_V^τ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.0378 ± 0.0020 OUR FIT				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
-0.0384 ± 0.0078	25k	127 ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
-0.038 ± 0.005	45.1k	128 BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
-0.037 ± 0.008	7441	129 ADRIANI	93M L3	Repl. by ACCIARRI 94
-0.039 ± 0.006		127 BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
127 The τ polarization result has been included.				
128 BUSKULIC 94 use the added constraint of τ polarization.				
129 ADRIANI 93M use their measurement of the τ polarization in addition to forward-backward lepton asymmetries.				

g_V^l

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.0377 ± 0.0007 OUR FIT				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
-0.039 ± 0.004	50.3k	¹³⁰ ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV
$-0.0378^{+0.0045}_{-0.0042}$	97k	¹³¹ ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
-0.034 ± 0.004	146k	¹³⁰ AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV
-0.038 ± 0.004	137.3k	¹³⁰ BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
-0.027 ± 0.008	58k	¹³⁰ ACTON	93D OPAL	Repl. by AKERS 94
$-0.040^{+0.006}_{-0.005}$		¹³¹ ADRIANI	93M L3	Repl. by ACCIARRI 94
$-0.034^{+0.004}_{-0.003}$		¹³¹ BUSKULIC	93J ALEP	Repl. by BUSKULIC 94

¹³⁰ Using forward-backward lepton asymmetries.¹³¹ The τ polarization result has been included.

Z AXIAL-VECTOR COUPLINGS TO CHARGED LEPTONS

These quantities are the effective axial-vector couplings of the Z to charged leptons. Their magnitude is derived from a measurement of the Z lineshape and the forward-backward lepton asymmetries as a function of energy around the Z mass. The relative sign among the vector to axial-vector couplings is obtained from a measurement of the Z asymmetry parameters, A_e and A_τ , or ν_e scattering. The fit values quoted below correspond to global nine- or five-parameter fits to lineshape, lepton forward-backward asymmetry, and A_e and A_τ measurements. See "Note on the Z boson" for details.

 g_A^e

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.5007 ± 0.0009 OUR FIT				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
-0.4977 ± 0.0045		¹³² ABE	95J SLD	$E_{cm}^{ee} = 91.31$ GeV
-0.4998 ± 0.0016	38k	¹³³ ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
-0.503 ± 0.002	45.8k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
-0.4980 ± 0.0021		¹³³ ADRIANI	93M L3	Repl. by ACCIARRI 94
-0.5029 ± 0.0018		¹³³ BUSKULIC	93J ALEP	Repl. by BUSKULIC 94

¹³² ABE 95J obtain this result combining polarized Bhabha results with the A_{LR} measurement of ABE 94C. The Bhabha results alone give $-0.4968 \pm 0.0039 \pm 0.0027$.¹³³ The τ -polarization constraint has been included. g_A^μ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.5015 ± 0.0012 OUR FIT				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$-0.4987^{+0.0030}_{-0.0026}$	34k	¹³⁴ ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
-0.501 ± 0.002	46.4k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
$-0.4968^{+0.0050}_{-0.0037}$		¹³⁴ ADRIANI	93M L3	Repl. by ACCIARRI 94
-0.5014 ± 0.0029		¹³⁴ BUSKULIC	93J ALEP	Repl. by BUSKULIC 94

¹³⁴ The τ -polarization constraint has been included.

g_A^τ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

–0.5009±0.0013 OUR FIT

••• We do not use the following data for averages, fits, limits, etc. •••

–0.5014±0.0029	25k	¹³⁵ ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
–0.502 ±0.003	45.1k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
–0.5032±0.0038	7441	¹³⁵ ADRIANI	93M L3	Repl. by ACCIARRI 94
–0.5016±0.0033		¹³⁵ BUSKULIC	93J ALEP	Repl. by BUSKULIC 94

¹³⁵ The τ -polarization constraint has been included.

 g_A^l

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

–0.5008±0.0008 OUR FIT

••• We do not use the following data for averages, fits, limits, etc. •••

–0.4999±0.0014	71k	ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV
–0.4998±0.0014	97k	¹³⁶ ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
–0.500 ±0.001	146k	AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV
–0.502 ±0.001	137k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
–0.4998±0.0016	58k	ACTON	93D OPAL	Repl. by AKERS 94
–0.4986±0.0015		¹³⁶ ADRIANI	93M L3	Repl. by ACCIARRI 94
–0.5022±0.0015		¹³⁶ BUSKULIC	93J ALEP	Repl. by BUSKULIC 94

¹³⁶ The τ -polarization constraint has been included.

Z COUPLINGS TO NEUTRAL LEPTONS

These quantities are the effective couplings of the Z to neutral leptons. $\nu_e e$ and $\nu_\mu e$ scattering results are combined with g_A^e and g_V^e measurements at the Z mass to obtain $g^{\nu e}$ and $g^{\nu\mu}$ following NOVIKOV 93C.

 $g^{\nu e}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.528±0.085	¹³⁷ VILAIN	94 CHM2	From $\nu_\mu e$ and $\nu_e e$ scattering

¹³⁷ VILAIN 94 derive this value from their value of $g^{\nu\mu}$ and their ratio $g^{\nu e}/g^{\nu\mu} = 1.05^{+0.15}_{-0.18}$.

 $g^{\nu\mu}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.502±0.017	¹³⁸ VILAIN	94 CHM2	From $\nu_\mu e$ scattering

¹³⁸ VILAIN 94 derive this value from their measurement of the couplings $g_A^{e\nu\mu} = -0.503 \pm 0.017$ and $g_V^{e\nu\mu} = -0.035 \pm 0.017$ obtained from $\nu_\mu e$ scattering. We have re-evaluated this value using the current PDG values for g_A^e and g_V^e .

Z ASYMMETRY PARAMETERS

For each fermion-antifermion pair coupling to the Z these quantities are defined as

$$A_f = \frac{2g_V^f g_A^f}{(g_V^f)^2 + (g_A^f)^2}$$

where g_V^f and g_A^f are the effective vector and axial-vector couplings. For their relation to the various lepton asymmetries see the 'Note on the Z Boson.'

 A_e

Using polarized beams, this quantity can also be measured as $(\sigma_L - \sigma_R)/(\sigma_L + \sigma_R)$, where σ_L and σ_R are the e^+e^- production cross sections for Z bosons produced with left-handed and right-handed electrons respectively.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.1519 ± 0.0034 OUR AVERAGE				
0.162 ± 0.041 ± 0.014	89838	139 ABE	97 SLD	$E_{cm}^{ee} = 91.27$ GeV
0.1543 ± 0.0039	93644	140 ABE	97E SLD	$E_{cm}^{ee} = 91.27$ GeV
0.152 ± 0.012		141 ABE	97N SLD	$E_{cm}^{ee} = 91.27$ GeV
0.129 ± 0.014 ± 0.005	89075	142 ALEXANDER	96U OPAL	$E_{cm}^{ee} = 88-94$ GeV
0.202 ± 0.038 ± 0.008		143 ABE	95J SLD	$E_{cm}^{ee} = 91.31$ GeV
0.136 ± 0.027 ± 0.003		144 ABREU	95I DLPH	$E_{cm}^{ee} = 88-94$ GeV
0.129 ± 0.016 ± 0.005	33000	145 BUSKULIC	95Q ALEP	$E_{cm}^{ee} = 88-94$ GeV
0.157 ± 0.020 ± 0.005	86000	144 ACCIARRI	94E L3	$E_{cm}^{ee} = 88-94$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.122 ± 0.030 ± 0.012	30663	144 AKERS	95 OPAL	Repl. by ALEXANDER 96U
0.1656 ± 0.0071 ± 0.0028	49392	146 ABE	94C SLD	Repl. by ABE 97E
0.097 ± 0.044 ± 0.004	10224	147 ABE	93 SLD	Repl. by ABE 97E
0.120 ± 0.026		144 BUSKULIC	93P ALEP	Repl. by BUSKULIC 95Q

139 ABE 97 obtain this result from a measurement of the observed left-right charge asymmetry, $A_Q^{obs} = 0.225 \pm 0.056 \pm 0.019$, in hadronic Z decays. If they combine this value of A_Q^{obs} with their earlier measurement of A_{LR}^{obs} they determine A_e to be $0.1574 \pm 0.0197 \pm 0.0067$ independent of the beam polarization.

140 ABE 97E measure the left-right asymmetry in hadronic Z production. This value (statistical and systematic errors added in quadrature) leads to $\sin^2\theta_W^{eff} = 0.23060 \pm 0.00050$.

141 ABE 97N obtain this direct measurement using the left-right cross section asymmetry and the left-right forward-backward asymmetry in leptonic decays of the Z boson obtained with a polarized electron beam.

142 ALEXANDER 96U measure the τ -lepton polarization and the forward-backward polarization asymmetry.

143 ABE 95J obtain this result from polarized Bhabha scattering.

144 Derived from the measurement of forward-backward τ polarization asymmetry.

145 BUSKULIC 95Q obtain this result fitting the τ polarization as a function of the polar τ production angle.

146 ABE 94C measured the left-right asymmetry in Z production. This value leads to $\sin^2\theta_W = 0.2292 \pm 0.0009 \pm 0.0004$.

147 ABE 93 measured the left-right asymmetry in Z production.

A_μ

This quantity is directly extracted from a measurement of the left-right forward-backward asymmetry in $\mu^+\mu^-$ production at SLC using a polarized electron beam. This double asymmetry eliminates the dependence on the Z - e - e coupling parameter A_e .

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.102±0.034	3788	148 ABE	97N SLD	$E_{cm}^{ee} = 91.27$ GeV

148 ABE 97N obtain this direct measurement using the left-right cross section asymmetry and the left-right forward-backward asymmetry in $\mu^+\mu^-$ decays of the Z boson obtained with a polarized electron beam.

 A_τ

The LEP Collaborations derive this quantity from the measurement of the average τ polarization in $Z \rightarrow \tau^+\tau^-$. The SLD Collaboration directly extracts this quantity from its measured left-right forward-backward asymmetry in $Z \rightarrow \tau^+\tau^-$ produced using a polarized e^- beam. This double asymmetry eliminates the dependence on the Z - e - e coupling parameter A_e .

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.143±0.008 OUR AVERAGE				

0.195±0.034		149 ABE	97N SLD	$E_{cm}^{ee} = 91.27$ GeV
-------------	--	---------	---------	---------------------------

0.134±0.009±0.010	89075	150 ALEXANDER	96U OPAL	$E_{cm}^{ee} = 88-94$ GeV
-------------------	-------	---------------	----------	---------------------------

0.148±0.017±0.014		ABREU	95I DLPH	$E_{cm}^{ee} = 88-94$ GeV
-------------------	--	-------	----------	---------------------------

0.136±0.012±0.009	33000	151 BUSKULIC	95Q ALEP	$E_{cm}^{ee} = 88-94$ GeV
-------------------	-------	--------------	----------	---------------------------

0.150±0.013±0.009	86000	ACCIARRI	94E L3	$E_{cm}^{ee} = 88-94$ GeV
-------------------	-------	----------	--------	---------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.153±0.019±0.013	30663	AKERS	95 OPAL	Repl. by ALEXANDER 96U
-------------------	-------	-------	---------	------------------------

0.132±0.033	10732	ADRIANI	93M L3	Repl. by ACCIARRI 94E
-------------	-------	---------	--------	-----------------------

0.143±0.023		BUSKULIC	93P ALEP	Repl. by BUSKULIC 95Q
-------------	--	----------	----------	-----------------------

0.24 ±0.07	2021	ABREU	92N DLPH	Repl. by ABREU 95I
------------	------	-------	----------	--------------------

149 ABE 97N obtain this direct measurement using the left-right cross section asymmetry and the left-right forward-backward asymmetry in $\tau^+\tau^-$ decays of the Z boson obtained with a polarized electron beam.

150 ALEXANDER 96U measure the τ -lepton polarization and the forward-backward polarization asymmetry.

151 BUSKULIC 95Q obtain this result fitting the τ polarization as a function of the polar τ production angle.

 A_c

This quantity is directly extracted from a measurement of the left-right forward-backward asymmetry in $c\bar{c}$ production at SLC using polarized electron beam. This double asymmetry eliminates the dependence on the Z - e - e coupling parameter A_e .

VALUE	DOCUMENT ID	TECN	COMMENT
0.59±0.19 OUR AVERAGE			

0.37±0.23±0.21	152 ABE	95L SLD	$E_{cm}^{ee} = 91.26$ GeV
----------------	---------	---------	---------------------------

0.73±0.22±0.10	153 ABE,K	95 SLD	$E_{cm}^{ee} = 91.26$ GeV
----------------	-----------	--------	---------------------------

152 ABE 95L tag b and c quarks through their semileptonic decays into electrons and muons. A maximum likelihood fit is performed to extract A_b and A_c .

153 ABE,K 95 tag $Z \rightarrow c\bar{c}$ events using D^{*+} and D^+ meson production. To take care of the $b\bar{b}$ contamination in their analysis they use $A_b^D = 0.64 \pm 0.11$ (which is A_b from

D^*/D tagging). This is obtained by starting with a Standard Model value of 0.935, assigning it an estimated error of ± 0.105 to cover LEP and SLD measurements, and finally taking into account $B\text{-}\bar{B}$ mixing ($1-2\chi_{\text{mix}} = 0.72 \pm 0.09$). Combining with ABE 95L they quote 0.59 ± 0.19 .

A_b

This quantity is directly extracted from a measurement of the left-right forward-backward asymmetry in $b\bar{b}$ production at SLC using polarized electron beam. This double asymmetry eliminates the dependence on the $Z\text{-}e\text{-}e$ coupling parameter A_e .

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.89\pm0.11 OUR AVERAGE				
0.87 \pm 0.11 \pm 0.09	4032	154 ABE	95K SLD	$E_{\text{cm}}^{ee} = 91.26$ GeV
0.91 \pm 0.14 \pm 0.07		155 ABE	95L SLD	$E_{\text{cm}}^{ee} = 91.26$ GeV

- 154 ABE 95K obtain an enriched sample of $b\bar{b}$ events tagging with the impact parameter. A momentum-weighted charge sum is used to identify the charge of the underlying b quark.
- 155 ABE 95L tag b and c quarks through their semileptonic decays into electrons and muons. A maximum likelihood fit is performed to extract A_b and A_c . Combining with ABE 95K, they quote $0.89 \pm 0.09 \pm 0.06$.

TRANSVERSE SPIN CORRELATIONS IN $Z \rightarrow \tau^+\tau^-$

The correlations between the transverse spin components of $\tau^+\tau^-$ produced in Z decays may be expressed in terms of the vector and axial-vector couplings:

$$C_{TT} = \frac{|g_A^\tau|^2 - |g_V^\tau|^2}{|g_A^\tau|^2 + |g_V^\tau|^2}$$

$$C_{TN} = -2 \frac{|g_A^\tau| |g_V^\tau|}{|g_A^\tau|^2 + |g_V^\tau|^2} \sin(\Phi_{g_V^\tau} - \Phi_{g_A^\tau})$$

C_{TT} refers to the transverse-transverse (within the collision plane) spin correlation and C_{TN} refers to the transverse-normal (to the collision plane) spin correlation.

The longitudinal τ polarization $P_\tau (= -A_\tau)$ is given by:

$$P_\tau = -2 \frac{|g_A^\tau| |g_V^\tau|}{|g_A^\tau|^2 + |g_V^\tau|^2} \cos(\Phi_{g_V^\tau} - \Phi_{g_A^\tau})$$

Here Φ is the phase and the phase difference $\Phi_{g_V^\tau} - \Phi_{g_A^\tau}$ can be obtained using both the measurements of C_{TN} and P_τ .

C_{TT}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.01\pm0.12 OUR AVERAGE				
0.87 \pm 0.20 $^{+0.10}_{-0.12}$	9.1K	ABREU	97G DLPH	$E_{\text{cm}}^{ee} = 91.2$ GeV
1.06 \pm 0.13 \pm 0.05	120K	BARATE	97D ALEP	$E_{\text{cm}}^{ee} = 91.2$ GeV

C_{TN}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.08 \pm 0.13 \pm 0.04$	120K	¹⁵⁶ BARATE	97D ALEP	$E_{cm}^{ee} = 91.2$ GeV
¹⁵⁶ BARATE 97D combine their value of C_{TN} with the world average $P_\tau = -0.140 \pm 0.007$ to obtain $\tan(\Phi_{g_V^T} - \Phi_{g_A^T}) = -0.57 \pm 0.97$.				

 $A_{FB}^{(0,e)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow e^+e^-$

For the Z peak, we report the pole asymmetry defined by $(3/4)A_e^2$ as determined by the nine-parameter fit to cross-section and lepton forward-backward asymmetry data. For details see the "Note on the Z boson."

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
1.51 ± 0.40 OUR FIT				
1.5 ± 0.4 OUR AVERAGE				
2.5 ± 0.9		91.2	ABREU	94 DLPH
1.04 ± 0.92		91.2	ACCIARRI	94 L3
0.62 ± 0.80		91.2	AKERS	94 OPAL
1.85 ± 0.66		91.2	BUSKULIC	94 ALEP

 $A_{FB}^{(0,\mu)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow \mu^+\mu^-$

For the Z peak, we report the pole asymmetry defined by $(3/4)A_e A_\mu$ as determined by the nine-parameter fit to cross-section and lepton forward-backward asymmetry data. For details see the "Note on the Z boson."

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
1.33 ± 0.26 OUR FIT				
1.34 ± 0.24 OUR AVERAGE				
1.4 ± 0.5		91.2	ABREU	94 DLPH
1.79 ± 0.61		91.2	ACCIARRI	94 L3
0.99 ± 0.42		91.2	AKERS	94 OPAL
1.46 ± 0.48		91.2	BUSKULIC	94 ALEP

• • • We do not use the following data for averages, fits, limits, etc. • • •

9 ± 30	-2	20	¹⁵⁷ ABREU	95M DLPH
7 ± 26	-10	40	¹⁵⁷ ABREU	95M DLPH
-11 ± 33	-25	57	¹⁵⁷ ABREU	95M DLPH
-62 ± 17	-45	69	¹⁵⁷ ABREU	95M DLPH
-56 ± 10	-58	79	¹⁵⁷ ABREU	95M DLPH
-13 ± 5	-23	87.5	¹⁵⁷ ABREU	95M DLPH
-29.0 ± 5.0 - 4.8 ± 0.5	-32.1	56.9	¹⁵⁸ ABE	90I VNS
- 9.9 $\pm 1.5 \pm 0.5$	-9.2	35	HEGNER	90 JADE
0.05 ± 0.22	0.026	91.14	¹⁵⁹ ABRAMS	89D MRK2
-43.4 ± 17.0	-24.9	52.0	¹⁶⁰ BACALA	89 AMY
-11.0 ± 16.5	-29.4	55.0	¹⁶⁰ BACALA	89 AMY
-30.0 ± 12.4	-31.2	56.0	¹⁶⁰ BACALA	89 AMY
-46.2 ± 14.9	-33.0	57.0	¹⁶⁰ BACALA	89 AMY
-29 ± 13	-25.9	53.3	ADACHI	88C TOPZ

+ 5.3 ± 5.0 ±0.5	-1.2	14.0	ADEVA	88	MRKJ
-10.4 ± 1.3 ±0.5	-8.6	34.8	ADEVA	88	MRKJ
-12.3 ± 5.3 ±0.5	-10.7	38.3	ADEVA	88	MRKJ
-15.6 ± 3.0 ±0.5	-14.9	43.8	ADEVA	88	MRKJ
- 1.0 ± 6.0	-1.2	13.9	BRAUNSCH...	88D	TASS
- 9.1 ± 2.3 ±0.5	-8.6	34.5	BRAUNSCH...	88D	TASS
-10.6 ⁺ 2.2 ₋ 2.3 ±0.5	-8.9	35.0	BRAUNSCH...	88D	TASS
-17.6 ⁺ 4.4 ₋ 4.3 ±0.5	-15.2	43.6	BRAUNSCH...	88D	TASS
- 4.8 ± 6.5 ±1.0	-11.5	39	BEHREND	87C	CELL
-18.8 ± 4.5 ±1.0	-15.5	44	BEHREND	87C	CELL
+ 2.7 ± 4.9	-1.2	13.9	BARTEL	86C	JADE
-11.1 ± 1.8 ±1.0	-8.6	34.4	BARTEL	86C	JADE
-17.3 ± 4.8 ±1.0	-13.7	41.5	BARTEL	86C	JADE
-22.8 ± 5.1 ±1.0	-16.6	44.8	BARTEL	86C	JADE
- 6.3 ± 0.8 ±0.2	-6.3	29	ASH	85	MAC
- 4.9 ± 1.5 ±0.5	-5.9	29	DERRICK	85	HRS
- 7.1 ± 1.7	-5.7	29	LEVI	83	MRK2
-16.1 ± 3.2	-9.2	34.2	BRANDELIK	82C	TASS

¹⁵⁷ ABREU 95M perform this measurement using radiative muon-pair events associated with high-energy isolated photons.

¹⁵⁸ ABE 90I measurements in the range $50 \leq \sqrt{s} \leq 60.8$ GeV.

¹⁵⁹ ABRAMS 89D asymmetry includes both 9 $\mu^+\mu^-$ and 15 $\tau^+\tau^-$ events.

¹⁶⁰ BACALA 89 systematic error is about 5%.

$A_{FB}^{(0,\tau)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow \tau^+\tau^-$

For the Z peak, we report the pole asymmetry defined by $(3/4)A_e A_\tau$ as determined by the nine-parameter fit to cross-section and lepton forward-backward asymmetry data. For details see the "Note on the Z boson."

<u>ASYMMETRY (%)</u>	<u>STD. MODEL</u>	<u>\sqrt{s} (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
2.12 ± 0.32 OUR FIT				
2.13 ± 0.31 OUR AVERAGE				
2.2 ± 0.7		91.2	ABREU	94 DLPH
2.65 ± 0.88		91.2	ACCIARRI	94 L3
2.05 ± 0.52		91.2	AKERS	94 OPAL
1.97 ± 0.56		91.2	BUSKULIC	94 ALEP

• • • We do not use the following data for averages, fits, limits, etc. • • •

-32.8 ⁺ 6.4 ₋ 6.2 ±1.5	-32.1	56.9	¹⁶¹ ABE	90I	VNS
- 8.1 ± 2.0 ±0.6	-9.2	35	HEGNER	90	JADE
-18.4 ±19.2	-24.9	52.0	¹⁶² BACALA	89	AMY
-17.7 ±26.1	-29.4	55.0	¹⁶² BACALA	89	AMY
-45.9 ±16.6	-31.2	56.0	¹⁶² BACALA	89	AMY
-49.5 ±18.0	-33.0	57.0	¹⁶² BACALA	89	AMY
-20 ±14	-25.9	53.3	ADACHI	88C	TOPZ
-10.6 ± 3.1 ±1.5	-8.5	34.7	ADEVA	88	MRKJ

$-8.5 \pm 6.6 \pm 1.5$	-15.4	43.8	ADEVA	88	MRKJ
$-6.0 \pm 2.5 \pm 1.0$	8.8	34.6	BARTEL	85F	JADE
$-11.8 \pm 4.6 \pm 1.0$	14.8	43.0	BARTEL	85F	JADE
$-5.5 \pm 1.2 \pm 0.5$	-0.063	29.0	FERNANDEZ	85	MAC
-4.2 ± 2.0	0.057	29	LEVI	83	MRK2
-10.3 ± 5.2	-9.2	34.2	BEHREND	82	CELL
-0.4 ± 6.6	-9.1	34.2	BRANDELIK	82C	TASS

¹⁶¹ ABE 90I measurements in the range $50 \leq \sqrt{s} \leq 60.8$ GeV.

¹⁶² BACALA 89 systematic error is about 5%.

$A_{FB}^{(0,\ell)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow \ell^+\ell^-$

For the Z peak, we report the pole asymmetry defined by $(3/4)A_\ell^2$ as determined by the five-parameter fit to cross-section and lepton forward-backward asymmetry data assuming lepton universality. For details see the "Note on the Z boson."

<u>ASYMMETRY (%)</u>	<u>STD. MODEL</u>	<u>\sqrt{s} (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
1.59±0.18 OUR FIT				
1.60±0.18 OUR AVERAGE				
1.77±0.37		91.2	ABREU 94	DLPH
1.84±0.45		91.2	ACCIARRI 94	L3
1.28±0.30		91.2	AKERS 94	OPAL
1.71±0.33		91.2	BUSKULIC 94	ALEP

$A_{FB}^{(0,u)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow u\bar{u}$

<u>ASYMMETRY (%)</u>	<u>STD. MODEL</u>	<u>\sqrt{s} (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
4.0±7.3 OUR EVALUATION				
4.0 ± 6.7 ± 2.8	6	91.2	¹⁶³ ACKERSTAFF 97T	OPAL

¹⁶³ ACKERSTAFF 97T measure the forward-backward asymmetry of various fast hadrons made of light quarks. Then using SU(2) isospin symmetry and flavor independence for down and strange quarks authors solve for the different quark types.

$A_{FB}^{(0,s)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow s\bar{s}$

The s -quark asymmetry is derived from measurements of the forward-backward asymmetry of fast hadrons containing an s quark.

<u>ASYMMETRY (%)</u>	<u>STD. MODEL</u>	<u>\sqrt{s} (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
9.9±3.1 OUR AVERAGE	Error includes scale factor of 1.2.			
6.8±3.5±1.1	10	91.2	¹⁶⁴ ACKERSTAFF 97T	OPAL
13.1±3.5±1.3		91.2	¹⁶⁵ ABREU 95G	DLPH

¹⁶⁴ ACKERSTAFF 97T measure the forward-backward asymmetry of various fast hadrons made of light quarks. Then using SU(2) isospin symmetry and flavor independence for down and strange quarks authors solve for the different quark types. The value reported here corresponds then to the forward-backward asymmetry for "down-type" quarks.

¹⁶⁵ ABREU 95G require the presence of a high-momentum charged kaon or Λ^0 to tag the s quark. An unresolved s - and d -quark asymmetry of $(11.2 \pm 3.1 \pm 5.4)\%$ is obtained by tagging the presence of a high-energy neutron or neutral kaon in the hadron calorimeter.

$A_{FB}^{(0,c)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow c\bar{c}$

OUR FIT, which is obtained by a simultaneous fit to several c - and b -quark measurements as explained in the "Note on the Z boson," refers to the Z pole asymmetry. As a cross check we have also performed a weighted average of the "near peak" measurements taking into account the various common systematic errors. We have assumed that the smallest common systematic error is fully correlated. Applying to this combined "peak" measurement QCD, QED, and energy-dependence corrections, our weighted average gives a pole asymmetry of $(7.20 \pm 0.64)\%$.

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
7.32 ± 0.58 OUR FIT				
6.3 ± 1.2 ± 0.6		91.22	166 ALEXANDER	97C OPAL
6.00 ± 0.67 ± 0.52		91.24	167 ALEXANDER	96 OPAL
7.7 ± 2.9 ± 1.2		91.27	168 ABREU	95E DLPH
8.3 ± 2.2 ± 1.6		91.27	169 ABREU	95K DLPH
6.99 ± 2.05 ± 1.02		91.24	170 BUSKULIC	95I ALEP
9.9 ± 2.0 ± 1.7		91.24	171 BUSKULIC	94G ALEP
8.3 ± 3.8 ± 2.7	5.6	91.24	172 ADRIANI	92D L3
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3.9 ± 5.1 ± 0.9		89.45	166 ALEXANDER	97C OPAL
15.8 ± 4.1 ± 1.1		93.00	166 ALEXANDER	97C OPAL
− 7.5 ± 3.4 ± 0.6	− 3.5	89.52	167 ALEXANDER	96 OPAL
14.1 ± 2.8 ± 0.9	12.0	92.94	167 ALEXANDER	96 OPAL
6.8 ± 4.2 ± 0.9		91.25	173 BUSKULIC	94J ALEP
1.4 ± 3.0 ± 2.0	5.6	91.24	174 ACTON	93K OPAL
3.8 ± 4.4 ± 1.0	5.4	91.28	175 AKERS	93D OPAL
− 12.9 ± 7.8 ± 5.5	− 13.6	35	BEHREND	90D CELL
7.7 ± 13.4 ± 5.0	− 22.1	43	BEHREND	90D CELL
− 12.8 ± 4.4 ± 4.1	− 13.6	35	ELSEN	90 JADE
− 10.9 ± 12.9 ± 4.6	− 23.2	44	ELSEN	90 JADE
− 14.9 ± 6.7	− 13.3	35	OULD-SAADA	89 JADE

¹⁶⁶ ALEXANDER 97C identify the b and c events using a D/D^* tag.

¹⁶⁷ ALEXANDER 96 tag heavy flavors using one or two identified leptons. This allows the simultaneous fitting of the b and c quark forward-backward asymmetries as well as the average $B^0-\bar{B}^0$ mixing.

¹⁶⁸ ABREU 95E require the presence of a $D^{*\pm}$ to identify c and b quarks.

¹⁶⁹ ABREU 95K identify c and b quarks using both electron and muon semileptonic decays.

¹⁷⁰ BUSKULIC 95I require the presence of a high momentum $D^{*\pm}$ to have an enriched sample of $Z \rightarrow c\bar{c}$ events.

¹⁷¹ BUSKULIC 94G perform a simultaneous fit to the p and p_T spectra of both single and dilepton events.

¹⁷² ADRIANI 92D use both electron and muon semileptonic decays.

¹⁷³ BUSKULIC 94J Identify the b and c decays using D^* . Replaced by BUSKULIC 95I.

¹⁷⁴ ACTON 93K use the lepton tagging technique. Replaced by ALEXANDER 96.

¹⁷⁵ AKERS 93D identify the b and c decays using D^* . Replaced by ALEXANDER 97C.

$A_{FB}^{(0,b)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow b\bar{b}$

OUR FIT, which is obtained by a simultaneous fit to several c - and b -quark measurements as explained in the "Note on the Z boson," refers to the Z pole asymmetry. As a cross check we have also performed a weighted average of the "near peak" measurements taking into account the various common systematic errors. We have assumed that the smallest common systematic error is fully correlated. Applying to this combined "peak" measurement QCD, QED, and energy-dependence corrections, our weighted average gives a pole asymmetry of $(10.07 \pm 0.32)\%$. For the jet-charge measurements (where the QCD corrections are already included since they represent an inherent part of the analysis), we subtract the QCD correction before combining.

<u>ASYMMETRY (%)</u>	<u>STD. MODEL</u>	<u>\sqrt{s} (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
10.02 ± 0.28 OUR FIT				
9.94 ± 0.52 ± 0.44		91.21	176 ACKERSTAFF	97P OPAL
9.4 ± 2.7 ± 2.2		91.22	177 ALEXANDER	97C OPAL
9.06 ± 0.51 ± 0.23		91.24	178 ALEXANDER	96 OPAL
9.65 ± 0.44 ± 0.26		91.21	179 BUSKULIC	96Q ALEP
5.9 ± 6.2 ± 2.4		91.27	180 ABREU	95E DLPH
10.4 ± 1.3 ± 0.5		91.27	181 ABREU	95K DLPH
11.5 ± 1.7 ± 1.0		91.27	182 ABREU	95K DLPH
8.7 ± 1.1 ± 0.4		91.3	183 ACCIARRI	94D L3
9.92 ± 0.84 ± 0.46		91.19	184 BUSKULIC	94I ALEP
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.1 ± 2.1 ± 0.2		89.44	176 ACKERSTAFF	97P OPAL
14.5 ± 1.7 ± 0.7		92.91	176 ACKERSTAFF	97P OPAL
– 8.6 ± 10.8 ± 2.9		89.45	177 ALEXANDER	97C OPAL
– 2.1 ± 9.0 ± 2.6		93.00	177 ALEXANDER	97C OPAL
5.5 ± 2.4 ± 0.3	5.5	89.52	178 ALEXANDER	96 OPAL
11.7 ± 2.0 ± 0.3	11.4	92.94	178 ALEXANDER	96 OPAL
– 3.4 ± 11.2 ± 0.7		88.38	179 BUSKULIC	96Q ALEP
5.3 ± 2.0 ± 0.2		89.38	179 BUSKULIC	96Q ALEP
8.9 ± 5.9 ± 0.4		90.21	179 BUSKULIC	96Q ALEP
3.8 ± 5.1 ± 0.2		92.05	179 BUSKULIC	96Q ALEP
10.3 ± 1.6 ± 0.4		92.94	179 BUSKULIC	96Q ALEP
8.8 ± 7.5 ± 0.5		93.90	179 BUSKULIC	96Q ALEP
6.2 ± 3.4 ± 0.2		89.52	185 AKERS	95S OPAL
9.63 ± 0.67 ± 0.38		91.25	185 AKERS	95S OPAL
17.2 ± 2.8 ± 0.7		92.94	185 AKERS	95S OPAL
8.7 ± 1.4 ± 0.2		91.24	186 BUSKULIC	94G ALEP
7.1 ± 5.4 ± 0.7	5.2	89.66	187 ACTON	93K OPAL
9.2 ± 1.8 ± 0.8	8.5	91.24	187 ACTON	93K OPAL

13.1 ± 4.7 ± 1.3	10.8	92.75	187	ACTON	93K	OPAL
13.9 ± 9.7 ± 4.9	9.4	91.28	188	AKERS	93D	OPAL
16.1 ± 6.0 ± 2.1		91.2	189	ABREU	92H	DLPH
8.6 ± 1.5 ± 0.7	8.2	91.24	190	ADRIANI	92D	L3
2.5 ± 5.1 ± 0.7	5.3	89.67	191	ADRIANI	92D	L3
9.7 ± 1.7 ± 0.7	8.2	91.24	191	ADRIANI	92D	L3
6.2 ± 4.2 ± 0.7	10.8	92.81	191	ADRIANI	92D	L3
-71 ± 34 ± 7 - 8	-58	58.3		SHIMONAKA	91	TOPZ
-22.2 ± 7.7 ± 3.5	-26.0	35		BEHREND	90D	CELL
-49.1 ± 16.0 ± 5.0	-39.7	43		BEHREND	90D	CELL
-28 ± 11	-23	35		BRAUNSCH...	90	TASS
-16.6 ± 7.7 ± 4.8	-24.3	35		ELSEN	90	JADE
-33.6 ± 22.2 ± 5.2	-39.9	44		ELSEN	90	JADE
3.4 ± 7.0 ± 3.5	-16.0	29.0		BAND	89	MAC
-72 ± 28 ± 13	-56	55.2		SAGAWA	89	AMY

176 ACKERSTAFF 97P tag b quarks using lifetime. The quark charge is measured using both jet charge and vertex charge, a weighted sum of the charges of tracks in a jet which contains a tagged secondary vertex.

177 ALEXANDER 97C identify the b and c events using a D/D^* tag.

178 ALEXANDER 96 tag heavy flavors using one or two identified leptons. This allows the simultaneous fitting of the b and c quark forward-backward asymmetries as well as the average $B^0-\bar{B}^0$ mixing.

179 BUSKULIC 96Q tag b -quark flavor and charge using high transverse momentum leptons. The asymmetry value at the Z peak is obtained using a charm charge asymmetry of 6.17%.

180 ABREU 95E require the presence of a $D^{*\pm}$ to identify c and b quarks.

181 ABREU 95K identify c and b quarks using both electron and muon semileptonic decays. The systematic error includes an uncertainty of ± 0.3 due to the mixing correction ($\chi = 0.115 \pm 0.011$).

182 ABREU 95K tag b quarks using lifetime; the quark charge is identified using jet charge. The systematic error includes an uncertainty of ± 0.3 due to the mixing correction ($\chi = 0.115 \pm 0.011$).

183 ACCIARRI 94D use both electron and muon semileptonic decays.

184 BUSKULIC 94I use the lifetime tag method to obtain a high purity sample of $Z \rightarrow b\bar{b}$ events and the hemisphere charge technique to obtain the jet charge.

185 AKERS 95S tag b quarks using lifetime; the quark charge is measured using jet charge. These asymmetry values are obtained using $R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons}) = 0.216$. For a value of R_b different from this by an amount ΔR_b , the change in the asymmetry values is given by $-K\Delta R_b$, where $K = 0.082, 0.471, \text{ and } 0.855$ for \sqrt{s} values of 89.52, 91.25, and 92.94 GeV respectively. Replaced by ACKERSTAFF 97P.

186 BUSKULIC 94G perform a simultaneous fit to the p and p_T spectra of both single and dilepton events. Replaced by BUSKULIC 96Q.

187 ACTON 93K use the lepton tagging technique. The systematic error includes the uncertainty on the mixing parameter. Replaced by ALEXANDER 96.

188 AKERS 93D identify the b and c decays using D^* . Replaced by ALEXANDER 97C.

189 B tagging via its semimuonic decay. Experimental value corrected using average LEP $B^0-\bar{B}^0$ mixing parameter $\chi = 0.143 \pm 0.023$.

190 ADRIANI 92D use both electron and muon semileptonic decays. For this measurement ADRIANI 92D average over all \sqrt{s} values to obtain a single result.

191 ADRIANI 92D use both electron and muon semileptonic decays. The quoted systematic error is common to all measurements. The peak value is superseded by ACCIARRI 94D.

CHARGE ASYMMETRY IN $e^+e^- \rightarrow q\bar{q}$

Summed over five lighter flavors.

Experimental and Standard Model values are somewhat event-selection dependent. Standard Model expectations contain some assumptions on $B^0-\bar{B}^0$ mixing and on other electroweak parameters.

<u>ASYMMETRY (%)</u>	<u>STD. MODEL</u>	<u>\sqrt{s} (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
– 0.76 ± 0.12 ± 0.15		91.2	¹⁹² ABREU	92I DLPH
4.0 ± 0.4 ± 0.63	4.0	91.3	¹⁹³ ACTON	92L OPAL
9.1 ± 1.4 ± 1.6	9.0	57.9	ADACHI	91 TOPZ
– 0.84 ± 0.15 ± 0.04		91	DECAMP	91B ALEP
8.3 ± 2.9 ± 1.9	8.7	56.6	STUART	90 AMY
11.4 ± 2.2 ± 2.1	8.7	57.6	ABE	89L VNS
6.0 ± 1.3	5.0	34.8	GREENSHAW	89 JADE
8.2 ± 2.9	8.5	43.6	GREENSHAW	89 JADE

¹⁹² ABREU 92I has 0.14 systematic error due to uncertainty of quark fragmentation.

¹⁹³ ACTON 92L use the weight function method on 259k selected $Z \rightarrow$ hadrons events.

The systematic error includes a contribution of 0.2 due to $B^0-\bar{B}^0$ mixing effect, 0.4 due to Monte Carlo (MC) fragmentation uncertainties and 0.3 due to MC statistics.

ACTON 92L derive a value of $\sin^2\theta_W^{\text{eff}}$ to be $0.2321 \pm 0.0017 \pm 0.0028$.

CHARGE ASYMMETRY IN $p\bar{p} \rightarrow Z \rightarrow e^+e^-$

<u>ASYMMETRY (%)</u>	<u>STD. MODEL</u>	<u>\sqrt{s} (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5.2 ± 5.9 ± 0.4		91	ABE	91E CDF

Z REFERENCES

ABE	98D PRL 80 660	K. Abe+	(SLD Collab.)
ACKERSTAFF	98E EPJ C1 439	K. Ackerstaff+	(OPAL Collab.)
ABE	97 PRL 78 17	+Abe, Abt, Akagi, Allen+	(SLD Collab.)
ABE	97E PRL 78 2075	+Abe, Abt, Akagi, Allen+	(SLD Collab.)
ABE	97N PRL 79 804	K. Abe+	(SLD Collab.)
ABREU	97C ZPHY C73 243	+Adam, Adye, Ajinenko+	(DELPHI Collab.)
ABREU	97G PL B404 194	P. Abreu+	(DELPHI Collab.)
ACCIARRI	97D PL B393 465	+Adriani, Aguilar-Benitez, Ahlen+	(L3 Collab.)
ACCIARRI	97J PL B407 351	M. Acciarri+	(L3 Collab.)
ACCIARRI	97K PL B407 361	M. Acciarri+	(L3 Collab.)
ACCIARRI	97L PL B407 389	M. Acciarri+	(L3 Collab.)
ACCIARRI	97R PL B413 167	M. Acciarri+	(L3 Collab.)
ACKERSTAFF	97C PL B391 221	+Alexander, Allison, Altekamp, Ametewee+	(OPAL Collab.)
ACKERSTAFF	97K ZPHY C74 1	K. Ackerstaff+	(OPAL Collab.)
ACKERSTAFF	97M ZPHY C74 413	K. Ackerstaff+	(OPAL Collab.)
ACKERSTAFF	97P ZPHY C75 385	K. Ackerstaff+	(OPAL Collab.)
ACKERSTAFF	97S PL B412 210	K. Ackerstaff+	(OPAL Collab.)
ACKERSTAFF	97T ZPHY C76 387	K. Ackerstaff+	(OPAL Collab.)
ACKERSTAFF	97W ZPHY C76 425	K. Ackerstaff+	(OPAL Collab.)
ALEXANDER	97C ZPHY C73 379	+Allison, Altekamp, Ametewee+	(OPAL Collab.)
ALEXANDER	97D ZPHY C73 569	+Allison, Altekamp, Ametewee+	(OPAL Collab.)
ALEXANDER	97E ZPHY C73 587	+Allison, Altekamp, Ametewee+	(OPAL Collab.)
BARATE	97D PL B405 191	R. Barate+	(ALEPH Collab.)
BARATE	97E PL B401 150	R. Barate+	(ALEPH Collab.)

BARATE	97F	PL B401 163	R. Barate+	(ALEPH Collab.)
BARATE	97J	ZPHY C74 451	R. Barate+	(ALEPH Collab.)
ABE	96E	PR D53 1023	+Abt, Ahn, Akagi, Allen+	(SLD Collab.)
ABREU	96	ZPHY C70 531	+Adam, Adye+	(DELPHI Collab.)
ABREU	96C	PL B379 309	+Adam, Adye+	(DELPHI Collab.)
ABREU	96R	ZPHY C72 31	+Adam, Adye+	(DELPHI Collab.)
ABREU	96S	PL B389 405	+Adam, Adye, Ajinenko+	(DELPHI Collab.)
ABREU	96U	ZPHY C73 61	+Adam, Adye, Agasi, Ajinenko+	(DELPHI Collab.)
ACCIARRI	96	PL B371 126	+Adam, Adriani+	(L3 Collab.)
ACCIARRI	96B	PL B370 195	+Adam, Adriani+	(L3 Collab.)
ADAM	96	ZPHY C69 561	+Adye, Agasi, Ajinenko+	(DELPHI Collab.)
ADAM	96B	ZPHY C70 371	+Adye, Agasi+	(DELPHI Collab.)
ALEXANDER	96	ZPHY C70 357	+Allison, Altekamp+	(OPAL Collab.)
ALEXANDER	96B	ZPHY C70 197	+Allison, Altekamp+	(OPAL Collab.)
ALEXANDER	96F	PL B370 185	+Allison, Altekamp+	(OPAL Collab.)
ALEXANDER	96N	PL B384 343	+Allison, Altekamp, Ametewee+	(OPAL Collab.)
ALEXANDER	96R	ZPHY C72 1	+Allison, Altekamp+	(OPAL Collab.)
ALEXANDER	96U	ZPHY C72 365	+Allison, Altekamp, Ametewee+	(OPAL Collab.)
ALEXANDER	96X	PL B376 232	G. Alexander+	(OPAL Collab.)
BUSKULIC	96D	ZPHY C69 393	+Casper, De Bonis, Decamp+	(ALEPH Collab.)
BUSKULIC	96H	ZPHY C69 379	+Casper, De Bonis+	(ALEPH Collab.)
BUSKULIC	96Q	PL B384 414	+De Bonis, Decamp, Ghez+	(ALEPH Collab.)
ABE	95J	PRL 74 2880	+Abt, Ahn, Akagi+	(SLD Collab.)
ABE	95K	PRL 74 2890	+Abt, Ahn, Akagi+	(SLD Collab.)
ABE	95L	PRL 74 2895	+Abt, Ahn, Akagi+	(SLD Collab.)
ABE,K	95	PRL 75 3609	K. Abe, Abt, Ahn, Akagi+	(SLD Collab.)
ABREU	95	ZPHY C65 709	erratum +Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	95D	ZPHY C66 323	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	95E	ZPHY C66 341	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	95F	NP B444 3	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	95G	ZPHY C67 1	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	95I	ZPHY C67 183	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	95J	ZPHY C65 555	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	95K	ZPHY C65 569	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	95L	ZPHY C65 587	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	95M	ZPHY C65 603	+Adam, Adye, Agasi, Ajinenko+	(DELPHI Collab.)
ABREU	95O	ZPHY C67 543	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	95R	ZPHY C68 353	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	95W	PL B361 207	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	95X	ZPHY C69 1	+Adam, Adye, Agasi+	(DELPHI Collab.)
ACCIARRI	95B	PL B345 589	+Adam, Adriani, Aguilar-Benitez+	(L3 Collab.)
ACCIARRI	95C	PL B345 609	+Adam, Adriani, Aguilar-Benitez+	(L3 Collab.)
ACCIARRI	95G	PL B353 136	+Adam, Adriani, Aguilar-Benitez, Ahlen+	(L3 Collab.)
AKERS	95	ZPHY C65 1	+Alexander, Allison+	(OPAL Collab.)
AKERS	95B	ZPHY C65 17	+Alexander, Allison+	(OPAL Collab.)
AKERS	95C	ZPHY C65 47	+Alexander, Allison+	(OPAL Collab.)
AKERS	95O	ZPHY C67 27	+Alexander, Allison+	(OPAL Collab.)
AKERS	95S	ZPHY C67 365	+Alexander, Allison+	(OPAL Collab.)
AKERS	95U	ZPHY C67 389	+Alexander, Allison+	(OPAL Collab.)
AKERS	95W	ZPHY C67 555	+Alexander, Allison+	(OPAL Collab.)
AKERS	95X	ZPHY C68 1	+Alexander, Allison+	(OPAL Collab.)
AKERS	95Z	ZPHY C68 203	+Alexander, Allison+	(OPAL Collab.)
ALEXANDER	95D	PL B358 162	+Allison, Altekamp+	(OPAL Collab.)
BUSKULIC	95I	PL B352 479	+Casper, De Bonis+	(ALEPH Collab.)
BUSKULIC	95Q	ZPHY C69 183	+Casper, De Bonis+	(ALEPH Collab.)
BUSKULIC	95R	ZPHY C69 15	+Casper, De Bonis, Decamp+	(ALEPH Collab.)
MIYABAYASHI	95	PL B347 171	+Adachi, Fujii+	(TOPAZ Collab.)
ABE	94C	PRL 73 25	+Abt, Ash, Aston, Bacchetta, Baird+	(SLD Collab.)
ABREU	94	NP B418 403	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	94B	PL B327 386	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	94P	PL B341 109	+Adam, Adye, Agasi, Ajinenko+	(DELPHI Collab.)
ACCIARRI	94	ZPHY C62 551	+Adam, Adriani, Aguilar-Benitez+	(L3 Collab.)
ACCIARRI	94B	PL B328 223	+Adam, Adriani, Aguilar-Benitez+	(L3 Collab.)
ACCIARRI	94D	PL B335 542	+Adam, Adriani, Aguilar-Benitez, Ahlen+	(L3 Collab.)
ACCIARRI	94E	PL B341 245	+Adam, Adriani+	(L3 Collab.)
AKERS	94	ZPHY C61 19	+Alexander, Allison+	(OPAL Collab.)
AKERS	94P	ZPHY C63 181	+Alexander, Allison+	(OPAL Collab.)
BUSKULIC	94	ZPHY C62 539	+Casper, De Bonis, Decamp, Ghez, Goy+	(ALEPH Collab.)
BUSKULIC	94G	ZPHY C62 179	+Casper, De Bonis, Decamp, Ghez+	(ALEPH Collab.)
BUSKULIC	94I	PL B335 99	+Casper, De Bonis+	(ALEPH Collab.)

BUSKULIC	94J	ZPHY C62 1	+De Bonis, Decamp+	(ALEPH Collab.)
BUSKULIC	94K	ZPHY C64 361	+De Bonis, Decamp+	(ALEPH Collab.)
VILAIN	94	PL B320 203	+Wilquet, Beyer+	(CHARM II Collab.)
ABE	93	PRL 70 2515	+Abt, Acton+	(SLD Collab.)
ABREU	93	PL B298 236	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	93I	ZPHY C59 533	+Adam, Adye, Agasi+	(DELPHI Collab.)
Also	95	ZPHY C65 709 erratum	Abreu, Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	93L	PL B318 249	+Adam, Adami, Adye+	(DELPHI Collab.)
ACTON	93	PL B305 407	+Alexander, Allison+	(OPAL Collab.)
ACTON	93D	ZPHY C58 219	+Alexander, Allison+	(OPAL Collab.)
ACTON	93E	PL B311 391	+Akers, Alexander+	(OPAL Collab.)
ACTON	93F	ZPHY C58 405	+Alexander, Allison+	(OPAL Collab.)
ACTON	93K	ZPHY C60 19	+Akers, Alexander+	(OPAL Collab.)
ADRIANI	93	PL B301 136	+Aguilar-Benitez, Ahlen+	(L3 Collab.)
ADRIANI	93E	PL B307 237	+Aguilar-Benitez, Ahlen+	(L3 Collab.)
ADRIANI	93F	PL B309 451	+Aguilar-Benitez, Ahlen+	(L3 Collab.)
ADRIANI	93I	PL B316 427	+Aguilar-Benitez, Ahlen+	(L3 Collab.)
ADRIANI	93J	PL B317 467	+Aguilar-Benitez, Ahlen, Alcaraz+	(L3 Collab.)
ADRIANI	93M	PRPL 236 1	+Aguilar-Benitez, Ahlen, Alcaraz, Aloisio+	(L3 Collab.)
AKERS	93B	ZPHY C60 199	+Alexander, Allison, Anderson, Arcelli+	(OPAL Collab.)
AKERS	93D	ZPHY C60 601	+Alexander, Allison+	(OPAL Collab.)
BUSKULIC	93J	ZPHY C60 71	+Decamp, Goy, Lees, Minard+	(ALEPH Collab.)
BUSKULIC	93L	PL B313 520	+De Bonis, Decamp+	(ALEPH Collab.)
BUSKULIC	93N	PL B313 549	+De Bonis, Decamp+	(ALEPH Collab.)
BUSKULIC	93P	ZPHY C59 369	+Decamp, Goy+	(ALEPH Collab.)
NOVIKOV	93C	PL B298 453	+Okun, Vysotsky	(ITEP)
ABREU	92	ZPHY C53 567	+Adam, Adami, Adye+	(DELPHI Collab.)
ABREU	92G	PL B275 231	+Adam, Adami, Adye+	(DELPHI Collab.)
ABREU	92H	PL B276 536	+Adam, Adami, Adye+	(DELPHI Collab.)
ABREU	92I	PL B277 371	+Adam, Adami, Adye+	(DELPHI Collab.)
ABREU	92K	PL B281 383	+Adam, Adami, Adye+	(DELPHI Collab.)
ABREU	92M	PL B289 199	+Adam, Adye, Agasi, Alekseev+	(DELPHI Collab.)
ABREU	92N	ZPHY C55 555	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	92O	PL B295 383	+Adam, Adami, Adye+	(DELPHI Collab.)
ACTON	92B	ZPHY C53 539	+Alexander, Allison, Allport+	(OPAL Collab.)
ACTON	92J	PL B291 503	+Alexander, Allison, Allport+	(OPAL Collab.)
ACTON	92L	PL B294 436	+Alexander, Allison, Allport+	(OPAL Collab.)
ACTON	92N	PL B295 357	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
ACTON	92O	ZPHY C56 521	+Alexander, Allison+	(OPAL Collab.)
ADEVA	92	PL B275 209	+Adriani, Aguilar-Benitez+	(L3 Collab.)
ADRIANI	92B	PL B288 404	+Aguilar-Benitez, Ahlen, Akbari, Alcaraz+	(L3 Collab.)
ADRIANI	92D	PL B292 454	+Aguilar-Benitez, Ahlen, Akbari+	(L3 Collab.)
ADRIANI	92E	PL B292 463	+Aguilar-Benitez, Ahlen, Akbari, Alcaraz+	(L3 Collab.)
ALITTI	92B	PL B276 354	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
BUSKULIC	92D	PL B292 210	+Decamp, Goy, Lees+	(ALEPH Collab.)
BUSKULIC	92E	PL B294 145	+Decamp, Goy, Lees, Minard+	(ALEPH Collab.)
DECAMP	92	PRPL 216 253	+Deschizeaux, Goy, Lees, Minard+	(ALEPH Collab.)
DECAMP	92B	ZPHY C53 1	+Deschizeaux, Goy, Lees, Minard+	(ALEPH Collab.)
LEP	92	PL B276 247	+ALEPH, DELPHI, L3, OPAL	(LEP Collabs.)
ABE	91E	PRL 67 1502	+Amidei, Apollinari+	(CDF Collab.)
ABREU	91H	ZPHY C50 185	+Adam, Adami, Adye+	(DELPHI Collab.)
ACTON	91B	PL B273 338	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
ADACHI	91	PL B255 613	+Anazawa, Doser, Enomoto+	(TOPAZ Collab.)
ADEVA	91I	PL B259 199	+Adriani, Aguilar-Benitez, Akbari+	(L3 Collab.)
AKRAWY	91F	PL B257 531	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
DECAMP	91B	PL B259 377	+Deschizeaux, Goy+	(ALEPH Collab.)
DECAMP	91J	PL B266 218	+Deschizeaux, Goy, Lees+	(ALEPH Collab.)
DECAMP	91K	PL B273 181	+Deschizeaux, Goy, Lees, Minard+	(ALEPH Collab.)
JACOBSEN	91	PRL 67 3347	+Koetke, Adolphsen, Fujino+	(Mark II Collab.)
SHIMONAKA	91	PL B268 457	+Fujii, Miyamoto+	(TOPAZ Collab.)
ABE	90I	ZPHY C48 13	+Amako, Arai, Asano, Chiba+	(VENUS Collab.)
ABRAMS	90	PRL 64 1334	+Adolphsen, Averill, Ballam+	(Mark II Collab.)
ADACHI	90F	PL B234 525	+Doser, Enomoto, Fujii+	(TOPAZ Collab.)
AKRAWY	90J	PL B246 285	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
BEHREND	90D	ZPHY C47 333	+Criegiee, Field, Franke, Jung+	(CELLO Collab.)
BRAUNSCH...	90	ZPHY C48 433	Braunschweig, Gerhards, Kirschfink+	(TASSO Collab.)
ELSEN	90	ZPHY C46 349	+Allison, Ambrus, Barlow, Bartel+	(JADE Collab.)
HEGNER	90	ZPHY C46 547	+Naroska, Schroth, Allison+	(JADE Collab.)
KRAL	90	PRL 64 1211	+Abrams, Adolphsen, Averill, Ballam+	(Mark II Collab.)

STUART	90	PRL 64 983	+Breedon, Kim, Ko, Lander, Maeshima+	(AMY Collab.)
ABE	89	PRL 62 613	+Amidei, Apollinari, Ascori, Atac+	(CDF Collab.)
ABE	89C	PRL 63 720	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ABE	89L	PL B232 425	+Amako, Arai, Asano, Chiba+	(VENUS Collab.)
ABRAMS	89B	PRL 63 2173	+Adolphsen, Averill, Ballam, Barish+	(Mark II Collab.)
ABRAMS	89D	PRL 63 2780	+Adolphsen, Averill, Ballam, Barish+	(Mark II Collab.)
ALBAJAR	89	ZPHY C44 15	+Albrow, Allkofer, Arnison, Astbury+	(UA1 Collab.)
BACALA	89	PL B218 112	+Malchow, Sparks, Imlay, Kirk+	(AMY Collab.)
BAND	89	PL B218 369	+Camporesi, Chadwick, Delfino, Desangro+	(MAC Collab.)
GREENSHAW	89	ZPHY C42 1	+Warming, Allison, Ambrus, Barlow+	(JADE Collab.)
OULD-SAAD	89	ZPHY C44 567	+Allison, Ambrus, Barlow, Bartel+	(JADE Collab.)
SAGAWA	89	PRL 63 2341	+Lim, Abe, Fujii, Higashi+	(AMY Collab.)
ADACHI	88C	PL B208 319	+Aihara, Dijkstra, Enomoto, Fujii+	(TOPAZ Collab.)
ADEVA	88	PR D38 2665	+Anderhub, Ansari, Becker+	(Mark-J Collab.)
BRAUNSCH...	88D	ZPHY C40 163	Braunschweig, Gerhards, Kirschfink+	(TASSO Collab.)
ANSARI	87	PL B186 440	+Bagnaia, Banner, Battiston+	(UA2 Collab.)
BEHREND	87C	PL B191 209	+Buerger, Criegee, Dainton+	(CELLO Collab.)
BARTEL	86C	ZPHY C30 371	+Becker, Cords, Felst, Haidt+	(JADE Collab.)
Also	85B	ZPHY C26 507	Bartel, Becker, Bowdery, Cords+	(JADE Collab.)
Also	82	PL 108B 140	Bartel, Cords, Dittmann, Eichler+	(JADE Collab.)
ASH	85	PRL 55 1831	+Band, Blume, Camporesi+	(MAC Collab.)
BARTEL	85F	PL 161B 188	+Becker, Cords, Felst+	(JADE Collab.)
DERRICK	85	PR D31 2352	+Fernandez, Fries, Hyman+	(HRS Collab.)
FERNANDEZ	85	PRL 54 1624	+Ford, Qi, Read+	(MAC Collab.)
LEVI	83	PRL 51 1941	+Blocker, Strait+	(Mark II Collab.)
BEHREND	82	PL 114B 282	+Chen, Fenner, Field+	(CELLO Collab.)
BRANDELIK	82C	PL 110B 173	+Braunschweig, Gather	(TASSO Collab.)