

# CHARMED BARYONS ( $C=+1$ )

$$\begin{aligned}\Lambda_c^+ &= u d c, & \Sigma_c^{++} &= u u c, & \Sigma_c^+ &= u d c, & \Sigma_c^0 &= d d c, \\ \Xi_c^+ &= u s c, & \Xi_c^0 &= d s c, & \Omega_c^0 &= s s c\end{aligned}$$

 **$\Lambda_c^+$** 

$$I(J^P) = 0(\frac{1}{2}^+)$$

$J$  not confirmed;  $\frac{1}{2}^+$  is the quark model prediction.

Mass  $m = 2284.9 \pm 0.6$  MeV

Mean life  $\tau = (0.206 \pm 0.012) \times 10^{-12}$  s

$c\tau = 61.8 \mu\text{m}$

### Decay asymmetry parameters

$$\Lambda\pi^+ \quad \alpha = -0.98 \pm 0.19$$

$$\Sigma^+\pi^0 \quad \alpha = -0.45 \pm 0.32$$

$$\Lambda\ell^+\nu_\ell \quad \alpha = -0.82^{+0.11}_{-0.07}$$

Nearly all branching fractions of the  $\Lambda_c^+$  are measured relative to the  $pK^-\pi^+$  mode, but there are no model-independent measurements of this branching fraction. We explain how we arrive at our value of  $B(\Lambda_c^+ \rightarrow pK^-\pi^+)$  in a Note at the beginning of the branching-ratio measurements, in the Listings. When this branching fraction is eventually well determined, all the other branching fractions will slide up or down proportionally as the true value differs from the value we use here.

$\Lambda_c^+$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level (MeV/c)
<b>Hadronic modes with a <math>p</math> and one <math>\bar{K}</math></b>		
$p\bar{K}^0$	( 2.5 $\pm$ 0.7 ) %	872
$pK^-\pi^+$	[k] ( 5.0 $\pm$ 1.3 ) %	822
$p\bar{K}^*(892)^0$	[l] ( 1.8 $\pm$ 0.6 ) %	681
$\Delta(1232)^{++} K^-$	( 8 $\pm$ 5 ) $\times 10^{-3}$	709
$\Lambda(1520)\pi^+$	[l] ( 4.5 $\pm$ 2.5 ) $\times 10^{-3}$	626
$pK^-\pi^+$ nonresonant	( 2.8 $\pm$ 0.9 ) %	822
$p\bar{K}^0\eta$	( 1.3 $\pm$ 0.4 ) %	567

$p\bar{K}^0\pi^+\pi^-$	( 2.4 ± 1.1 ) %	753
$pK^-\pi^+\pi^0$	seen	758
$pK^*(892)^-\pi^+$	[I] ( 1.1 ± 0.6 ) %	579
$p(K^-\pi^+)_{\text{nonresonant}}\pi^0$	( 3.6 ± 1.2 ) %	758
$\Delta(1232)\bar{K}^*(892)$	seen	416
$pK^-\pi^+\pi^+\pi^-$	( 1.1 ± 0.8 ) × 10 <sup>-3</sup>	670
$pK^-\pi^+\pi^0\pi^0$	( 8 ± 4 ) × 10 <sup>-3</sup>	676
$pK^-\pi^+\pi^0\pi^0\pi^0$	( 5.0 ± 3.4 ) × 10 <sup>-3</sup>	573

**Hadronic modes with a  $p$  and zero or two  $K$ 's**

$p\pi^+\pi^-$	( 3.5 ± 2.0 ) × 10 <sup>-3</sup>	926
$p f_0(980)$	[I] ( 2.8 ± 1.9 ) × 10 <sup>-3</sup>	621
$p\pi^+\pi^+\pi^-\pi^-$	( 1.8 ± 1.2 ) × 10 <sup>-3</sup>	851
$pK^+K^-$	( 2.3 ± 0.9 ) × 10 <sup>-3</sup>	615
$p\phi$	[I] ( 1.2 ± 0.5 ) × 10 <sup>-3</sup>	589

**Hadronic modes with a hyperon**

$\Lambda\pi^+$	( 9.0 ± 2.8 ) × 10 <sup>-3</sup>	863
$\Lambda\pi^+\pi^0$	( 3.6 ± 1.3 ) %	843
$\Lambda\rho^+$	< 5 %	CL=95% 638
$\Lambda\pi^+\pi^+\pi^-$	( 3.3 ± 1.0 ) %	806
$\Lambda\pi^+\eta$	( 1.7 ± 0.6 ) %	690
$\Sigma(1385)^+\eta$	[I] ( 8.5 ± 3.3 ) × 10 <sup>-3</sup>	569
$\Lambda K^+\bar{K}^0$	( 6.0 ± 2.1 ) × 10 <sup>-3</sup>	441
$\Sigma^0\pi^+$	( 9.9 ± 3.2 ) × 10 <sup>-3</sup>	824
$\Sigma^+\pi^0$	( 1.00 ± 0.34 ) %	826
$\Sigma^+\eta$	( 5.5 ± 2.3 ) × 10 <sup>-3</sup>	712
$\Sigma^+\pi^+\pi^-$	( 3.4 ± 1.0 ) %	803
$\Sigma^+\rho^0$	< 1.4 %	CL=95% 578
$\Sigma^-\pi^+\pi^+$	( 1.8 ± 0.8 ) %	798
$\Sigma^0\pi^+\pi^0$	( 1.8 ± 0.8 ) %	802
$\Sigma^0\pi^+\pi^+\pi^-$	( 1.1 ± 0.4 ) %	762
$\Sigma^+\pi^+\pi^-\pi^0$	—	766
$\Sigma^+\omega$	[I] ( 2.7 ± 1.0 ) %	568
$\Sigma^+\pi^+\pi^+\pi^-\pi^-$	( 3.0 ± 4.1 ) × 10 <sup>-3</sup>	707
$\Sigma^+K^+K^-$	( 3.5 ± 1.2 ) × 10 <sup>-3</sup>	346
$\Sigma^+\phi$	[I] ( 3.5 ± 1.7 ) × 10 <sup>-3</sup>	292
$\Sigma^+K^+\pi^-$	( 7 ± 6 ) × 10 <sup>-3</sup>	668
$\Xi^0K^+$	( 3.9 ± 1.4 ) × 10 <sup>-3</sup>	652
$\Xi^-K^+\pi^+$	( 4.9 ± 1.7 ) × 10 <sup>-3</sup>	564
$\Xi(1530)^0K^+$	[I] ( 2.6 ± 1.0 ) × 10 <sup>-3</sup>	471

**Semileptonic modes**

$\Lambda \ell^+ \nu_\ell$	[m]	( 2.0 $\pm$ 0.6 ) %	—
$\Lambda e^+ \nu_e$		( 2.1 $\pm$ 0.6 ) %	—
$\Lambda \mu^+ \nu_\mu$		( 2.0 $\pm$ 0.7 ) %	—
$e^+$ anything		( 4.5 $\pm$ 1.7 ) %	—
$p e^+$ anything		( 1.8 $\pm$ 0.9 ) %	—
$\Lambda e^+$ anything		—	—
$\Lambda \mu^+$ anything		—	—
$\Lambda \ell^+ \nu_\ell$ anything		—	—

**Inclusive modes**

$p$ anything		(50 $\pm$ 16) %	—
$p$ anything (no $\Lambda$ )		(12 $\pm$ 19) %	—
$p$ hadrons		—	—
$n$ anything		(50 $\pm$ 16) %	—
$n$ anything (no $\Lambda$ )		(29 $\pm$ 17) %	—
$\Lambda$ anything		(35 $\pm$ 11) %	S=1.4
$\Sigma^\pm$ anything	[n]	(10 $\pm$ 5) %	—

 **$\Delta C = 1$  weak neutral current (*C1*) modes, or  
Lepton number (*L*) violating modes**

$p \mu^+ \mu^-$	<i>C1</i>	< 3.4	$\times 10^{-4}$	CL=90%	936
$\Sigma^- \mu^+ \mu^+$	<i>L</i>	< 7.0	$\times 10^{-4}$	CL=90%	811

---

**$\Lambda_c(2593)^+$** 

$$I(J^P) = 0(\frac{1}{2}^-)$$

The spin-parity follows from the fact that  $\Sigma_c(2455)\pi$  decays, with little available phase space, are dominant.

Mass  $m = 2593.9 \pm 0.8$  MeV

$m - m_{\Lambda_c^+} = 308.9 \pm 0.6$  MeV (S = 1.1)

Full width  $\Gamma = 3.6^{+2.0}_{-1.3}$  MeV

$\Lambda_c^+\pi\pi$  and its submode  $\Sigma_c(2455)\pi$  — the latter just barely — are the only strong decays allowed to an excited  $\Lambda_c^+$  having this mass; and the  $\Lambda_c^+\pi^+\pi^-$  mode seems to be largely via  $\Sigma_c^{++}\pi^-$  or  $\Sigma_c^0\pi^+$ .

<b><math>\Lambda_c(2593)^+</math> DECAY MODES</b>	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+\pi^+\pi^-$	[o] $\approx 67$ %	124
$\Sigma_c(2455)^{++}\pi^-$	$24 \pm 7$ %	17
$\Sigma_c(2455)^0\pi^+$	$24 \pm 7$ %	23
$\Lambda_c^+\pi^+\pi^-$ 3-body	$18 \pm 10$ %	124
$\Lambda_c^+\pi^0$	not seen	261
$\Lambda_c^+\gamma$	not seen	290

 **$\Lambda_c(2625)^+$** 

$$I(J^P) = 0(?^?)$$

$J^P$  is expected to be  $3/2^-$ .

Mass  $m = 2626.6 \pm 0.8$  MeV (S = 1.2)

$m - m_{\Lambda_c^+} = 341.7 \pm 0.6$  MeV (S = 1.6)

Full width  $\Gamma < 1.9$  MeV, CL = 90%

$\Lambda_c^+\pi\pi$  and its submode  $\Sigma(2455)\pi$  are the only strong decays allowed to an excited  $\Lambda_c^+$  having this mass.

<b><math>\Lambda_c(2625)^+</math> DECAY MODES</b>	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+\pi^+\pi^-$	seen	184
$\Sigma_c(2455)^{++}\pi^-$	small	100
$\Sigma_c(2455)^0\pi^+$	small	101
$\Lambda_c^+\pi^+\pi^-$ 3-body	large	184
$\Lambda_c^+\pi^0$	not seen	293
$\Lambda_c^+\gamma$	not seen	319

**$\Sigma_c(2455)$** 

$$I(J^P) = 1(\frac{1}{2}^+)$$

$J^P$  not confirmed;  $\frac{1}{2}^+$  is the quark model prediction.

$$\Sigma_c(2455)^{++}\text{mass } m = 2452.8 \pm 0.6 \text{ MeV}$$

$$\Sigma_c(2455)^+\text{ mass } m = 2453.6 \pm 0.9 \text{ MeV}$$

$$\Sigma_c(2455)^0\text{ mass } m = 2452.2 \pm 0.6 \text{ MeV}$$

$$m_{\Sigma_c^{++}} - m_{\Lambda_c^+} = 167.87 \pm 0.19 \text{ MeV}$$

$$m_{\Sigma_c^+} - m_{\Lambda_c^+} = 168.7 \pm 0.6 \text{ MeV}$$

$$m_{\Sigma_c^0} - m_{\Lambda_c^+} = 167.30 \pm 0.20 \text{ MeV}$$

$$m_{\Sigma_c^{++}} - m_{\Sigma_c^0} = 0.57 \pm 0.23 \text{ MeV}$$

$$m_{\Sigma_c^+} - m_{\Sigma_c^0} = 1.4 \pm 0.6 \text{ MeV}$$

$\Lambda_c^+ \pi$  is the only strong decay allowed to a  $\Sigma_c$  having this mass.

 **$\Sigma_c(2455)$  DECAY MODES**Fraction ( $\Gamma_i/\Gamma$ ) $p$  (MeV/c)

$$\Lambda_c^+ \pi$$

 $\approx 100 \text{ \%}$ 

90

 **$\Sigma_c(2520)$** 

$$I(J^P) = 1(?^?)$$

$$\Sigma_c(2520)^{++}\text{mass } m = 2519.4 \pm 1.5 \text{ MeV}$$

$$\Sigma_c(2520)^0\text{ mass } m = 2517.5 \pm 1.4 \text{ MeV}$$


$$m_{\Sigma_c(2520)^{++}} - m_{\Lambda_c^+} = 234.5 \pm 1.4 \text{ MeV}$$

$$m_{\Sigma_c(2520)^0} - m_{\Lambda_c^+} = 232.6 \pm 1.3 \text{ MeV}$$

$$m_{\Sigma_c(2520)^{++}} - m_{\Sigma_c(2520)^0} = 1.9 \pm 1.9 \text{ MeV}$$

$$\Sigma_c(2520)^{++}\text{full width } \Gamma = 18 \pm 5 \text{ MeV}$$

$$\Sigma_c(2520)^0\text{ full width } \Gamma = 13 \pm 5 \text{ MeV}$$



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$


$I(J^P)$  not confirmed;  $\frac{1}{2}(\frac{1}{2}^+)$  is the quark model prediction.

Mass  $m = 2465.6 \pm 1.4$  MeV

Mean life  $\tau = (0.35^{+0.07}_{-0.04}) \times 10^{-12}$  s

$$c\tau = 106 \text{ } \mu\text{m}$$

$\Xi_c^+$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda K^- \pi^+ \pi^+$	seen	784
$\Lambda \bar{K}^*(892)^0 \pi^+$	not seen	601
$\Sigma(1385)^+ K^- \pi^+$	not seen	676
$\Sigma^+ K^- \pi^+$	seen	808
$\Sigma^+ \bar{K}^*(892)^0$	seen	653
$\Sigma^0 K^- \pi^+ \pi^+$	seen	733
$\Xi^0 \pi^+$	seen	875
$\Xi^- \pi^+ \pi^+$	seen	850
$\Xi(1530)^0 \pi^+$	not seen	748
$\Xi^0 \pi^+ \pi^0$	seen	854
$\Xi^0 \pi^+ \pi^+ \pi^-$	seen	817
$\Xi^0 e^+ \nu_e$	seen	882



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$I(J^P)$  not confirmed;  $\frac{1}{2}(\frac{1}{2}^+)$  is the quark model prediction.

Mass  $m = 2470.3 \pm 1.8$  MeV ( $S = 1.3$ )

$m_{\Xi_c^0} - m_{\Xi_c^+} = 4.7 \pm 2.1$  MeV ( $S = 1.2$ )

Mean life  $\tau = (0.098^{+0.023}_{-0.015}) \times 10^{-12}$  s

$$c\tau = 29 \text{ } \mu\text{m}$$

$\Xi_c^0$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda \bar{K}^0$	seen	864
$\Xi^- \pi^+$	seen	875
$\Xi^- \pi^+ \pi^+ \pi^-$	seen	816
$p K^- \bar{K}^*(892)^0$	seen	406
$\Omega^- K^+$	seen	522
$\Xi^- e^+ \nu_e$	seen	882
$\Xi^- \ell^+ \text{anything}$	seen	—

**$\Xi_c(2645)$** 

$I(J^P) = ?(?^?)$

 $\Xi_c(2645)^+ \text{ mass } m = 2644.6 \pm 2.1 \text{ MeV } (S = 1.2)$  $\Xi_c(2645)^0 \text{ mass } m = 2643.8 \pm 1.8 \text{ MeV}$  $m_{\Xi_c(2645)^+} - m_{\Xi_c^0} = 174.3 \pm 1.1 \text{ MeV}$  $m_{\Xi_c(2645)^0} - m_{\Xi_c^+} = 178.2 \pm 1.1 \text{ MeV}$  $\Xi_c(2645)^+ \text{ full width } \Gamma < 3.1 \text{ MeV, CL} = 90\%$  $\Xi_c(2645)^0 \text{ full width } \Gamma < 5.5 \text{ MeV, CL} = 90\%$  $\Xi_c \pi$  is the only strong decay allowed to a  $\Xi_c$  resonance having this mass. **$\Xi_c(2645)$  DECAY MODES**Fraction ( $\Gamma_i/\Gamma$ ) $p$  (MeV/c) $\Xi_c^0 \pi^+$ 

seen

101

 $\Xi_c^+ \pi^-$ 

seen

107

 **$\Omega_c^0$** 

$I(J^P) = 0(\frac{1}{2}^+)$

 $I(J^P)$  not confirmed;  $0(\frac{1}{2}^+)$  is the quark model prediction.Mass  $m = 2704 \pm 4$  MeV  $(S = 1.8)$ Mean life  $\tau = (0.064 \pm 0.020) \times 10^{-12}$  s $c\tau = 19 \mu\text{m}$  **$\Omega_c^0$  DECAY MODES**Fraction ( $\Gamma_i/\Gamma$ ) $p$  (MeV/c) $\Sigma^+ K^- K^- \pi^+$ 

seen

697

 $\Xi^- K^- \pi^+ \pi^+$ 

seen

838

 $\Omega^- \pi^+$ 

seen

827

 $\Omega^- \pi^- \pi^+ \pi^+$ 

seen

759