

$\Xi(1690)$ $I(J^P) = \frac{1}{2}(\text{??})$ Status: ***

DIONISI 78 sees a threshold enhancement in both the neutral and negatively charged $\Sigma\bar{K}$ mass spectra in $K^- p \rightarrow (\Sigma\bar{K})K\pi$ at 4.2 GeV/c. The data from the $\Sigma\bar{K}$ channels alone cannot distinguish between a resonance and a large scattering length. Weaker evidence at the same mass is seen in the corresponding $\Lambda\bar{K}$ channels, and a coupled-channel analysis yields results consistent with a new Ξ .

BIAGI 81 sees an enhancement at 1700 MeV in the diffractively produced ΛK^- system. A peak is also observed in the $\Lambda\bar{K}^0$ mass spectrum at 1660 MeV that is consistent with a 1720 MeV resonance decaying to $\Sigma^0\bar{K}^0$, with the γ from the Σ^0 decay not detected.

BIAGI 87 provides further confirmation of this state in diffractive dissociation of Ξ^- into ΛK^- . The significance claimed is 6.7 standard deviations.

ADAMOVICH 98 sees a peak of 1400 ± 300 events in the $\Xi^-\pi^+$ spectrum produced by 345 GeV/c Σ^- -nucleus interactions.

 $\Xi(1690)$ MASSES**MIXED CHARGES**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1690 ± 10 OUR ESTIMATE				This is only an educated guess; the error given is larger than the error on the average of the published values.
1686 \pm 4	1400	ADAMOVICH 98	WA89	Σ^- nucleus, 345 GeV/c
1699 \pm 5	175	¹ DIONISI 78	HBC	$K^- p$ 4.2 GeV/c
1684 \pm 5	183	² DIONISI 78	HBC	$K^- p$ 4.2 GeV/c

 $\Xi(1690)^0$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1686 \pm 4	1400	ADAMOVICH 98	WA89	Σ^- nucleus, 345 GeV/c
1699 \pm 5	175	¹ DIONISI 78	HBC	$K^- p$ 4.2 GeV/c
1684 \pm 5	183	² DIONISI 78	HBC	$K^- p$ 4.2 GeV/c

 $\Xi(1690)^-$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1691.1 \pm 1.9 \pm 2.0	104	BIAGI 87	SPEC	Ξ^- Be 116 GeV
1700 \pm 10	150	³ BIAGI 81	SPEC	Ξ^- H 100, 135 GeV
1694 \pm 6	45	⁴ DIONISI 78	HBC	$K^- p$ 4.2 GeV/c

$\Xi(1690)$ WIDTHS

MIXED CHARGES

VALUE (MeV)	EVTS	DOCUMENT ID
<30 OUR ESTIMATE		

$\Xi(1690)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
10 ± 6	1400	ADAMOVICH 98	WA89	Σ^- nucleus, 345 GeV/c	■
44 ± 23	175	¹ DIONISI 78	HBC	$K^- p$ 4.2 GeV/c	
20 ± 4	183	² DIONISI 78	HBC	$K^- p$ 4.2 GeV/c	

$\Xi(1690)^-$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 8	90	104	BIAGI 87	SPEC	Ξ^- Be 116 GeV
47 ± 14		150	³ BIAGI 81	SPEC	Ξ^- H 100, 135 GeV
26 ± 6		45	⁴ DIONISI 78	HBC	$K^- p$ 4.2 GeV/c

$\Xi(1690)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \Lambda \bar{K}$	seen
$\Gamma_2 \Sigma \bar{K}$	seen
$\Gamma_3 \Xi \pi$	seen
$\Gamma_4 \Xi^- \pi^+ \pi^0$	
$\Gamma_5 \Xi^- \pi^+ \pi^-$	possibly seen
$\Gamma_6 \Xi(1530) \pi$	

$\Xi(1690)$ BRANCHING RATIOS

$\Gamma(\Lambda \bar{K})/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
seen	104	BIAGI 87	SPEC	—	Ξ^- Be 116 GeV	

$\Gamma(\Sigma \bar{K})/\Gamma(\Lambda \bar{K})$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_1
2.7 ± 0.9	DIONISI 78	HBC	0	$K^- p$ 4.2 GeV/c	
3.1 ± 1.4	DIONISI 78	HBC	—	$K^- p$ 4.2 GeV/c	

$\Gamma(\Xi \pi)/\Gamma(\Sigma \bar{K})$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ_2
<0.09	DIONISI 78	HBC	0	$K^- p$ 4.2 GeV/c	

$\Gamma(\Xi \pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
seen	ADAMOVICH 98	WA89	Σ^- nucleus, 345 GeV/c	■

$\Gamma(\Xi^-\pi^+\pi^0)/\Gamma(\Sigma\bar{K})$

<u>VALUE</u>
<0.04

$\Gamma(\Xi^-\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>EVTS</u>
possibly seen	4

$\Gamma(\Xi^-\pi^+\pi^-)/\Gamma(\Sigma\bar{K})$

<u>VALUE</u>
<0.03

$\Gamma(\Xi(1530)\pi)/\Gamma(\Sigma\bar{K})$

<u>VALUE</u>
<0.06

Γ_4/Γ_2

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
DIONISI	78	HBC	$K^- p$ 4.2 GeV/c

Γ_5/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
BIAGI	87	SPEC	$\Xi^- \text{Be}$ 116 GeV

Γ_5/Γ_2

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
DIONISI	78	HBC	$K^- p$ 4.2 GeV/c

Γ_6/Γ_2

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
DIONISI	78	HBC	$K^- p$ 4.2 GeV/c

$\Xi(1690)$ FOOTNOTES

¹ From a fit to the $\Sigma^+ K^-$ spectrum.

² From a coupled-channel analysis of the $\Sigma^+ K^-$ and $\Lambda\bar{K}^0$ spectra.

³ A fit to the inclusive spectrum from $\Xi^- N \rightarrow \Lambda K^- X$.

⁴ From a coupled-channel analysis of the $\Sigma^0 K^-$ and ΛK^- spectra.

$\Xi(1690)$ REFERENCES

ADAMOVICH 98 EPJ C5 621
 BIAGI 87 ZPHY C34 15
 BIAGI 81 ZPHY C9 305
 DIONISI 78 PL 80B 145

M.I. Adamovich+ (CERN WA89 Collab.)
 + (BRIS, CERN, GEVA, HEIDP, LAUS, LOQM, RAL) I
 + (BRIS, CAVE, GEVA, HEIDP, LAUS, LOQM, RHEL)
 + Diaz, Armenteros+ (CERN, AMST, NIJM, OXF) I